

# TOPOLOGICAL FINITENESS PROPERTIES OF MONOIDS PART 2: SPECIAL MONOIDS, ONE-RELATOR MONOIDS, AMALGAMATED FREE PRODUCTS, AND HNN EXTENSIONS

ROBERT D. GRAY<sup>1</sup> and BENJAMIN STEINBERG<sup>2</sup>

**ABSTRACT.** We show how topological methods developed in a previous article can be applied to prove new results about topological and homological finiteness properties of monoids. A monoid presentation is called special if the right-hand side of each relation is equal to 1. We prove results which relate the finiteness properties of a monoid defined by a special presentation with those of its group of units. Specifically we show that the monoid inherits the finiteness properties  $F_n$  and  $FP_n$  from its group of units. We also obtain results which relate the geometric and cohomological dimensions of such a monoid to those of its group of units. We apply these results to prove a Lyndon's Identity Theorem for one-relator monoids of the form  $\langle A \mid r = 1 \rangle$ . In particular we show that all such monoids are of type  $F_\infty$  (and  $FP_\infty$ ), and that when  $r$  is not a proper power, then the monoid has geometric and cohomological dimension at most 2. The first of these results, resolves an important case of a question of Kobayashi from 2000 on homological finiteness properties of one-relator monoids. We also show how our topological approach can be used to prove results about the closure properties of various homological and topological finiteness properties for amalgamated free products and HNN-extensions of monoids. To prove these results we introduce new methods for constructing equivariant classifying spaces for monoids, as well as developing a Bass–Serre theory for free constructions of monoids.

## 1. INTRODUCTION

Topological methods play an important role in the modern study of infinite discrete groups. Recall that an Eilenberg–Mac Lane complex of type  $K(G, 1)$  is an aspherical CW complex with fundamental group  $G$ . For any group  $G$  a  $K(G, 1)$  complex exists, and it is unique up to homotopy equivalence. While the existence of such spaces is elementary, it is often a much harder problem to find a  $K(G, 1)$  complex which is suitably ‘nice’ to be used for doing calculations. This is important if one wants to compute homology and cohomology groups. This is part of the motivation for the study of higher order topological finiteness properties of groups, a topic which goes back to pioneering work of Wall [61] and Serre [57]. We recall that a group is of type  $F_n$  if there is a  $K(G, 1)$ -complex with a finite  $n$ -skeleton. The property  $F_1$  is equivalent to finite generation, while a group is of type  $F_2$  if and only if it is finitely presented, so  $F_n$  gives a natural higher dimensional analogue of these two fundamental finiteness properties. The geometric dimension of a group  $G$ , denoted  $\text{gd}(G)$ , is the minimum dimension of a  $K(G, 1)$  complex. The topological finiteness property  $F_n$  and geometric dimension correspond, respectively, to the homological finiteness property  $FP_n$  and the cohomological dimension of the group. The study of topological and homological finiteness properties is an active area of

---

2010 *Mathematics Subject Classification.* 20M50, 20M05, 20J05, 57M07, 20F10, 20F65.

*Key words and phrases.* Equivariant CW-complex, homological finiteness property, classifying space, cohomological dimension, Hochschild cohomological dimension, geometric dimension, Bass–Serre tree, monoid, special monoid, one-relator monoid, free product with amalgamation, HNN extension.

This work was supported by the EPSRC grants EP/N033353/1 ‘Special inverse monoids: subgroups, structure, geometry, rewriting systems and the word problem’ and EP/V032003/1 ‘Algorithmic, topological and geometric aspects of infinite groups, monoids and inverse semigroups’. The second author was supported by NSA MSP #H98230-16-1-0047 and a PSC-CUNY award.

<sup>1</sup>School of Mathematics, University of East Anglia, Norwich NR4 7TJ, England. Email [Robert.D.Gray@uea.ac.uk](mailto:Robert.D.Gray@uea.ac.uk).

<sup>2</sup>Department of Mathematics, City College of New York, Convent Avenue at 138th Street, New York, New York 10031, USA. Email [bsteinberg@ccny.cuny.edu](mailto:bsteinberg@ccny.cuny.edu).

research. We refer the reader to [11, Chapter 8], [24, Chapters 6-9] and [12] for more background on this topic.

The homological finiteness properties  $FP_n$  and cohomological dimension have also been extensively studied more generally for monoids. One major motivation for studying homological finiteness properties of monoids comes from important connections with the theory of rewriting systems, and the word problem for finitely presented monoids. It is well known that there are finitely presented monoids with undecidable word problem. Given that the word problem is undecidable in general, a central theme running through the development of geometric and combinatorial group and monoid theory has been to identify and study classes of finitely presented monoids all of whose members have solvable word problem. A finite complete rewriting system is a finite presentation for a monoid of a particular form (both confluent and Noetherian) which gives a solution of the word problem for the monoid; see [7]. Complete rewriting systems are also of interest because of their close connection with the theory of Gröbner–Shirshov bases; see [60]. The connection between complete rewriting systems and homological finiteness properties is given by the Anick–Groves–Squier Theorem which shows that a monoid that admits such a presentation must be of type  $FP_\infty$ ; see [4, 58] and [10]. The property  $FP_n$  for monoids also arises in the study of Bieri–Neumann–Strebel–Renz invariants of groups; see [6].

A number of other interesting homological and homotopical finiteness properties have been studied in relation to monoids defined by complete rewriting systems; see [3, 28, 55]. The cohomological dimension of monoids has also received attention in the literature; see for example [13, 27, 47]. In fact, for monoids these properties depend on whether one works with left  $\mathbb{Z}M$ -modules or right  $\mathbb{Z}M$ -modules, giving rise to the notions of both left- and right- $FP_n$ , and left and right cohomological dimension. In general these are independent of each other; see [15, 27, 53]. Working with bimodule resolutions of the  $(\mathbb{Z}M, \mathbb{Z}M)$ -bimodule  $\mathbb{Z}M$  one obtains the notion bi- $FP_n$  introduced and studied in [36]. This property is of interest from the point of view of Hochschild cohomology, which is the standard notion of cohomology for rings; see [30, 51]. For more background on the study of homological finiteness properties in monoid theory, and the connections with the theory of string rewriting systems, see [10, 16, 52].

While homological finiteness properties of monoids have been extensively studied, in contrast, until recently there was no corresponding theory of topological finiteness properties of monoids. The results in this paper are part of a research programme of the authors, initiated in [25], aimed at developing such a theory. A central theme of this work is that the topological approach allows for less technical, and more conceptual, proofs than had previously been possible using only algebraic means. Other recent results in the literature where topological methods have been usefully applied in the study of monoids include, e.g. [8, 46, 50].

This paper is the sequel to the article [25] where we set out the foundations of  $M$ -equivariant homotopy theory for monoids acting on CW complexes, and the corresponding study of topological finiteness properties of monoids. In that paper we introduced the notion of a left equivariant classifying space for a monoid, which is a contractible projective  $M$ -CW complex. A left equivariant classifying space always exists, for any monoid  $M$ , and it is unique up to  $M$ -homotopy equivalence. We then define the corresponding finiteness conditions left- $F_n$  and left geometric dimension in the obvious natural way in terms of the existence of a left equivariant classifying space satisfying appropriate finiteness properties. It follows easily from the definitions that left- $F_n$  implies left- $FP_n$ , and that the left geometric dimension is an upper bound on the left cohomological dimension of the monoid. There are obvious dual definitions and statements working with right actions. We also developed a two-sided analogue of this theory in [25], with two-sided  $M$  actions, defining the notion of a bi-equivariant classifying space for a monoid, and the resulting finiteness properties bi- $F_n$  and geometric dimension. It follows from the definitions that bi- $F_n$  implies bi- $FP_n$  (in the sense of [36]) and that the geometric dimension is an upper bound for the Hochschild cohomological dimension. See Section 2 below for full details and formal definitions of all of these notions.

The aim of this paper is to apply the ideas and results from [25] to solve some open problems concerning homological finiteness properties of monoids that seemed resistant to algebraic techniques. Let us begin with some history. An important open problem is whether every one-relator monoid has decidable word problem. While the question is open in general, it has been solved in a number of special cases; see Adjan [1] and Adjan and Oganesyan [2]. Related to this is another open question which asks whether every one-relator monoid admits a presentation by a finite complete rewriting system. Of course, a positive answer to this question would imply a positive solution to the word problem. In light of the Anick-Groves-Squier Theorem which states that monoids which admit finite complete presentations are of type right- and left- $\text{FP}_\infty$ , it is natural to ask whether all one-relator monoids are of type  $\text{FP}_\infty$ . This question was posed by Kobayashi in [34, Problem 1]. The question is also natural given the fact that all one-relator groups are all of type  $\text{FP}_\infty$ , as a consequence of Lyndon's Identity Theorem for one-relator groups; see Lyndon [42].

The first positive result concerning the word problem for one-relator monoids dealt with the case of, so-called, special one-relator monoids [1]. A *special* monoid is one defined by a finite presentation of the form  $\langle A \mid w_1 = 1, \dots, w_k = 1 \rangle$ . They were first studied in the sixties by Adjan [1] and Makanin [44]. Adjan proved that the group of units of a one-relator special monoid is a one-relator group and reduced the word problem of the monoid to that of the group, which has a decidable word problem by Magnus's theorem [43]. Makanin proved more generally that the group of units of a  $k$ -relator special monoid is a  $k$ -relator group and reduced the word problem of the monoid to that of the group. See [63] for a modern approach to these results. Thus there is a much closer connection for special monoids between the group of units and the monoid than is customary.

One of the main results of this paper is that if  $M = \langle A \mid w_1 = 1, \dots, w_k = 1 \rangle$ , and if  $G$  is the group of units of  $M$ , then if  $G$  is of type  $\text{FP}_n$  with  $1 \leq n \leq \infty$ , then  $M$  is also of type left- and right- $\text{FP}_n$ . Moreover, we prove that both the left and right cohomological dimensions of  $M$  are bounded below by  $\text{cd } G$ , and are bounded above by  $\max\{2, \text{cd } G\}$ . We shall also prove the topological analogues of these results, obtaining the corresponding statements with right and left- $\text{F}_n$  and geometric dimension. These results are obtained by proving new results about the geometry of Cayley digraphs of special monoids, including the observation that the quotient of the Cayley digraph by its strongly connected components is a regular rooted tree on which the monoid acts by simplicial maps. We use this to show how one can construct a left equivariant classifying space for a special monoid from an equivariant classifying space for its group of units.

We shall then go on to apply these results to prove a Lyndon's Identity Theorem [42] for one-relator monoids of the form  $\langle A \mid w = 1 \rangle$ . Specifically, we show that our results can be applied to construct equivariant classifying spaces for one-relator monoids of this form, which have finitely many orbits of cells in each dimension, and have dimension at most 2 unless the monoid has torsion. We apply this to give a positive answer to Kobayashi's question [34, Problem 1] on homological finiteness properties of one-relator monoids, in the case of one-relator monoids of the form  $\langle A \mid w = 1 \rangle$ , by proving that all such monoids are of type left- and right- $\text{F}_\infty$  and  $\text{FP}_\infty$ . We also show that if  $M = \langle A \mid w = 1 \rangle$  with  $w$  not a proper power then the left and right cohomological dimension of  $M$  are bounded above by 2, and if  $w$  is a proper power then they are both equal to  $\infty$ . The analogous topological result for the left and right geometric dimension of a one-relator special monoid is also obtained. In fact, it will follow from our results that when  $w$  is not a proper power then the Cayley complex of the one-relator monoid  $M$  is an equivariant classifying space for  $M$  of dimension at most 2. This is the analogue, for one-relator special monoids, of the fact that the presentation complex of a torsion-free one-relator group is aspherical and is thus a  $K(G, 1)$  complex for the group of dimension at most 2; see [14, 22]. These results on special monoids, and one-relator monoids, will be given in Section 3.

The results we obtain in this paper for special one-relator monoids form an important infinite family of base cases for the main result in our article [26] where we prove a Lyndon's Identity Theorem for arbitrary one-relator monoids  $\langle A \mid u = v \rangle$ . Applying this result, in [26] we give a

positive answer to Kobayashi's question by showing that every one-relator monoid  $\langle A \mid u = v \rangle$  is of type left- and right- $\text{FP}_\infty$ .

In Section 4 below we prove several new results about the preservation of topological and homological finiteness properties for amalgamated free products of monoids. Monoid amalgamated products are far more complicated than group ones. For example, an amalgamated free product of finite monoids can have an undecidable word problem, and the factors do not necessarily embed, or intersect, in the base monoid; see [56]. In particular there are no normal form results at our disposal when working with monoid amalgamated free products. We give a method for constructing an equivariant classifying spaces for an amalgamated free product of monoids  $L = M_1 *_W M_2$  from equivariant classifying spaces of the monoids  $M_1$ ,  $M_2$  and  $W$ . To do this, we use homological ideas of Dicks [18] on derivations to construct a Bass–Serre tree  $T$  for the amalgam  $L$ . We also develop an analogous theory in the two-sided case. These constructions are used to prove several results about the closure properties of  $\text{F}_n$ ,  $\text{FP}_n$ , and geometric and cohomological dimension.

Finally, in Section 5 we consider HNN extensions construction for monoids, in the sense of Otto and Pride [54], and those defined by Howie [32]. As in the case of amalgamated free products, we give constructions of equivariant classifying spaces, and apply these to deduce results about the closure properties of topological and homological finiteness properties. This also involves constructing appropriate Bass–Serre trees. As special cases of our results we recover generalisations of a number of results of Otto and Pride from [54] and [55].

## 2. PRELIMINARIES

In this section we recall some of the relevant background from [25] needed for the rest of the article. For full details, and proofs of the statements made here we refer the reader to [25, Sections 2–4]. For additional general background on algebraic topology, and topological methods in group theory, we refer the reader to [48] and [24].

**2.1. The category of  $M$ -sets.** Let  $M$  be a monoid. A *left  $M$ -set* consists of a set  $X$  and a mapping  $M \times X \rightarrow X$  written  $(m, x) \mapsto mx$  called a *left action*, such that  $1x = x$  and  $m(nx) = (mn)x$  for all  $m, n \in M$  and  $x \in X$ . Right  $M$ -sets are defined dually, they are the same thing as left  $M^{op}$ -sets, where  $M^{op}$  is the *opposite* of the monoid  $M$  which is the monoid with the same underlying set  $M$  and multiplication given by  $x \cdot y = yx$ . A *bi- $M$ -set* is an  $M \times M^{op}$ -set. A mapping  $f: X \rightarrow Y$  between  $M$ -sets is  *$M$ -equivariant* if  $f(mx) = mf(x)$  for all  $x \in X$ ,  $m \in M$ , and  $M$ -sets together with  $M$ -equivariant mappings form a category.

If  $X$  is an  $M$ -set and  $A \subseteq X$ , then  $A$  is said to be a *free basis for  $X$*  if and only if each element of  $X$  can be uniquely expressed as  $ma$  with  $m \in M$  and  $a \in A$ . The free left  $M$ -set on  $A$  exists and can be realised as the set  $M \times A$  with action  $m(m', a) = (mm', a)$ . Note that if  $G$  is a group, then a left  $G$ -set  $X$  is free if and only if  $G$  acts freely on  $X$ , that is, each element of  $X$  has trivial stabilizer. In this case, any set of orbit representatives is a basis. An  $M$ -set  $P$  is *projective* if any  $M$ -equivariant surjective mapping  $f: X \rightarrow P$  has an  $M$ -equivariant section  $s: P \rightarrow X$  with  $f \circ s = 1_P$ . Every free  $M$ -set is projective, and an  $M$ -set is projective if and only if it is a retract of a free one. Each projective  $M$ -set  $P$  is isomorphic to an  $M$ -set of the form  $\coprod_{a \in A} Me_a$  (disjoint union, which is the coproduct in the category of  $M$ -sets) with  $e_a \in E(M)$ , where  $E(M)$  denotes the set of idempotents of the monoid  $M$ . In particular, projective  $G$ -sets are the same thing as free  $G$ -sets for a group  $G$ .

If  $A$  is a right  $M$ -set and  $B$  is a left  $M$ -set, then  $A \otimes_M B$  is the quotient of  $A \times B$  by the least equivalence relation  $\sim$  such that  $(am, b) \sim (a, mb)$  for all  $a \in A$ ,  $b \in B$  and  $m \in M$ . We write  $a \otimes b$  for the class of  $(a, b)$  and note that the mapping  $(a, b) \mapsto a \otimes b$  is universal for mappings  $f: A \times B \rightarrow X$  with  $X$  a set and  $f(am, b) = f(a, mb)$ . If  $M$  happens to be a group, then  $M$  acts on  $A \times B$  via  $m(a, b) = (am^{-1}, mb)$  and  $A \otimes_M B$  is just the set of orbits of this action. The tensor product  $A \otimes_M ()$  preserves all colimits because it is a left adjoint to the functor  $X \mapsto X^A$ .

If  $B$  is a left  $M$ -set there is a natural preorder relation  $\leq$  on  $B$  where  $x \leq y$  if and only if  $Mx \subseteq My$ . We write  $x \approx y$  if there is a sequence  $z_1, z_2, \dots, z_n$  of elements of  $B$  such that for each  $0 \leq i \leq n-1$  either  $z_i \leq z_{i+1}$  or  $z_i \geq z_{i+1}$ . This is clearly an equivalence relation and we call the  $\approx$ -classes of  $B$  the *weak orbits* of the  $M$ -set. This corresponds to the notion of the weakly connected components in a directed graph. If  $B$  is a right  $M$ -set then we use  $B/M$  to denote the set of weak orbits of the  $M$ -set while if  $B$  is a left  $M$ -set we use  $M \backslash B$  to denote the set of weak orbits. Note that if  $1$  denotes the trivial right  $M$ -set and  $B$  is a left  $M$ -set, then we have  $M \backslash B = 1 \otimes_M B$ . Let  $M, N$  be monoids. An  $M$ - $N$ -biset is an  $M \times N^{op}$ -set. If  $A$  is an  $M$ - $N$ -biset and  $B$  is a left  $N$ -set, then the equivalence relation defining  $A \otimes_N B$  is left  $M$ -invariant and so  $A \otimes_N B$  is a left  $M$ -set with action  $m(a \otimes b) = ma \otimes b$ .

**2.2. Projective  $M$ -CW complexes.** A *left  $M$ -space* is a topological space  $X$  with a continuous left action  $M \times X \rightarrow X$  where  $M$  has the discrete topology. A *right  $M$ -space* is the same thing as a left  $M^{op}$ -space and a *bi- $M$ -space* is an  $M \times M^{op}$ -space. Each  $M$ -set can be viewed as a discrete  $M$ -space. Colimits in the category of  $M$ -spaces are formed by taking colimits in the category of spaces and observing that the result has a natural  $M$ -action.

Our main interest in this article will be in  $M$ -spaces  $X$  where  $X$  is a CW complex. Following [25] we define a (projective)  *$M$ -cell* of dimension  $n$  to be an  $M$ -space of the form  $Me \times B^n$  where  $e \in E(M)$  is an idempotent and  $B^n$  has the trivial action. In the special case  $e = 1$ , we call it a *free  $M$ -cell*. We then define a projective  $M$ -CW complex in an inductive fashion by imitating the usual definition of a CW complex but by attaching  $M$ -cells  $Me \times B^n$  via  $M$ -equivariant maps from  $Me \times S^{n-1}$  to the  $(n-1)$ -skeleton. Formally, a *projective (left) relative  $M$ -CW complex* is a pair  $(X, A)$  of  $M$ -spaces such that  $X = \varinjlim X_n$  with  $i_n: X_n \rightarrow X_{n+1}$  inclusions,  $X_{-1} = A$ ,  $X_0 = P_0 \cup A$  with  $P_0$  a projective  $M$ -set and where  $X_n$  is obtained as a pushout of  $M$ -spaces

$$\begin{array}{ccc} P_n \times S^{n-1} & \longrightarrow & X_{n-1} \\ \downarrow & & \downarrow \\ P_n \times B^n & \longrightarrow & X_n \end{array} \quad (2.1)$$

with  $P_n$  a projective  $M$ -set and  $B^n$  having a trivial  $M$ -action for  $n \geq 1$ . The set  $X_n$  is the  *$n$ -skeleton* of  $X$  and if  $X_n = X$  and  $P_n \neq \emptyset$ , then  $X$  is said to have *dimension  $n$* . Since  $P_n$  is isomorphic to a coproduct of  $M$ -sets of the form  $Me$  with  $e \in E(M)$ , we are indeed attaching  $M$ -cells at each step. If  $A = \emptyset$ , we call  $X$  a *projective  $M$ -CW complex*. Note that a projective  $M$ -CW complex is a CW complex and the  $M$ -action is cellular (in fact, takes  $n$ -cells to  $n$ -cells). We can define projective right  $M$ -CW complexes and projective bi- $M$ -CW complexes by replacing  $M$  with  $M^{op}$  and  $M \times M^{op}$ , respectively. We say that  $X$  is a *free  $M$ -CW complex* if each  $P_n$  is a free  $M$ -set. A projective  $M$ -CW complex  $X$  is of  *$M$ -finite type* if  $P_n$  is a finitely generated projective  $M$ -set for each  $n$ , and we say that  $X$  is  *$M$ -finite* if it is finite dimensional and of  $M$ -finite type (i.e.,  $X$  is constructed from finitely many  $M$ -cells). The degree  $n$  component of the cellular chain complex for the projective  $M$ -CW complex  $X$  is isomorphic to  $\mathbb{Z}P_n$  as a  $\mathbb{Z}M$ -module, and hence is projective.

A *projective  $M$ -CW subcomplex* of  $X$  is an  $M$ -invariant subcomplex  $A \subseteq X$  which is a union of  $M$ -cells of  $X$ . If  $X$  is a projective  $M$ -CW complex then so is  $Y = X \times I$  where  $I$  is given the trivial action. If we retain the above notation, then  $Y_0 = X_0 \times \partial I \cong X_0 \coprod X_0$ . The  $n$ -cells for  $n \geq 1$  are obtained from attaching  $P_n \times B^n \times \partial I \cong (P_n \coprod P_n) \times B^n$  and  $P_{n-1} \times B^{n-1} \times I$ . Notice that  $X \times \partial I$  is a projective  $M$ -CW subcomplex of  $X \times I$ . An  *$M$ -homotopy* between  $M$ -equivariant continuous maps  $f, g: X \rightarrow Y$  between  $M$ -spaces  $X$  and  $Y$  is an  $M$ -equivariant mapping  $H: X \times I \rightarrow Y$  with  $H(x, 0) = f(x)$  and  $H(x, 1) = g(x)$  for  $x \in X$  where  $I$  is viewed as having the trivial  $M$ -action. We write  $f \simeq_M g$  in this case. We say that  $X, Y$  are  *$M$ -homotopy equivalent*, written  $X \simeq_M Y$ , if there are  $M$ -equivariant continuous mappings (called  *$M$ -homotopy equivalences*)  $f: X \rightarrow Y$  and  $g: Y \rightarrow X$  such that  $gf \simeq_M 1_X$  and  $fg \simeq_M 1_Y$ . Every  $M$ -equivariant continuous mapping of projective  $M$ -CW

complexes is  $M$ -homotopy equivalent to a cellular one. This is the *cellular approximation theorem*; see [25, Theorem 2.8].

If  $X$  is a left  $M$ -space and  $A$  is a right  $M$ -set, then  $A \otimes_M X$  is a topological space with the quotient topology. The following base change result will be used frequently below.

**Proposition 2.1.** [25, Proposition 3.1 and Corollary 3.2] *If  $A$  is an  $M$ - $N$ -biset that is projective (free) as an  $M$ -set and  $X$  is a projective (free)  $N$ -CW complex, then  $A \otimes_N X$  is a projective (free)  $M$ -CW complex. If  $A$  is in addition finitely generated as an  $M$ -set and  $X$  is of  $N$ -finite type, then  $A \otimes_N X$  is of  $M$ -finite type. Moreover,  $\dim A \otimes_N X = \dim X$ .*

*Remark 2.2.* We shall use the observation that if  $X$  is a free right  $M$ -set on  $A$ , then  $A$  is in bijection with  $X/M$  and hence  $X \cong X/M \times M$  as a right  $M$ -set where  $M$  acts trivially on  $X/M$ . Hence if  $Y$  is a projective  $M$ -CW complex, then  $X \otimes_M Y \cong \coprod_A Y \cong X/M \times Y$  where  $X/M$  has the discrete topology. Moreover, these homeomorphisms come from isomorphisms of the CW structure.

### 2.3. Equivariant classifying spaces and topological finiteness properties for monoids.

A (left) *equivariant classifying space*  $X$  for a monoid  $M$  is a projective  $M$ -CW complex which is contractible. A right equivariant classifying space for  $M$  will be a left equivariant classifying space for  $M^{op}$ . In some cases, an equivariant classifying space for a monoid may be constructed using the Cayley digraph of the monoid as the 1-skeleton. Recall that if  $M$  is a monoid and  $A \subseteq M$ , then the (right) *Cayley digraph*  $\Gamma(M, A)$  of  $M$  with respect to  $A$  is the graph with vertex set  $M$  and with edges in bijection with  $M \times A$  where the directed edge (arc) corresponding to  $(m, a)$  starts at  $m$  and ends at  $ma$ . Note that  $\Gamma(M, A)$  is a free  $M$ -graph and is  $M$ -finite if and only if  $A$  is finite (see Section 4 below for the definition of  $M$ -graph).

Equivariant classifying spaces of monoids are unique up to  $M$ -homotopy equivalence; see [25, Theorem 6.3 & Corollary 6.5]. The definition of equivariant classifying spaces for monoids leads naturally to the definitions of the following topological finiteness properties. A monoid  $M$  is of type *left- $F_n$*  (for a non-negative integer  $n$ ) if there is an equivariant classifying space  $X$  for  $M$  such that  $X_n$  is  $M$ -finite, i.e., such that  $M \backslash X$  has finite  $n$ -skeleton. We say that  $M$  is of type *left- $F_\infty$*  if  $M$  has an equivariant classifying space  $X$  that is of  $M$ -finite type, i.e.,  $M \backslash X$  is of finite type. The monoid  $M$  is defined to have type *right- $F_n$*  if  $M^{op}$  is of type *left- $F_n$*  for  $0 \leq n \leq \infty$ . The *left geometric dimension* of  $M$  is defined to be the minimum dimension of a left equivariant classifying space for  $M$ . The right geometric dimension is defined dually.

The homological analogue of *left- $F_n$*  is the finiteness property *left- $FP_n$* , where a monoid  $M$  is said to be of type *left- $FP_n$*  if there is a projective resolution  $P = (P_i)_{i \geq 0}$  of the trivial left  $\mathbb{Z}M$ -module  $\mathbb{Z}$  such that  $P_i$  is finitely generated for  $i \leq n$ . There is a dual notion of *right- $FP_n$* , and we say a monoid is of type *FP $_n$*  if it is both of type *left- $FP_n$*  and *right- $FP_n$* . For any monoid  $M$ , if  $M$  is of type *left- $F_n$*  for some  $0 \leq n \leq \infty$  then it is of type *left- $FP_n$* . Indeed, if  $X$  is an equivariant classifying space for  $M$  then the augmented cellular chain complex of  $X$  gives a projective  $\mathbb{Z}M$ -resolution of the trivial  $\mathbb{Z}M$ -module  $\mathbb{Z}$  with the desired finiteness properties. If  $M$  is a monoid of type *left- $F_2$* , then  $M$  is of type *left- $F_n$*  if and only if  $M$  is of type *left- $FP_n$*  for  $0 \leq n \leq \infty$ . In particular, for finitely presented monoids the conditions *left- $F_n$*  and *left- $FP_n$*  are equivalent. In the special case that the monoid  $M$  is a group, the definition of *left- $F_n$*  above is easily seen to agree with the usual definition of  $F_n$  for groups. The left geometric dimension is clearly an upper bound on the left cohomological dimension, denoted *left cd*  $M$ , of a monoid  $M$  where the left *cohomological dimension* of  $M$  is the shortest length of a projective resolution of the trivial left  $\mathbb{Z}M$ -module  $\mathbb{Z}$ .

To define the bilateral notion of a classifying space, first recall that  $M$  is an  $M \times M^{op}$ -set via the action  $(m_L, m_R)m = m_L m m_R$ . We say that a projective  $M \times M^{op}$ -CW complex  $X$  is a *bi-equivariant classifying space* for  $M$  if  $\pi_0(X) \cong M$  as an  $M \times M^{op}$ -set and each component of  $X$  is contractible; equivalently,  $X$  has an  $M \times M^{op}$ -equivariant homotopy equivalence to the discrete  $M \times M^{op}$ -set  $M$ . We can augment the cellular chain complex of  $X$  via the canonical surjection  $\varepsilon: C_0(X) \rightarrow H_0(X) \cong \mathbb{Z}\pi_0(X) \cong \mathbb{Z}M$ . Since each component of  $X$  is contractible,

this gives a projective bimodule resolution of  $\mathbb{Z}M$ . A bi-equivariant classifying space may be constructed for any monoid [25, Corollary 7.4]. As in the one-sided case, bi-equivariant classifying spaces are unique up to  $M \times M^{op}$ -homotopy equivalence; see [25, Theorem 7.2].

A monoid  $M$  is said to be of type  $bi\text{-}F_n$  if there is a bi-equivariant classifying space  $X$  for  $M$  such that  $X_n$  is  $M \times M^{op}$ -finite, i.e.,  $M \backslash X / M$  has finite  $n$ -skeleton. We say that  $M$  is of type  $bi\text{-}F_\infty$  if  $M$  has a bi-equivariant classifying space  $X$  that is of  $M \times M^{op}$ -finite type, i.e.,  $M \backslash X / M$  is of finite type. We define the *geometric dimension* of  $M$  to be the minimum dimension of a bi-equivariant classifying space for  $M$ . The homological analogue of  $bi\text{-}F_n$  is the property  $bi\text{-}FP_n$  (in the sense of [36]), where a monoid is said to be of type  $bi\text{-}FP_n$  if there is a projective resolution

$$\cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \mathbb{Z}M \rightarrow 0$$

of the  $(\mathbb{Z}M, \mathbb{Z}M)$ -bimodule  $\mathbb{Z}M$ , where  $P_0, P_1, \dots, P_n$  are finitely generated projective  $(\mathbb{Z}M, \mathbb{Z}M)$ -bimodules. For  $0 \leq n \leq \infty$ , if  $M$  is of type  $bi\text{-}F_n$ , then it is of type  $bi\text{-}FP_n$ . If  $M$  is of type  $bi\text{-}F_n$  for  $0 \leq n \leq \infty$ , then  $M$  is of type  $left\text{-}F_n$  and type  $right\text{-}F_n$ . If  $M$  is a monoid of type  $bi\text{-}F_2$ , then  $M$  is of type  $bi\text{-}F_n$  if and only if  $M$  is of type  $bi\text{-}FP_n$  for  $0 \leq n \leq \infty$ ; see [25, Theorem 7.15]. In particular, for finitely presented monoids  $bi\text{-}F_n$  and  $bi\text{-}FP_n$  are equivalent. The *Hochschild cohomological dimension* of  $M$ , written  $\dim M$ , is the length of a shortest projective resolution of  $\mathbb{Z}M$  as a  $\mathbb{Z}[M \times M^{op}]$ -module. The Hochschild cohomological dimension bounds both the left and right cohomological dimension and the geometric dimension bounds the Hochschild cohomological dimension. The geometric dimension also bounds both the left and right geometric dimensions because if  $X$  is a bi-equivariant classifying space for  $M$  of dimension  $n$ , then  $X/M$  is an equivariant classifying space of dimension  $n$ .

**2.4. A theorem of Brown.** We end this section by recalling a result of Brown which will be useful for proofs of results about homological finiteness properties of monoids. Unless otherwise stated, all modules considered here are left modules. Let us say that a module  $V$  over a (unital) ring  $R$  is of type  $FP_n$  if it has a projective resolution that is finitely generated through degree  $n$ ; this is equivalent to having a free resolution that is finitely generated through degree  $n$ ; see [11, Proposition 4.3]. We say that  $V$  is of type  $FP_\infty$  if it has a projective (equivalently, free) resolution that is finitely generated in all degrees. So a monoid is of type  $left\text{-}FP_n$  if and only if the trivial left module is of type  $FP_n$ . One says that  $V$  has *projective dimension* at most  $d$  if it has a projective resolution of length  $d$ . Note that the left cohomological dimension of a monoid is the projective dimension of the trivial left module. Notice also that both the class of modules of type  $FP_n$  and the class of modules having projective dimension at most  $d$  are closed under direct sum.

The following is lemma of K. Brown [9]. Recall that a morphism of chain complexes is a *weak equivalence* if it induces an isomorphism on homology.

**Lemma 2.3.** [9, Lemma 1.5] *Let  $R$  be a ring and  $C = (C_i)$  a chain complex of (left)  $R$ -modules and, for each  $i$ , let  $(P_{ij})_{j \geq 0}$  be a projective resolution of  $C_i$ . Then one can find a chain complex  $Q = (Q_n)$  with  $Q_n = \bigoplus_{i+j=n} P_{ij}$  such that there is a weak equivalence  $f: Q \rightarrow C$ .*

**Corollary 2.4.** *Suppose that  $R$  is a ring and*

$$C_n \longrightarrow C_{n-1} \longrightarrow \cdots \longrightarrow C_0 \longrightarrow V$$

*is a partial resolution of an  $R$ -module  $V$ .*

- (1) *If  $C_i$  is of type  $FP_{n-i}$ , for  $0 \leq i \leq n$ , then  $V$  is of type  $FP_n$ .*
- (2) *Let  $d \geq n$  and suppose that  $C_n \rightarrow C_{n-1}$  is injective. If  $C_i$  has a projective dimension of at most  $d - i$ , for  $0 \leq i \leq n$ , then  $V$  has a projective dimension at most  $d$ .*

*Proof.* To prove the first item, put  $C = (C_i)$  and let  $(P_{ij})_{j \geq 0}$  be a projective resolution of  $C_i$  by finitely generated projectives that is finitely generated through degree  $n - i$ . Then the chain complex  $Q$  from Lemma 2.3 is a complex of projectives with  $Q_k$  finitely generated, for

$0 \leq k \leq n$ , with  $H_0(Q) \cong H_0(C) = V$  and  $H_q(Q) \cong H_q(C) = 0$  for  $0 < q < n$ . Thus if we augment

$$Q_n \longrightarrow Q_{n-1} \longrightarrow \cdots \longrightarrow Q_0$$

by the natural epimorphism  $Q_0 \rightarrow H_0(Q) \cong V$ , we obtain a partial projective resolution of  $V$  of length  $n$  by finitely generated projectives.

For the second item, again let  $C = (C_i)$  and let  $(P_{ij})_{j \geq 0}$  be a projective resolution of  $C_i$  of length at most  $d - i$ . Then the chain complex  $Q$  from Lemma 2.3 is a complex of projectives of length at most  $d$  with  $H_0(Q) \cong H_0(C) \cong V$  and  $H_q(Q) = H_q(C) = 0$  for  $q > 0$ . Thus if we augment  $Q$  by the canonical epimorphism  $Q_0 \rightarrow H_0(Q) \cong V$ , we obtain a projective resolution of  $V$  of length at most  $d$ .  $\square$

Next we show that projective dimension and  $\text{FP}_n$  are stable under flat base extension.

**Lemma 2.5.** *Suppose that  $\varphi: R \rightarrow S$  is a ring homomorphism and that  $S$  is flat as a right  $R$ -module. Let  $V$  be a left  $R$ -module.*

- (1) *If  $V$  is of type  $\text{FP}_n$ , then  $S \otimes_R V$  is of type  $\text{FP}_n$  as an  $S$ -module.*
- (2) *If  $V$  has projective dimension at most  $d$ , then  $S \otimes_R V$  has projective dimension at most  $d$  over  $S$ .*

*Proof.* Since  $S \otimes_R R \cong S$  and tensor products preserve direct sums and retracts, it follows that if  $P$  is a (finitely generated) projective  $R$ -module, then  $S \otimes_R P$  is a (finitely generated) projective  $S$ -module. If  $(P_i)$  is a projective resolution of  $V$ , then by flatness of  $S$  and the preceding observation, we obtain that  $(S \otimes_R P_i)$  is a projective resolution of  $S \otimes_R V$  with  $S \otimes_R P_i$  finitely generated whenever  $P_i$  is. The result follows.  $\square$

A typical way to apply Corollary 2.4 in order to prove that a monoid  $M$  is of type  $\text{FP}_n$  is to find an action of  $M$  by cellular mappings on a contractible CW complex  $X$  such that the  $i^{\text{th}}$ -cellular chain group  $C_i(X)$  is of type  $\text{FP}_{n-i}$  as a  $\mathbb{Z}M$ -module for  $0 \leq i \leq n$ .

### 3. SPECIAL MONOIDS AND ONE-RELATOR MONOIDS

Let  $M$  be the monoid defined by the finite presentation  $\langle A \mid w_1 = 1, \dots, w_k = 1 \rangle$ . Presentations of this form are called *special*, and monoids which admit such presentations are called *special monoids*. Special presentations were first studied by Adjan [1] and Makanin [44]. The main aim of this section is to prove some results which relate the topological and homological finiteness properties of special monoids to the corresponding properties holding in their group of units. By specialising to the case of one-relator monoids and combining with results of Adjan [1] and Lyndon [42] we then obtain a result characterising homological and cohomological finiteness properties of special one-relator monoids. These results answer an important case of the open problem of Kobayashi [34] which asks whether all one-relator monoids are of type right and left- $\text{FP}_\infty$ . As discussed in the introduction to this paper, additional motivation for this question comes from its connection to the question of whether one-relator monoids admit presentations by finite complete rewriting systems which, in turn, relates to the longstanding open problem of whether such monoids have decidable word problem.

For rewriting systems we follow [31, Chapter 12]. We recall some basic definitions and notation here. Let  $A$  be a non-empty set, known as an alphabet, and let  $A^*$  denote the free monoid of all words over  $A$ . If  $w = a_1 a_2 \dots a_n \in A^*$ , with  $a_i \in A$  for  $1 \leq i \leq n$ , then we write  $|w| = n$  and call this the *length* of the word  $w$ . A *rewriting system*  $\mathfrak{R}$  over  $A$  is a subset of  $A^* \times A^*$ . The pair  $\langle A \mid \mathfrak{R} \rangle$  is called a *monoid presentation*. The elements of  $\mathfrak{R}$  are called *rewrite rules*. For words  $u, v \in A^*$  we write  $u \rightarrow_{\mathfrak{R}} v$  if there are words  $\alpha, \beta \in A^*$  and a rewrite rule  $(l, r)$  in  $\mathfrak{R}$  such that  $u = \alpha l \beta$  and  $v = \alpha r \beta$ . We use  $\rightarrow_{\mathfrak{R}}^*$  to denote the reflexive transitive closure of  $\rightarrow_{\mathfrak{R}}$ , while  $\leftrightarrow_{\mathfrak{R}}^*$  denotes the symmetric closure of  $\rightarrow_{\mathfrak{R}}^*$ . The relation  $\leftrightarrow_{\mathfrak{R}}^*$  defines a congruence on  $A^*$  and the quotient  $A^* / \leftrightarrow_{\mathfrak{R}}^*$  is called the *monoid defined by the presentation*  $\langle A \mid \mathfrak{R} \rangle$ . For any word  $w \in A^*$  we use  $[w]_{\mathfrak{R}}$  to denote the  $\leftrightarrow_{\mathfrak{R}}^*$ -class of the word  $w$ . So for words  $u, v \in A^*$



when we write  $u = v$  it means that  $u$  and  $v$  are equal as words in  $A^*$ , while  $[u]_{\mathfrak{R}} = [v]_{\mathfrak{R}}$  means that  $u$  and  $v$  represent the same element of the monoid defined by the presentation. We also sometimes write  $u =_{\mathfrak{R}} v$  to mean the  $[u]_{\mathfrak{R}} = [v]_{\mathfrak{R}}$ . When the set of rewrite rules with respect to which we are working with is clear from context, we shall often omit the subscript  $\mathfrak{R}$  and simply write  $[u]$ ,  $\rightarrow$ ,  $\rightarrow^*$  and  $\leftrightarrow^*$ .

A word  $u$  is called *irreducible* if no rewrite rule can be applied to it, that is, there is no word  $v$  such that  $u \rightarrow v$ . We use  $\text{Irr}(\mathfrak{R})$  to denote the set of irreducible words of the system  $\mathfrak{R}$ . The rewriting system  $\mathfrak{R}$  is *Noetherian* if there is no infinite chain of words  $u_i \in A^*$  with  $u_i \rightarrow u_{i+1}$  for all  $i \geq 1$ . The system is *confluent* if whenever  $u \rightarrow^* u_1$  and  $u \rightarrow^* u_2$  there is a word  $v \in A^*$  such that  $u_1 \rightarrow^* v$  and  $u_2 \rightarrow^* v$ . A rewriting system that is both Noetherian and confluent is called *complete*. If  $\mathfrak{R}$  is a complete rewriting system then each  $\leftrightarrow^*$  equivalence class contains a unique irreducible word. Thus in this situation,  $\text{Irr}(\mathfrak{R})$  provides a set of normal forms for the elements of the monoid defined by the presentation  $\langle A \mid \mathfrak{R} \rangle$ .

Let  $M = \langle A \mid w_1 = 1, \dots, w_k = 1 \rangle = \langle A \mid T \rangle$  be the finitely presented special monoid defined above. The symbol  $M$  will be used to denote this monoid for the remainder of this section. We call  $w_1, w_2, \dots, w_k$  the *defining relators* of this presentation. Let  $\Gamma(M, A)$  denote the right Cayley graph of  $M$  with respect to  $A$ . The strongly connected components of  $\Gamma(M, A)$  are called the *Schützenberger graphs* of  $M$ . Here we say that two vertices  $u$  and  $v$  of a directed graph belong to the same strongly connected component if and only if there is a directed path from  $u$  to  $v$ , and also a directed path from  $v$  to  $u$ . Our aim is to prove that any two Schützenberger graphs of  $M$  are isomorphic to each other and that, modulo the Schützenberger graphs, the Cayley graph of  $M$  has a tree-like structure. We begin by summarising some results of Zhang [63] on special monoids that will be used extensively below.

Let  $G$  be the group of units of  $M$ . By [63, Theorem 3.7], we have that  $G$  has a group presentation with  $k$  defining relations. Let  $R$  be the submonoid of right invertible elements. Then  $R$  is isomorphic to a free product of  $G$  with a finitely generated free monoid by [63, Theorem 4.4].

In more detail, we say that a word  $u \in A^*$  is invertible if  $[u] \in M$  is invertible. Let  $u \in A^+$  be a non-empty invertible word. We say that the invertible word  $u$  is *indecomposable* if no non-empty proper prefix of  $u$  is invertible. Every non-empty invertible word  $v$  has a unique decomposition  $v = v_1 v_2 \dots v_l$  where each  $v_i$  is indecomposable. To obtain this decomposition, first write  $v = v_1 u_1$  where  $v_1$  is the shortest non-empty invertible prefix of  $v$ . Since  $v$  and  $v_1$  are invertible it follows that  $u_1$  is invertible. If  $u_1$  is non-empty we repeat this process writing  $u_1 = v_2 u_2$  where  $v_2$  is the shortest non-empty invertible prefix of  $u_1$ . Continuing in this way gives the decomposition  $v = v_1 v_2 \dots v_l$ . It is unique since if  $v'_1 v'_2 \dots v'_k$  were some other such decomposition then  $v_1 v_2 \dots v_l = v'_1 v'_2 \dots v'_k$ , neither  $v_1$  nor  $v'_1$  can be a proper prefix of the other, hence  $v_1 = v'_1$ , and then inductively we see that  $v_i = v'_i$  for all  $i$ . We call  $u \in A^+$  a *minimal invertible word* if it is indecomposable and invertible and the length of  $u$  does not exceed the length of any of the relators in  $T$ . Each relation word  $w_i$  in  $T$  represents the identity of  $M$  and thus is invertible. Therefore each relation word  $w_i$  has a unique decomposition  $w_i = w_{i,1} w_{i,2} \dots w_{i,n_i}$  into indecomposable invertible words. The words  $w_{i,j}$  for  $1 \leq i \leq n$ ,  $1 \leq j \leq n_j$  are called the *minimal factors* of the relators of the presentation. Each minimal factor is clearly a minimal invertible word.

Let  $\Delta$  be the set of all minimal invertible words  $\delta \in A^*$  such that  $\delta$  is equal in  $M$  to at least one of the minimal factors  $w_{i,j}$  of the relators. Clearly  $\Delta$  is a finite set of words over  $A$ . It is also immediate from the definition that  $\Delta$  contains in particular all of the minimal factors  $w_{i,j}$  of the relators. It is also a consequence of the definitions that no non-empty proper prefix of a word from  $\Delta$  can be equal to a non-empty proper suffix of a word from  $\Delta$ . On the other hand, a word from  $\Delta$  can, in general, arise as a subword of a word from  $\Delta$  (and there are examples where this happens). It also follows from the definitions that  $\Delta$  is a prefix code, meaning that no word from  $\Delta$  is a prefix of any other word from  $\Delta$ . It follows that  $\Delta$  freely generates a free submonoid of  $A^*$ .

The elements represented by the words from  $\Delta$  give a finite generating set for the group of units  $G$  of the monoid  $M$ . Indeed, it may be shown that every indecomposable invertible word  $v$  is equal in  $M$  to some word from  $\Delta$ ; see [63, Lemma 3.4], and every invertible word can be written as a product of indecomposable invertible words.

A finite presentation for the group of units  $G$  of  $M$ , with respect to the finite generating set  $\Delta$  may be constructed in the following way. We partition the finite set of words  $\Delta$  as the disjoint union  $\Delta = \Delta_1 \cup \Delta_2 \cup \dots \cup \Delta_m$  of non-empty sets where two words belong to the same set  $\Delta_j$  if and only if they represent the same element of the monoid  $M$ . Note that two distinct factors  $w_{i,j}$  could well represent the same element of  $M$  even if they are not equal as words. Set  $B = \{b_1, b_2, \dots, b_m\}$  and define a map  $\phi$  from  $\Delta$  to  $B$  which maps every word from the set  $\Delta_j$  to the letter  $b_j$ . Extend this to a surjective homomorphism  $\phi: \Delta^* \rightarrow B^*$ . Note that for any word  $v \in A^*$ , if  $v \in \Delta^*$  then as observed above  $v$  has a unique decomposition  $v = v_1 v_2 \dots v_l$  where each  $v_i \in \Delta^*$  and thus the mapping  $\phi$  is well-defined on the subset  $\Delta^*$  of  $A^*$ . Let  $T_0$  be the rewriting system over the alphabet  $B$  given by applying  $\phi$  to each of the relators from the presentation  $\langle A \mid T \rangle$  (recall that each  $w_j \in \Delta^*$ ) to obtain

$$T_0 = \{(s, 1) : s \text{ is some cyclic permutation of some } \phi(w_j)\}.$$

This means for each relator  $w_j$  from  $T$ , we decompose  $w_j$  into its minimal factors, then read the factors recording the sets  $\Delta_i$  to which each of them belongs, and then write down the corresponding word over  $B$ , and all of its cyclic conjugates.

**Theorem 3.1.** [63, Theorem 3.7] *Let  $M$  be the monoid defined by a finite special presentation  $\langle A \mid T \rangle$ . Then  $\langle B \mid T_0 \rangle$  is a finite monoid presentation for the group of units  $G$  of  $M$ .*

It follows that  $\langle B \mid \phi(w_1) = 1, \dots, \phi(w_k) = 1 \rangle$  is a group presentation for the group of units of  $M$  with the same number of defining relations as the presentation of  $M$ .

Choose and fix some order on the finite alphabet  $A$ , and for words  $x, y \in A^*$  write  $x < y$  if  $x$  precedes  $y$  in the resulting shortlex ordering [31, Definition 2.60]. Now define a rewriting system  $S = S(T)$  over  $A$  as follows:

$$S = \{(u, v) \mid u, v \in \Delta^* : \phi(u) =_{T_0} \phi(v) \text{ \& } u > v\}.$$

In fact, it follows from the results of Zhang that the condition  $\phi(u) =_{T_0} \phi(v)$  is equivalent to saying that  $u =_T v$ , i.e. that  $u$  and  $v$  represent the same element of the group of units of the monoid  $M$ . So the condition  $\phi(u) =_{T_0} \phi(v)$  could be replaced by the condition  $u =_T v$  in the definition of  $S$ .

**Theorem 3.2.** [63, Proposition 3.2] *The infinite presentation  $\langle A \mid S \rangle$  is Noetherian, confluent and defines the monoid  $M$ . In fact, the rewriting systems  $T$  and  $S = S(T)$  are equivalent, that is,  $\leftrightarrow_S^* = \leftrightarrow_T^*$ .*

We shall prove statements about  $M$  by working with the irreducible words  $\text{Irr}(S)$  associated with this infinite complete rewriting system. For the rest of this section, when we say a word over the alphabet  $A$  is irreducible, we mean that it is irreducible with respect to the rewriting system  $S$ .

The submonoid of right units  $R$  is generated by the prefixes of the words from  $\Delta$ . Indeed, let  $I$  be the set of non-empty prefixes of words from  $\Delta$ , that is,

$$I = \{x \in A^+ \mid xy \in \Delta \text{ for some } y \in A^*\}.$$

Clearly all words in the set  $I$  represent right invertible elements of  $M$ . Conversely, we have the following result.

**Lemma 3.3.** [63, Lemma 3.3] *Let  $u \in A^*$  be irreducible modulo  $S = S(T)$ . If  $[u]_T$  is right invertible, then  $u \in I^*$ .*

It follows from this lemma that  $I$  constitutes a finite generating set for the submonoid  $R$  of right units of the monoid  $M$  (that is, the submonoid of all right invertible elements). Furthermore, in [63, Theorem 4.4] Zhang proves the following result which describes the structure of the submonoid of right units of the monoid  $M$ .

**Theorem 3.4.** [63, Theorem 4.4] *Let  $M$  be a finitely presented special monoid. The submonoid of right units  $R$  of  $M$  is a free product of the group of units  $G$  and a finitely generated free monoid.*

Monoid free products will be formally defined in Section 4 below. Our next goal is to show that the Cayley graph of a special monoid has a tree-like structure. The action of the monoid on the corresponding tree will be used to construct a free resolution of the trivial module.

Let  $\mathcal{T}$  be the set of irreducible words in  $A^*$  with no suffix in  $I$ .

**Lemma 3.5.** *Let  $w \in \mathcal{T}$  and let  $u \in A^*$  be irreducible. Then  $wu$  is irreducible.*

*Proof.* If  $wu$  is not irreducible, then since both  $w$  and  $u$  are irreducible it follows that  $w = xy$  and  $u = zw$  with  $yz$  a left-hand side of a rewrite rule and  $y, z$  both non-empty. But every left hand side of a rewrite rule is in  $\Delta^*$  and so  $y$  has a non-empty suffix  $v$  that is a prefix of an element of  $\Delta$ . But then  $v \in I$ , contradicting that  $w \in \mathcal{T}$ .  $\square$

We recall the definition of the pre-order  $\leq_{\mathcal{R}}$  on the monoid  $M$ . For all  $m, n \in M$  we write  $m \leq_{\mathcal{R}} n$  if and only if  $mM \subseteq nM$ , and write  $m \mathcal{R} n$  if  $m \leq_{\mathcal{R}} n$  and  $n \leq_{\mathcal{R}} m$ . Obviously  $\mathcal{R}$  is an equivalence relation on  $M$ , usually called Green's  $\mathcal{R}$ -relation, and  $M/\mathcal{R}$  is a poset with the order induced by  $\leq_{\mathcal{R}}$ . In terms of the right Cayley graph  $\Gamma(M, A)$  of  $M$  we have  $m \leq_{\mathcal{R}} n$  if and only if there is a directed path from  $n$  to  $m$ , while the  $\mathcal{R}$ -classes are the vertex sets of the Schützenberger graphs of the monoid.

Let  $\mathcal{L}$  be a subset of  $A^*$  containing the empty word. For any two words  $\alpha, \beta \in \mathcal{L}$  write  $\alpha \prec \beta$  if and only if  $\beta$  is a prefix of  $\alpha$ . This defines a poset which we denote by  $P_{\mathcal{L}}$ . This poset is the reversal of the prefix order on the set of words  $\mathcal{L}$ . This poset is countable since  $A$  is finite. The empty word is the unique maximal element of the poset. This poset is *locally-finite* in the sense that every interval  $[x, y]$  in this poset contains finitely many elements. In fact the principal filter of every element in this poset is finite since a word admits only finitely many prefixes. Recall that if  $s$  and  $t$  are elements of a poset  $P$  then we say  $s$  *covers*  $t$  if  $s < t$  and  $[s, t] = \{s, t\}$ . A locally finite poset is completely determined by its cover relations. The *Hasse diagram* of a poset  $P$  is a graph whose edges are the cover relations. Hasse diagrams are drawn in such a way that if  $s < t$  then  $t$  is drawn with a higher vertical coordinate than  $s$ .

**Proposition 3.6.** *Let  $\mathcal{L} \subseteq A^*$  contain the empty word. Then the Hasse diagram of  $P_{\mathcal{L}}$  is a rooted tree (with root the empty word).*

*Proof.* For  $n \geq 0$ , let  $\mathcal{L}_n$  consist of those words from  $\mathcal{L}$  of length at most  $n$ . Let  $\Lambda$  (respectively,  $\Lambda_n$ ) be the Hasse diagram of  $P_{\mathcal{L}}$  (respectively,  $P_{\mathcal{L}_n}$ ). Then  $\Lambda = \varinjlim \Lambda_n$  and hence, since a direct limit of trees is a tree, it suffices to handle the case that  $\mathcal{L}$  is finite. We proceed by induction on  $|\mathcal{L}|$ . If  $|\mathcal{L}| = 1$ , then  $\Lambda$  consists of a single vertex and there is nothing to prove. Assume true for languages with at most  $n$  elements and suppose that  $\mathcal{L}$  has  $n + 1$  elements. Suppose that  $w \in \mathcal{L}$  has maximum length. Let  $v$  be the longest proper prefix of  $w$  belonging to  $\mathcal{L}$  (it could be the empty word). Let  $\Lambda'$  be the Hasse diagram of  $P_{\mathcal{L} \setminus \{w\}}$ ; it is a rooted tree with root the empty word by induction. Then there is an edge between  $v$  to  $w$  in  $\Lambda$  and that is the only edge incident on  $w$ . Hence  $\Lambda$  and  $\Lambda'$  have the same Euler characteristic and so  $\Lambda$  is a tree (as  $\Lambda'$  was).  $\square$

It is possible for an element of  $P_{\mathcal{L}}$  to cover infinitely many distinct elements of  $P_{\mathcal{L}}$ . For example, if  $\mathcal{L} = \{\epsilon, ab, aab, aaab, aaaab, \dots\}$  then  $\epsilon$  covers all the other words in this set.

The following fact is essentially established in [63, Lemma 5.2] and the discussion afterwards.

**Proposition 3.7.** *Every element  $m \in M$  can uniquely be expressed in the form  $m = [w_m]u_m$  with  $w_m \in \mathcal{T}$  and  $u_m \in R$ . Moreover, the irreducible word  $v \in A^*$  representing  $m$  is  $w_mt$  where  $t \in I^*$  is the longest suffix of  $v$  in  $I^*$  and  $[t] = u_m$ . Furthermore, if  $m, n \in M$ , then  $m \leq_{\mathcal{R}} n$  if and only if  $w_n$  is a prefix of  $w_m$ . Hence the Hasse diagram of  $M/\mathcal{R}$  is a tree rooted at 1.*

*Proof.* Let  $v \in A^*$  be the irreducible word with  $[v] = m$ . Then  $v = v'v''$  where  $v''$  is the longest suffix in  $I^*$ . It follows that  $v' \in \mathcal{T}$  and  $v''$  represents an element of  $R$ . This shows the existence of such a factorization. For uniqueness, let  $w \in \mathcal{T}$  and  $x \in A^*$  be an irreducible word representing an element of  $R$ . By [63, Lemma 3.3], we have that  $x \in I^*$ . Then  $wx$  is irreducible by Lemma 3.5. Thus  $wx = v'v''$ . By choice of  $v''$ , we must have  $|x| \leq |v''|$ . If  $|x| < |v''|$ , then some non-empty prefix of  $v''$  is a suffix of  $w$ . As  $I$  is prefix-closed, whence so is  $I^*$ , this contradicts that  $w \in \mathcal{T}$ . Thus  $x = v''$  and hence  $w = v'$ . This establishes the uniqueness of the decomposition.

Suppose now that  $m = nn'$  with  $n' \in M$ . Let  $z$  be a right inverse of  $u_m$  and let  $v$  be an irreducible word representing  $u_n n' z$ . Then  $w_n v$  is an irreducible word representing  $nn'z = mz = [w_m]u_m z = [w_m]$  by Lemma 3.5. Thus  $w_m = w_n v$  and so  $w_n$  is a prefix of  $w_m$ . Conversely, suppose that  $w_n$  is a prefix of  $w_m$ . Clearly,  $[w_n] \mathcal{R} n$  and  $[w_m] \mathcal{R} m$  as  $u_m, u_n$  are right invertible. So it suffices to observe that  $[w_m] \leq_{\mathcal{R}} [w_n]$ .

The final statement follows from Proposition 3.6.  $\square$

Retaining the notation of Proposition 3.7 we obtain the following immediate corollary.

**Corollary 3.8.** *The action of  $R$  on the right of  $M$  is free with transversal  $\mathcal{T} = \{[w] \mid w \in \mathcal{T}\}$ . Furthermore,  $M/R \cong M/\mathcal{R}$ .*

Another corollary is that all principal right ideals of  $M$  are isomorphic as right  $M$ -sets.

**Corollary 3.9.** *Let  $n \in M$ . Then the mapping  $\varphi_n: M \rightarrow nM$  given by  $\varphi_n(m) = [w_n]m$  is an isomorphism of right  $M$ -sets.*

*Proof.* As  $nM = [w_n]M$ , the map  $\varphi_n$  is clearly a surjective homomorphism of right  $M$ -sets. To see that is an isomorphism, suppose that  $\varphi_n(m) = \varphi_n(m')$ . Let  $v, v' \in A^*$  be irreducible words representing  $m, m'$ , respectively. Then  $w_n v$  and  $w_n v'$  are irreducible by Lemma 3.5. As they represent the same element of  $M$ , we deduce that  $v = v'$  and so  $m = m'$ .  $\square$

We now generalize Corollary 3.9 to show that every right ideal of  $M$  is a free  $M$ -set.

**Theorem 3.10.** *Let  $M$  be a special monoid. Then every right ideal of  $M$  is a free right  $M$ -set and dually every left ideal of  $M$  is a free left  $M$ -set.*

*Proof.* Let  $X$  be a right ideal of  $M$  and let  $X' = \{w \in \mathcal{T} \mid [w] \in X\}$ . Let  $U'$  be the set of elements  $w \in X'$  with no proper prefix in  $X'$ . We claim that  $X$  is freely generated as an  $M$ -set by  $U = \{[w] \mid w \in U'\}$ . By Proposition 3.7 if  $s, t \in U$  are distinct, then  $sM \cap tM = \emptyset$ . Indeed, if  $m \in sM \cap tM$ , then  $w_m$  has both  $w_s$  and  $w_t$  as prefixes and hence either  $w_s$  is a prefix of  $w_t$ , or vice versa, contradicting the definition of  $U'$ . Also, by Corollary 3.9, for each  $s \in U$ , we have that  $sM \cong M$  as a right  $M$ -set. It follows that  $U$  freely generates a sub- $M$ -subset  $Y$  of  $X$ . We show that  $Y = X$ .

If  $m \in X$ , then  $m = [w_m]u_m$  with  $w_m \in \mathcal{T}$  and  $u_m \in R$ . Then  $[w_m] \in X$  as  $[w_m] \mathcal{R} m$ . Let  $w \in \mathcal{T}$  be the shortest prefix of  $w_m$  with  $[w] \in X$ . Then  $w \in U'$  and  $m \in [w]M \subseteq Y$ . This completes the proof.  $\square$

*Remark 3.11.* Note that if  $X$  is a free right  $M$ -set on a subset  $B$  and if  $X$  has a finite generating set, then  $B$  is finite. Indeed, if  $C$  is a finite generating set for  $X$ , then there is a finite subset  $B' \subseteq B$  such that  $C \subseteq B'M$ . But then  $B \subseteq B'M$  and hence  $B = B'$  by freeness of the action.

Let  $\Gamma(M, A)$  be the Cayley graph of  $M$  with respect to  $A$ . Let  $\Gamma(M, A, m)$  denote the strongly connected component of  $m$  (also called the Schützenberger graph of  $m$ ). An immediate geometric consequence of Corollary 3.9 is the following.

**Corollary 3.12.** *Let  $n \in M$ . Then there is an isomorphism of  $A$ -labeled graphs  $\Gamma(M, A, 1) \rightarrow \Gamma(M, A, n)$  sending 1 to  $[w_n]$ . If  $\Gamma_n$  is the induced subgraph of  $\Gamma(M, A)$  consisting of all vertices accessible from  $n$ , then  $\Gamma(M, A)$  is isomorphic to  $\Gamma_n$  as an  $A$ -labeled graph via an isomorphism taking 1 to  $[w_n]$ .*

Corollary 3.12 recovers as a special case the result [45, Theorem 4.6] that all the maximal subgroups of a special monoid are isomorphic to each other. This is because the Schützenberger group of a regular  $\mathcal{R}$ -class is isomorphic to the automorphism group of its labelled Schützenberger graph [59, Theorem 3].

Next we wish to show that there is a unique edge entering any strongly connected component of  $\Gamma(M, A)$  other than the strong component of 1, and that it ends at an element of  $\mathcal{T}$  (see Corollary 3.8 for the notation). Let us say that an edge of a digraph *enters* a strong component  $C$  of the graph if its initial vertex is not in  $C$  and its terminal vertex is in  $C$ .

**Proposition 3.13.** *Let  $n \in \mathcal{T} \setminus \{1\}$  (and so  $n = [w_n]$ ). Then if  $w_n = xa$  with  $a \in A$ , we have that  $[x] >_{\mathcal{R}} n$ ,  $[a] \notin R$  and  $[x] \xrightarrow{a} n$  is the unique edge entering  $\Gamma(M, A, n)$ .*

*Proof.* Note that  $x$  is irreducible. Let  $x = x'x''$  with  $x''$  the longest suffix of  $x$  in  $I^*$ . Then  $x' = w_{[x]}$  and  $w_n$  is not a prefix of  $x'$ . Thus  $[x] >_{\mathcal{R}} [w_n] = n$  by Proposition 3.7. It follows that  $[x] \xrightarrow{a} n$  enters  $\Gamma(M, A, n)$  and hence  $a \notin R$ .

Suppose that  $m \xrightarrow{b} m'$  enters  $\Gamma(M, A, n)$ . Let  $w$  be an irreducible word representing  $m$ . Then  $w = w_my$  where  $y \in I^*$  is the longest suffix of  $w$  in  $I^*$ . We claim that  $wb$  has no suffix in  $I$ . Indeed, if it did, then since  $I$  is prefix-closed and  $w_m$  has no suffix in  $I$ , we must have that  $yb$  has a suffix in  $I$ . Then  $yb = rs$  where  $s \in I$ . Since  $r$  is a prefix of  $y$  and  $I$  (and hence  $I^*$ ) is prefix-closed, we obtain that  $yb = rs \in I^*$ . Thus  $yb$  represents an element of  $R$  and so

$$m' = [w_myb] \mathcal{R} [w_m] \mathcal{R} m$$

a contradiction. Thus  $wb$  has no suffix in  $I$ .

We claim that  $wb$  is irreducible. Suppose that  $wb$  is not irreducible. Then since  $w$  is irreducible, each left-hand side in the rewriting system belongs to  $\Delta^*$  and  $\Delta \subseteq I$ , we must have that  $wb$  has a suffix in  $I$ , a contradiction.

Putting it all together, we deduce that  $wb \in \mathcal{T}$  and so  $wb = w_n$  by Proposition 3.7. It follows that  $b = a$  and  $w = x$ , completing the proof.  $\square$

Let  $\Gamma$  be the directed graph obtained from  $\Gamma(M, A)$  by collapsing each strongly connected component (and its internal edges) to a point. So the vertex set of  $\Gamma$  is  $M/\mathcal{R}$  and there is an edge  $(m, a)$  from the  $\mathcal{R}$ -class  $R_m$  of  $m$  to the  $\mathcal{R}$ -class  $R_{ma}$  of  $ma$  if  $m \in M$ ,  $a \in A$  and  $R_m \neq R_{ma}$ . We aim to show that  $\Gamma$  is a regular rooted tree isomorphic to the Hasse diagram of  $M/\mathcal{R}$ . Note that this tree can be of infinite degree.

**Theorem 3.14.** *The graph  $\Gamma$  is isomorphic as a digraph to the Hasse diagram of  $M/\mathcal{R}$  ordered by  $\geq_{\mathcal{R}}$ . This graph is a regular rooted tree with root the strong component of 1.*

*Proof.* We retain the above notation. Suppose first that  $w, w' \in \mathcal{T}$  and there is an edge from  $\Gamma(M, A, [w'])$  to  $\Gamma(M, A, [w])$ ; it is unique by Proposition 3.13. Then, by Proposition 3.13, we have that if  $w = xa$  with  $a \in A$ , then  $[x] \mathcal{R} [w']$ . Thus if  $x'$  is the longest suffix of  $x$  belonging to  $I^*$ , then  $x = w'x'$  and  $w = w'x'a$ . Since  $I$  is prefix-closed, it follows that if  $y$  is any non-empty prefix of  $x'$ , then  $w'y$  has a suffix in  $I$  and hence does not belong to  $\mathcal{T}$ . Thus in the prefix order on  $\mathcal{T}$ , there is no element between  $w'$  and  $w$ . It follows from Proposition 3.7 that in the Hasse diagram of  $M/\mathcal{R}$  with respect to  $\geq_{\mathcal{R}}$ , there is an edge from  $R_{[w']}$  to  $R_{[w]}$ .

Conversely, suppose that there is an edge in the Hasse diagram from  $R_{[w']}$  to  $R_{[w]}$  with  $w, w' \in \mathcal{T}$ . Then  $w'$  is a proper prefix of  $w$  by Proposition 3.7 and so  $w = w'y$  with  $y \in A^*$  irreducible and non-empty. Let  $a \in A$  be the last letter of  $y$ , so  $y = y'a$ . Then  $[w'] \leq_{\mathcal{R}} [w'y'] \leq_{\mathcal{R}} [w]$  and so one of these inequalities is an equality. Since  $w$  is not a prefix of  $w'y'$ , it follows from Proposition 3.7 (or by [63, Lemma 5.2]) that the second inequality is strict. Thus  $[w'y']$

belongs to the strong component of  $[w']$  and the image of the edge  $[w'y'] \xrightarrow{a} [w]$  connects the strong component of  $[w']$  to the strong component of  $[w]$  in  $\Gamma$  (and is the only such edge by Proposition 3.13).

Since the reverse prefix order on any set of words containing the empty word is a rooted tree, it follows that  $\Gamma$  is a rooted tree with root the strong component of 1. By construction of  $\Gamma$  and Corollary 3.12 it follows that all vertices have the same cardinality set of children.  $\square$

Note that in general if  $M$  is a monoid generated by a finite set  $A$ , and if  $R'$  and  $R''$  are  $\mathcal{R}$ -classes of  $M$  such that  $R'$  covers  $R''$  in the poset  $M/\mathcal{R}$ , then there must exist elements  $x \in R'$  and  $y \in R''$  and a generator  $a \in A$  such that  $xa = y$  in  $M$ . The second part of the proof of the above theorem shows that in a finitely generated special monoid in this situation there are unique elements  $x \in R', y \in R''$  and  $a \in A$  satisfying these properties.

We note that the left action of  $M$  on  $\Gamma(M, A)$  induces a left action of  $M$  on  $\Gamma$  by cellular mappings since strong components are mapped into strong components. However, elements of  $m$  can collapse edges to a point. In fact,  $\Gamma$  (being a tree) is a simplicial graph (1-dimensional simplicial complex) and  $M$  acts by simplicial mappings. For example, consider the bicyclic monoid  $B = \langle a, b \mid ab = 1 \rangle$ . Then since  $a \mathcal{R} 1$  left multiplication by  $a$  collapses the vertices corresponding to the strong components of 1 and  $b$  and hence collapses the edge between these components.

We can view the vertex set of  $\Gamma$  as  $M/R$  and so if we use the simplicial chain complex for  $\Gamma$ , we have  $C_0(\Gamma) \cong \mathbb{Z}[M/R] \cong \mathbb{Z}M \otimes_{\mathbb{Z}R} \mathbb{Z}$  as a  $\mathbb{Z}M$ -module. We can identify  $C_1(\Gamma)$  as a  $\mathbb{Z}M$ -module with the quotient  $C_1(\Gamma(M, A))/N$  where  $N$  is the  $\mathbb{Z}M$ -submodule generated as an abelian group by edges  $m \xrightarrow{a} ma$  with  $a \in A$  and  $m \mathcal{R} ma$ . Note that  $C_1(\Gamma(M, A))$  is a free  $\mathbb{Z}M$ -module of rank  $|A|$ . We shall show that  $N$  is a free  $\mathbb{Z}M$ -module of finite rank, as well. It will then follow that  $C_1(\Gamma)$  is of type  $\text{FP}_\infty$  with projective dimension at most 1.

Note that  $N$  is the direct sum over all  $a \in A$  of the submodules  $N_a$  spanned by edges  $m \xrightarrow{a} ma$  with  $m \mathcal{R} ma$  and so it suffices to show that each of these submodules  $N_a$  is a finitely generated free  $\mathbb{Z}M$ -module.

**Proposition 3.15.** *Let  $a \in A$ . Then  $N_a$  is a finitely generated free  $\mathbb{Z}M$ -module. Consequently,  $N$  is a finitely generated free  $\mathbb{Z}M$ -module.*

*Proof.* Let  $L = \{m \in M \mid m \mathcal{R} ma\}$ . Then  $L$  is a left ideal of  $M$  and  $N_a \cong \mathbb{Z}L$ . First observe that if  $a \in R$ , then  $L = M$  and there is nothing to prove. So assume that  $a \in A \setminus R$ . By Theorem 3.10 we have that  $L$  is a free left  $M$ -set. By Remark 3.11 it suffices to prove that  $L$  is finitely generated.

We claim that  $L$  is generated by  $I' = \{[w] \in L \mid w \in I\}$ , which is finite as  $I$  is finite. Let  $m \in L$  and let  $w \in A^*$  be irreducible with  $[w] = m$ . There are two cases. Assume first that  $wa$  is irreducible. Then since  $ma \mathcal{R} m$ , it follows from Proposition 3.7 that  $wa \notin \mathcal{T}$  (as  $wa$  is not a prefix of  $w$ ) and so  $wa = sxa$  with  $xa \in I$ . Since  $a \notin R$ , we must have that  $x$  is non-empty. Since  $I$  is prefix-closed,  $x \in I$ . Thus  $[x], [xa] \in R$  and hence  $[x] \mathcal{R} [x]a$ . Then  $m = [s][x]$  and  $[x] \in I'$ . So  $m \in MI'$ .

Next assume that  $wa$  is not irreducible. Then  $wa = sxa$  with  $xa \in \Delta$ , as  $w$  is irreducible. But  $a \notin R$  and so  $x$  is non-empty. Thus  $x \in I$ . Also  $xa \in \Delta \subseteq I$ . Thus  $[x], [xa] \in R$  and so  $[x] \mathcal{R} [x]a$ . Also,  $m = [s][x]$  with  $[x] \in I'$  and so  $m \in MI'$ . This completes the proof.  $\square$

Now all is in place to prove the first main result of this section.

**Theorem 3.16.** *Let  $M$  be a finitely presented special monoid with group of units  $G$ .*

- (1) *If  $G$  is of type  $\text{FP}_n$  with  $1 \leq n \leq \infty$ , then  $M$  is of type left- $\text{FP}_n$  and of type right- $\text{FP}_n$ .*
- (2)  *$\text{cd } G \leq \text{left cd } M \leq \max\{2, \text{cd } G\}$  and  $\text{cd } G \leq \text{right cd } M \leq \max\{2, \text{cd } G\}$ .*

*Proof.* We retain the above notation. We prove the results for left- $\text{FP}_n$  and left cohomological dimension (the other results are dual). First note that if  $L$  denotes the submonoid of left invertible elements, then  $M$  is a free left  $L$ -set by the dual of Proposition 3.7. If  $B$  is the basis

of  $M$  as a left  $L$ -set, then each element  $m \in M$  can be expressed uniquely as  $u_m b_m$  with  $b_m \in B$  and  $u_m \in L$ . But then if  $g \in G$  with  $gm = m$ , we must have  $gu_m b_m = u_m b_m$ . It follows that  $gu_m = u_m$  by uniqueness. But since  $L$  is a free product of  $G$  with a finitely generated free monoid by [63, Theorem 4.4], it follows that  $G$  acts freely on the left of  $L$  and so  $g = 1$ . Thus  $\mathbb{Z}M$  is a free left  $\mathbb{Z}G$ -module and so  $\text{cd } G \leq \text{left cd } M$  as any projective resolution of  $\mathbb{Z}$  over  $\mathbb{Z}M$  is a projective resolution over  $\mathbb{Z}G$ .

The graph  $\Gamma$  is a tree with a simplicial action by  $M$  described above. So we have an exact sequence of  $\mathbb{Z}M$ -modules

$$0 \longrightarrow C_1(\Gamma) \longrightarrow C_0(\Gamma) \longrightarrow \mathbb{Z} \longrightarrow 0.$$

We have identified  $C_1(\Gamma) \cong C_1(\Gamma(M, A))/N$  where  $C_1(\Gamma(M, A))$  is free of rank  $|A|$  and  $N$  is a finitely generated free module by Proposition 3.15. Thus  $C_1(\Gamma)$  is of type  $\text{FP}_\infty$  and has projective dimension at most 1.

On the other hand,  $C_0(\Gamma) \cong \mathbb{Z}[M/R] \cong \mathbb{Z}M \otimes_{\mathbb{Z}R} \mathbb{Z}$ . By Zhang's theorem [63, Theorem 4.4],  $R = G * C^*$  where  $C$  is a finite alphabet, and hence  $R$  is of type  $\text{FP}_n$  whenever  $G$  is, and,  $\text{left cd } R \leq \max\{1, \text{cd } G\}$  by [17, Theorem 5.5] (or see Corollaries 4.6 and 4.9 below). Note that a finitely generated free monoid is of type  $\text{FP}_\infty$  and of cohomological dimension 1 because its Cayley graph is a tree and a free  $M$ -CW complex of finite type of dimension 1.

As  $\mathbb{Z}M$  is a free, and hence flat, right  $\mathbb{Z}R$ -module by Corollary 3.8, it follows from Lemma 2.5 that  $C_0(\Gamma) \cong \mathbb{Z}M \otimes_{\mathbb{Z}R} \mathbb{Z}$  is of type  $\text{FP}_n$  and of projective dimension at most  $\max\{1, \text{cd } G\}$ .

The result now follows from an application of Corollary 2.4.  $\square$

In general, the left- and right-cohomological dimensions of a monoid are not equal. In fact they are completely independent of each other; see [27]. One immediate corollary of the above result is that if  $M$  is a finitely presented special monoid with left- and right-cohomological dimensions both at least equal to 2, then the left cohomological dimension of  $M$  is equal to its right cohomological dimension.

As an application of Theorem 3.16 we now show how it can be used to prove that all special one-relator monoids are of type  $\text{FP}_\infty$ , answering a case of a question of Kobayashi. We also recover Kobayashi's result (see [33, Theorem 7.2] and [34, Corollary 7.5]) that if the relator is not a proper power then the cohomological dimension is at most 2.

A word  $u \in A^*$  is called *primitive* if it is not a proper power in  $A^*$ .

**Lemma 3.17.** [41, Corollary 4.2] *For every nonempty word  $w \in A^*$  there is a unique primitive word  $p$  and a unique integer  $k \geq 1$  such that  $w = p^k$ .*

The following lemma is well known. We include it here for completeness.

**Lemma 3.18.** *Let  $M = \langle A \mid w = 1 \rangle$ . Write  $w = p^k$  where  $p$  is a primitive word and  $k \geq 1$ . The group of units  $G$  of  $M$  is a one-relator group with torsion if and only if  $k > 1$ .*

*Proof.* Since it is a prefix and suffix of  $w$ , it follows that  $p$  is invertible in  $M$ . Therefore, the decomposition of  $w$  into indecomposable invertible factors has the form  $w = (p_1 p_2 \dots p_l)^k$  where  $p_1 p_2 \dots p_l$  is the decomposition of  $p$  into indecomposable invertible factors. Let  $P = \{p_i : 1 \leq i \leq l\} \subseteq A^*$ . Let  $X = \{x_p : p \in P\}$  be an alphabet in bijection with the set of words  $P$ , so distinct words  $p_i$  and  $p_j$  from  $P$  correspond to distinct letters  $x_{p_i}$  and  $x_{p_j}$  from the alphabet  $X$ . It follows from [1, Lemma 96] that the group of units of the monoid  $M$  is isomorphic to the group defined by the group presentation  $\text{Gp}\langle X \mid (x_{p_1} x_{p_2} \dots x_{p_l})^k = 1 \rangle$ . Observe that  $x_{p_1} x_{p_2} \dots x_{p_l} \in X^*$ , i.e. this is a positive word over the alphabet  $X$ . In particular the word  $(x_{p_1} x_{p_2} \dots x_{p_l})^k$  is cyclically reduced. Since the word  $p_1 p_2 \dots p_l$  is primitive by assumption it follows that the word  $x_{p_1} x_{p_2} \dots x_{p_l} \in X^*$  is also primitive. Hence  $(x_{p_1} x_{p_2} \dots x_{p_l})^k$  is a proper power if and only if  $k > 1$ . But then by a well-known result of Karrass, Magnus and Solitar characterising elements of finite order in one-relator groups [43, Theorem 5.2] it follows that the group of units of  $M$  is a one-relator group with torsion if and only if  $k > 1$ .  $\square$

Well-written accounts of the result [1, Lemma 96] of Adjan used in the previous proof may be found in [38, Section 1] and [39, Section 2]. The following result gives a positive answer to Kobayashi's question [34, Problem 1] in the case of special one-relator monoids

**Corollary 3.19.** *Let  $M$  be the one-relator monoid  $\langle A \mid w = 1 \rangle$ . Then  $M$  is of type left- and right- $\text{FP}_\infty$ . Moreover, if  $w$  is not a proper power then  $\text{left cd } M \leq 2$  and  $\text{right cd } M \leq 2$ , and otherwise  $\text{left cd } M = \text{right cd } M = \infty$ .*

*Proof.* We prove the results for  $\text{left-FP}_\infty$  and left cohomological dimension (the other results are dual). The group of units  $G$  of  $M$  is a one-relator group by Adjan's theorem [1, Lemma 96] (this also follows from the results of Zhang described above), and hence of type  $\text{FP}_\infty$  by Lyndon's theorem [42]. This proves the first statement in light of Theorem 3.16. The second statement follows since by Lemma 3.18 the group  $G$  is a one-relator group whose defining relator is not a proper power in the first case and is a proper power in the second. By a theorem of Lyndon [42]  $G$  has cohomological dimension at most 2 in the first case and has infinite cohomological dimension in the second. The result now follows from Theorem 3.16.  $\square$

We now turn our attention to proving the topological analogue of Theorem 3.16. We do this by showing how an equivariant classifying space for a special monoid may be constructed from an equivariant classifying space for its group of units.

Note that while for finitely presented monoids it follows from [25] that the properties  $\text{left-FP}_n$  and  $\text{left-F}_n$  are equivalent, in contrast it is not known whether  $\text{left cd}(M)$  and  $\text{left gd}(M)$  coincide (this is even open for groups). Therefore, the second part of the following theorem is not an immediate consequence of Theorem 3.16.

**Theorem 3.20.** *Let  $M$  be a finitely presented special monoid with group of units  $G$ .*

- (1) *If  $G$  is of type  $\text{F}_n$  with  $1 \leq n \leq \infty$ , then  $M$  is of type left- and right- $\text{F}_n$ .*
- (2)  *$\text{gd } G \leq \text{left gd } M \leq \max\{2, \text{gd } G\}$  and  $\text{gd } G \leq \text{right gd } M \leq \max\{2, \text{gd } G\}$ .*

*Proof.* We prove the results for  $\text{left-F}_n$  and left geometric dimension. The other results are dual. It is proved in [25, Section 6] for finitely presented monoids the properties  $\text{left-F}_n$  and  $\text{left-FP}_n$  coincide. Now part (1) of the theorem follows from the first part of Theorem 3.16. (One can also see this directly from the construction below.)

To prove part (2), first note that we showed that  $M$  was a free left  $G$ -set at the beginning of the proof of Theorem 3.16. Hence any free  $M$ -CW complex is a free  $G$ -CW complex. Also note that Theorem 3.10 implies that every projective  $M$ -set is free, as  $Me$  is a left ideal for any idempotent  $e$ . Thus any projective  $M$ -CW complex  $X$  is a free  $M$ -CW complex and so it follows that  $G \backslash X$  is  $K(G, 1)$ -space. The inequality  $\text{gd } G \leq \text{right gd } M$  follows.

We shall now explain how to construct an equivariant classifying space for  $M$  of dimension  $\max\{2, \text{gd}(G)\}$ .

Let  $X_G$  be an equivariant classifying space for the group  $G$ . Since  $G$  is a group it follows that the projective  $G$ -CW complex  $X_G$  is a free  $G$ -CW complex. By Zhang's theorem [63, Theorem 4.4], the submonoid of right units  $R$  of  $M$  is isomorphic to the monoid free product  $G * C^*$  where  $C^*$  is a free monoid over a finite alphabet  $C$ . The right Cayley graph  $\Gamma(C^*)$  of  $C^*$  with respect to the generating set  $C$  is a tree and thus is a free equivariant classifying space for the monoid  $C^*$ . In particular  $C^*$  is of geometric dimension at most 1. Let  $X$  be the left equivariant classifying space for  $R \cong G * C^*$  given by the construction in the proof of Theorem 4.5 in Section 4 below. From the construction it follows that  $X$  is a free  $R$ -CW complex and an equivariant classifying space for  $R$ . (If  $X_G$  has a  $G$ -finite  $n$ -skeleton, then  $X$  has an  $R$ -finite  $n$ -skeleton.) It also follows from the construction of  $X$  that  $\dim X \leq \max\{1, \dim X_G\}$  (compare with Theorem 4.8).

Now  $M$  is an  $M$ - $R$ -biset, which is free as a left  $M$ -set and is also free as a right  $R$ -set by Corollary 3.8, and  $X$  is a free left  $R$ -CW complex. It follows from Proposition 2.1 that  $M \otimes_R X$  is a free left  $M$ -CW complex with  $\dim M \otimes_R X = \dim X$ . (It will have  $M$ -finite  $n$ -skeleton if  $X$  has  $R$ -finite  $n$ -skeleton.) The complex  $M \otimes_R X$  is a disjoint union of copies of  $X$ , one for each



$\mathcal{R}$ -class of  $M$  by Remark 2.2. To make this concrete, take the transversal  $\mathcal{T}$  of the  $\mathcal{R}$ -classes of  $M$  defined above, which is a basis for  $M$  as a free right  $R$ -set. Then each element of  $M \otimes_R X$  can be uniquely written in the form  $t \otimes x$  with  $t \in \mathcal{T}$  and  $x \in X$  and  $M \otimes_R X = \coprod_{t \in \mathcal{T}} t \otimes X$ . We say that two elements  $m \otimes x$  and  $m' \otimes x'$  of  $M \otimes_R X$  belong to the same copy of  $X$  in  $M \otimes_R X$  if and only if  $m \mathcal{R} m'$ .

Fix a basepoint  $x_0 \in \mathcal{Q} \subseteq X_0$ . Next we connect the space  $M \otimes_R X$  by attaching edges  $m \otimes x_0 \rightarrow ma \otimes x_0$  for each  $m \in M$  and  $a \in A$ . This is the same as attaching a free  $M$ -cell  $M \times B^1$  of dimension 1 based at  $1 \otimes x_0 \rightarrow a \otimes x_0$  for each  $a \in A$ . Let  $Y$  denote the resulting free  $M$ -CW complex. The  $\mathcal{R}$ -order in the monoid  $M$  induces in a natural way an order on the copies of  $X$  in  $Y$ , and there is an edge joining two distinct copies of  $X$  in  $Y$  if and only if there is an edge in the right Cayley graph of  $M$  joining the corresponding  $\mathcal{R}$ -classes. Moreover, it follows from the definition of  $Y$ , and Proposition 3.13, that there is at most one edge joining any pair of distinct copies of  $X$  in  $Y$ . It follows that if we contract each of the copies of  $X$  in  $Y$  we obtain the graph  $\Gamma$  in Theorem 3.14, which is a regular rooted tree, together with possibly infinitely many loops at each vertex. These loops arise from the edges  $m \otimes x_0 \rightarrow ma \otimes x_0$  where  $m \mathcal{R} ma$  added in the construction of  $Y$ . (Notice that if  $M \otimes_R X$  has  $M$ -finite  $n$ -skeleton, then so does  $Y$ .)

To turn  $Y$  into an equivariant classifying space for  $M$  we add 2-cells to deal with these loops, in the following way. It follows from Proposition 3.15 that for each  $a \in A$ , the set  $L = \{m \in M \mid m \mathcal{R} ma\}$  is a free left  $M$ -set generated by a finite set  $F_a \subseteq L$  with  $F_a \subseteq R$ . For each  $r \in F_a$ , choose a path  $p_r$  in  $1 \otimes X$  from  $1 \otimes x_0$  to  $1 \otimes rx_0$ , choose a path  $q_r$  in  $1 \otimes X$  from  $1 \otimes x_0$  to  $1 \otimes rax_0$ , and let  $e_r$  denote the edge in  $Y$  labelled by  $a$  from  $1 \otimes rx_0$  to  $1 \otimes rax_0$ . Note that since  $r \in F_a \subseteq L$  it follows that  $r \in R$  and  $ra \in R$  and so  $1 \otimes rx_0 = r \otimes x_0$  and  $1 \otimes rax_0 = ra \otimes x_0$  and hence  $e_r$  is indeed one of the edges that was added during the construction of  $Y$ . Now for each  $a \in A$  attach a free 2-cell  $M \times B^2$  to  $Y$  by attaching a 2-cell at  $1 \otimes x_0$  with boundary path  $p_r e_r q_r^{-1}$  and all of its translates under the action of  $M$ . We do this for each  $a \in A$  and call the resulting complex  $Z$ . Now if we contract the copies of  $X$  in  $Z$ , we obtain the tree  $\Gamma$ , together with loops at each vertex each of which bounds a single disk. Thus  $Z$  is homotopy equivalent to the tree  $\Gamma$ , and hence is contractible. This shows that  $Z$  is an equivariant classifying space for the monoid  $M$ . (Note that if  $Y$  has  $M$ -finite  $n$ -skeleton, then so does  $Z$  hence giving an alternative proof that if  $G$  is of type  $F_n$ , then  $M$  is of type left- $F_n$ .)

To complete the proof, since the free  $M$ -CW complex  $Z$  was constructed from  $M \otimes_R X$  by attaching 1-cells and 2-cells, and since we have already observed that  $\dim M \otimes_R X = \dim X \leq \max\{1, \dim X_G\}$ , it follows that  $\dim Z \leq \max\{2, \dim X_G\}$  and hence  $\text{left gd}(M) \leq \max\{2, \text{gd}(G)\}$ .  $\square$

For special one-relator monoids we obtain the following corollary which is the topological analogue of Corollary 3.19.

**Corollary 3.21.** *Let  $M$  be the one-relator monoid  $\langle A \mid w = 1 \rangle$ . Then  $M$  is of type left- and right- $F_\infty$ . Moreover, if  $w$  is not a proper power then  $\text{left gd } M \leq 2$  and  $\text{right gd } M \leq 2$ , and otherwise  $\text{left gd } M = \text{right gd } M = \infty$ .*

In particular this results says that for every special one-relator monoid whose defining relator is not a proper power admits an equivariant classifying space of dimension at most 2. In fact, in this case it turns out that the Cayley complex of the monoid gives an equivariant classifying space of dimension at most 2, as the following result demonstrates.

**Theorem 3.22.** *Let  $M = \langle A \mid w = 1 \rangle$  such that  $w$  is not a proper power. Let  $X$  be the 2-complex obtained by filling in each loop labeled by  $w$  in the Cayley graph  $\Gamma(M, A)$  of  $M$ . Then  $X$  is left equivariant classifying space for  $M$  with dimension at most 2.*

*Proof.* It follows from the proof of [25, Theorem 6.14] that  $X$  is an  $M$ -finite simply connected free  $M$ -CW complex of dimension at most 2. It is shown in [34, Corollary 7.5] that the presentation  $\langle A \mid w = 1 \rangle$  is strictly aspherical in the sense defined in [33, Section 2]. The cellular chain

complex of  $X$  gives a free resolution displayed in Equation (7.2) in [33, Theorem 7.2]. This shows that  $X$  is acyclic. Since  $X$  is acyclic and simply connected it follows from the Whitehead and Hurewicz theorems that  $X$  is contractible, and hence  $X$  is a left equivariant classifying space for the monoid  $M$ .  $\square$

The analogous result to Theorem 3.22 is also known to hold for one-relator groups. This was first observed in [14] and is a consequence of Lyndon's Identity Theorem [42]. A more topological proof is given in [22].

We currently do not know whether the two-sided analogues of the results proved in this section, for bi- $F_n$  and (two-sided) geometric dimension, hold. One way to establish these results might be to seek a better understanding of the two-sided Cayley graphs of special monoids.

As mentioned in the introduction, building on the ideas presented in this section, in [26] we have extended these results to arbitrary one-relator monoids. In particular in [26] we give a positive answer to Kobayashi's question [34, Problem 1] by showing that every one-relator monoid  $\langle A \mid u = v \rangle$  is of type left- and right- $F_\infty$  and  $FP_\infty$ .

#### 4. AMALGAMATED FREE PRODUCTS

For graph of groups, including free products with amalgamation and HNN extensions, there are well-established methods for constructing a  $K(G, 1)$  from  $K(G, 1)$ s of the vertex and edge groups; see for example [29, page 92]. This can then be used to prove results for groups about the behaviour of the properties  $F_n$  and geometric dimension for amalgamated free products and HNN extensions. In this section, and the two sections that follow it, we use topological methods to investigate the behaviour of topological and homological finiteness properties of monoids, for free products with amalgamation, and HNN extension constructions.

A *monoid amalgam* is a triple  $[M_1, M_2; W]$  where  $M_1, M_2$  are monoids with a common submonoid  $W$ . The *amalgamated free product* is then the pushout in the diagram

$$\begin{array}{ccc} W & \longrightarrow & M_1 \\ \downarrow & & \downarrow \\ M_2 & \longrightarrow & M_1 *_W M_2 \end{array} \quad (4.1)$$

in the category of monoids. Monoid amalgamated products are **much** more complicated than group ones. For instance, the amalgamated free product of finite monoids can have an undecidable word problem, and the factors do not have to embed or intersect in the base monoid; see [56]. So there are no normal forms available in complete generality that allow one construct a Bass–Serre tree. We use instead the homological ideas of Dicks. For more details about these methods we refer the reader to [19, Chapter 1, Sections 4–7].

An  $M$ -graph  $X$  is a one-dimensional CW complex with a cellular action by  $M$  sending edges to edges. Given an  $M$ -graph  $X$  we use  $V$  to denote its set of 0-cells and  $E$  to denote its set of 1-cells. Given any  $M$ -graph, if we choose some orientation for the edges, then the attaching maps of the 1-cells define functions  $\iota, \tau$  from  $E$  to  $V$  where in  $X$  each oriented edge  $e$  starts at  $\iota e$  and ends at  $\tau e$ . We call  $V$  and  $E$  the vertex set, and edge set respectively, of the  $M$ -graph  $X$ . We shall assume that the monoid action preserves the orientation. It shall sometimes be useful to think of an  $M$ -graph as given by a tuple  $(X, V, E, \iota, \tau)$  where  $X$  is an  $M$ -set,  $X = V \cup E$  a disjoint union where each of  $V$  and  $E$  is closed under the action of  $M$ , and  $\iota, \tau: E \rightarrow V$  are  $M$ -equivariant maps.

Let  $M$  be a monoid and let  $X$  be an  $M$ -graph. Let  $\mathbb{Z}V$  and  $\mathbb{Z}E$  denote the free abelian groups on  $V$  and  $E$ , respectively. The cellular boundary map of  $X$  is the  $M$ -linear map  $\partial: \mathbb{Z}E \rightarrow \mathbb{Z}V$  with  $\partial(e) = \tau e - \iota e$  for all  $e \in E$ . The sequence

$$\mathbb{Z}E \xrightarrow{\partial} \mathbb{Z}V \xrightarrow{\epsilon} \mathbb{Z} \longrightarrow 0$$

is the augmented cellular chain complex of  $X$ , where  $\epsilon$  is the augmentation map sending  $\sum_{v \in V} n_v v$  to  $\sum_{v \in V} n_v$  (i.e., each element of the basis  $V$  is mapped to 1). Throughout this section we shall frequently be confronted with the task of showing that a given  $M$ -graph is a tree or a forest. To do this, it is useful to recall that the  $M$ -graph  $X$  is a forest if and only if  $\partial: \mathbb{Z}E \rightarrow \mathbb{Z}V$  is injective; see [19, Lemmas 6.4], i.e.,

$$0 \longrightarrow \mathbb{Z}E \xrightarrow{\partial} \mathbb{Z}V \xrightarrow{\epsilon} \mathbb{Z} \longrightarrow 0$$

is exact.

The results in this section improve, and give simpler proofs of, several results of Cremanns and Otto [17] on the behaviour of  $\text{FP}_n$  under free products and certain rather restricted free products of monoids with amalgamation. The proofs in Cremanns and Otto are quite long and technical, as is often the case for results in this area. The results in this section demonstrate the type of result our topological methods were introduced to prove. They show that the topological approach may be used to prove more general results in a less technical and more conceptual way. Our results also generalise and simplify proofs of some results of Kobayashi [35] on preservation of left-, right- and bi- $\text{FP}_n$  under free products (see for example [35, Proposition 4.1]). There are no bi- $\text{FP}_n$  analogues in the literature of the two-sided results we obtain below on the behaviour of bi- $\text{F}_n$  and geometric dimension for free products with amalgamation. Also, as far as we are aware, the results that we obtain here are the first to appear in the literature on cohomological dimension of amalgamated free products of monoids.

A monoid presentation is said to have finite homological type, abbreviated to FHT, if the, so-called, homotopy bimodule of the given presentation is finitely generated. The homotopy bimodule is a  $\mathbb{Z}M$ -bimodule constructed from a complex of  $\mathbb{Z}A^*$ -bimodules defined using the set of defining relations  $R$  of the presentation  $\langle A \mid \mathfrak{R} \rangle$  of the monoid  $M$ , and a particular family of disjoint circuits in the derivation graph associated with the presentation. The property FHT was originally introduced by Wang and Pride [62]. We refer the reader to that paper, or to [36, Section 3], for full details of the definition of FHT. It was proved in [37] that for finitely presented monoids FHT and bi- $\text{FP}_3$  (equivalently bi- $\text{F}_3$ ) are equivalent. So some of the results below also have an interpretation in terms of FHT.

**4.1. The one-sided setting.** Let us define a tree  $T$  for a pushout diagram (4.1). Let us assume that  $f_i: W \rightarrow M_i$ , for  $i = 1, 2$ , is the homomorphism in the diagram and put  $L = M_1 *_W M_2$  for the pushout. The right multiplicative actions of  $M_1$ ,  $M_2$  and  $W$  give three different partitions of  $L$  into weak orbits. Since  $W \leq M_i$  the  $W$ -orbits give a finer partition than both the  $M_1$ - and  $M_2$ -orbits. We can then define a directed bipartite graph  $T$  with one part given by the  $M_1$ -orbits and the other part given by the  $M_2$ -orbits. When an  $M_1$ -orbit intersects an  $M_2$ -orbit, that intersection will be a union of  $W$ -orbits, and in this case we draw directed edges from the  $M_1$ -orbit to the  $M_2$ -orbit labelled by the  $W$ -orbits in this intersection.

In more detail, let  $T$  be the  $L$ -graph with vertex set

$$V = L/M_1 \amalg L/M_2$$

and edge set

$$E = L/W$$

where  $M_1, M_2, W$  act on the right of  $L$  by first applying the canonical map to the pushout and then right multiplying. We write  $[x]_K$  for the class of  $x \in L$  in  $L/K$ . The edge  $[x]_W$  connects  $[x]_{M_1}$  with  $[x]_{M_2}$  (and we usually think of it as oriented in this direction). The incidence here is easily seen to be well defined and the action of  $L$  on the left of these sets is by cellular mappings sending edges to edges and preserving orientation. Hence  $T$  is an  $L$ -graph.

**Lemma 4.1.** *The graph  $T$  is connected.*

*Proof.* The pushout  $L$ , being a quotient of the free product  $M_1 * M_2$ , is generated by the images of  $M_1$  and  $M_2$  under the natural maps (which we omit from the notation even though they need

not be injective). We define the length of  $x \in L$  to be the minimum  $k$  such that  $x = x_1 \cdots x_k$  with  $x_i \in M_1 \cup M_2$ . We prove by induction on the length of  $x$  that there is a path in  $T$  from  $[1]_{M_1}$  to  $[x]_{M_1}$ . If  $x = 1$ , this is trivial, so assume the statement is true for length  $k$  and  $x = x_1 \cdots x_{k+1}$ . Let  $p$  be a path from  $[1]_{M_1}$  to  $[x_2 \cdots x_{k+1}]_{M_1}$ . Then  $x_1 p$  is a path from  $[x_1]_{M_1}$  to  $[x]_{M_1}$ . If  $x_1 \in M_1$ , then  $[x_1]_{M_1} = [1]_{M_1}$  and so  $x_1 p$  is a path from  $[1]_{M_1}$  to  $[x]_{M_1}$ . If  $x_1 \in M_2$ , then  $[x_1]_W$  is an edge connecting  $[x_1]_{M_1}$  and  $[x_1]_{M_2} = [1]_{M_2}$  and  $[1]_W$  is an edge connecting  $[1]_{M_1}$  with  $[1]_{M_2}$  and so there is a path from  $[1]_{M_1}$  to  $[x]_{M_1}$ . Finally, if  $x \in L$ , then  $[x]_{M_2}$  is connected by  $[x]_W$  to  $[x]_{M_1}$ , which in turn is connected by a path to  $[1]_{M_1}$ . Thus  $T$  is connected.  $\square$

We aim to prove that  $T$  is a tree by showing that the cellular boundary map  $\partial: \mathbb{Z}E \rightarrow \mathbb{Z}V$  is injective. To prove this we shall make use of semidirect products of monoids and the concept of a derivation. An account of this theory for groups may be found in [19] where it is applied to show that the standard graph of the fundamental group of a graph of groups is a tree; see [19, Theorem 7.6].

Let  $M$  be a monoid and let  $A$  be a left  $\mathbb{Z}M$ -module. Then we can form the semidirect product  $A \rtimes M$ , of the abelian group  $A$  and the monoid  $M$ , with elements  $A \times M$  and multiplication given by

$$(a, m)(a', m') = (a + ma', mm').$$

The natural projection  $\pi: A \rtimes M \rightarrow M$ ,  $(a, m) \mapsto m$  is clearly a monoid homomorphism. A *splitting* of this projection is a monoid homomorphism  $\sigma: M \rightarrow A \rtimes M$  such that  $\pi(\sigma(m)) = m$  for all  $m \in M$ . Associated to any splitting  $\sigma$  of  $\pi$  is a mapping  $d: M \rightarrow A$  defined as the unique function satisfying

$$\sigma(m) = (d(m), m)$$

for all  $m \in M$ . It follows from the fact that  $\sigma$  is a homomorphism that the function  $d: M \rightarrow A$  must satisfy

$$d(mm') = d(m) + md(m') \quad (4.2)$$

for all  $m, m' \in M$ . Any function  $d: M \rightarrow A$  satisfying (4.2) is called a *derivation*. A derivation is called *inner* if it is of the form  $d(m) = ma - a$  for some  $a \in A$ . It is easy to check that a mapping  $d: M \rightarrow A$  is a derivation if and only if  $m \mapsto (d(m), m)$  provides a splitting of the semidirect product projection  $A \rtimes M \rightarrow M$ .

**Lemma 4.2.** *The graph  $T$  is a tree.*

*Proof.* Since  $T$  is connected by Lemma 4.1, it suffices to show that the cellular boundary map  $\partial: \mathbb{Z}E \rightarrow \mathbb{Z}V$  is injective. To show this, we define a left inverse  $\beta: \mathbb{Z}V \rightarrow \mathbb{Z}E$ . In what follows, we abuse notation by identifying an element of  $M_1$ ,  $M_2$  or  $W$  with its image in  $L$ .

First define  $\varphi_1: M_1 \rightarrow \mathbb{Z}E \rtimes L$  by  $\varphi_1(m_1) = (0, m_1)$ . Then  $\varphi_1$  is clearly a monoid homomorphism. Define  $\varphi_2: M_2 \rightarrow \mathbb{Z}E \rtimes L$  by  $\varphi_2(m_2) = ([1]_W - [m_2]_W, m_2)$ . Notice that  $m_2 \mapsto [1]_W - [m_2]_W$  is the inner derivation of the  $\mathbb{Z}M_2$ -module  $\mathbb{Z}E$  associated to  $-[1]_W \in \mathbb{Z}E$  and hence  $\varphi_2$  is a homomorphism. Next, we observe that  $\varphi_1 f_1 = \varphi_2 f_2$ . Indeed, if  $w \in W$ , then  $\varphi_1 f_1(w) = (0, w)$  and  $\varphi_2 f_2(w) = ([1]_W - [w]_W, w) = (0, w)$  as  $[1]_W = [w]_W$ . Thus there is a well defined homomorphism  $\varphi: L \rightarrow \mathbb{Z}E \rtimes L$  extending  $\varphi_1, \varphi_2$  by the universal property of a pushout. This map must split the semidirect product projection by construction of  $\varphi_1, \varphi_2$ . Indeed, for all  $m_1 \in L$  in the image of  $M_1$  we have  $\varphi(m_1) = \varphi_1(m_1) = (0, m_1)$  and for all  $m_2 \in L$  in the image of  $M_2$  we have

$$\varphi(m_2) = \varphi_2(m_2) = ([1]_W - [m_2]_W, m_2).$$

It follows that for all  $m_1 \in L$  in the image of  $M_1$  we have  $\pi(\varphi(m_1)) = m_1$ , and for all  $m_2 \in L$  in the image of  $M_2$  we have  $\pi(\varphi(m_2)) = m_2$ . Since, as already observed above,  $L$  is generated by the images of  $M_1$  and  $M_2$  under the natural maps, and since  $\pi$  and  $\varphi$  are homomorphisms, we conclude that  $\pi(\varphi(l)) = l$  for all  $l \in L$ , as required. It follows that  $\varphi(x) = (d(x), x)$  for some

derivation  $d: L \rightarrow \mathbb{Z}E$  with the property that  $d(m_1) = 0$  for  $m_1 \in M_1$  and  $d(m_2) = [1]_W - [m_2]_W$  for  $m_2 \in M_2$ .

Define  $\beta: \mathbb{Z}V \rightarrow \mathbb{Z}E$  by  $\beta([x]_{M_1}) = d(x)$  and  $\beta([x]_{M_2}) = d(x) + [x]_W$  for  $x \in L$ . We must show that this is well defined. First suppose that  $x \in L$  and  $m_1 \in M_1$ . Then  $d(xm_1) = xd(m_1) + d(x) = d(x)$  because  $d$  vanishes on the image of  $M_1$ . If  $x \in L$  and  $m_2 \in M_2$ , then

$$\begin{aligned} d(xm_2) + [xm_2]_W &= xd(m_2) + d(x) + [xm_2]_W \\ &= x([1]_W - [m_2]_W) + d(x) + [xm_2]_W = d(x) + [x]_W. \end{aligned}$$

It follows that  $\beta$  is well defined.

We now compute

$$\beta\partial([x]_W) = \beta([x]_{M_2}) - \beta([x]_{M_1}) = d(x) + [x]_W - d(x) = [x]_W$$

for  $x \in L$ . Thus  $\beta\partial = 1_{\mathbb{Z}E}$  and so  $\partial$  is injective. This completes the proof that  $T$  is a tree.  $\square$

Since  $T$  is a tree we obtain an exact sequence of  $\mathbb{Z}L$ -modules

$$0 \longrightarrow \mathbb{Z}E \xrightarrow{\partial} \mathbb{Z}V \xrightarrow{\epsilon} \mathbb{Z} \longrightarrow 0$$

where  $E, V$  are the edge and vertex sets of  $T$ , respectively. See [19, Theorem 6.6]. The exactness of this cellular chain complex of  $T$  can be reformulated in the following manner.

**Corollary 4.3.** *There is an exact sequence of  $\mathbb{Z}L$ -modules*

$$0 \longrightarrow \mathbb{Z}L \otimes_{\mathbb{Z}W} \mathbb{Z} \longrightarrow (\mathbb{Z}L \otimes_{\mathbb{Z}M_1} \mathbb{Z}) \oplus (\mathbb{Z}L \otimes_{\mathbb{Z}M_2} \mathbb{Z}) \longrightarrow \mathbb{Z} \longrightarrow 0$$

where  $L = M_1 *_W M_2$  is the pushout.

*Proof.* This follows from the definition of  $T$ , the fact that  $T$  is a tree, and the observation that  $\mathbb{Z}[L/K] \cong \mathbb{Z}L \otimes_{\mathbb{Z}K} \mathbb{Z}$  for  $K = M_1, M_2, W$ .  $\square$

We call  $T$  the *Bass–Serre tree* of the pushout.

If  $f: X \rightarrow Y$  and  $g: X \rightarrow Z$  are continuous mappings of topological spaces, the *homotopy pushout* of  $f, g$  is the space obtained by attaching  $X \times I$  to  $Y \amalg Z$  by the mapping  $h: X \times \partial I \rightarrow Y \amalg Z$  with  $h(x, 0) = f(x)$  and  $h(x, 1) = g(x)$ . If  $X, Y, Z$  are CW complexes and  $f, g$  are cellular mappings, then  $h$  is cellular and so the homotopy pushout  $U$  of  $f$  and  $g$  is a CW complex. If, in addition,  $X, Y, Z$  are projective  $M$ -CW complexes and  $f, g$  are cellular and  $M$ -equivariant, then  $U$  is a projective  $M$ -CW complex by [25, Lemma 2.1]. Moreover, by the description of the cells coming from the proof of [25, Lemma 2.1], if  $Y, Z$  have  $M$ -finite  $n$ -skeleton and  $X$  has  $M$ -finite  $(n - 1)$ -skeleton (whence  $X \times I$  has  $M$ -finite  $n$ -skeleton), then  $U$  has  $M$ -finite  $n$ -skeleton.

The homotopy pushout construction is functorial with respect to commutative diagrams

$$\begin{array}{ccccc} Y & \xrightarrow{f_1} & X_1 & & \\ f_2 \downarrow & \searrow r & \searrow s & & \\ X_2 & & Y' & \xrightarrow{g_1} & X'_1 \\ & \searrow t & \downarrow g_2 & & \\ & & X'_2 & & \end{array}$$

Moreover, if  $r, s, t$  are homotopy equivalences, then it is well known that the induced mapping of homotopy pushouts is a homotopy equivalence; see for example [20, Theorem 4.2.1], or [21, page 19] where it is observed that homotopy colimits have the strong homotopy equivalence property.

For the reader's convenience, we shall prove a special case of this fact that will be crucial in what follows. Recall that if  $Y$  is a space, the *suspension* of  $Y$  is the space  $\Sigma Y = Y \times I / (Y \times \{0\} \cup Y \times \{1\})$ . If  $Y$  is contractible, then the mapping  $\Sigma Y \rightarrow I$  induced by the projection  $Y \times I \rightarrow I$  is a homotopy equivalence.

**Lemma 4.4.** *Let  $M$  be a monoid and  $X_1, X_2, Y$  locally path connected  $M$ -spaces. Assume that the natural mappings  $r_i: X_i \rightarrow \pi_0(X_i)$ , for  $i = 1, 2$ , and  $r: Y \rightarrow \pi_0(Y)$  are homotopy equivalences (where the set of path components is given the discrete topology). Let  $f_i: Y \rightarrow X_i$  be continuous mappings, for  $i = 1, 2$ , and let  $Z$  be the homotopy pushout of  $X_1, X_2$  along  $Y$ , which is naturally an  $M$ -space. Let  $\Gamma$  be the  $M$ -graph with vertex set  $\pi_0(X_1) \amalg \pi_0(X_2)$  and edge set  $\pi_0(Y)$  where the edge corresponding to  $C \in \pi_0(Y)$  connects the component of  $f_1(C)$  to the component of  $f_2(C)$ ; this is the homotopy pushout of  $\pi_0(X_1)$  and  $\pi_0(X_2)$  along  $\pi_0(Y)$ . Then the natural  $M$ -equivariant mapping  $h: Z \rightarrow \Gamma$  is a homotopy equivalence.*

*Proof.* The mapping  $h$  takes an element of  $X_i$  to its path component and an element  $(y, t) \in Y \times I$  to  $(C, t)$  where  $C$  is the component of  $y$ . This is well defined, by construction of the homotopy pushout, and is  $M$ -equivariant. As the connected components of  $X_i$ , for  $i = 1, 2$ , are disjoint and contractible subcomplexes,  $Z$  is homotopy equivalent to the space obtained by contracting each of these subcomplexes to a point. Then  $Z$  has the homotopy type of the CW complex obtained by adjunction of  $\coprod_{C \in \pi_0(Y)} \Sigma C$  to the discrete set  $\pi_0(X_1) \amalg \pi_0(X_2)$  where  $\Sigma C$  is attached via the mapping sending  $(y, 0)$  to the component of  $f_1(C)$  and  $(y, 1)$  to the component of  $f_2(C)$ . Since the mapping  $\Sigma C \rightarrow I$  induced by the projection  $C \times I \rightarrow I$  is a homotopy equivalence by contractibility of  $C$ , it follows that  $h$  is a homotopy equivalence. This completes the proof.  $\square$

We now prove some preservation results for amalgamated free products. We shall apply the observation in Remark 2.2 without comment.

**Theorem 4.5.** *Let  $[M_1, M_2; W]$  be an amalgam of monoids such that  $M_1, M_2$  are free as right  $W$ -sets. If  $M_1, M_2$  are of type left- $F_n$  and  $W$  is of type left- $F_{n-1}$ , then  $M_1 *_W M_2$  is of type left- $F_n$ .*

*Proof.* Let  $X_i$  be an equivariant classifying space for  $M_i$  with  $M_i$ -finite  $n$ -skeleton, for  $i = 1, 2$ , and let  $Y$  be an equivariant classifying space for  $W$  with  $W$ -finite  $(n-1)$ -skeleton. By [25, Lemma 6.2] and the cellular approximation theorem [25, Theorem 2.8], we can find  $W$ -equivariant cellular mappings  $f_i: Y \rightarrow X_i$ , for  $i = 1, 2$ . Let  $L = M_1 *_W M_2$ . By McDuff [49],  $L$  is a free right  $M_i$ -set, for  $i = 1, 2$ , and a free right  $W$ -set. Then  $X'_i = L \otimes_{M_i} X_i$ , for  $i = 1, 2$ , is a projective  $L$ -CW complex with  $L$ -finite  $n$ -skeleton and  $Y' = L \otimes_W Y$  is a projective  $L$ -CW complex with  $L$ -finite  $(n-1)$ -skeleton by Proposition 2.1. Let  $\tilde{f}_i: Y' \rightarrow X'_i$  be the map induced by  $f_i$ , for  $i = 1, 2$ , and let  $Z$  be the homotopy pushout of  $\tilde{f}_1, \tilde{f}_2$ . It is a projective  $L$ -CW complex. We claim that  $Z$  is an equivariant classifying space for  $L$ . Note that  $Z$  has an  $L$ -finite  $n$ -skeleton by construction.

Our goal is to show that  $Z$  is homotopy equivalent to the Bass–Serre tree  $T$ . By [25, Proposition 3.4], we have that  $\pi_0(X'_i) \cong L \otimes_{M_i} \pi_0(X_i) \cong L/M_i$  and  $\pi_0(Y') \cong L \otimes_W \pi_0(Y) \cong L/W$  and  $f_i$  induces the natural mapping  $L/W \rightarrow L/M_i$  under these identifications, for  $i = 1, 2$ . As  $X'_i \cong L/M_i \times X_i$  and  $Y' \cong L/W \times Y$  (by freeness of  $L$  as a right  $K$ -set for  $K = M_1, M_2, W$ ) and  $X_i$ , for  $i = 1, 2$ , and  $Y$  are contractible, the projections  $X'_i \rightarrow \pi_0(X'_i)$ , for  $i = 1, 2$ , and  $Y' \rightarrow \pi_0(Y')$  are homotopy equivalences. It follows that  $Z$  is homotopy equivalent to  $T$ , by Lemma 4.4, and hence contractible. This completes the proof.  $\square$

Note that we do not assume that the monoids  $M_1$  and  $M_2$  are finitely generated, or finitely presented, in the above result. Recall that a monoid can be of type left- $F_2$  without being finitely presented, and can be of type left- $F_1$  without being finitely generated; see [25, Section 6]. The hypotheses of Theorem 4.5 hold if  $W$  is trivial or if  $M_1, M_2$  are left cancellative and  $W$  is a group. As another example, if we consider  $\mathbb{N}$ , then, for any  $k > 0$ ,  $\mathbb{N}$  is a free  $k\mathbb{N}$ -set with basis  $\{0, 1, \dots, k-1\}$ . Since  $k\mathbb{N} \cong \mathbb{N}$ , it follows from Theorem 4.5 that  $\mathbb{N} *_k \mathbb{N} \mathbb{N}$  is of type left- $F_\infty$ , as  $\mathbb{N}$  is of type left- $F_\infty$ , for any  $k, m > 0$ . As a special case of Theorem 4.5 we obtain the following result as a corollary.

**Corollary 4.6.** *A free product  $M * N$  of monoids of type  $\text{left-F}_n$  is of type  $\text{left-F}_n$ . If  $M, N$  are finitely presented monoids, then  $M * N$  is of type  $\text{left-F}_n$  if and only if  $M$  and  $N$  both are of type  $\text{left-F}_n$ .*

*Proof.* If  $M$  and  $N$  are of type  $\text{left-F}_n$ , then  $M * N$  is of type  $\text{left-F}_n$  by Theorem 4.5 as  $M, N$  are free  $\{1\}$ -sets. Conversely, if  $M, N$  are finitely presented, then so is  $M * N$  and hence  $\text{left-F}_n$  is equivalent to  $\text{left-FP}_n$  for these monoids. A result of Pride [53] says that a retract of a  $\text{left-FP}_n$  monoid is  $\text{left-FP}_n$ . As  $M, N$  are retracts of  $M * N$ , the converse follows.  $\square$

The fact that for finitely presented monoids  $M, N$  of type  $\text{left-FP}_n$ , the free product  $M * N$  is of type  $\text{left-FP}_n$  was first proved in [17, Theorem 5.5].

The following corollary is classical.

**Corollary 4.7.** *If  $[G_1, G_2; H]$  is an amalgam of groups with  $G_1, G_2$  of type  $\text{left-F}_n$  and  $H$  of type  $\text{left-F}_{n-1}$ , then  $G_1 *_H G_2$  is of type  $\text{left-F}_n$ .*

*Proof.* Since  $G_1, G_2$  are free left  $H$ -sets, this follows from Theorem 4.5.  $\square$

The homotopy pushout construction in the proof of Theorem 4.5 also serves to establish the following.

**Theorem 4.8.** *Let  $[M_1, M_2; W]$  be an amalgam of monoids such that  $M_1, M_2$  are free as right  $W$ -sets. Suppose that  $d_i$  is the left geometric dimension of  $M_i$ , for  $i = 1, 2$ , and  $d$  is the left geometric dimension of  $W$ . Then the left geometric dimension of  $M_1 *_W M_2$  is bounded above by  $\max\{d_1, d_2, d + 1\}$ .*

**Corollary 4.9.** *Let  $M$  and  $N$  be monoids of left geometric dimension at most  $n$ . Then  $M * N$  has left geometric dimension at most  $\max\{n, 1\}$ .*

We now wish to prove a homological analogue of Theorem 4.5.

**Theorem 4.10.** *Let  $[M_1, M_2; W]$  be an amalgam of monoids such that  $\mathbb{Z}L$  is flat as a right  $\mathbb{Z}M_1$ -,  $\mathbb{Z}M_2$ - and  $\mathbb{Z}W$ -module, where  $L = M_1 *_W M_2$ . If  $M_1, M_2$  are of type  $\text{left-FP}_n$  and  $W$  is of type  $\text{left-FP}_{n-1}$ , then  $M_1 *_W M_2$  is of type  $\text{left-FP}_n$ .*

*Proof.* By Lemma 2.5 and the hypotheses, we deduce that  $\mathbb{Z}L \otimes_{\mathbb{Z}M_i} \mathbb{Z}$  is of type  $\text{FP}_n$ , for  $i = 1, 2$ , and  $\mathbb{Z}L \otimes_{\mathbb{Z}W} \mathbb{Z}$  is of type  $\text{FP}_{n-1}$ . The result now follows by applying Corollary 2.4 to the exact sequence in Corollary 4.3.  $\square$

*Remark 4.11.* It is reasonable to consider whether it might be possible to weaken the hypothesis of Theorem 4.10 to just assuming that  $\mathbb{Z}M_1$  and  $\mathbb{Z}M_2$  are flat as  $\mathbb{Z}W$ -modules. In [23, Lemma 5.2(a)], Fiedorowicz claims that if  $[M_1, M_2; W]$  is an amalgam of monoids such that  $\mathbb{Z}M_1$  and  $\mathbb{Z}M_2$  are flat as left  $\mathbb{Z}W$ -modules, then  $\mathbb{Z}L$  (where  $L = M_1 *_W M_2$ ) is flat as a left  $\mathbb{Z}M_i$ -module, for  $i = 1, 2$ , and as a left  $\mathbb{Z}W$ -module. Unfortunately, his result is not correct. The following counterexample to [23, Lemma 5.2(a)] is due to Tyler Lawson (see [40]), whom we thank for allowing us to reproduce it. Let

$$M_1 = \langle a, a^{-1} \mid aa^{-1} = 1, a^{-1}a = 1 \rangle, \quad W = \{b\}^*, \quad \text{and} \quad M_2 = \{c, d\}^*.$$

So  $M_1$  is isomorphic to the infinite cyclic group, and  $W$  and  $M_2$  are the free monoids of ranks 1 and 2, respectively. Let  $f_1: W \rightarrow M_1$  be the homomorphism which maps  $b \mapsto a$ , let  $f_2: W \rightarrow M_2$  be the homomorphism which maps  $b \mapsto c$ , and let  $L$  be the monoid amalgam  $[M_1, M_2; W]$  with respect to the embeddings  $f_1$  and  $f_2$ . Then  $L$  is isomorphic to the monoid with presentation

$$\langle a, a^{-1}, d \mid aa^{-1} = 1, a^{-1}a = 1 \rangle,$$

that is, to  $\mathbb{Z} * \{d\}^*$ .

As the commutative ring  $\mathbb{Z}M_1$  is a localization of  $\mathbb{Z}W$ , it is clearly flat as a left  $\mathbb{Z}W$ -module. Since  $W$  is a free factor in  $M_2$ , we have that  $M_2$  is a free left  $W$ -set and hence  $\mathbb{Z}M_2$  is a free

left  $\mathbb{Z}W$ -module (and thus flat). On the other hand,  $\mathbb{Z}L$  is not flat as a left  $\mathbb{Z}M_2$ -module. This may be shown by considering the exact sequence of  $\mathbb{Z}M_2$ -modules

$$0 \rightarrow \mathbb{Z}M_2 \oplus \mathbb{Z}M_2 \rightarrow \mathbb{Z}M_2 \rightarrow \mathbb{Z} \rightarrow 0,$$

where the first map sends  $(u, v)$  to  $uc + vd$ , and the second sends  $c$  and  $d$  to zero. Here  $\mathbb{Z}$  is made a left  $\mathbb{Z}M_2$ -module by having  $c$  and  $d$  annihilate it rather than via the trivial module structure. Tensoring this sequence over  $\mathbb{Z}M_2$  on the left by  $\mathbb{Z}L$  gives the sequence

$$0 \rightarrow \mathbb{Z}L \oplus \mathbb{Z}L \rightarrow \mathbb{Z}L \rightarrow 0 \rightarrow 0,$$

which is not left exact since the first factor of the direct sum is taken isomorphically to the middle term by invertibility of  $a$ . Hence  $\mathbb{Z}L$  is not flat as a  $\mathbb{Z}M_2$ -module. A nearly identical proof was given by Bergman to show that universal localization does not preserve flatness in the non-commutative setting [5, Page 70].

Since [23, Lemma 5.2(a)] does not hold, it cannot be used to weaken the hypothesis of Theorem 4.10 to assuming only that  $\mathbb{Z}M_1$  and  $\mathbb{Z}M_2$  are flat as  $\mathbb{Z}W$ -modules. Similarly [23, Lemma 5.2(a)] cannot be used to weaken the hypotheses of any of Theorems 4.14, 4.28 or 4.29.

It follows from results of McDuff [49] that the hypotheses of Theorem 4.5 are satisfied when  $M_1$  and  $M_2$  are free as  $W$ -sets which gives the following corollary.

**Corollary 4.12.** *Let  $[M_1, M_2; W]$  be an amalgam of monoids such that  $M_1, M_2$  are free as right  $W$ -sets. If  $M_1, M_2$  are of type left-FP $_n$  and  $W$  is of type left-FP $_{n-1}$ , then  $M_1 *_W M_2$  is of type left-FP $_n$ .*

Corollary 4.12 applies, in particular, when  $W$  is trivial. Thus we obtain the following improvement on [17, Theorem 5.5] in which we do not need to assume the factors are finitely presented.

**Corollary 4.13.** *Let  $M_1, M_2$  be monoids of type left-FP $_n$ . Then  $M_1 * M_2$  is of type left-FP $_n$ .*

**Theorem 4.14.** *Let  $[M_1, M_2; W]$  be an amalgam of monoids such that  $\mathbb{Z}L$ , where  $L = M_1 *_W M_2$ , is flat as a right  $\mathbb{Z}M_1$ -,  $\mathbb{Z}M_2$ - and  $\mathbb{Z}W$ -module. If  $M_1, M_2$  have left cohomological dimension at most  $d$  and  $W$  has left cohomological dimension at most  $d - 1$ , then  $M_1 *_W M_2$  has left cohomological dimension at most  $d$ .*

*Proof.* By Lemma 2.5 and the hypotheses, we deduce that  $\mathbb{Z}L \otimes_{\mathbb{Z}M_i} \mathbb{Z}$  is of cohomological dimension at most  $d$ , for  $i = 1, 2$ , and  $\mathbb{Z}L \otimes_{\mathbb{Z}A} \mathbb{Z}$  is of cohomological dimension  $d - 1$ . We deduce the theorem by applying Corollary 2.4 to the exact sequence in Corollary 4.3.  $\square$

Again, combining this with results of McDuff [49] gives the following.

**Corollary 4.15.** *Let  $[M_1, M_2; W]$  be an amalgam of monoids such that  $M_1, M_2$  are free as right  $W$ -sets. Suppose that  $d_i$  is the left cohomological dimension of  $M_i$ , for  $i = 1, 2$ , and  $d$  is the left cohomological dimension of  $W$ . Then the left cohomological dimension of  $M_1 *_W M_2$  is bounded above by  $\max\{d_1, d_2, d + 1\}$ .*

**4.2. The two-sided setting.** We need some preliminary properties of tensor products before investigating amalgams in the two-sided context.

**Proposition 4.16.** *If  $f: M \rightarrow N$  is a monoid homomorphism, then there is an  $N \times N^{op}$  isomorphism  $F: N \otimes_M M \otimes_M N \rightarrow N \otimes_M N$  defined by  $F(n \otimes m \otimes n') = nm \otimes n'$ .*

*Proof.* The mapping  $h: N \times M \times N \rightarrow N \otimes_M N$  given by  $(n, m, n') \mapsto nm \otimes n'$  is  $N \times N^{op}$ -equivariant and satisfies  $(nm', m, m''n') \mapsto nm'm \otimes m''n' = nm'mm'' \otimes n' = h(n, m'mm'', n')$  and so the mapping  $F$  is well defined. The mapping  $k: N \times N \rightarrow N \otimes_M M \otimes_M N$  given by  $(n, n') \mapsto n \otimes 1 \otimes n'$  satisfies  $k(nm, n') = nm \otimes 1 \otimes n' = n \otimes m \otimes n' = n \otimes 1 \otimes mn' = k(n, mn')$  for  $m \in M$  and hence induces a mapping  $N \otimes_M N \rightarrow N \otimes_M M \otimes_M N$ . Clearly,  $h$  and  $k$  induce inverse mappings as  $nm \otimes 1 \otimes n' = n \otimes m \otimes n'$  for  $m \in M$ .  $\square$



The next proposition will frequently be used to decongest notation.

**Proposition 4.17.** *Let  $A$  be right  $M$ -set,  $B$  a left  $M$ -set and  $C$  a left  $M \times M^{op}$ -set. Then  $A \otimes_M C \otimes_M B$  is naturally isomorphic to  $(A \times B) \otimes_{M \times M^{op}} C$  in the category of sets where we view  $A \times B$  as a right  $M \times M^{op}$ -set via the action  $(a, b)(m, m') = (am, m'b)$ .*

*Proof.* Define  $f: A \times C \times B \rightarrow (A \times B) \otimes_{M \times M^{op}} C$  by  $f(a, c, b) = (a, b) \otimes c$ . Then  $f(am, c, m'b) = (am, m'b) \otimes c = (a, b) \otimes mcm'$  and so  $f$  induces a well-defined mapping  $A \otimes_M C \otimes_M B \rightarrow (A \times B) \otimes_{M \times M^{op}} C$ . Define  $g: A \times B \times C \rightarrow A \otimes_M C \otimes_M B$  by  $g(a, b, c) = a \otimes c \otimes b$ . Then  $g(am, m'b, c) = am \otimes c \otimes m'b = a \otimes mcm' \otimes b = g(a, b, mcm')$  and so  $g$  induces a well-defined mapping  $(A \times B) \otimes_{M \times M^{op}} C \rightarrow A \otimes_M C \otimes_M B$ . The maps induced by  $f$  and  $g$  are clearly mutually inverse and natural in  $A, B, C$ .  $\square$

*Remark 4.18.* A nearly identical proof shows that if  $A$  is a right  $\mathbb{Z}M$ -module,  $B$  is a left  $\mathbb{Z}M$ -module and  $C$  is an  $\mathbb{Z}M$ -bimodule, then we have that  $A \otimes_{\mathbb{Z}M} C \otimes_{\mathbb{Z}M} B \cong (A \otimes B) \otimes_{\mathbb{Z}M \otimes \mathbb{Z}M^{op}} C$  as abelian groups and the isomorphism is natural.

**Proposition 4.19.** *Suppose that  $A$  is a free right  $M$ -set,  $B$  is a free left  $M$ -set and  $C$  is an  $M$ - $M$ -biset. Then  $A \otimes_M C \otimes_M B$  is naturally isomorphic to  $A/M \times C \times M \setminus B$  in the category of sets.*

*Proof.* By freeness,  $A \otimes_M C \cong A/M \times C$  via  $a \otimes c \mapsto ([a], c)$  where  $[a]$  is the class of  $a$  and, moreover, this is a right  $M$ -set isomorphism. Therefore,  $A \otimes_M C \otimes_M B \cong (A/M \times C) \otimes_M B \cong A/M \times C \times M \setminus B$  because  $B$  is a free left  $M$ -set on  $M \setminus B$ . The isomorphism is clearly natural in  $A, B, C$ .  $\square$

We now wish to consider a pushout diagram (4.1) in the bimodule setting. Let us assume that  $f_i: W \rightarrow M_i$  is the homomorphism in the diagram, for  $i = 1, 2$ , and we continue to use  $L$  to denote the pushout. Let us proceed to define a forest  $T$ . The vertex set of  $T$  will be

$$V = (L \otimes_{M_1} L) \coprod (L \otimes_{M_2} L)$$

and the edge set will be

$$E = L \otimes_W L.$$

We shall write  $[x, y]_K$  for the tensor  $x \otimes y$  in  $L \otimes_K L$  for  $K = M_1, M_2, W$ . The edge  $[x, y]_W$  will connect  $[x, y]_{M_1}$  to  $[x, y]_{M_2}$ , and we think of it as oriented in this direction. Note that  $T$  is an  $L \times L^{op}$ -graph. Note that  $[x, y]_K \mapsto xy$  is well defined for any of  $K = M_1, M_2, W$ .

**Lemma 4.20.** *There is an  $L \times L^{op}$ -equivariant isomorphism  $\pi_0(T) \rightarrow L$  induced by the multiplication map on vertices.*

*Proof.* As an edge  $[x, y]_W$  connects  $[x, y]_{M_1}$  to  $[x, y]_{M_2}$ , we have that multiplication  $[x, y]_{M_i} \mapsto xy$  on vertices induces an  $L \times L^{op}$ -equivariant surjective mapping  $\pi_0(T) \rightarrow L$ . To prove the injectivity, we first claim that  $[1, x]_{M_1}$  is connected by an edge path to  $[x, 1]_{M_1}$  for all  $x \in L$  by induction on the length of  $x$ . If  $x = 1$ , there is nothing to prove. So assume the claim for length  $k$  and let  $x = x_1 \cdots x_{k+1}$  with  $x_i \in M_1 \cup M_2$  (again abusing notation as  $M_i$  need not embed in  $L$ ). Let  $p$  be a path from  $[1, x_2 \cdots x_{k+1}]_{M_1}$  to  $[x_2 \cdots x_{k+1}, 1]_{M_1}$ . Then  $x_1 p_1$  is a path from  $[x_1, x_2 \cdots x_{k+1}]_{M_1}$  to  $[x, 1]_{M_1}$ . If  $x_1 \in M_1$ , then  $[x_1, x_2 \cdots x_{k+1}]_{M_1} = [1, x]_{M_1}$  and we are done. If  $x_1 \in M_2$ , then  $[x_1, x_2 \cdots x_{k+1}]_W$  is an edge between  $[x_1, x_2 \cdots x_{k+1}]_{M_1}$  and  $[1, x]_{M_2}$ . But  $[1, x]_W$  is an edge from  $[1, x]_{M_1}$  to  $[1, x]_{M_2}$  and so we are again done in this case.

If  $x = x_1 x_2$  with  $x_1, x_2 \in L$ , there is a path  $p$  from  $[1, x_1]_{M_1}$  to  $[x_1, 1]_{M_1}$  by the above claim. Then  $p x_2$  is a path from  $[1, x]_{M_1}$  to  $[x_1, x_2]_{M_1}$ . Thus any two vertices  $[u, v]_{M_1}$  and  $[u', v']_{M_1}$  with  $uv = u'v'$  are connected in  $T$ . But  $[u, v]_W$  connects  $[u, v]_{M_2}$  to  $[u, v]_{M_1}$  and hence any two vertices  $[u, v]_{M_i}$  and  $[u', v']_{M_j}$  with  $uv = u'v'$  are connected for all  $i, j \in \{1, 2\}$ . This completes the proof.  $\square$

Next we prove that  $T$  is a forest. Note that  $\mathbb{Z}E$  is a  $\mathbb{Z}L$ -bimodule.

If  $A$  is a bimodule over a monoid ring  $\mathbb{Z}K$  then we can form the two-sided semidirect product  $A \bowtie K$ , of the abelian group  $A$  and the monoid  $K$ , with elements  $A \times K$  and multiplication given by

$$(a, k)(a', k') = (ak' + ka', kk').$$

A *splitting*  $\sigma$  of the projection  $\pi: A \bowtie K \rightarrow K$  is a monoid homomorphism  $\sigma: K \rightarrow A \bowtie K$  such that  $\pi(\sigma(k)) = k$  for all  $k \in K$ . A mapping  $d: K \rightarrow A$  is a *derivation* if

$$d(kk') = kd(k') + d(k)k'$$

for all  $k, k' \in K$ . A derivation is *inner* if  $d(k) = ka - ak$  for some  $a \in A$ . Derivations correspond to splittings of the two-sided semidirect product projection  $A \bowtie K \rightarrow K$ , each splitting being of the form  $k \mapsto (d(k), k)$  with  $d$  a derivation.

**Lemma 4.21.** *The graph  $T$  is a forest.*

*Proof.* A graph with vertex set  $V$  and edge set  $E$  is a forest if and only if the cellular boundary map  $\partial: \mathbb{Z}E \rightarrow \mathbb{Z}V$  is injective. We again use derivations to construct a left inverse to  $\partial$ . As usual, we identify elements of  $M_1$ ,  $M_2$  and  $W$  with their images in  $L$  (abusing notation).

Define  $\varphi_1: M_1 \rightarrow \mathbb{Z}E \bowtie L$  by  $\varphi_1(m_1) = (0, m_1)$ ; this is clearly a homomorphism. Next define  $\varphi_2: M_2 \rightarrow \mathbb{Z}E \bowtie L$  by  $\varphi_2(m_2) = ([1, m_2]_W - [m_2, 1]_W, m_2)$ . Note that  $m_2 \mapsto [1, m_2]_W - [m_2, 1]_W$  is the inner derivation of the  $\mathbb{Z}M_2$ -bimodule  $\mathbb{Z}E$  associated to the element  $-[1, 1]_W$  and hence  $\varphi_2$  is a homomorphism. If  $w \in W$ , then

$$\varphi_2 f_2(w) = ([1, w]_W - [w, 1]_W, w) = (0, w) = \varphi_1 f_1(w)$$

as  $[1, w]_W = [w, 1]_W$  for  $w \in W$ . Therefore, there is a homomorphism  $\varphi: L \rightarrow \mathbb{Z}E \bowtie L$  extending  $\varphi_1, \varphi_2$ , which is a splitting of the projection by construction. Thus  $\varphi(x) = (d(x), x)$  for some derivation  $d: L \rightarrow \mathbb{Z}E$  satisfying  $d(m_1) = 0$  for  $m_1 \in M_1$  and  $d(m_2) = [1, m_2]_W - [m_2, 1]_W$  for  $m_2 \in M_2$ .

We now define  $\beta: \mathbb{Z}V \rightarrow \mathbb{Z}E$  by  $\beta([x, y]_{M_1}) = d(x)y$  and  $\beta([x, y]_{M_2}) = d(x)y + [x, y]_W$ . To show that this is well defined, we need that if  $m_1 \in M_1$ , then  $[xm_1, y]_{M_1}$  and  $[x, m_1y]_{M_1}$  are sent to the same element and if  $m_2 \in M_2$ , then  $[xm_2, y]_{M_2}$  and  $[x, m_2y]_{M_2}$  are sent to the same element. But  $d(xm_1)y = xd(m_1)y + d(x)m_1y = d(x)m_1y$  because  $d(m_1) = 0$ . Also, we compute

$$\begin{aligned} d(xm_2)y + [xm_2, y]_W &= xd(m_2)y + d(x)m_2y + [xm_2, y]_W \\ &= x([1, m_2]_W - [m_2, 1]_W)y + d(x)m_2y + [xm_2, y]_W \\ &= d(x)m_2y + [x, m_2y]_W. \end{aligned}$$

We then obtain

$$\beta\partial([x, y]_W) = \beta([x, y]_{M_2}) - \beta([x, y]_{M_1}) = d(x)y + [x, y]_W - d(x)y = [x, y]_W.$$

Thus  $\beta\partial = 1_{\mathbb{Z}E}$  and hence  $\partial$  is injective. This completes the proof that  $T$  is a forest.  $\square$

We call  $T$  the *Bass–Serre forest* of the pushout. Since  $H_0(T) \cong \mathbb{Z}\pi_0(T) \cong \mathbb{Z}L$  as an  $L \times L^{op}$ -bimodule (by Lemma 4.20), Lemma 4.21 has the following reinterpretation.

**Corollary 4.22.** *There is an exact sequence of  $L \times L^{op}$ -modules*

$$0 \longrightarrow \mathbb{Z}L \otimes_{\mathbb{Z}W} \mathbb{Z}L \longrightarrow (\mathbb{Z}L \otimes_{\mathbb{Z}M_1} \mathbb{Z}L) \oplus (\mathbb{Z}L \otimes_{\mathbb{Z}M_2} \mathbb{Z}L) \longrightarrow \mathbb{Z}L \longrightarrow 0$$

where  $L = M_1 *_W M_2$  is the pushout.

*Proof.* This follows by consideration of the cellular chain complex of the forest  $T$  and using that  $\mathbb{Z}V/\partial\mathbb{Z}E = H_0(T) \cong \mathbb{Z}L$ , as observed before the corollary.  $\square$

**Theorem 4.23.** *Let  $[M_1, M_2; W]$  be an amalgam of monoids such that  $M_1, M_2$  are free as both left and right  $W$ -sets. If  $M_1, M_2$  are of type  $bi\text{-}F_n$  and  $W$  is of type  $bi\text{-}F_{n-1}$ , then  $M_1 *_W M_2$  is of type  $bi\text{-}F_n$ .*

*Proof.* Let  $X_i$  be a bi-equivariant classifying space for  $M_i$  with  $M_i \times M_i^{op}$ -finite  $n$ -skeleton, for  $i = 1, 2$ , and  $Y$  a bi-equivariant classifying space for  $W$  with  $W \times W^{op}$ -finite  $(n - 1)$ -skeleton. Fix bi-equivariant isomorphisms  $r_i: M_i \rightarrow \pi_0(X_i)$  and  $r: W \rightarrow \pi_0(Y)$ . By [25, Lemma 7.1] and the cellular approximation theorem [25, Theorem 2.8], we can find  $W \times W^{op}$ -equivariant cellular mappings  $f_i: Y \rightarrow X_i$ , for  $i = 1, 2$ , such that the composition of  $r$  with the composition of the mapping induced by  $f_i$  with  $r_i^{-1}$  is the inclusion, for  $i = 1, 2$ . Let  $L = M_1 *_W M_2$ . By McDuff [49],  $L$  is a free as both a left and a right  $M_i$ -set, for  $i = 1, 2$ , and as a left and right  $W$ -set.

For  $i = 1, 2$ ,  $X'_i = L \otimes_{M_i} X_i \otimes_{M_i} L \cong (L \times L^{op}) \otimes_{L \times L^{op}} X_i$  (the isomorphism by Proposition 4.17) is a projective  $L \times L^{op}$ -CW complex with  $L \times L^{op}$ -finite  $n$ -skeleton and  $Y' = L \otimes_W Y \otimes_W L \cong (L \times L^{op}) \otimes_{L \times L^{op}} Y$  is a projective  $L \times L^{op}$ -CW complex with  $L \times L^{op}$ -finite  $(n - 1)$ -skeleton by Proposition 2.1. Let  $F_i: Y' \rightarrow X'_i$  be the mapping induced by  $f_i$ , for  $i = 1, 2$ , and let  $Z$  be the homotopy pushout of  $F_1, F_2$ ; it is a projective  $L \times L^{op}$ -CW complex. We claim that  $Z$  is a bi-equivariant classifying space for  $L$ . Note that  $Z$  has an  $L \times L^{op}$ -finite  $n$ -skeleton by construction.

Our goal is to show that  $Z$  is homotopy equivalent to the Bass–Serre forest  $T$  via an  $L \times L^{op}$ -equivariant homotopy equivalence. By [25, Proposition 3.4] and Proposition 4.16 we have that  $\pi_0(X'_i) \cong L \otimes_{M_i} M \otimes_{M_i} L \cong L \otimes_{M_i} L$ , for  $i = 1, 2$ , and  $\pi_0(Y') \cong L \otimes_W W \otimes_W L \cong L \otimes_W L$  and, moreover,  $F_i$  induces the natural mapping  $L \otimes_W L \rightarrow L \otimes_{M_i} L$ , for  $i = 1, 2$  (by construction). Thus, by Lemma 4.4, it suffices to show that the projections  $X'_i \rightarrow \pi_0(X'_i)$ , for  $i = 1, 2$ , and  $Y' \rightarrow \pi_0(Y')$  are homotopy equivalences.

Since  $L$  is free as a left and right  $M_i$ -set, for  $i = 1, 2$ , and as a left and right  $W$ -set, we have by Proposition 4.19 that  $X'_i \cong L/M_i \times X_i \times M_i/L$  (for  $i = 1, 2$ ) and  $Y' \cong L/W \times Y \times W/L$ . As  $X_1, X_2, Y$  are homotopy equivalent to their sets of path components via the canonical projection, we deduce that the projections to path components are, indeed, homotopy equivalences for  $X'_1, X'_2, Y'$ . This completes the proof.  $\square$

The hypotheses of Theorem 4.23, of course, hold if  $W$  is trivial. It also holds if we amalgamate two copies of  $\mathbb{N}$  along cyclic submonoids. So  $\mathbb{N} *_k \mathbb{N} = m\mathbb{N}$  is of type  $\text{bi-F}_\infty$  for any  $m, k > 0$ .

**Corollary 4.24.** *A free product  $M * N$  of monoids of type  $\text{bi-F}_n$  is of type  $\text{bi-F}_n$ . If  $M, N$  are finitely presented monoids, then  $M * N$  is of type  $\text{bi-FP}_n$  if and only if  $M$  and  $N$  both are of type  $\text{bi-FP}_n$ .*

*Proof.* The first statement follows from Theorem 4.23. The second follows from the equivalence of  $\text{bi-F}_n$  and  $\text{bi-FP}_n$  for finitely presented monoids and the result of Pride [53] that the class of monoids of type  $\text{bi-FP}_n$  is closed under retracts.  $\square$

The hypotheses of Theorem 4.23 also hold if  $M_1, M_2$  are cancellative and  $W$  is a group. The homotopy pushout construction in the proof of Theorem 4.23 yields the following theorem.

**Theorem 4.25.** *Let  $[M_1, M_2; W]$  be an amalgam of monoids such that  $M_1, M_2$  are free as left and right  $W$ -sets. Suppose that  $d_i$  is the geometric dimension of  $M_i$ , for  $i = 1, 2$  and  $d$  is the geometric dimension of  $W$ . Then the geometric dimension of  $M_1 *_W M_2$  is bounded above by  $\max\{d_1, d_2, d + 1\}$ .*

Since only the trivial monoid has geometric dimension 0, we obtain the following special case.

**Corollary 4.26.** *Let  $M$  and  $N$  be monoids of geometric dimension at most  $n$ . Then  $M * N$  has geometric dimension at most  $n$ .*

Next we wish to consider the homological analogue.

**Proposition 4.27.** *Suppose that  $A$  is a flat right  $\mathbb{Z}M$ -module and  $B$  is a flat left  $\mathbb{Z}M$ -module. Then  $A \otimes B$  is a flat right  $\mathbb{Z}M \otimes \mathbb{Z}M^{op}$ -module (with respect to the structure  $(a \otimes b)(m, m') = am \otimes m'b$ ).*

*Proof.* If  $0 \rightarrow J \rightarrow K \rightarrow L \rightarrow 0$  is a short exact sequences of  $M$ -bimodules, then  $0 \rightarrow A \otimes_{\mathbb{Z}M} J \rightarrow A \otimes_{\mathbb{Z}M} K \rightarrow A \otimes_{\mathbb{Z}M} L \rightarrow 0$  is exact by flatness of  $A$ . Therefore,

$$0 \rightarrow A \otimes_{\mathbb{Z}M} J \otimes_{\mathbb{Z}M} B \rightarrow A \otimes_{\mathbb{Z}M} K \otimes_{\mathbb{Z}M} B \rightarrow A \otimes_{\mathbb{Z}M} L \otimes_{\mathbb{Z}M} B \rightarrow 0$$

is exact by flatness of  $B$ . The result now follows by Remark 4.18.  $\square$

**Theorem 4.28.** *Let  $[M_1, M_2; W]$  be an amalgam of monoids such that  $\mathbb{Z}L$  is flat as both a left and right  $\mathbb{Z}M_i$ -module and  $\mathbb{Z}W$ -module, for  $i = 1, 2$ , where  $L = M_1 *_W M_2$ . If  $M_1, M_2$  are of type bi-FP $_n$  and  $W$  is of type bi-FP $_{n-1}$ , then  $M_1 *_W M_2$  is of type bi-FP $_n$ .*

*Proof.* Note that  $\mathbb{Z}[L \times L^{op}] \cong \mathbb{Z}L \otimes \mathbb{Z}L^{op}$  is a flat right  $\mathbb{Z}[M_i \times M_i^{op}]$ -module, for  $i = 1, 2$ , and a flat right- $\mathbb{Z}[W \times W^{op}]$ -module by Proposition 4.27. By Lemma 2.5 and the hypotheses, we deduce that  $\mathbb{Z}[L \times L^{op}] \otimes_{\mathbb{Z}[M_i \times M_i^{op}]} \mathbb{Z}M_i$  is of type FP $_n$ , for  $i = 1, 2$ , and  $\mathbb{Z}[L \times L^{op}] \otimes_{\mathbb{Z}[W \times W^{op}]} \mathbb{Z}W$  is of type FP $_{n-1}$ . The result now follows by applying Corollary 2.4 to the exact sequence in Corollary 4.22, in light of Proposition 4.16 and Proposition 4.17.  $\square$

**Theorem 4.29.** *Suppose that  $[M_1, M_2; W]$  is an amalgam of monoids such that  $M_i$  has Hochschild cohomological dimension at most  $d$ , for  $i = 1, 2$ ,  $W$  has Hochschild cohomological dimension at most  $d - 1$ , and  $\mathbb{Z}L$  is flat as both a left and right  $\mathbb{Z}M_i$ -module and  $\mathbb{Z}W$ -module, for  $i = 1, 2$ , where  $L = M_1 *_W M_2$ . Then  $M_1 *_W M_2$  has Hochschild cohomological dimension at most  $d$ .*

As with the one-sided results, combining these results with results of McDuff [49] gives the following corollaries.

**Corollary 4.30.** *Let  $[M_1, M_2; W]$  be an amalgam of monoids such that  $M_1, M_2$  are free as both left and right  $W$ -sets. If  $M_1, M_2$  are of type bi-FP $_n$  and  $W$  is of type bi-FP $_{n-1}$ , then  $M_1 *_W M_2$  is of type bi-FP $_n$ . This applies, in particular, to free products.*

**Corollary 4.31.** *Let  $[M_1, M_2; W]$  be an amalgam of monoids such that  $M_1, M_2$  are free as left and right  $W$ -sets. Suppose that  $d_i$  is the Hochschild cohomological dimension of  $M_i$ , for  $i = 1, 2$  and  $d$  is the Hochschild cohomological dimension of  $W$ . Then the Hochschild cohomological dimension of  $M_1 *_W M_2$  is bounded above by  $\max\{d_1, d_2, d + 1\}$ .*

We remark that the results of this section and the previous section have analogues for the amalgamation of a finite family of monoids over a common submonoid.

## 5. HNN EXTENSIONS

In this section we shall present several new theorems about the behaviour of homological and topological finiteness properties for HNN extensions of monoids. Several natural HNN extension definitions for monoids have arisen in the literature in different contexts.

First in this section we consider a generalization of a construction of Otto and Pride, which they used to distinguish finite derivation type from finite homological type [54]. Let  $M$  be a monoid,  $A$  a submonoid and  $\varphi: A \rightarrow M$  a homomorphism. The free monoid generated by a set  $A$  is denoted by  $A^*$ . Then the *Otto-Pride extension* of  $M$  with base monoid  $A$  is the quotient  $L$  of the free product  $M * \{t\}^*$  by the smallest congruence such that  $at = t\varphi(a)$  for  $a \in A$ , i.e.,  $L = \langle M, t \mid at = t\varphi(a), a \in A \rangle$ . For example, if  $A = M$  and  $\varphi$  is the trivial homomorphism, then the Otto-Pride extension is the monoid  $M \cup \overline{M}$  where  $\overline{M}$  is an adjoined set of right zeroes in bijection with  $M$ . Otto and Pride have considered Otto-Pride extensions of groups where  $\varphi$  is injective, in [54] and [55].

**5.1. The one-sided case.** The following model for  $L$  will be useful for constructing normal forms and for proving flatness results.

**Proposition 5.1.** *View  $M$  as a right  $A$ -set via right multiplication and as a left  $A$ -set via the action  $a \odot m = \varphi(a)m$  for  $a \in A$ . Then  $L$  is isomorphic to the monoid with underlying set  $R = \coprod_{i=0}^{\infty} R_i$ , where  $R_0 = M$  and  $R_{i+1} = R_i \otimes_A M$ , and with multiplication defined by*

$$(m_1 \otimes \cdots \otimes m_k)(m'_1 \otimes \cdots \otimes m'_\ell) = m_1 \otimes \cdots \otimes m_{k-1} \otimes m_k m'_1 \otimes m'_2 \otimes \cdots \otimes m'_\ell.$$

In particular,  $M$  and  $t^*$  embed in  $L$  (where  $t$  is identified with  $1 \otimes 1 \in R_1$ ).

*Proof.* It is a straightforward exercise to verify that  $R$  is a monoid with identity  $1 \in R_0 = M$ . Define  $f: M \cup \{t\} \rightarrow R$  by  $f(m) = m$  and  $f(t) = 1 \otimes 1$ . Then if  $a \in A$ , we have that  $f(a)f(t) = a \otimes 1 = 1 \otimes \varphi(a) = f(t)f(\varphi(a))$  and so  $f$  induces a homomorphism  $f: L \rightarrow R$ . Note that  $f$  is surjective. Indeed,  $R_0$  is in the image of  $f$  by construction. Assume that  $R_i$  is in the image of  $f$  and let  $m_1 \otimes \cdots \otimes m_{i+1} \in R_i$ . If  $f(x) = m_1 \otimes \cdots \otimes m_i$  (by induction), then  $f(xtm_{i+1}) = m_1 \otimes \cdots \otimes m_i \otimes m_{i+1}$ . Now define  $g: R \rightarrow L$  by  $g(m_1 \otimes \cdots \otimes m_i) = m_1 t m_2 t \cdots t m_i$ . It is easy to verify that this is well defined using the defining relations of  $L$  and trivially  $g$  is a homomorphism. Now  $gf(m) = m$  for  $m \in M$  and  $gf(t) = g(1 \otimes 1) = t$ . Therefore,  $gf = 1_L$  and so  $f$  is injective. This concludes the proof that  $f$  is an isomorphism.  $\square$

As a corollary, we can deduce a normal form theorem for  $L$  if  $M$  is free as a right  $A$ -set.

**Corollary 5.2.** *Let  $\varphi: A \rightarrow M$  be a homomorphism with  $A$  a submonoid of  $M$ . Let  $L = \langle M, t \mid at = t\varphi(a), a \in A \rangle$  be the Otto-Pride extension. Suppose that  $M$  is a free right  $A$ -set with basis  $C$  containing  $1$ . Then every element of  $M$  can be uniquely written in the form  $c_0 c_1 t c_2 \cdots t c_k a$  with  $k \geq 0$ ,  $c_i \in C$  and  $a \in A$ . Consequently,  $L$  is free both as a right  $M$ -set and a right  $A$ -set.*

*Proof.* Since  $M$  is free as a right  $A$ -set on  $C$ , retaining the notation of Proposition 5.1, we have that  $R_i \cong C^{i+1} \times A$  via the mapping  $(c_0, \dots, c_i, a) \mapsto c_0 \otimes c_1 \otimes \cdots \otimes c_i a$ . Composing this mapping with the isomorphism  $g$  in the proof of Proposition 5.1 provides the desired normal form. Clearly,  $L$  is a free right  $M$ -set on the normal forms with  $c_k = 1 = a$  and  $L$  is a free right  $A$ -set on the normal forms with  $a = 1$ . This completes the proof.  $\square$

Note that if  $M$  is left cancellative and  $A$  is a group, then  $M$  is a free right  $A$ -set.

**Corollary 5.3.** *Let  $M$  be a monoid,  $A$  a submonoid and  $\varphi: A \rightarrow M$  be a homomorphism. Let  $L = \langle M, t \mid at = t\varphi(a), a \in A \rangle$  be the Otto-Pride extension. Suppose that  $\mathbb{Z}M$  is flat as a right  $\mathbb{Z}A$ -module. Then  $\mathbb{Z}L$  is flat both as a right  $\mathbb{Z}M$ -module and a right  $\mathbb{Z}A$ -module.*

*Proof.* Put  $V_0 = \mathbb{Z}M$  and  $V_{i+1} = V_i \otimes_{\mathbb{Z}A} \mathbb{Z}M$ . Then by Proposition 5.1, we have that as a right  $\mathbb{Z}M$ -module,  $\mathbb{Z}L \cong \bigoplus_{i \geq 0} V_i$  so it suffices to show that  $V_i$  is flat as both a right  $\mathbb{Z}M$ -module and a right  $\mathbb{Z}A$ -module. We prove this by induction. As  $V_0$  is a free right  $\mathbb{Z}M$ -module and a flat  $\mathbb{Z}A$ -module, by assumption, this case is handled. Assume that  $V_i$  is flat both as a right  $\mathbb{Z}M$ -module and a right  $\mathbb{Z}A$ -module. Let  $h: U \rightarrow W$  be an injective homomorphism of  $\mathbb{Z}M$ -modules (respectively,  $\mathbb{Z}A$ -modules). Then the induced mapping  $\mathbb{Z}M \otimes_{\mathbb{Z}M} U \rightarrow \mathbb{Z}M \otimes_{\mathbb{Z}M} W$  (respectively,  $\mathbb{Z}M \otimes_{\mathbb{Z}A} U \rightarrow \mathbb{Z}M \otimes_{\mathbb{Z}A} W$ ) is injective since  $\mathbb{Z}M$  is flat as a right module over both  $\mathbb{Z}M$  and  $\mathbb{Z}A$ . Then tensoring these injective mappings on the left with  $V_i$  over  $\mathbb{Z}A$  results in an injective mapping by flatness of  $V_i$ . Thus we see that  $V_{i+1}$  is flat as a right  $\mathbb{Z}M$ -module and as a right  $\mathbb{Z}A$ -module.  $\square$

We now construct a Bass-Serre tree for Otto-Pride extensions. Again fix a monoid  $M$  together with a homomorphism  $\varphi: A \rightarrow M$  from a submonoid  $A$  and let  $L$  be the Otto-Pride extension. We define a graph  $T$  with vertex set  $V = L/M$  and edge set  $E = L/A$ . An edge  $[x]_A$  connects  $[x]_M$  to  $[xt]_M$  (oriented in this way), where  $[x]_K$  denotes the class of  $x$  in  $L/K$ . This is well defined because if  $a \in A$ , then  $[xa]_M = [x]_M$  and  $[xat]_M = [xt\varphi(a)]_M = [xt]_M$ . Clearly, the left action of  $L$  is by cellular mappings sending edges to edges and so  $T$  is an  $L$ -graph. We aim to prove that  $T$  is a tree.

**Lemma 5.4.** *The graph  $T$  is connected.*

*Proof.* The monoid  $L$  is generated by  $M \cup \{t\}$ . The length of an element  $x$  is its shortest expression as a product in these generators. We prove by induction on length that there is a path from  $[1]_M$  to  $[x]_M$ . If  $x = 1$ , there is nothing to prove. Assume that  $x = yz$  with  $y \in M \cup \{t\}$  and  $z$  of length one shorter. Let  $p$  be a path from  $[1]_M$  to  $[z]_M$ . Then  $yp$  is a path from  $[y]_M$  to  $[x]_M$ . If  $y \in M$ , then  $[y]_M = [1]_M$  and we are done. If  $y = t$ , then since  $[1]_A$

connects  $[1]_M$  with  $[t]_M = [y]_M$  and so we are done in this case, as well. It follows that  $T$  is connected.  $\square$

Next we use derivations to prove that  $T$  is a tree.

**Lemma 5.5.** *The graph  $T$  is a tree.*

*Proof.* We prove that  $\partial: \mathbb{Z}E \rightarrow \mathbb{Z}V$  is injective. It will then follow that  $T$  is a tree as it was already shown to be connected in Lemma 5.4. Define  $\gamma: M \cup \{t\} \rightarrow \mathbb{Z}E \rtimes L$  by  $\gamma(m) = (0, m)$  for  $m \in M$  and  $\gamma(t) = ([1]_A, t)$ . Then if  $a \in A$ , we have that  $\gamma(a)\gamma(t) = (0, a)([1]_A, t) = ([a]_A, at) = ([1]_A, t\varphi(a)) = ([1]_A, t)(0, \varphi(a)) = \gamma(t)\gamma(\varphi(a))$ . Therefore,  $\gamma$  extends to a homomorphism  $\gamma: L \rightarrow \mathbb{Z}E \rtimes L$  splitting the semidirect product. Thus  $\gamma(x) = (d(x), x)$  for some derivation  $d: L \rightarrow \mathbb{Z}E$  with  $d(m) = 0$  for  $m \in M$  and  $d(t) = [1]_A$ .

Define  $\beta: \mathbb{Z}V \rightarrow \mathbb{Z}E$  by  $\beta([x]_M) = d(x)$ . This is well defined because if  $m \in M$ , then  $d(xm) = xd(m) + d(x) = d(x)$  as  $d(m) = 0$ . Now we compute that  $\beta\partial([x]_A) = \beta([xt]_M) - \beta([x]_M) = d(xt) - d(x) = xd(t) + d(x) - d(x) = x[1]_A = [x]_A$ . Therefore,  $\beta\partial = 1_{\mathbb{Z}E}$  and hence  $\partial$  is injective. We conclude that  $T$  is a tree.  $\square$

We call  $T$  the *Bass–Serre tree* of the extension. Lemma 5.5 can be restated in terms of exact sequences using that  $\mathbb{Z}[L/K] \cong \mathbb{Z}L \otimes_{\mathbb{Z}K} \mathbb{Z}$  for  $K = M, A$ .

**Corollary 5.6.** *There is an exact sequence*

$$0 \longrightarrow \mathbb{Z}L \otimes_{\mathbb{Z}A} \mathbb{Z} \longrightarrow \mathbb{Z}L \otimes_{\mathbb{Z}M} \mathbb{Z} \longrightarrow \mathbb{Z} \longrightarrow 0$$

*of left  $\mathbb{Z}L$ -modules.*

The analogue of the homotopy pushout that we shall need in this context is the homotopy coequalizer. If  $f, g: Y \rightarrow X$  are continuous mappings, then the *homotopy coequalizer*  $M(f, g)$  is the space obtained by gluing  $Y \times I$  to  $X$  via the mapping  $h: Y \times \partial I \rightarrow X$  given by  $h(y, 0) = f(y)$  and  $h(y, 1) = g(y)$ . If  $X$  and  $Y$  are CW complexes and  $f, g$  are cellular, then  $M(f, g)$  is a CW complex. If  $X, Y$  are projective  $M$ -CW complexes and  $f, g$  are  $M$ -equivariant and cellular, then  $M(f, g)$  is a projective  $M$ -CW complex by [25, Lemma 2.1]. Moreover, if  $X$  has  $M$ -finite  $n$ -skeleton and  $Y$  has  $M$ -finite  $(n-1)$ -skeleton, then  $M(f, g)$  has  $M$ -finite  $n$ -skeleton.

Homotopy coequalizers like homotopy pushouts, are examples of homotopy colimits. If  $f', g': Y' \rightarrow X'$  are continuous mappings and  $r: Y \rightarrow Y'$  and  $s: X \rightarrow X'$  are continuous such that

$$\begin{array}{ccc} Y & \xrightarrow[f]{g} & X \\ r \downarrow & & \downarrow s \\ Y' & \xrightarrow[f']{g'} & X' \end{array}$$

commutes, then there is an induced continuous mapping  $t: M(f, g) \rightarrow M(f', g')$  (which will be  $M$ -equivariant if all spaces are  $M$ -spaces and all maps are  $M$ -equivariant). Moreover, if  $r, s$  are homotopy equivalences, then so is  $t$ ; see [21, page 19]. For example, the graph  $T$  is the homotopy coequalizer of  $i, j: L/A \rightarrow L/M$  given by  $i([x]_A) = [x]_A$  and  $j([x]_A) = [xt]_A$  (where these sets are viewed as discrete spaces).

**Theorem 5.7.** *Let  $M$  be a monoid,  $A$  a submonoid and  $\varphi: A \rightarrow M$  be a homomorphism. Let  $L = \langle M, t \mid at = t\varphi(a), a \in A \rangle$  be the Otto–Pride extension. Suppose that  $M$  is free as a right  $A$ -set. If  $M$  is of type  $\text{left-}\mathcal{F}_n$  and  $A$  is of type  $\text{left-}\mathcal{F}_{n-1}$ , then  $L$  is of type  $\text{left-}\mathcal{F}_n$ .*

*Proof.* Let  $X$  be an equivariant classifying space for  $M$  with  $M$ -finite  $n$ -skeleton and let  $Y$  be an equivariant classifying space for  $A$  with  $A$ -finite  $(n-1)$ -skeleton. Using [25, Lemma 6.2] and the cellular approximation theorem [25, Theorem 2.8], we can find continuous cellular mappings  $f, g: Y \rightarrow X$  such that  $f(ay) = af(y)$  and  $g(ay) = \varphi(a)g(y)$  for all  $a \in A$  and  $y \in Y$ . To construct  $g$ , we view  $X$  as an  $A$ -space via the action  $a \odot x = \varphi(a)x$  for  $a \in A$ . Let

$X' = L \otimes_M X$  and  $Y' = L \otimes_A Y$ . These are projective  $L$ -CW complexes by Proposition 2.1 and  $X'$  has  $L$ -finite  $n$ -skeleton,  $Y'$  has  $L$ -finite  $(n-1)$ -skeleton.

Let  $F: Y' \rightarrow X'$  be the mapping induced by  $f$  and define  $G: Y' \rightarrow X'$  by  $G(u \otimes y) = ut \otimes g(y)$ . The latter is well defined since if  $a \in A$ , then  $uat \otimes g(y) = ut\varphi(a) \otimes g(y) = ut \otimes \varphi(a)g(y) = ut \otimes g(ay)$ . Clearly,  $G$  is  $L$ -equivariant, continuous and cellular. Let  $Z = M(F, G)$  be the homotopy coequalizer. Then  $Z$  is a projective  $L$ -CW complex with  $L$ -finite  $n$ -skeleton. We aim to show that  $Z$  is homotopy equivalent to  $T$  and hence contractible.

By [25, Proposition 3.4] we have that  $\pi_0(Y') \cong L \otimes_A \pi_0(Y) \cong L/A$  and  $\pi_0(X') \cong L \otimes_M \pi_0(X) \cong L/M$  as  $X, Y$  are connected. By construction  $F$  and  $G$  induce the mappings  $[u]_A \mapsto [u]_M$  and  $[u]_A \mapsto [ut]_M$ , respectively, on path components under these identifications. As the tree  $T$  is the homotopy coequalizer of these two mappings, it suffices to show that the projections  $X' \rightarrow \pi_0(X')$  and  $Y' \rightarrow \pi_0(Y')$  are homotopy equivalences. Then  $Z$  will be homotopy equivalent to  $T$ .

Since  $L$  is free as a right  $M$ -set and as a right  $A$ -set, we have that  $X' \cong L/M \times X$  and  $Y' \cong L/A \times Y$  as  $L$ -CW complexes. As  $X$  and  $Y$  are contractible and  $L/M$  and  $L/A$  are discrete, we deduce that the projections to connected components are homotopy equivalences in both cases. This completes the proof.  $\square$

The proof of Theorem 5.7 can be used to show that if  $M$  is free as a right  $A$ -set,  $M$  has left geometric dimension  $d$  and  $A$  has left geometric dimension  $d'$ , then  $L$  has left geometric dimension at most  $\max\{d, d'+1\}$ . The hypothesis of Theorem 5.7 applies if  $M$  is left cancellative and  $A$  is a group or if  $M = \mathbb{N}$  and  $A$  is a cyclic submonoid.

Next we prove the homological analogue of Theorem 5.7 under the weaker assumption of flatness.

**Theorem 5.8.** *Let  $M$  be a monoid and let  $\varphi: A \rightarrow M$  be a homomorphism from a submonoid  $A$  of  $M$ . Let  $L = \langle M, t \mid at = t\varphi(a), a \in A \rangle$  be the Otto-Pride extension. Suppose that  $\mathbb{Z}M$  is flat as a right  $\mathbb{Z}A$ -module. If  $M$  is of type  $\text{left-FP}_n$  and  $A$  is of type  $\text{left-FP}_{n-1}$ , then  $L$  is of type  $\text{left-FP}_n$ .*

*Proof.* By Corollary 5.3,  $\mathbb{Z}L$  is flat as a right  $\mathbb{Z}M$ -module and as a right  $\mathbb{Z}A$ -module. It follows from Lemma 2.5 and the hypotheses that  $\mathbb{Z}L \otimes_{\mathbb{Z}M} \mathbb{Z}$  is of type  $\text{FP}_n$  and  $\mathbb{Z}L \otimes_{\mathbb{Z}A} \mathbb{Z}$  is of type  $\text{FP}_{n-1}$ . The result now follows by applying Corollary 2.4 to the exact sequence in Corollary 5.6.  $\square$

One can prove similarly the following theorem.

**Theorem 5.9.** *Let  $M$  be a monoid and  $\varphi: A \rightarrow M$  a homomorphism from a submonoid  $A$  of  $M$ . Let  $L = \langle M, t \mid at = t\varphi(a), a \in A \rangle$  be the Otto-Pride extension. Suppose that  $\mathbb{Z}M$  is flat as a right  $\mathbb{Z}A$ -module. If  $M$  has left cohomological dimension at most  $d$  and  $A$  has left cohomological dimension at most  $d-1$ , then  $L$  has left cohomological dimension at most  $d$ .*

**5.2. The two-sided case.** It turns out that in the two-sided setting we shall need to consider Otto-Pride extensions corresponding to injective monoid homomorphisms  $\varphi: A \rightarrow M$  from a submonoid  $A$  of  $M$  in order to make the construction left-right dual. Putting  $B = \varphi(A)$ , we have that  $B$  is isomorphic to  $A$ . Otto and Pride considered the special case when  $M$  and  $A$  are groups (and hence so is  $B$ ). We shall call an Otto-Pride extension *HNN-like* if  $\varphi$  is injective. Let  $L$  be the Otto-Pride extension. It is straightforward to check  $L = \langle M, t \mid tb = \varphi^{-1}(b)t, b \in B \rangle$  and hence left/right duals of Proposition 5.1 and Corollary 5.2 are valid with  $B$  in the role of  $A$  and using left sets instead of right sets. Note that an HNN-like Otto-Pride extension of groups, which is the case considered by Otto and Pride, embeds as a submonoid of the corresponding group HNN extension (note that the Otto-Pride extension does not contain  $t^{-1}$  and hence is a monoid, not a group). Our results give geometric proofs of a number of the results of [54] and [55].

In what follows, we shall always view  $L$  as a right  $A$ -set via left multiplication and as a left  $A$ -set via  $a \odot x = \varphi(a)x$ . Therefore, we view  $L \times L^{op}$  as a right  $A \times A^{op}$ -set via  $(x, y)(a, a') = (xa, \varphi(a')y)$ .

**Proposition 5.10.** *There is an isomorphism*

$$L \otimes_A L \cong L \otimes_A A \otimes_A L \cong (L \times L^{op}) \otimes_{A \times A^{op}} A$$

of left  $L \times L^{op}$ -sets.

*Proof.* The first isomorphism is given by  $x \otimes y \mapsto x \otimes 1 \otimes y$  with inverse  $x \otimes a \otimes y \mapsto xa \otimes y$  (the reader should check that these are well defined and equivariant). The second isomorphism sends  $x \otimes a \otimes y$  to  $(x, y) \otimes a$  with inverse mapping  $(x, y) \otimes a$  to  $x \otimes a \otimes y$ . The reader should again check that this is well defined and equivariant.  $\square$

We now associate a Bass–Serre forest  $T$  to an HNN-like Otto-pride extension. The vertex set of  $T$  is  $V = L \otimes_M L$  and the edge set is  $E = L \otimes_A L$ . Again, we write  $[x, y]_K$  for the tensor  $x \otimes y$  of  $L \otimes_K L$ , for  $K = M, A$ . With this notation, the edge  $[x, y]_A$  connects  $[x, ty]_M$  to  $[xt, y]_M$  (which we think of as oriented in this way). To check that this is well defined, observe that if  $x, y \in L$  and  $a \in A$ , then  $[xa, y]_A = [x, \varphi(a)y]_A$  and  $[xa, ty]_M = [x, aty]_M = [x, t\varphi(a)y]_M$  and  $[xat, y]_M = [xt\varphi(a), y]_M = [xt, \varphi(a)y]_M$ . By construction,  $T$  is an  $L \times L^{op}$ -graph.

It is immediate from the definition of the incidences in  $T$  that the multiplication mapping  $L \otimes_M L \rightarrow L$  induces an  $L \times L^{op}$ -equivariant surjection  $\pi_0(T) \rightarrow L$ . We aim to show that it is an isomorphism.

**Lemma 5.11.** *The multiplication mapping  $L \otimes_M L \rightarrow L$  induces an  $L \times L^{op}$ -equivariant isomorphism of  $\pi_0(T)$  with  $L$ .*

*Proof.* We first prove by induction on the length of  $x$  as a product of elements of  $M \cup \{t\}$  that there is a path from  $[1, x]_M$  to  $[x, 1]_M$ . If  $x = 1$ , there is nothing to prove. Otherwise, assume  $x = uy$  with  $u \in M \cup \{t\}$  and  $y$  of shorter length. Let  $p$  be a path from  $[1, y]_M$  to  $[y, 1]_M$ . Then  $up$  is a path from  $[u, y]_M$  to  $[x, 1]_M$ . If  $u \in M$ , then  $[u, y]_M = [1, x]_M$  and we are done. If  $u = t$ , then  $[1, y]_A$  is an edge connecting  $[1, x]_M = [1, ty]_M$  to  $[t, y]_M = [u, y]_M$  and so we are again done.

Now if  $x = uv$  in  $L$ , then by the above, there is a path  $p$  from  $[1, u]_M$  to  $[u, 1]_M$ . Then  $pv$  is a path from  $[1, x]_M$  to  $[u, v]_M$ . It follows that all vertices  $[u', v']_M$  with  $u'v' = x$  are in a single connected component and hence the multiplication map induces an isomorphism from  $\pi_0(T)$  to  $L$ .  $\square$

Next we use derivations to prove that  $T$  is a forest.

**Lemma 5.12.** *The graph  $T$  is a forest.*

*Proof.* It suffices to prove that the cellular boundary map  $\partial: \mathbb{Z}E \rightarrow \mathbb{Z}V$  is injective. Define a mapping  $\gamma: M \cup \{t\} \rightarrow \mathbb{Z}E \rtimes L$  by  $\gamma(m) = (0, m)$  for  $m \in M$  and  $\gamma(t) = ([1, 1]_A, t)$ . If  $a \in A$ , then we compute  $\gamma(a)\gamma(t) = ([a, 1]_A, at) = ([1, \varphi(a)]_A, t\varphi(a)) = \gamma(t)\gamma(\varphi(a))$  and hence  $\gamma$  extends to a homomorphism  $\gamma: L \rightarrow \mathbb{Z}E \rtimes L$  splitting the two-sided semidirect product projection. Thus  $\gamma(x) = (d(x), x)$  for some derivation  $d: L \rightarrow \mathbb{Z}E$  such that  $d(m) = 0$  for  $m \in M$  and  $d(t) = [1, 1]_A$ . Define  $\beta: \mathbb{Z}V \rightarrow \mathbb{Z}E$  by  $\beta([x, y]_M) = d(x)y$ . We must verify that  $\beta$  is well defined. If  $m \in M$ , then  $d(xm)y = xd(m)y + d(x)my = d(x)my$  because  $d(m) = 0$ . This shows that  $\beta$  is well defined. Next we compute that

$$\begin{aligned} \beta\partial([x, y]_A) &= \beta([xt, y]_M) - \beta([x, ty]_M) = d(xt)y - d(x)ty \\ &= xd(t)y + d(x)ty - d(x)ty = x[1, 1]_Ay = [x, y]_A \end{aligned}$$

as  $d(t) = [1, 1]_A$ . This establishes that  $\beta\partial = 1_{\mathbb{Z}E}$  and hence  $T$  is a forest.  $\square$



We call  $T$  the *Bass–Serre forest* for  $L$ .

The exactness of the sequence

$$0 \longrightarrow \mathbb{Z}E \longrightarrow \mathbb{Z}V \longrightarrow H_0(T) \longrightarrow 0,$$

coming from  $T$  being a forest, together with the isomorphism  $\mathbb{Z}L \cong \mathbb{Z}\pi_0(L) \cong H_0(T)$  coming from Lemma 5.11, yields the following exact sequence.

**Corollary 5.13.** *Let  $L$  be the HNN-like Otto–Pride extension associated to a monomorphism  $\varphi: A \rightarrow M$  with  $A$  a submonoid of  $M$ . Then there is an exact sequence*

$$0 \longrightarrow \mathbb{Z}L \otimes_{\mathbb{Z}A} \mathbb{Z}L \longrightarrow \mathbb{Z}L \otimes_{\mathbb{Z}M} \mathbb{Z}L \longrightarrow \mathbb{Z}L \longrightarrow 0$$

where  $\mathbb{Z}L$  is viewed as a right  $\mathbb{Z}A$ -module via the inclusion and as a left  $\mathbb{Z}A$ -module via  $\varphi$ .

Suppose that we have an HNN-like Otto–Pride extension  $L$  with base monoid  $A$  and monomorphism  $\varphi: A \rightarrow M$ . Put  $B = \varphi(A)$ .

**Proposition 5.14.** *If  $M$  is free as a right  $A$ -set and as a left  $B$ -set, then  $L$  is free as both a right and a left  $M$ -set. Moreover,  $L$  is free as a right  $A$ -set and a left  $B$ -set. Hence  $L$  is free as a left  $A$ -set via the action  $a \odot x = \varphi(a)x$  for  $a \in A$  and  $x \in L$ .*

*Proof.* This follows from Corollary 5.2 and its dual.  $\square$

The flat version is the following.

**Proposition 5.15.** *If  $\mathbb{Z}M$  is a flat right  $\mathbb{Z}A$ -module and a flat left  $\mathbb{Z}B$ -module, then  $\mathbb{Z}L$  is flat as both a right and a left  $\mathbb{Z}M$ -module. Furthermore,  $\mathbb{Z}L$  is flat as a right  $\mathbb{Z}A$ -module and a left  $\mathbb{Z}B$ -module. Thus  $\mathbb{Z}L$  is flat as a left  $\mathbb{Z}A$ -module via the  $\mathbb{Z}A$ -module structure coming from  $\varphi$ .*

*Proof.* This follows from Corollary 5.3 and its dual.  $\square$

We can now investigate the two-sided topological and homological finiteness of HNN-like Otto–Pride extensions. The following theorem generalises [54, Theorem 1] and [55, Theorem 5].

**Theorem 5.16.** *Let  $L$  be an HNN-like Otto–Pride extension of  $M$  with respect to an injective homomorphism  $\varphi: A \rightarrow M$  and put  $B = \varphi(A)$ . Suppose that  $M$  is free as a right  $A$ -set and as a left  $B$ -set. Then if  $M$  is of type  $\text{bi-F}_n$  and  $A$  is of type  $\text{bi-F}_{n-1}$ , then  $L$  is of type  $\text{bi-F}_n$ .*

*Proof.* Let  $X$  be a bi-equivariant classifying space for  $M$  with  $M$ -finite  $n$ -skeleton and  $Y$  a bi-equivariant classifying space for  $A$  with  $A$ -finite  $(n-1)$ -skeleton. Let  $r: M \rightarrow \pi_0(X)$  and  $r': A \rightarrow \pi_0(Y)$  be equivariant isomorphisms. By [25, Lemma 7.1] and the cellular approximation theorem [25, Theorem 2.8], we can find cellular mappings  $f_1, f_2: Y \rightarrow X$  such that  $f_1(aya') = af_1(y)a'$  and  $f_2(aya') = \varphi(a)f_2(y)\varphi(a')$  for  $a, a' \in A$  and  $y \in Y$  and, moreover,  $r^{-1}(f_1)_*r'$  is the inclusion and  $r^{-1}(f_2)_*r' = \varphi$  where  $(f_i)_*$  is the induced mapping on the set of path components, for  $i = 1, 2$ .

In what follows, we view  $L$  as a (free) right  $A$ -set via the inclusion and a (free) left  $A$ -set via  $\varphi$ . Put  $X' = L \otimes_M X \otimes_M L$  and  $Y' = L \otimes_A Y \otimes_A L$ . They are projective  $L \times L^{op}$ -CW complexes with  $L \times L^{op}$ -finite  $n$ -,  $(n-1)$ -skeletons, respectively, by Proposition 2.1 and 4.17. Define  $F_1, F_2: Y' \rightarrow X'$  by  $F_1(u \otimes y \otimes v) = u \otimes f_1(y) \otimes tv$  and  $F_2(u \otimes y \otimes v) = ut \otimes f_2(y) \otimes v$ . Let us verify that this is well defined. If  $a, a' \in A$ , then we have that  $ua \otimes f_1(y) \otimes t\varphi(a')v = ua \otimes f_1(y) \otimes a'tv = u \otimes f_1(aya') \otimes tv$  and so  $F_1$  is well defined. Also, we have that  $uat \otimes f_2(y) \otimes \varphi(a')v = ut\varphi(a) \otimes f_2(y) \otimes \varphi(a')v = ut\varphi(a)f_2(y)\varphi(a') \otimes v = ut\varphi(a)f_2(y)\varphi(a') \otimes v$  and so  $F_2$  is well defined. Clearly,  $F_1, F_2$  are continuous  $L \times L^{op}$ -equivariant cellular mappings. Let  $Z = M(F_1, F_2)$  be the homotopy coequalizer. It is a projective  $L \times L^{op}$ -CW complex with  $L \times L^{op}$ -finite  $n$ -skeleton by construction. We prove that  $Z$  is a bi-equivariant classifying space for  $Z$ . To do this it suffices to construct an  $L \times L^{op}$ -equivariant homotopy equivalence to the Bass–Serre forest  $T$ .

First note, by [25, Proposition 3.4], that  $\pi_0(X') \cong L \otimes_M M \otimes_M L \cong L \otimes_M L$  (by Proposition 4.16) and  $\pi_0(Y') \cong L \otimes_A A \otimes_A L \cong L \otimes_A L$  (by Proposition 5.10). The mapping  $L \otimes_A L \rightarrow L \otimes_M L$  induced by  $F_1$  is  $u \otimes v \mapsto u \otimes tv$  and the mapping induced by  $F_2$  is

$u \otimes v \mapsto ut \otimes v$ . As  $T$  is the homotopy coequalizer of these two mappings of discrete sets  $L \otimes_A L \rightarrow L \otimes_M L$ , to complete the proof it suffices to show that  $X'$  and  $Y'$  are homotopy equivalent to their sets of path components (via the natural projections). But this follows because  $X$  and  $Y$  are homotopy equivalent to their respective sets of path components and the isomorphisms  $X' \cong L/M \times X \times M \setminus L$  and  $Y' \cong L/A \times Y \times B \setminus L$  coming from  $L$  being free as both a left and right  $M$ -set and as a right  $A$ -set and left  $B$ -set (cf. Proposition 5.14).  $\square$

The hypotheses of Theorem 5.16 hold if  $M$  and  $A$  are groups or, more generally, if  $M$  is cancellative and  $A$  is a group. It also holds if  $M = \mathbb{N}$  and  $A$  is a cyclic submonoid. The proof of Theorem 5.16 shows that if  $M$  is free as a right  $A$ -set and a left  $B$ -set,  $M$  has geometric dimension  $d$  and  $A$  has geometric dimension  $d'$ , then  $L$  has geometric dimension at most  $\max\{d, d' + 1\}$ .

The flat homological analogue of Theorem 5.16 has a similar proof.

**Theorem 5.17.** *Let  $L$  be an HNN-like Otto-Pride extension of  $M$  with respect to a monomorphism  $\varphi: A \rightarrow M$  and put  $B = \varphi(A)$ . Assume that  $\mathbb{Z}M$  is flat as a right  $\mathbb{Z}A$ -module and as a left  $\mathbb{Z}B$ -module. If  $M$  is of type  $\text{bi-FP}_n$  and  $A$  is of type  $\text{bi-FP}_{n-1}$ , then  $L$  is of type  $\text{bi-FP}_n$ .*

*Proof.* We have that  $\mathbb{Z}L$  is flat as both a right and a left  $\mathbb{Z}A$ -module and as a right and a left  $\mathbb{Z}M$ -module by Proposition 5.15 (viewing  $L$  as a left  $A$ -module via  $\varphi$ ). Therefore,  $\mathbb{Z}[L \times L^{op}] \cong \mathbb{Z}L \otimes \mathbb{Z}L^{op}$  is flat as both a right  $\mathbb{Z}[M \times M^{op}]$ -module and as a right- $\mathbb{Z}[A \times A^{op}]$ -module by Proposition 4.27. Applying Lemma 2.5 and the hypotheses, we conclude that  $\mathbb{Z}[L \times L^{op}] \otimes_{\mathbb{Z}[M \times M^{op}]} \mathbb{Z}M$  is of type  $\text{FP}_n$  and  $\mathbb{Z}[L \times L^{op}] \otimes_{\mathbb{Z}[A \times A^{op}]} \mathbb{Z}A$  is of type  $\text{FP}_{n-1}$ . The result now follows by applying Corollary 2.4 to the exact sequence in Corollary 5.13, taking into account Proposition 4.16, Proposition 4.17 and Proposition 5.10.  $\square$

As an example, if  $M$  is any group containing a copy of  $\mathbb{Z}$  and  $A = \mathbb{N}$ , viewed as a submonoid of  $M$ , then since  $\mathbb{Z}M$  is free as a module over the group ring of  $\mathbb{Z}$ , which in turn is flat over the monoid ring of  $\mathbb{N}$ , being a localization, we conclude that  $\mathbb{Z}M$  is flat over the monoid ring of  $\mathbb{N}$ .

One can similarly prove that if  $L$  is an HNN-like Otto-Pride extension of  $M$  with respect to a monomorphism  $\varphi: A \rightarrow M$  and  $\mathbb{Z}M$  is flat as a right  $\mathbb{Z}A$ -module and as a left  $\mathbb{Z}B$ -module, where  $B = \varphi(A)$ , then if  $M$  has Hochschild cohomological dimension at most  $d$  and  $A$  has Hochschild cohomological dimension at most  $d - 1$ , then  $L$  has Hochschild cohomological dimension at most  $d$ .

We end this section by briefly explaining what happens for a different HNN extensions of monoids construction of the sort considered by Howie [32]. Suppose that  $M$  is a monoid and  $A, B$  are isomorphic submonoids via an isomorphism  $\varphi: A \rightarrow B$ . Let  $C$  be an infinite cyclic group generated by  $t$ . The HNN extension of  $M$  with base monoids  $A, B$  is the quotient  $L$  of the free product  $M * C$  by the congruence generated by the relations  $at = t\varphi(a)$  for  $a \in A$ . In other words,  $L = \langle M, t, t^{-1} \mid tt^{-1} = 1 = t^{-1}t, at = t\varphi(a), \forall a \in A \rangle$ . The following results may be proved in a similar way to Theorems 5.7 and Theorem 5.16, respectively, using suitably modified definition of Bass–Serre tree, and Bass–Serre forest, for these contexts.

**Theorem 5.18.** *Let  $L$  be an HNN extension of  $M$  with base monoids  $A, B$ . Suppose that, furthermore,  $M$  is free as both a right  $A$ -set and a right  $B$ -set. If  $M$  is of type  $\text{left-F}_n$  and  $A$  is of type  $\text{left-F}_{n-1}$ , then  $L$  is of type  $\text{left-F}_n$ .*

**Theorem 5.19.** *Let  $L$  be an HNN extension of  $M$  with base monoids  $A, B$ . Suppose that, furthermore,  $M$  is free as both a right and a left  $A$ -set (via the inclusion) and as a right and a left  $B$ -set. If  $M$  is of type  $\text{left-F}_n$  and  $A$  is of type  $\text{bi-F}_{n-1}$ , then  $L$  is of type  $\text{bi-F}_n$ .*

Theorem 5.18 recovers the usual topological finiteness result for HNN extensions of groups. It also applies if  $M$  is left cancellative and  $A$  is a group. The analogue of Theorem 5.18 for left geometric dimensions states that if  $M$  is free as both a right  $A$ -set and a right  $B$ -set,  $M$  has left geometric dimension at most  $d$  and  $A$  has geometric dimension at most  $d - 1$ , then  $L$  has geometric dimension at most  $d$ . Theorem 5.19 applies if  $M$  is cancellative and  $A$  is a group.

Similarly, if  $M$  is free as both a right and a left  $A$ -set and as a right and a left  $B$ -set, then if  $M$  has geometric dimension at most  $d$  and  $A$  has geometric dimension at most  $d - 1$ , then  $L$  has geometric dimension at most  $d$ .

## REFERENCES

- [1] S. I. Adjan. Defining relations and algorithmic problems for groups and semigroups. *Trudy Mat. Inst. Steklov.*, 85:123, 1966.
- [2] S. I. Adjan and G. U. Oganessian. On the word and divisibility problems for semigroups with one relation. *Mat. Zametki*, 41(3):412–421, 458, 1987.
- [3] J. M. Alonso and S. M. Hermiller. Homological finite derivation type. *Internat. J. Algebra Comput.*, 13(3):341–359, 2003.
- [4] D. J. Anick. On the homology of associative algebras. *Trans. Amer. Math. Soc.*, 296(2):641–659, 1986.
- [5] G. M. Bergman. Coproducts and some universal ring constructions. *Trans. Amer. Math. Soc.*, 200:33–88, 1974.
- [6] R. Bieri, W. D. Neumann, and R. Strebel. A geometric invariant of discrete groups. *Invent. Math.*, 90(3):451–477, 1987.
- [7] R. V. Book and F. Otto. *String-rewriting systems*. Texts and Monographs in Computer Science. Springer-Verlag, New York, 1993.
- [8] M. Brittenham, S. W. Margolis, and J. Meakin. Subgroups of the free idempotent generated semigroups need not be free. *J. Algebra*, 321(10):3026–3042, 2009.
- [9] K. S. Brown. Complete Euler characteristics and fixed-point theory. *J. Pure Appl. Algebra*, 24(2):103–121, 1982.
- [10] K. S. Brown. The geometry of rewriting systems: a proof of the Anick-Groves-Squier theorem. In *Algorithms and classification in combinatorial group theory (Berkeley, CA, 1989)*, volume 23 of *Math. Sci. Res. Inst. Publ.*, pages 137–163. Springer, New York, 1992.
- [11] K. S. Brown. *Cohomology of groups*, volume 87 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994. Corrected reprint of the 1982 original.
- [12] K. S. Brown. Lectures on the cohomology of groups. In *Cohomology of groups and algebraic K-theory*, volume 12 of *Adv. Lect. Math. (ALM)*, pages 131–166. Int. Press, Somerville, MA, 2010.
- [13] C. C.-a. Cheng and J. Shapiro. Cohomological dimension of an abelian monoid. *Proc. Amer. Math. Soc.*, 80(4):547–551, 1980.
- [14] W. H. Cockcroft. On two-dimensional aspherical complexes. *Proc. London Math. Soc. (3)*, 4:375–384, 1954.
- [15] D. E. Cohen. A monoid which is right  $FP_\infty$  but not left  $FP_1$ . *Bull. London Math. Soc.*, 24(4):340–342, 1992.
- [16] D. E. Cohen. String rewriting and homology of monoids. *Math. Structures Comput. Sci.*, 7(3):207–240, 1997.
- [17] R. Cremanns and F. Otto.  $FP_\infty$  is undecidable for finitely presented monoids with word problems decidable in polynomial time. *Mathematische Schriften Kassel 11/98, Universitat Kassel.*, 1998.
- [18] W. Dicks. *Groups, trees and projective modules*, volume 790 of *Lecture Notes in Mathematics*. Springer, Berlin, 1980.
- [19] W. Dicks and M. J. Dunwoody. *Groups acting on graphs*, volume 17 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1989.
- [20] T. Dieck. *Algebraic topology*. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2008.
- [21] W. G. Dwyer and H.-W. Henn. *Homotopy theoretic methods in group cohomology*. Advanced Courses in Mathematics. CRM Barcelona. Birkhäuser Verlag, Basel, 2001.
- [22] E. Dyer and A. T. Vasquez. Some small aspherical spaces. *J. Austral. Math. Soc.*, 16:332–352, 1973. Collection of articles dedicated to the memory of Hanna Neumann, III.
- [23] Z. Fiedorowicz. Classifying spaces of topological monoids and categories. *Amer. J. Math.*, 106(2):301–350, 1984.
- [24] R. Geoghegan. *Topological methods in group theory*, volume 243 of *Graduate Texts in Mathematics*. Springer, New York, 2008.
- [25] R. D. Gray and B. Steinberg. Topological finiteness properties of monoids I: Foundations. *Algebr. Geom. Topol.*, 22(7) 3083–3170, 2022.
- [26] R. D. Gray and B. Steinberg. A Lyndon’s identity theorem for one-relator monoids. *Sel. Math. New Ser.* 28, 59 (2022).
- [27] V. S. Guba and S. J. Pride. On the left and right cohomological dimension of monoids. *Bull. London Math. Soc.*, 30(4):391–396, 1998.
- [28] Y. Guiraud and P. Malbos. Higher-dimensional normalisation strategies for acyclicity. *Adv. Math.*, 231(3-4):2294–2351, 2012.
- [29] A. Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002.
- [30] G. Hochschild. On the cohomology groups of an associative algebra. *Ann. of Math. (2)*, 46:58–67, 1945.

- [31] D. F. Holt, B. Eick, and E. A. O'Brien. *Handbook of computational group theory*. Discrete Mathematics and its Applications (Boca Raton). Chapman & Hall/CRC, Boca Raton, FL, 2005.
- [32] J. M. Howie. Embedding theorems for semigroups. *Quart. J. Math. Oxford Ser. (2)*, 14:254–258, 1963.
- [33] Y. Kobayashi. Homotopy reduction systems for monoid presentations: asphericity and low-dimensional homology. *J. Pure Appl. Algebra*, 130(2):159–195, 1998.
- [34] Y. Kobayashi. Finite homotopy bases of one-relator monoids. *J. Algebra*, 229(2):547–569, 2000.
- [35] Y. Kobayashi. The homological finiteness properties left-, right-, and bi-FP<sub>n</sub> of monoids. *Comm. Algebra*, 38(11):3975–3986, 2010.
- [36] Y. Kobayashi and F. Otto. On homotopical and homological finiteness conditions for finitely presented monoids. *Internat. J. Algebra Comput.*, 11(3):391–403, 2001.
- [37] Y. Kobayashi and F. Otto. For finitely presented monoids the homological finiteness conditions FHT and bi-FP<sub>3</sub> coincide. *J. Algebra*, 264(2):327–341, 2003.
- [38] G. Lallement. On monoids presented by a single relation. *J. Algebra*, 32:370–388, 1974.
- [39] G. Lallement. Some algorithms for semigroups and monoids presented by a single relation. In *Semigroups, theory and applications (Oberwolfach, 1986)*, volume 1320 of *Lecture Notes in Math.*, pages 176–182. Springer, Berlin, 1988.
- [40] T. Lawson. A flatness result of fiedorwicz for amalgamated free products of monoids in connection with classifying spaces of monoids. *MathOverflow*.
- [41] R. C. Lyndon and M. P. Schützenberger. The equation  $a^M = b^N c^P$  in a free group. *Michigan Math. J.*, 9:289–298, 1962.
- [42] R. C. Lyndon. Cohomology theory of groups with a single defining relation. *Ann. of Math. (2)*, 52:650–665, 1950.
- [43] R. C. Lyndon and P. E. Schupp. *Combinatorial group theory*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1977 edition.
- [44] G. S. Makanin. On the identity problem in finitely defined semigroups. *Dokl. Akad. Nauk SSSR*, 171:285–287, 1966.
- [45] A. Malheiro. Complete rewriting systems for codified submonoids. *Internat. J. Algebra Comput.*, 15(2):207–216, 2005.
- [46] S. Margolis, J. Rhodes, and P. V. Silva. On the topology of a boolean representable simplicial complex. *Internat. J. Algebra Comput.*, 27(1):121–156, 2017.
- [47] S. Margolis, F. Saliola, and B. Steinberg. Combinatorial topology and the global dimension of algebras arising in combinatorics. *J. Eur. Math. Soc. (JEMS)*, 17(12):3037–3080, 2015.
- [48] J. P. May. *A concise course in algebraic topology*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1999.
- [49] D. McDuff. On the classifying spaces of discrete monoids. *Topology*, 18(4):313–320, 1979.
- [50] J. Meakin and N. Szakács. Inverse monoids and immersions of 2-complexes. *Internat. J. Algebra Comput.*, 25(1-2):301–323, 2015.
- [51] B. Mitchell. Rings with several objects. *Advances in Math.*, 8:1–161, 1972.
- [52] F. Otto and Y. Kobayashi. Properties of monoids that are presented by finite convergent string-rewriting systems—a survey. In *Advances in algorithms, languages, and complexity*, pages 225–266. Kluwer Acad. Publ., Dordrecht, 1997.
- [53] S. J. Pride. Homological finiteness conditions for groups, monoids, and algebras. *Comm. Algebra*, 34(10):3525–3536, 2006.
- [54] S. J. Pride and F. Otto. For rewriting systems the topological finiteness conditions FDT and FHT are not equivalent. *J. London Math. Soc. (2)*, 69(2):363–382, 2004.
- [55] S. J. Pride and F. Otto. On higher order homological finiteness of rewriting systems. *J. Pure Appl. Algebra*, 200(1-2):149–161, 2005.
- [56] M. V. Sapir. Algorithmic problems for amalgams of finite semigroups. *J. Algebra*, 229(2):514–531, 2000.
- [57] J.-P. Serre. *Cohomologie des groupes discrets*. Princeton Univ. Press, Princeton, N.J., 1971.
- [58] C. C. Squier. Word problems and a homological finiteness condition for monoids. *J. Pure Appl. Algebra*, 49(1-2):201–217, 1987.
- [59] J. B. Stephen. The automorphism group of the graph of an  $\mathcal{R}$  class. *Semigroup Forum*, 53(3):387–389, 1996.
- [60] V. Ufnarowski. Introduction to noncommutative Gröbner bases theory. In *Gröbner bases and applications (Linz, 1998)*, volume 251 of *London Math. Soc. Lecture Note Ser.*, pages 259–280. Cambridge Univ. Press, Cambridge, 1998.
- [61] C. T. C. Wall. Finiteness conditions for CW-complexes. *Ann. of Math. (2)*, 81:56–69, 1965.
- [62] X. Wang and S. J. Pride. Second order Dehn functions of groups and monoids. *Internat. J. Algebra Comput.*, 10(4):425–456, 2000.
- [63] L. Zhang. Applying rewriting methods to special monoids. *Math. Proc. Cambridge Philos. Soc.*, 112(3):495–505, 1992.