

Input-output equations and identifiability of linear ODE models

Alexey Ovchinnikov, Gleb Pogudin, and Peter Thompson

Abstract—Structural identifiability is a property of a differential model with parameters that allows for the parameters to be determined from the model equations in the absence of noise. The method of input-output equations is one method for verifying structural identifiability. This method stands out in its importance because the additional insights it provides can be used to analyze and improve models. However, its complete theoretical grounds and applicability are still to be established. A subtlety and key for this method to work correctly is knowing whether the coefficients of these equations are identifiable.

In this paper, to address this, we prove identifiability of the coefficients of input-output equations for types of differential models that often appear in practice, such as linear models with one output and linear compartment models in which, from each compartment, one can reach either a leak or an input. This shows that checking identifiability via input-output equations for these models is legitimate and, as we prove, that the field of identifiable functions is generated by the coefficients of the input-output equations. Finally, we exploit a connection between input-output equations and the transfer function matrix to show that, for a linear compartment model with an input and strongly connected graph, the field of all identifiable functions is generated by the coefficients of the transfer function matrix even if the initial conditions are generic.

Index Terms—identifiable functions, input-output equations, linear compartment models, structural parameter identifiability

I. INTRODUCTION

A. Background

Structural global identifiability (in what follows, we will say just “identifiability” for simplicity) is a property of a differential model with parameters that allows for the parameters to be uniquely determined from the model equations, noiseless data and sufficiently exciting inputs (also known as the persistence of excitation, see [1]–[3]). Performing identifiability analysis is an important first step in evaluating and, if needed, adjusting the model. There are different approaches

to assessing identifiability (see [4]–[6] for descriptions of methods). If structural identifiability is established, one can assess practical identifiability before doing reliable parameter identification [7], [8].

There is a relaxed version of identifiability, namely, local identifiability. It refers to the possibility of determining finitely many feasible parameter values. There are efficient algorithms [9], [10] for checking whether a given function of parameters is locally identifiable. To the best of our knowledge, there are no complete and efficient algorithms for finding all locally identifiable functions of parameters (see [11, page 7] for a partial algorithm), a key to efficient model reparametrization for improving the model.

1) *How the errors that we prevent occur in existing methods:* One of the approaches, which is widely used, is based on **input-output equations** [12]–[23], and has appeared in software packages such as COMBOS, DAISY, and their successors. An existing challenge is to understand the a priori applicability of the method, as the above software packages make incorrect identifiability conclusions for some models. We address this challenge in the present paper.

We will now discuss this in more detail. Roughly speaking, input-output equations are “minimal” equations that depend only on the input and output variables and parameters (see [24] for applications other than identifiability). We will describe a typical algorithm based on this approach using the following linear compartment model as a running example:

$$\begin{cases} x_1' = -(a_{01} + a_{21})x_1 + a_{12}x_2 + u, \\ x_2' = a_{21}x_1 - a_{12}x_2, \\ y = x_2. \end{cases} \quad (1)$$

In the above system,

- x_1 and x_2 are the state variables;
- y is the output observed in the experiment;
- u is the input (control) function to be chosen by the experimenter;
- a_{01}, a_{12}, a_{21} are unknown scalar parameters.

The question is whether the values of the parameters a_{01}, a_{12}, a_{21} can be determined from y and u . A typical algorithm operates as follows:

- (1) Find input-output equations, writing them as (differential) polynomials in the input and output variables. For (1), a calculation shows that the input-output equation is

$$y'' + (a_{01} + a_{12} + a_{21})y' + a_{01}a_{12}y - a_{21}u = 0. \quad (2)$$

- (2) Use the following **Assumption (A)**:

This work was partially supported by the NSF grants CCF-1563942, CCF-1564132, CCF-1319632, DMS-1760448, CCF-1708884, DMS-1853650, DMS-1853482; NSA grant #H98230-18-1-0016; and CUNY grants PSC-CUNY #69827-0047, #60098-00 48.

A. Ovchinnikov is with CUNY Queens College, Department of Mathematics, 65-30 Kissena Blvd, Queens, NY 11367, USA and CUNY Graduate Center, Ph.D. Programs in Mathematics and Computer Science, 365 Fifth Avenue, New York, NY 10016, USA (e-mail: aovchinnikov@qc.cuny.edu).

G. Pogudin was with Courant Institute of Mathematical Sciences, New York University, New York, NY 10012, USA and the Higher School of Economics, Faculty of Computer Science, Moscow, 109028, Russia. He is now with LIX, CNRS, École Polytechnique, Institut Polytechnique de Paris, 1 rue Honoré d'Estienne d'Orves, 91120, Palaiseau, France (e-mail: gleb.pogudin@polytechnique.edu).

P. Thompson was with CUNY Graduate Center, Ph.D. Program in Mathematics, 365 Fifth Avenue, New York, NY 10016, USA (e-mail: peterthompson-math@gmail.com, peter.thompson@liu.se).

a function of parameters is identifiable if and only if it can be expressed as a rational function of the coefficients of the input-output equations.

In our example, this amounts to assuming that a function of parameters is identifiable if and only if it can be expressed as a rational function of $a_{01} + a_{12} + a_{21}$, $a_{01}a_{12}$, and a_{21} .

One possible rationale behind this assumption is the “solvability” condition from [13, Remark 3]: due to the “minimality” of the input-output equations, one would expect that there exist N and $t_1, \dots, t_N \in \mathbb{R}$ such that the linear system

$$\begin{cases} y''(t_1) + c_1 y'(t_1) + c_2 y(t_1) + c_3 u(t_1) = 0 \\ \vdots \\ y''(t_N) + c_1 y'(t_N) + c_2 y(t_N) + c_3 u(t_N) = 0 \end{cases} \quad (3)$$

in c_1, c_2, c_3 has a unique solution in terms of $y(t_i), y'(t_i), y''(t_i), u(t_i)$, $1 \leq i \leq N$, so the coefficients of (2) are identifiable. However, the assumption is not always satisfied and, consequently, such N and $t_1, \dots, t_N$ might not exist at all. This is a reason, e.g., why DAISY may miss the non-identifiability of some of the parameters in those systems. An example is given in Section IV-A.1 (see also [5, Example 2.14] and [25, Sections 5.2 and 5.3]).

- (3) Set up a system of polynomial equations in the parameters setting the coefficients of (2) equal to new variables,

$$\begin{cases} a_{01} + a_{12} + a_{21} = c_1 \\ a_{01}a_{12} = c_2 \\ -a_{21} = c_3, \end{cases} \quad (4)$$

and verify if (4) as a system in the a ’s with coefficients in the field $\mathbb{C}(c_1, c_2, c_3)$ has a unique solution. This can be done, e.g., using Gröbner bases. Alternatively, for (4), one can see that $a_{21} = -c_3$ can be uniquely recovered, but the values of a_{01} and a_{12} are known only up to exchange due to the symmetry of (4) with respect to a_{01} and a_{12} .

2) *Importance of the IO-equation method: finds all identifiable combinations and helps with reparametrization:* Even though there are complete algorithms (that is, not relying on any assumption like [Assumption \(A\)](#) above) for assessing structural identifiability (see, e.g., [26]), establishing when the input-output equation method is valid is important because:

- This method can produce *all identifiable functions* (also referred to as “true parameters” in [24, Remark 2]), not just assess identifiability of specific parameters. More precisely, [27, Corollary 5.8] shows that the field generated by the coefficients of the input-output equations contains all of the identifiable functions.

In example (1), the field of identifiable functions is generated by the coefficients of (2), so it is equal to

$$\mathbb{C}(a_{01} + a_{12} + a_{21}, a_{01}a_{12}, a_{21}) = \mathbb{C}(a_{01} + a_{12}, a_{01}a_{12}, a_{21}).$$

Generators of the field of identifiable functions can be used to reparametrize the model [12], [28], [29].

- This method can be used for proving general theorems about classes of models [14], [15].

- For a large class of linear compartment models, there are efficient methods for computing their input-output equations [14], [15], [22].

B. The problem

As was described above, the approach to assessing identifiability via input-output equations has been used much in the last three decades and has its own distinctive features. However, it heavily relies on [Assumption \(A\)](#), which is not always true (see [5, Example 2.14] and [25, Section 5.2]). It can be verified by an algorithm [30, Section 4.1] and [31, Section 3.4] but is not verified in any implementation we have seen (including [17], [20]). **The general problem** studied in this paper is:

to determine classes of ODE models that satisfy [Assumption \(A\)](#) a priori; consequently, the approach via input-output equations gives correct result for these models.

Discrepancy between different notions of identifiability is not unusual given the wide range of experimental setups and mathematical tools involved. We refer the reader to a recent review [32] (see also [3]) presenting a number of notions of identifiability together with some known (in)equivalences between them. Our work clarifies this big picture by giving explicit and easy to check (unlike [27]) conditions for equivalence of different ways to assess identifiability.

C. Our results

The first part of our results shows that [Assumption \(A\)](#) is a priori satisfied for the following classes of models often appearing in practice [7], [20], [33]–[38]:

- linear models with one output (Main Result 1);
- linear compartment models such that, from every vertex of the graph of the model, at least one leak or input is reachable (Main Result 2).

Checking whether the model is of one of these types can be done just by visual inspection. For instance, as we will see in Example 1, each of these theorems is applicable to model (1). Main Result 1 cannot be strengthened to more than one output if all linear models are allowed, see Section IV-A.1 and, for non-linear systems, see [39, Lemma 5.1].

The second part is devoted to relaxing the “minimality” condition on the input-output equations. For linear compartment models, elegant relations involving only parameters, inputs, and outputs were proposed in [14, Theorem 2] based on Cramer’s rule (see also [15, Proposition 2.3]). In general, using these equations instead of the “minimal” relations in the algorithm above would give incorrect results [15, Remark 3.11].

However, in Main Result 3, we show that, for linear compartment models with an input and whose graph is strongly connected, one can use these equations as the input-output equations and obtain the full field of identifiable functions.

Furthermore, we apply Main Result 3 to the transfer function method [40, page 444]. It is known that, in case of multiple outputs, using only the coefficients of the transfer function matrix (as opposed to the full output transforms) may

lead to incorrect identifiability conclusions [40, Example 10.6]. As a corollary of our results (Corollary 3), we show that this is not the case for such linear compartment models.

We state the consequences of our results for algorithms for computing identifiable functions in Section II-C and illustrate the conditions in our main results in Section IV.

D. Structure of the paper

Basic notions and notation from differential algebra, identifiability, and linear compartment models are given in Section II. The main results in a brief form are stated in Section III and then stated and proved in Section V. In Section IV, we illustrate our main results with examples, e.g., showing how existing identifiability approaches could fail. The appendix has results we use relating the notions used in the paper for linear models to the corresponding notions for nonlinear systems.

II. PRELIMINARIES

In this section, we recall the notation/notions found in the literature and introduce our own notation/notions to state our main results in Section III. All fields have characteristic zero.

A. Identifiability of linear models

Fix positive integers λ , n , m , and κ for the remainder of the paper. Let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_\lambda)$, $\mathbf{x} = (x_1, \dots, x_n)$, $\mathbf{y} = (y_1, \dots, y_m)$, and $\mathbf{u} = (u_1, \dots, u_\kappa)$. Consider a system of ODEs

$$\Sigma = \begin{cases} \mathbf{x}' = \mathbf{f}(\mathbf{x}, \boldsymbol{\mu}, \mathbf{u}), \\ \mathbf{y} = \mathbf{g}(\mathbf{x}, \boldsymbol{\mu}, \mathbf{u}), \\ \mathbf{x}(0) = \mathbf{x}^*, \end{cases} \quad (5)$$

where $\mathbf{f} = (f_1, \dots, f_n)$ and $\mathbf{g} = (g_1, \dots, g_m)$ are tuples of polynomials in $\mathbf{x}, \mathbf{u}$ over $\mathbb{C}(\boldsymbol{\mu})$ of degree at most one.

For a rational function $h(\boldsymbol{\mu}) \in \mathbb{C}(\boldsymbol{\mu})$, we will define two notions of identifiability: *identifiability* and *IO-identifiability*. The former is meaningful from the modeling standpoint; the latter is what the algorithm outlined in the introduction checks.

1) *Identifiability*: We fix notation to give rigorous definitions:

Notation 1 (Auxiliary analytic notation):

- (a) Let $\mathbb{C}^\infty(0)$ denote the set of all functions that are complex analytic in some neighborhood of $t = 0$.
- (b) Let $\Omega \subset \mathbb{C}^\lambda$ be the complement to the set where at least one of the denominators of the coefficients of (5) in $\mathbb{C}(\boldsymbol{\mu})$ vanishes.
- (c) For every $h \in \mathbb{C}(\boldsymbol{\mu})$, we set

$$\Omega_h := \mathbb{C}^n \times \{\hat{\boldsymbol{\mu}} \in \Omega \mid h(\hat{\boldsymbol{\mu}}) \text{ well-defined}\} \times (\mathbb{C}^\infty(0))^\kappa.$$

- (d) For $(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}})$ such that $\hat{\boldsymbol{\mu}} \in \Omega$, let $X(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}})$ and $Y(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}})$ denote the unique solution over $\mathbb{C}^\infty(0)$ of the instance of Σ with $\mathbf{x}^* = \hat{\mathbf{x}}^*$, $\boldsymbol{\mu} = \hat{\boldsymbol{\mu}}$, and $\mathbf{u} = \hat{\mathbf{u}}$ (see [41, Theorem 2.2.2]).
- (e) For any positive integer s , a subset $U \subset \mathbb{C}^s$ is called *Zariski open* if there exists a polynomial P on $\mathbb{C}^s$ such that U is the complement to the zero set of P .
- (f) For any positive integer s , a subset $U \subset (\mathbb{C}^\infty(0))^s$ is called *Zariski open* if there exists a polynomial P in $z_1, \dots, z_s$ and their derivatives such that

$$U = \{\hat{\mathbf{z}} \in (\mathbb{C}^\infty(0))^s \mid P(\hat{\mathbf{z}})|_{t=0} \neq 0\}.$$

- (g) For any positive integer s and $X = \mathbb{C}^s$ or $(\mathbb{C}^\infty(0))^s$, the set of all nonempty Zariski open subsets of X will be denoted by $\tau(X)$.

Definition 1 (Identifiability, see [5, Definition 2.5]): We say that $h(\boldsymbol{\mu}) \in \mathbb{C}(\boldsymbol{\mu})$ is *identifiable* if

$$\exists \Theta \in \tau(\mathbb{C}^n \times \mathbb{C}^\lambda) \exists U \in \tau((\mathbb{C}^\infty(0))^\kappa)$$

$$\forall (\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}}) \in (\Theta \times U) \cap \Omega_h \quad |S_h(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}})| = 1,$$

$$\text{where } S_h(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}}) := \{h(\tilde{\boldsymbol{\mu}}) \mid \exists (\tilde{\mathbf{x}}^*, \tilde{\boldsymbol{\mu}}, \tilde{\mathbf{u}}) \in \Omega_h$$

$$\text{such that } Y(\tilde{\mathbf{x}}^*, \tilde{\boldsymbol{\mu}}, \tilde{\mathbf{u}}) = Y(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}})\}.$$

The field $\{h \in \mathbb{C}(\boldsymbol{\mu}) \mid h \text{ is identifiable}\}$ will be called *the field of identifiable functions*.

2) *Input-output identifiability*: The notion of IO-identifiability can be defined for systems with rational right-hand side (see Section A from the Appendix). Here we give a specialization of the general definition to the linear case (the equivalence of Definition 3 and Definition 7 restricted to the linear case is established in Proposition 1). For this, we will first recall several standard notions from differential algebra:

Notation 2 (Differential rings and ideals):

- (a) A *differential ring* (R, δ) is a commutative ring with a derivation $\delta: R \rightarrow R$, that is, a map such that, for all $a, b \in R$, $(a+b)' = a' + b'$ and $(ab)' = a'b + ab'$.
- (b) The *ring of differential polynomials* in the variables $x_1, \dots, x_n$ over a field K is the ring $K[x_j^{(i)} \mid i \geq 0, 1 \leq j \leq n]$ with a derivation defined on the ring by $(x_j^{(i)})' := x_j^{(i+1)}$. This differential ring is denoted by $K\{x_1, \dots, x_n\}$.
- (c) For a differential polynomial $P \in K\{x_1, \dots, x_n\}$ and $1 \leq i \leq n$, the *order of P with respect to x_i* is the order of the highest derivative of x_i appearing in P ($-\infty$ if x_i does not appear in P). It is denoted by $\text{ord}_{x_i} P$.
- (d) An ideal I of a differential ring (R, δ) is called a *differential ideal* if, for all $a \in I$, $\delta(a) \in I$. For $F \subset R$, the smallest differential ideal containing set F is denoted by $[F]$.
- (e) For Σ as in (5), let $I_\Sigma = [\mathbf{x}' - \mathbf{f}, \mathbf{y} - \mathbf{g}] \subset \mathbb{C}(\boldsymbol{\mu})\{\mathbf{x}, \mathbf{y}, \mathbf{u}\}$ be the differential ideal of Σ . Informally, I_Σ is the ideal of all relations among components of a generic solution of Σ .

Definition 2 (a full set of input-output equations): For

Σ as in (5), a tuple $(p_1, \dots, p_m)$ of differential polynomials from $\mathbb{C}(\boldsymbol{\mu})\{\mathbf{y}, \mathbf{u}\}$ is called a *full set of input-output equations* if there is an ordering of the output variables, which we will assume to be $y_1 < y_2 < \dots < y_m$ to simplify notation, such that

- (1) p_1 is the linear differential polynomial in y_1 and $\mathbf{u}$ in I_Σ of the smallest possible order in y_1 such that the coefficient of the highest derivative of y_1 is one.
- (2) For every $\ell > 1$, p_ℓ is the linear differential polynomial in $y_1, \dots, y_\ell$ and $\mathbf{u}$ in I_Σ such that
 - $\text{ord}_{y_j} p_\ell < \text{ord}_{y_j} p_j$ for every $1 \leq j < \ell$;
 - the coefficient of the highest derivative of y_ℓ in p_ℓ is 1;
 - $\text{ord}_{y_\ell} p_\ell$ is the smallest possible.

Definition 3 (IO-identifiable function): For a system Σ , consider a full set E of input-output equations. Then the subfield k of $\mathbb{C}(\boldsymbol{\mu})$ generated by the coefficients of E over $\mathbb{C}$ is called *the field of input-output identifiable (IO-identifiable) functions*. We call $h \in \mathbb{C}(\boldsymbol{\mu})$ *IO-identifiable* if $h \in k$.

Remark 1: Proposition 1 establishes the equivalence of this definition to Definition 7, which is applicable to a general

rational ODE systems. Proposition 1 also implies that the field of input-output identifiable functions does not depend on the choice of a full set of input-output equations.

For examples of input-output equations and IO-identifiable functions, see Section IV.

3) Comparison of identifiability and IO-identifiability:

Remark 2 (Meaning of IO-identifiability): One can see that the field of IO-identifiable functions is exactly what will be computed by the first two steps of the algorithm outlined in the introduction (see also Algorithm II.1). The **general problem** as stated in Section I-B can be restated as:

*Determine classes of ODE models for which
identifiable $\iff$ IO-identifiable.*

[27, Theorem 4.2] together with [5, Example 2.14] (see also Section IV-A.1 and [25, Sections 5.2 and 5.3] with non-constant dynamics and outputs) imply that:

$$\boxed{\text{Identifiable}} \subsetneq \boxed{\text{IO-identifiable}}. \quad (6)$$

B. Linear compartment models

In this section, we discuss linear compartment models [42]. Such a model consists of a set of compartments in which material is transferred from some compartments to other compartments. We also allow for leakage of material from some compartments out of the system, and for input of material into some compartments from outside the system.

We use the notation of [14, Section 2] but our construction will be slightly more general (allowing scaling for inputs and outputs). Let G be a simple directed graph with n vertices V and edges E . Let In, Out, and Leak be subsets of V . The coefficients of material transfer are

$$\{a_{ji} \mid j \leftarrow i \in E\} \quad \text{and} \quad \{a_{0i} \mid i \in \text{Leak}\},$$

and there may be some additional parameters, we will denote all the parameters by μ as before. For $i = 1, \dots, n$, let x_i be the quantity of material in compartment i . If $i \in \text{In}$, let $b_i(\mu)u_i$ be the rate at which the experimenter inputs material into the i -th compartment, where $b_i \in \mathbb{C}(\mu) \setminus \{0\}$. If $i \in \text{Out}$, let $y_i = c_i(\mu)x_i$, where $c_i \in \mathbb{C}(\mu) \setminus \{0\}$. Without loss of generality, we assume

$$\text{Out} = \{1, \dots, m\}.$$

Now the system of equations governing the dynamics of $x_1, \dots, x_n$ is given by

$$\Sigma = \begin{cases} \mathbf{x}' = A(G)\mathbf{x} + \mathbf{u}, \\ y_i = c_i(\mu)x_i, \quad \text{for every } i \in \text{Out}, \end{cases} \quad (7)$$

where $\mathbf{x} = (x_1, \dots, x_n)^T$, $\mathbf{u}$ is the $n \times 1$ matrix whose i -th entry is $b_i(\mu)u_i$ if $i \in \text{In}$ and 0 otherwise, and $A(G)$ is the matrix (generalizing the Laplacian of the graph) defined by

$$A(G)_{ij} = \begin{cases} -a_{0i} - \sum_{k:i \rightarrow k \in E} a_{ki}, & i = j, i \in \text{Leak} \\ -\sum_{k:i \rightarrow k \in E} a_{ki}, & i = j, i \notin \text{Leak} \\ a_{ij}, & j \rightarrow i \in E \\ 0, & \text{otherwise.} \end{cases} \quad (8)$$

In the notation of (19), we have

$$\mathbf{x} = \{x_1, \dots, x_n\}, \quad \mathbf{y} = \{y_1, \dots, y_m\}, \quad \mathbf{u} = \{u_i \mid i \in \text{In}\}.$$

Definition 4: A system Σ is called a *linear compartment model* if there exists a simple directed graph G with edges E and vertices V , subsets In, Out, and Leak of V , and functions $b_i, c_j \in \mathbb{C}(\mu) \setminus \{0\}$ such that Σ has the form of (7).


It was observed in [14, Theorem 2] that, for a linear compartment model, one can obtain relations among inputs, outputs, and parameters as follows. Let ∂ be the differentiation operator. Let $M_{ji}(G)$ be the submatrix of $\partial I - A(G)$ obtained by deleting the j -th row and i -th column. Then [14, Theorem 2] yields that system (7) implies that for every $i \in \text{Out}$,

$$\det(\partial I - A)(y_i) - \frac{1}{c_i(\mu)} \sum_{j \in \text{In}} (-1)^{i+j} \det(M_{ji}(G)) (b_j(\mu)u_j) = 0. \quad (9)$$

[15, Theorem 3.8] gives a refined version of (9) coinciding with (9) for the cases we consider in our main results.

Definition 5 (Reachability): We say *vertex v is reachable from vertex w* or *one can reach vertex v from vertex w* if there exists a directed path from w to v . For example, in the graph $1 \rightarrow 2$, vertex 2 is reachable from vertex 1. We say a leak (resp. input) is reachable from w if there exists a vertex v in Leak (resp. In) such that v is reachable from w .

Example 1: Consider the graph



Here G is the graph given by

$$V = \{1, 2\} \quad \text{and} \quad E = \{1 \rightarrow 2, 2 \rightarrow 1\}.$$

The arrow leaving compartment 1 indicates that $\text{Leak} = \{1\}$, the arrow entering compartment 1 indicates that $\text{In} = \{1\}$, and the other decoration to compartment 2 indicates that $\text{Out} = \{2\}$. Note that the input and leak arrows, as well as the output decoration, are not considered part of the graph. One can see that the corresponding system of differential equations coincides with (1) and can be written as

$$\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}' = \begin{pmatrix} -(a_{01} + a_{21}) & a_{12} \\ a_{21} & -a_{12} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} u \\ 0 \end{pmatrix}, \quad y = x_2.$$

One can see that this system satisfies the conditions of Theorems 1, 2, and 3. A direct computation shows that the input-output equation (2) is a special case of (9).

C. Existing algorithms used in practice yet to be justified

In this section, we will present and justify (rephrasing our Main Results 1, 2, and 3) the correctness of two versions (Algorithms II.1 and II.2) of the algorithm outlines in Section I-A that were not previously fully justified. Algorithm II.1 is one of the key components of, e.g., DAISY [17], and Algorithm II.2 summarizes the approach from [15, Definition 3.9]. Our justifications are based on the assumptions stated

in Corollaries 1 and 2. Omitting some of the assumptions could lead to incorrect conclusions, as we show in Section IV-A.

Algorithm II.1 Computing identifiable functions

Input System Σ as in (19)

Output Generators of the field of identifiable functions of Σ (see Corollary 1)

- (Step 1) Compute a full set C of input-output equations of Σ .
 (Step 2) Return the coefficients of C considered as differential polynomials in $\mathbf{y}$ and $\mathbf{u}$.
-

Corollary 1: Assume that Σ satisfies one of the following:

- (1) Σ is as in (5) and has exactly one output;
- (2) Σ is a linear compartment model such that one can reach a leak or an input from every vertex.

Then Algorithm II.1 will produce a correct result for Σ .

Proof: Algorithm II.1 will compute generators of the field of IO-identifiable functions. Main Results 1 and 2 imply that, for Σ that we consider, the field of IO-identifiable functions coincides with the field of identifiable functions. ■

Algorithm II.2 Computing identifiable functions

Input System Σ as in (19) corresponding to a linear compartment model with graph G

Output Generators of the field of identifiable functions of Σ (see Corollary 2)

- (Step 1) For every $i \in \text{Out}$, compute an input-output equation p_i as in (9) (or a refined version from [15, Theorem 3.8]).
 (Step 2) Return the coefficients of $\{p_i \mid i \in \text{Out}\}$ considered as differential polynomials in $\mathbf{y}$ and $\mathbf{u}$.
-

Corollary 2: In the notation of Algorithm II.2, if graph G is strongly connected and has at least one input, then Algorithm II.2 will produce a correct result.

Proof: Follows from Main Result 3. ■

III. MAIN RESULTS

In this section, we will state our main results in a condensed form. For the detailed statements, see the corresponding theorems in Section V. In Section II-C, we show how our main results apply to justifying an algorithm computing all identifiable functions of an ODE model. In Section IV, we present examples (both of applied and of purely mathematical nature) illustrating the importance and use of the conditions in the statements of our main results. Note that, while the first result is restricted to MISO systems, the second and third are applicable to MIMO systems as well.

Main Result 1 (see Theorem 1): If system Σ as in (5) has exactly one output, then IO-identifiable functions coincide with identifiable functions.

Main Result 2 (see Theorem 2): If the graph of a linear compartment model is such that one can reach a leak or an input from every vertex, then IO-identifiable functions coincide with identifiable functions.

Problem 1: Will Main Result 2 remain true if the condition on the graph is removed or relaxed?

In other words, Main Results 1 and 2 provide classes of models for which the approach via input-output equations outlined in the introduction gives the correct result.

Main Result 3 (see Theorem 3): For a linear compartment model with at least one input and whose graph is strongly connected, the field of all identifiable functions is generated by the coefficients of equations (9).

This theorem combined with Lemma 5 yields:

Corollary 3: For a linear compartment model satisfying the assumptions of Main Result 3, the field of all identifiable functions is generated by the coefficients of the entries of the transfer function matrix (see Section C in Appendix).

IV. EXAMPLES

A. How identifiability methods could make mistakes

In this section, we will consider several examples to illustrate how the methods based on IO-equations, formula (17), and transfer functions may lead to incorrect conclusions about identifiability. This is to make the reader more aware of the conditions in our main results.

1) *Failure to detect non-identifiability with multiple outputs using IO-equations:* We will discuss a simple example of a linear system such that the classical method of IO-equations will not be able to decide on the (non-)identifiability of 2 of the 3 parameters. The example will also show that the condition of having only one output cannot be removed from our Main Result 1.

We begin with an ODE for radioactive decay $x' = -ax$, with a being an unknown decay rate. Suppose now that we have an unknown constant inflow b , and so $x' = -ax + b$. Consider the following output (e.g., the radiation level): $y = cx$, in which the unknown parameter c represents the properties of the medium between the observer and the radioactive species.

Suppose now that there is a known fixed outflow w (e.g., through a hole of fixed size), and so the ODE model becomes

$$\begin{cases} x' = -ax + b - w \\ w' = 0 \\ y_1 = cx, \quad y_2 = w \end{cases} \quad (10)$$

We then have

$$\begin{aligned} y_1' &= cx' = -cax + cb - cw = -ay_1 + cb - cy_2 \\ y_2' &= w' = 0, \end{aligned} \quad (11)$$

which can be shown to be a full set of input-output equations. To check the solvability condition, consider system (3):

$$\begin{cases} y_1'(t_1) = -ay_1(t_1) + cb - cy_2(t_1) \\ y_1'(t_2) = -ay_1(t_2) + cb - cy_2(t_2) \\ y_1'(t_3) = -ay_1(t_3) + cb - cy_2(t_3), \end{cases} \quad (12)$$

which we consider as a linear system in a , cb , and c . The matrix of the system is

$$A := \begin{pmatrix} -y_1(t_1) & 1 & -y_2(t_1) \\ -y_1(t_2) & 1 & -y_2(t_2) \\ -y_1(t_3) & 1 & -y_2(t_3) \end{pmatrix}$$

Since y_2 is a constant, the second and third columns of the matrix are proportional. Therefore, system (12) has infinitely


many solutions for the corresponding coefficients of the input-output equations, cb and b . Hence, the matrix is rank-deficient and the solvability condition is not satisfied. Therefore, proceeding further with trying to check the identifiability of b and c based just on (12) could (and will for this example, as we will see) cause an incorrect conclusion as the validity of this method is currently guaranteed under the solvability condition.

For example, the software DAISY (which is based on input-output equations) applied to this model concludes that all of a , b , and c are globally identifiable. However, neither b , nor c is even locally identifiable. This can be seen, e.g., by noticing that the following is an output-preserving transformation of system (10) for all non-zero k :

$$x \rightarrow kx, \quad c \rightarrow \frac{c}{k}, \quad b \rightarrow kb + w - kw.$$

Therefore, **Assumption (A)** is not satisfied. For an analogous example without constant states, see [5, Remark 2.15].

2) *The transfer function method and generic initial conditions* (see also [40, Example 10.6]): Consider the linear compartment model



in which an input function u is applied to compartment 1, the quantity in compartment 1 is measured, and material flows from compartment 1 to compartment 3 and from compartment 2 to compartment 3. The corresponding system is

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}' = \begin{pmatrix} -a_{31} & 0 & 0 \\ 0 & -a_{32} & 0 \\ a_{31} & a_{32} & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} + \begin{pmatrix} u \\ 0 \\ 0 \end{pmatrix} \quad (13)$$

$$y_1 = x_1.$$

Since the system satisfies the hypothesis of Theorem 1, we can find the field of identifiable functions, $\mathbb{C}(a_{31})$, by looking at the input-output equation:

$$y_1' + a_{31}y_1 - u = 0.$$

The transfer function (see Section C in Appendix) for the system is $\frac{1}{s+a_{31}}$, so the transfer function method gives the same correct result although the initial conditions are not zero but generic and the assumptions of Theorem 3 are not satisfied. However, using transfer function will lead to erroneous results for this model if we move the output to compartment 3 (that is, replace $y_1 = x_1$ with $y_1 = x_3$). In this case, the transfer function is $\frac{a_{31}}{s(s+a_{31})}$, indicating that a_{32} is not identifiable. However, it actually is identifiable. This can be shown again by using Algorithm 1 (as the assumption of Theorem 1 is still satisfied), that is, considering the following input-output equation:

$$y_1''' + (a_{31} + a_{32})y_1'' + a_{31}a_{32}y_1' - a_{31}a_{32}u - a_{31}u'.$$

Thus, the hypotheses of Corollary 3 cannot be omitted. Note that Algorithm II.2 will give a correct result for this case even though the assumptions of Theorem 3 are not satisfied.

B. Positive examples for applying our theory

Below we give examples from the literature satisfying at least one of the sufficient conditions from our main results.

1) *Kinetics of lead in humans and our results for one output.*: The following system of equations is used in [42, Section 4A] to model the kinetics of lead in the human body:

$$\begin{cases} x_1' = k_1x_1 + k_2x_2 + k_3x_3 + k_4 \\ x_2' = k_5x_1 + k_6x_2 \\ x_3' = k_7x_1 - k_3x_3 \\ y_1 = x_1 \end{cases}$$

A full set of input-output equations is unique in this case and consists of a single differential polynomial:

$$y_1''' - (k_1 + k_3 + k_6)y_1'' + (-k_1k_3 + k_1k_6 - k_2k_5 - k_3k_6 - k_3k_7)y_1' + (k_1k_3k_6 - k_2k_3k_5 + k_3k_6k_7)y_1 + k_3k_4k_6.$$

By Corollary 1 (condition (1)), the field of identifiable functions is generated by

$$k_1 + k_3 + k_6, \quad -k_1k_3 + k_1k_6 - k_2k_5 - k_3k_6 - k_3k_7, \\ k_3(k_1k_6 - k_2k_5 + k_6k_7), \quad k_3k_4k_6.$$

In other words, these parameter combinations are identifiable, and moreover any other identifiable combination of parameters can be written as a rational combination of these.

2) *Hepatobiliary kinetics of bromosulfophthalein*: The following linear compartment model is taken from [43, Section 6.3]:

$$\begin{cases} x_1' = -k_{31}x_1 + k_{13}x_3 + u \\ x_2' = -k_{42}x_2 + k_{24}x_4 \\ x_3' = k_{31}x_1 - (k_{03} + k_{13} + k_{43})x_3 \\ x_4' = k_{42}x_2 + k_{43}x_3 - (k_{04} + k_{24})x_4 \\ y_1 = x_1, \\ y_2 = x_2. \end{cases}$$

A full set of IO-equations is too large to display here but their coefficients are

$$k_{13}, \quad k_{31}, \quad k_{04}k_{42}, \quad k_{24}k_{43}, \quad k_{03} + k_{43}, \quad k_{04} + k_{24} + k_{42}. \quad (14)$$

Hence, by Corollary 1, the field of identifiable functions is generated by (14). This refines the analysis performed in [44, Example 3], where it was shown that (14) generate the field of IO-identifiable functions (as we have seen, for some examples, there are IO-identifiable functions that are not identifiable).

3) *Cyclic model*: The following model can be obtained from [35, Model M] by adding extra leaks and an output for a better illustration of the computation in connection to our results:

$$\begin{cases} x_1' = a_{13}x_3 - a_{21}x_1 - a_{01}x_1 \\ x_2' = a_{21}x_1 - a_{32}x_2 - a_{02}x_2 + u \\ x_3' = a_{32}x_2 - a_{13}x_3 \\ y_1 = x_1, \quad y_2 = x_2 \end{cases}$$

Using the coefficients of a full set of IO-equations or of the transfer function matrix, we obtain by Corollaries 1 and 3 that the field of identifiable functions is generated by:

$$a_{21}, \quad (a_{01} + a_{21})a_{13}, \quad a_{01} + a_{13}, \quad a_{13}a_{32}, \quad a_{02} + a_{32}.$$

V. PROOFS

A. “Identifiability $\iff$ IO-identifiability” for linear systems with one output (proof of Theorem 1)

In this section, we prove one of the main results, Theorem 1, which shows that, for a linear system with one output, IO-identifiability and identifiability are equivalent. We begin with showing a preliminary result.

Lemma 1: Let K be a field. Consider

- the differential polynomial ring $K\{y, \mathbf{u}\}$ with derivation ∂ satisfying $\partial(K) = 0$,
- $P \in K\{y, \mathbf{u}\}$ of the form $P = D_P(y) + U_P$, where $D_P \in K[\partial]$ is a linear differential operator over K with leading coefficient 1 and $U_P \in K\{\mathbf{u}\}$.

Let W be the Wronskian of all the monomials of P except for the one of the highest order with respect to y . Then $W \notin [P]$.

Proof: Since the coefficients of P and W are in K , the membership $W \in [P]$ would be the same considered over K or its algebraic closure. Hence, replacing K with its algebraic closure if necessary, we assume that K is algebraically closed.

Consider a lexicographic monomial ordering induced by an ordering of the variables such that $y^{(i+1)} > y^{(i)}$ for every $i \geq 0$ and y is greater than any derivative of $\mathbf{u}$. Since for all r $P, P', \dots, P^{(r)}$ is a Gröbner basis for

$$[P] \cap K[y, y', \dots, y^{(r)}, \mathbf{u}, \mathbf{u}', \dots, \mathbf{u}^{(r)}],$$

it follows from [45, Lemma 1.5] that $P, P', \dots$ form a Gröbner basis of $[P]$ with respect to this ordering as defined by [45, Definition 1.4].

Since the leading terms of a Gröbner basis are linear, $[P]$ is a prime ideal. Thus, we can introduce $L := \text{Frac}(K\{y, \mathbf{u}\}/[P])$. Denote the field of constants of L by $C(L)$ and the images of y and $\mathbf{u}$ in L by $\bar{y}$ and $\bar{\mathbf{u}}$, respectively. Since none of derivatives of $\mathbf{u}$ appear in the leading terms of the Gröbner basis, $\bar{\mathbf{u}}$ and their derivatives are algebraically independent over K .

Assume that the statement of the lemma is not true. Due to [46, Theorem 3.7, p. 21], this implies that the images in L of the monomials of P except for the one of the highest order in y are linearly dependent over $C(L)$. Therefore, there exists a nonzero polynomial

$$Q = D_Q(y) + U_Q,$$

where $D_Q \in C(L)[\partial]$ is monic and $U_Q \in C(L)\{\mathbf{u}\}$, such that $Q(\bar{y}, \bar{\mathbf{u}}) = 0$ and $\text{ord} D_Q < \text{ord} D_P$. Let D_0 be the gcd of D_P and D_Q with the leading coefficient 1. Then $\text{ord} D_0 < \text{ord} D_P$.

If F is an algebraically closed field, $p \in F[X]$, and p is divisible by a $q \in E[X]$ with the leading coefficient 1, where E is an extension of F , then $q \in F[X]$. Hence, as D_0 divides D_P and K is algebraically closed, $D_0 \in K[\partial]$ and there is $D_1 \in K[\partial]$ such that $D_P = D_1 D_0$. There also are $A, B \in C(L)[\partial]$ such that

$$D_0 = AD_P + BD_Q.$$

Consider $R := A(P) + B(Q) = D_0(y) + U_R$, where $U_R = A(U_P) + B(U_Q)$. Then $R(\bar{y}, \bar{\mathbf{u}}) = 0$. Since $P - D_1(R) \in C(L)\{\mathbf{u}\}$ vanishes on $\bar{\mathbf{u}}$ and $\bar{\mathbf{u}}$ is differentially independent over $C(L)$, it follows that $P = D_1(R)$.

Considering a basis of $C(L)$ over K , we can write

$$U_R = U_0 + e_1 U_1 + \dots + e_N U_N,$$

where $U_0, \dots, U_N \in K\{\mathbf{u}\}$ and $1, e_1, e_2, \dots, e_N \in C(L)$ are linearly independent over K . Since $D_1(U_R) = U_P$ and $D_1 \in K[\partial]$, $U_1, \dots, U_N \in \ker D_1$, where we consider D_1 as a function from $C(L)\{y, \mathbf{u}\}$ to $C(L)\{y, \mathbf{u}\}$. There are two cases:

- D_1 is not divisible by ∂ . Then $\ker D_1 = \{0\}$. Hence,

$$U_1 = \dots = U_N = 0.$$

- D_1 is divisible by ∂ . Then $\ker D_1 = C(L)$. Thus, $U_1, \dots, U_N \in K$. However, since $U_P = D_1(U_R)$, U_P does not contain a term in K . Hence, U_Q does not contain a term in $C(L)$ and, consequently, U_R does not contain a term in $C(L)$. Thus,

$$U_1 = \dots = U_N = 0.$$

In both cases, we have shown that $U_R \in K\{\mathbf{u}\}$. Thus, $R \in K\{y, \mathbf{u}\}$ and $R \in [P]$. But this is impossible because $P, P', P'', \dots$ is a Gröbner basis of $[P]$ with respect to the monomial ordering introduced in the beginning of the proof, and $\text{ord} D_0 < \text{ord} D_P$, so R is not reducible with respect to this basis. ■

Theorem 1 (Main Result 1): For every Σ as in (5) with $m = 1$ (that is, single output), for all $h \in \mathbb{C}(\mu)$,

$$h \text{ is identifiable} \iff h \text{ is IO-identifiable}.$$

Proof: [27, Theorem 4.2] implies that identifiable functions are always IO-identifiable, so it remains to show the reverse inclusion. Consider a full set of input-output equations for Σ . Since $m = 1$, it will consist of a single linear differential polynomial $p \in \mathbb{C}(\mu)\{y, \mathbf{u}\}$. Then, Lemma 1 and [27, Lemma 4.6] imply that its coefficients are identifiable, so the reverse inclusion holds as well. ■

B. Sufficient condition for “identifiability $\iff$ IO-identifiability” for linear compartment models (proof of Theorem 2)

For the notation, see Section II-B.

Lemma 2: Let $F = \text{Frac}(\mathbb{C}(\mu)\{\mathbf{x}, \mathbf{y}, \mathbf{u}\}/I_\Sigma)$. The field of constants of F lies in the subfield of F generated by $\mathbb{C}$, μ and $\mathbf{x}$.

Proof: Observe that F as a field is generated by μ , $\mathbf{x}$, and all the derivatives of $\mathbf{u}$, and all these elements are algebraically independent. Assume that there exists $\ell \geq 0$ and $h \in \mathbb{C}(\mu, \mathbf{x}, \mathbf{u}, \dots, \mathbf{u}^{(\ell)})$ such that $h' = 0$ and, without loss of generality, $\frac{\partial}{\partial u_{\kappa}^{(\ell)}} h \neq 0$. Then we have

$$h' = \sum_{i=0}^{\ell} \sum_{r=1}^{\kappa} u_r^{(i+1)} \frac{\partial}{\partial u_r^{(i)}} h + \sum_{j=1}^n x_j' \frac{\partial}{\partial x_j} h = u_{\kappa}^{(\ell+1)} \frac{\partial}{\partial u_{\kappa}^{(\ell)}} h + a,$$

$$a \in \mathbb{C}(\mu, \mathbf{x}, \mathbf{u}, \dots, \mathbf{u}^{(\ell)}, u_1^{(\ell+1)}, \dots, u_{\kappa-1}^{(\ell+1)}).$$


Now $h' = 0$ yields a contradiction since $u_{\kappa}^{(\ell+1)}$ is transcendental over $\mathbb{C}(\mu, \mathbf{x}, \mathbf{u}, \dots, \mathbf{u}^{(\ell)}, u_1^{(\ell+1)}, \dots, u_{\kappa-1}^{(\ell+1)})$ and $\frac{\partial}{\partial u_{\kappa}^{(\ell)}} h \neq 0$. ■

Lemma 3: Consider a graph G such that, from every vertex, at least one leak can be reached. Then the eigenvalues of $A(G)$ are distinct and algebraically independent over $\mathbb{Q}$.


Proof: Let H be a directed spanning forest of G constructed by a breadth-first search (depth-first search would work as well) with the set Leak as the source such that, from every vertex, there is a path to some element of Leak. Relabeling vertices if necessary, $A(H)$ is upper triangular with algebraically independent diagonal entries. It is well known

that a breadth-first search on a graph will construct a spanning forest containing all vertices reachable from the source set (cf. [47, Section 22.2]).

We illustrate our procedure with an example. Let G be the graph shown below, with $\text{Leak} = \{1, 6\}$:



The steps of a breadth-first search with source set $\{1, 6\}$ are the first three upper left, upper right, and lower left graphs shown below. The fourth lower right graph is a relabeling of the third as described above.



Taking H to be the fourth graph, we have

$$A(H) = \begin{pmatrix} -a_{01} & a_{12} & & & & \\ & -a_{12} & & & & \\ & & -a_{03} & a_{34} & a_{35} & \\ & & & -a_{34} & & \\ & & & & -a_{35} & a_{56} \\ & & & & & -a_{56} \end{pmatrix}.$$

Since the diagonal entries are algebraically independent over $\mathbb{Q}$ and algebraic over the field extension of $\mathbb{Q}$ generated by the coefficients of the characteristic polynomial of $A(H)$, it follows that the coefficients of the characteristic polynomial of $A(H)$ are algebraically independent over $\mathbb{Q}$.

For all i, j , if the coefficients of the characteristic polynomial of $A(G)|_{a_{i,j}=0}$ are algebraically independent, then the coefficients of the characteristic polynomial of $A(G)$ are algebraically independent. Since $A(H)$ can be obtained from $A(G)$ by setting equal to 0 those $a_{i,j}$ such that H has no edge from j to i , it follows that the coefficients of the characteristic polynomial of $A(G)$ are non-zero and algebraically independent. Since these n coefficients belong to the field extension of $\mathbb{Q}$ generated by n eigenvalues, the eigenvalues must be algebraically independent as well. ■

Theorem 2 (Main Result 2): Let Σ be a linear compartment model with graph G such that, from every vertex of G , at least one leak or input is reachable. Then the fields of identifiable and IO-identifiable functions coincide.

Proof: Let $K := \text{Frac}(\mathbb{C}(\boldsymbol{\mu})\{\mathbf{x}, \mathbf{y}, \mathbf{u}\}/I_\Sigma)$. We will show that Σ does not have a rational first integral, that is $C(K) = \mathbb{C}(\boldsymbol{\mu})$. Then the theorem will follow from [27, Theorem 4.7]. Consider a model Σ^* with a graph G^* obtained from G by replacing every input with a leak (if there was a vertex with an input and a leak, we simply remove the input). The theorem will follow from the following two claims.

Claim: If Σ has a rational first integral, then Σ^* also does. Consider a first integral of Σ , that is, an element of $C(K) \setminus \mathbb{C}(\boldsymbol{\mu})$. Lemma 2 implies that there exists $R \in \mathbb{C}(\boldsymbol{\mu}, \mathbf{x}) \setminus \mathbb{C}$ such that c is the image of R in K . Since

$$\mathbb{C}[\boldsymbol{\mu}, \mathbf{x}]\{\mathbf{u}\} \cap I_\Sigma = 0$$

due to [5, Lemma 3.1] and the image of R in K is a constant, the Lie derivative of R with respect to Σ ,

$$\mathcal{L}_\Sigma(R) := \sum_{i=1}^n \frac{\partial R}{\partial x_i} f_i,$$

where $f_1, \dots, f_n$ are as in (19), is zero. If there exists $i \in \text{In}$ such that x_i appears in R , then $\mathcal{L}_\Sigma(R)$ will be of the form

$$\mathcal{L}_\Sigma(R) = \frac{\partial R}{\partial x_i} b_i(\boldsymbol{\mu}) u_i + (\text{something not involving } u_i) \neq 0.$$

Thus, R does not involve any x_i with $i \in \text{In}$. Then, due to the construction of G^* , $\mathcal{L}_{\Sigma^*}(R) = \mathcal{L}_\Sigma(R) = 0$, so Σ^* also has a rational first integral.

Claim: Σ^* does not have rational first integrals. Lemma 3 implies that the eigenvalues of $A(G^*)$ are algebraically independent. Then [48, Theorem 10.1.2, p. 118] implies that Σ^* does not have rational first integrals. ■

C. Using more convenient IO-equations (proof of Theorem 3)

For the notation, see Section II-B.

Lemma 4: Let K be a field. For all $a, b, c \in K[x]$ such that $\gcd(a, b) = 1$, there exists at most one pair (p, q) of elements of $K[x]$ such that $ap + bq = c$ and $\deg p < \deg b$.

Proof: Suppose (p, q) and (p_1, q_1) are distinct pairs satisfying the two properties above. It follows that

$$a(p - p_1) + b(q - q_1) = 0. \quad (15)$$

Since $(p, q) \neq (p_1, q_1)$, (15) implies that $p \neq p_1$. Since

$$\deg(p - p_1) < \deg b,$$

(15) implies $\gcd(a, b) \neq 1$, contradicting our hypothesis. ■

Corollary 4: Let K be a field containing $\mathbb{C}$ and $a, b, c \in K[x]$ with $\gcd(a, b) = 1$. If there is a pair of polynomials (p, q) with

$$ap + bq = c \quad \text{and} \quad \deg p < \deg b,$$

then the coefficients of p and q belong to the field extension of $\mathbb{C}$ generated by the coefficients of a , b , and c .

Proof: Suppose some coefficient of p or q does not belong to the field generated by the coefficients of a , b , and c . By [49, Theorem 9.29, p. 117], there is a field automorphism σ of $\bar{K}$ that fixes the field extension of $\mathbb{C}$ generated by the coefficients of a , b , and c and moves this coefficient.

We extend σ to $\bar{K}[x]$ by $\sigma(x) = x$. Applying σ to both sides of $ap + bq = c$ gives us $a\sigma(p) + b\sigma(q) = c$. Using $\bar{K}$ for K in Lemma 4, we arrive at a contradiction. ■

Theorem 3 (Main Result 3): Let Σ be a linear compartment model with a graph G . Let $A = A(G)$ and M_{ji} be the submatrix of $\partial I - A$ obtained by deleting the j -th row and the i -th column of $\partial I - A$. Recall that (see (9)), for every solution of Σ , we have for every $i \in \text{Out}$,

$$\det(\partial I - A)(y_i) = \frac{1}{c_i(\boldsymbol{\mu})} \sum_{j \in \text{In}} (-1)^{i+j} \det(M_{ji})(b_j(\boldsymbol{\mu})u_j).$$

If G is strongly connected and has at least one input, then the coefficients of these differential polynomials with respect to y 's and u 's generate the field of identifiable functions of Σ .

Proof: Without loss of generality, assume $\text{Out} = \{1, \dots, m\}$. We set, for $i = 1, \dots, m$,

$$h_i := \det(\partial I - A)(y_i) - \frac{1}{c_i(\boldsymbol{\mu})} \sum_{j \in \text{In}} (-1)^{i+j} \det(M_{ji}) b_j(\boldsymbol{\mu}) u_j. \quad (16)$$

Let also $D = \det(\partial I - A)$ and, for $i = 1, \dots, m$, let Q_i be the $1 \times n$ matrix of operators defined by

$$\begin{cases} (Q_i)_j = \frac{(-1)^{i+j}}{c_i(\boldsymbol{\mu})} \det(M_{ji}) b_j(\boldsymbol{\mu}), & j \in \text{In}, \\ (Q_i)_j = 0, & j \notin \text{In}. \end{cases} \quad (17)$$

Observe that, for $i = 1, \dots, m$ $h_i = D(y_i) - Q_i \cdot \mathbf{u}$, where $\mathbf{u}$ is the $n \times 1$ matrix defined by $\mathbf{u}_j = u_j$ if $j \in \text{In}$ and $\mathbf{u}_j = 0$ otherwise.

First we show that the coefficients of $h_1, \dots, h_m$ are IO-identifiable. Fix i . Consider an ordering of the outputs such that y_i is the smallest one. Let $p_1, \dots, p_m$ be a full set of input-output equations with respect to this ordering (see Definition 2) which exists due to Proposition 1. Then p_1 is of the form

$$E(y_i) + B \cdot \mathbf{u},$$

where E is a linear differential operator and B is a $1 \times n$ matrix of linear differential operators, both with coefficients in $\mathbb{C}(\boldsymbol{\mu})$. Since $h_i \in I_\Sigma$ and h_i involves only y_i and $\mathbf{u}$, the second part of Proposition 1 implies that $h_i \in [p_1]$, so there exists a differential operator $D_0 \in \mathbb{C}(\boldsymbol{\mu})[\partial]$ such that $h_i = D_0 p_1$. Since G is strongly connected and has an input, by [15, Proposition 3.19],

$$\gcd(D \cup \{(Q_i)_j \mid (Q_i)_j \neq 0\}) = 1.$$

Thus D_0 has order zero, so h_i and p_1 are proportional. Therefore, the coefficients of (16) are IO-identifiable.

Next, we show that the field generated by the coefficients of $h_1, \dots, h_m$ contains the field of IO-identifiable functions. Fix an ordering on the outputs $y_m > \dots > y_1$. We will show that the full set $p_1, \dots, p_m$ of input-output equations with respect to this ordering satisfies:

$$\text{ord}_{y_1} p_1 = n, \quad \text{ord}_{y_i} p_i = 0 \text{ for every } 2 \leq i \leq m. \quad (18)$$

The fact that $\text{ord}_{y_1} p_1 = n$ is implied by the previous paragraph. From (5), we see that the transcendence degree of

$$\mathbb{C}(\boldsymbol{\mu})\{\mathbf{x}, \mathbf{y}, \mathbf{u}\} / I_\Sigma$$

over $\mathbb{C}(\boldsymbol{\mu})\{\mathbf{u}\}$ is equal to n , so the transcendence degree of

$$\mathbb{C}(\boldsymbol{\mu})\{\mathbf{y}, \mathbf{u}\} / (I_\Sigma \cap \mathbb{C}(\boldsymbol{\mu})\{\mathbf{y}, \mathbf{u}\})$$

over $\mathbb{C}(\boldsymbol{\mu})\{\mathbf{u}\}$ is less than or equal to n . From the form of p_1 , we have that $y_1, y_1', \dots, y_1^{(n-1)}$ are algebraically independent over $\mathbb{C}(\boldsymbol{\mu})\{\mathbf{u}\}$, so for $i = 2, \dots, m$, the elements $y_i, y_1, y_1', \dots, y_1^{(n-1)}$ must be algebraically dependent over $\mathbb{C}(\boldsymbol{\mu})\{\mathbf{u}\}$. Hence, the equation for y_i has order 0 in y_i . Thus,

$$p_i = D(y_i) - Q_i \cdot \mathbf{u}$$

and, for every, $2 \leq i \leq m$, we can write

$$p_i = y_i + D_i(y_1) + P_i \cdot \mathbf{u},$$

where P_i is a $1 \times n$ matrix of linear differential operators and the order of operator D_i is at most $n - 1$.

We show that the coefficients of $p_1, \dots, p_m$ can be written in terms of the coefficients of $h_1, \dots, h_m$. Since h_1 equals $D(y_1) -$

$Q_1 \cdot \mathbf{u}$, this is true for the coefficients of D and Q_1 . It remains to show this for the coefficients of $D_2, \dots, D_m$ and $P_2, \dots, P_m$. Note that for all i and for $j \notin \text{In}$ we have $(P_i)_j = 0$, so we need only address the coefficients of $(P_i)_j$ for $j \in \text{In}$.

Fix $i > 1$ and let $g = y_i + D_i(y_1) + P_i(\mathbf{u})$. We have that

$$D(g) - D_i(h_1) = D(y_i) + (DP_i + D_i Q_1)(\mathbf{u}) \in I_\Sigma.$$

It follows that $D(y_i) + (DP_i + D_i Q_1)(\mathbf{u}) = h_i$, so, for all j ,

$$D(P_i)_j + D_i(Q_1)_j = -(Q_i)_j.$$

By the hypothesis of the theorem, $\text{In} \neq \emptyset$. Fix $j \in \text{In}$. We apply [15, Proposition 3.19] to the model obtained from Σ by deleting all the inputs except for j and obtain, using $D \neq 1$, that $\gcd(D, (Q_1)_j) = 1$ for every $j \in \text{In}$. By Corollary 4, the coefficients of $(P_i)_j$ and D_i belong to the field extension of $\mathbb{C}$ generated by the coefficients of D , $(Q_1)_j$, and $(Q_i)_j$. We showed that the field extension of $\mathbb{C}$ generated by the coefficients of $h_1, \dots, h_m$ is the field of IO-identifiable functions. By Theorem 2, this is the field of identifiable functions. ■

APPENDIX

A. General definition of identifiability

In this section, we will generalize the notions from Section II-A to ODE systems with rational right-hand side. Fix positive integers λ , n , m , and κ for the remainder of the appendix. Let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_\lambda)$, $\mathbf{x} = (x_1, \dots, x_n)$, $\mathbf{y} = (y_1, \dots, y_m)$, and $\mathbf{u} = (u_1, \dots, u_\kappa)$. Consider a system of ODEs

$$\Sigma = \begin{cases} \mathbf{x}' = \frac{\mathbf{f}(\mathbf{x}, \boldsymbol{\mu}, \mathbf{u})}{Q(\mathbf{x}, \boldsymbol{\mu}, \mathbf{u})}, \\ \mathbf{y} = \frac{\mathbf{g}(\mathbf{x}, \boldsymbol{\mu}, \mathbf{u})}{Q(\mathbf{x}, \boldsymbol{\mu}, \mathbf{u})}, \\ \mathbf{x}(0) = \mathbf{x}^*, \end{cases} \quad (19)$$

where $\mathbf{f} = (f_1, \dots, f_n)$ and $\mathbf{g} = (g_1, \dots, g_m)$ are tuples of elements of $\mathbb{C}[\boldsymbol{\mu}, \mathbf{x}, \mathbf{u}]$ and $Q \in \mathbb{C}[\boldsymbol{\mu}, \mathbf{x}, \mathbf{u}] \setminus \{0\}$.

Notation 3 (Ideal I_Σ):

(a) For an ideal I and element a in a ring R , we denote

$$I : a^\infty = \{r \in R \mid \exists \ell : a^\ell r \in I\}.$$

This set is also an ideal in R .

(b) Given Σ as in (19), we define the differential ideal of Σ :

$$I_\Sigma = [Q\mathbf{x}' - \mathbf{f}, Q\mathbf{y} - \mathbf{g}] : Q^\infty \subset \mathbb{C}(\boldsymbol{\mu})\{\mathbf{x}, \mathbf{y}, \mathbf{u}\}.$$

For the case of a linear system as in (5), this ideal coincides with the one from Notation 2.

Notation 4 (Auxiliary analytic notation):

(a) For every given $h \in \mathbb{C}(\mathbf{x}^*, \boldsymbol{\mu})$, let

$$\begin{aligned} \Omega &= \{(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}}) \in \mathbb{C}^n \times \mathbb{C}^\lambda \times (\mathbb{C}^\infty(0))^\kappa \mid Q(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}}(0)) \neq 0\} \\ \Omega_h &= \Omega \cap \{(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}) \in \mathbb{C}^{n+\lambda} \mid h(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}) \text{ well-defined}\} \\ &\quad \times (\mathbb{C}^\infty(0))^\kappa. \end{aligned}$$

(b) For $(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}}) \in \Omega$, let $X(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}})$ and $Y(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}})$ denote the unique solution over $\mathbb{C}^\infty(0)$ of the instance of Σ with $\mathbf{x}^* = \hat{\mathbf{x}}^*$, $\boldsymbol{\mu} = \hat{\boldsymbol{\mu}}$, and $\mathbf{u} = \hat{\mathbf{u}}$ (see [41, Theorem 2.2.2]).

Definition 6 (Identifiability, see [5, Definition 2.5]): We say that $h(\mathbf{x}^*, \boldsymbol{\mu}) \in \mathbb{C}(\mathbf{x}^*, \boldsymbol{\mu})$ is *identifiable* if

$$\begin{aligned} & \exists \Theta \in \tau(\mathbb{C}^n \times \mathbb{C}^\lambda) \exists U \in \tau((\mathbb{C}^\infty(0))^\kappa) \\ & \forall (\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}}) \in (\Theta \times U) \cap \Omega_h \quad |S_h(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}})| = 1, \\ & \text{where } S_h(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}}) := \{h(\tilde{\mathbf{x}}^*, \tilde{\boldsymbol{\mu}}) \mid (\tilde{\mathbf{x}}^*, \tilde{\boldsymbol{\mu}}, \hat{\mathbf{u}}) \in \Omega_h \\ & \quad Y(\hat{\mathbf{x}}^*, \hat{\boldsymbol{\mu}}, \hat{\mathbf{u}}) = Y(\tilde{\mathbf{x}}^*, \tilde{\boldsymbol{\mu}}, \hat{\mathbf{u}})\}. \end{aligned}$$

In this paper, we are interested in comparing identifiability and IO-identifiability (Definition 7), and the latter is defined for functions in $\boldsymbol{\mu}$, not in $\mathbf{x}^*$. Thus, just for the purpose of comparison, we will restrict ourselves to the field

$$\{h \in \mathbb{C}(\boldsymbol{\mu}) \mid h \text{ is identifiable}\},$$

which we will call *the field of identifiable functions*.

Definition 7 (IO-identifiability): The smallest field k such that $\mathbb{C} \subset k \subset \mathbb{C}(\boldsymbol{\mu})$ and $I_\Sigma \cap \mathbb{C}(\boldsymbol{\mu})\{\mathbf{y}, \mathbf{u}\}$ is generated (as an ideal or as a differential ideal) by $I_\Sigma \cap k\{\mathbf{y}, \mathbf{u}\}$ is called *the field of IO-identifiable functions*.

We call $h \in \mathbb{C}(\boldsymbol{\mu})$ *IO-identifiable* if $h \in k$.

B. Specialization to the linear case

Proposition 1: For every system Σ of the form (5):

- (1) for every ordering of output variables, there exists a unique full set of input-output equations with respect to this ordering;
- (2) if $p_1, \dots, p_m$ is the full set of input-output equations with respect to $y_1 < \dots < y_m$, then the derivatives of $p_1, \dots, p_m$ form a Gröbner basis of $I_\Sigma \cap \mathbb{C}(\boldsymbol{\mu})\{\mathbf{y}, \mathbf{u}\}$ with respect to any lexicographic monomial ordering such that
 - any derivative of any of $\mathbf{y}$'s is greater any derivative of any of $\mathbf{u}$'s;
 - $y_{i_1}^{(j_1)} > y_{i_2}^{(j_2)}$ iff $i_1 > i_2$ or $i_1 = i_2$ and $j_1 > j_2$.

An analogous statement holds for any ordering of outputs.

- (3) Definitions 7 and 3 define the same field. In particular, the field defined in Definition 3 does not depend on the choice of a full set of input-output equations.

Proof: We fix an ordering $y_1 < \dots < y_m$ of outputs. Assume that there are full sets of input-output equations $p_1, \dots, p_m$ and $q_1, \dots, q_m$ with respect to this ordering. Let ℓ be the smallest integer such that $p_\ell \neq q_\ell$. By the definition, $\text{ord}_{y_\ell} p_\ell = \text{ord}_{y_\ell} q_\ell$. Then $\text{ord}_{y_i}(p_\ell - q_\ell) < \text{ord}_{y_i} p_i$ for every $i \leq \ell$; this contradicts the definition of a full set of input-output equations. To finish the proof of part (1) of the proposition, we will show the existence of a full set of input-output equations.

Let $J := I_\Sigma \cap \mathbb{C}(\boldsymbol{\mu})\{\mathbf{y}, \mathbf{u}\}$. Consider the set of differential polynomials

$$S := \{\mathbf{x}' - \mathbf{f}, \mathbf{x}'' - \mathbf{f}', \dots, \mathbf{y} - \mathbf{g}, \mathbf{y}' - \mathbf{g}', \dots\}.$$

By the definition of I_Σ , S generates I_Σ . Since these generators are linear, I_Σ has a linear Gröbner basis (see [45, Definition 1.4]) with respect to any monomial ordering. Since J is an elimination ideal of I_Σ , it also has a linear Gröbner basis with respect to any monomial ordering. Consider any lexicographic monomial ordering on $\mathbb{C}(\boldsymbol{\mu})\{\mathbf{x}, \mathbf{y}, \mathbf{u}\}$ such that

- any derivative of any of $y_1, \dots, y_m$ is greater than any derivative of any of $x_1, \dots, x_n$;

- any derivative of any of $x_1, \dots, x_n$ is greater than any derivative of any of $u_1, \dots, u_\kappa$;
- for $a = x, y$, $a_{i_1}^{(j_1)} > a_{i_2}^{(j_2)}$ iff $i_1 > i_2$ or $i_1 = i_2$ and $j_1 > j_2$.

Observe that S is a Gröbner basis of I_Σ with respect to any such monomial ordering. Therefore, $\mathbf{u}$ and their derivatives are algebraically independent modulo I_Σ , and the transcendence degree of $\mathbb{C}(\boldsymbol{\mu})\{\mathbf{x}, \mathbf{y}, \mathbf{u}\}$ over $\mathbb{C}(\boldsymbol{\mu})\{\mathbf{u}\}$ modulo I_Σ is finite.

Consider the restriction of the ordering described above to $\mathbb{C}(\boldsymbol{\mu})\{\mathbf{y}, \mathbf{u}\}$. Consider the reduced Gröbner basis B of J with respect to this ordering. As we have shown, it is linear. Since the transcendence degree of $\mathbb{C}(\boldsymbol{\mu})\{\mathbf{y}, \mathbf{u}\}$ over $\mathbb{C}(\boldsymbol{\mu})\{\mathbf{u}\}$ modulo J is finite, for every $1 \leq i \leq m$, there is a derivative of y_i among the leading terms of B . Moreover, by differentiating the corresponding element of B , we see that all higher derivatives of y_i will appear as leading terms of B .

For each $1 \leq i \leq m$, we set p_i to be the element in B with the leading term being $y_i^{(j)}$ such that j is the smallest possible. Then the fact that $p_1, \dots, p_m$ are a part of the reduced Gröbner basis implies that they form a full set of input-output equations with respect to the ordering $y_1 < y_2 < \dots < y_m$. This finishes the proof of part (1) of the proposition.

To prove part (2) of the proposition, observe that the derivatives of $p_1, \dots, p_m$ form a Gröbner basis of $[p_1, \dots, p_m]$ with respect to the described ordering. Thus, it remains to show that $[p_1, \dots, p_m] = J$. Assume that there is $q \in J \setminus [p_1, \dots, p_m]$. By reducing it with respect to appropriate derivatives of $p_1, \dots, p_m$, we can assume that $\text{ord}_{y_i} q < \text{ord}_{y_i} p_i$ for every $1 \leq i \leq m$. But this would imply that $p_1, \dots, p_m$ is not a full set of input-output equations, proving part (2) of the proposition.

To prove part (3) of the proposition, note that, since a full set of input-output equations is a part of a reduced Gröbner basis of J , its coefficients are in the field of definition of J . On the other hand, since the set of all derivatives of $p_1, \dots, p_m$ forms a Gröbner basis of J and the coefficients of these derivatives are the same as the coefficients of $p_1, \dots, p_m$, the coefficients of $p_1, \dots, p_m$ generate the field of definition of J . ■

C. Input-output equations based on the Cramer's rule and the transfer function matrix

Recall [40, page 444] that the *transfer function matrix* of a linear system

$$\begin{cases} \mathbf{x}' = A(\boldsymbol{\mu})\mathbf{x} + B(\boldsymbol{\mu})\mathbf{u}, \\ \mathbf{y} = C(\boldsymbol{\mu})\mathbf{x}. \end{cases}$$

is defined by

$$H(\boldsymbol{\mu}, s) := C(\boldsymbol{\mu})(sI - A(\boldsymbol{\mu}))^{-1}B(\boldsymbol{\mu}), \quad (20)$$

where s is a new algebraic variable and I is the identity matrix. This matrix relates the Laplace transforms of $\mathbf{y}$ and $\mathbf{u}$ under the assumption that the initial conditions are zero [40, page 75]. The formulas (9) and (20) look similar and in fact are related. We give a connection we are interested in as Lemma 5 below, and refer for further connection to an upcoming paper [50].

For a rational function $f \in \mathbb{C}(\boldsymbol{\mu})(s)$ in s , by *the coefficients of f* , we will understand the union of the coefficients of the numerator and denominator in *the reduced form* if the denominator is taken to be monic.

Lemma 5: Consider a linear compartment model with at least one input and whose graph is strongly connected. Then the following sets generate the same subfield in $\mathbb{C}(\mu)$

- coefficients of the input-output equations (9);
- coefficients of the entries of the transfer function matrix.

Proof: Since each of the equations (9) involves only one output and each row in the matrix (20) corresponds to an output, proving the lemma for the single-output case will yield the general case by taking the union of the respective generators.

We write (9) as $p(\partial)y = q_1(\partial)u_1 + \dots + q_r(\partial)u_r$ for nonzero $p(s), q_1(s), \dots, q_r(s) \in \mathbb{C}(\mu)[s]$ such that $p(s)$ is monic. Let F_1 be the field generated by the coefficients of $p, q_1, \dots, q_r$. Since the graph is strongly connected and has an input, [15, Proposition 3.19] implies that $\gcd(p, q_1, \dots, q_r) = 1$. A direct computation shows that $H(\mu)$ defined by (20) is equal to

$$H(\mu) := (h_1(s), h_2(s), \dots, h_r(s)) = \left(\frac{q_1(s)}{p(s)}, \frac{q_2(s)}{p(s)}, \dots, \frac{q_r(s)}{p(s)} \right).$$

Let F_2 be the field generated by the coefficients of $h_1, \dots, h_r$. For all integers $n_1, \dots, n_r$, the coefficients of $n_1 h_1 + \dots + n_r h_r$ belong to F_2 . Since $q_1, \dots, q_r, p$ are coprime, there exist integers $n_1, \dots, n_r$ such that $n_1 q_1 + \dots + n_r q_r$ is coprime with p , so p is the denominator of $n_1 h_1 + \dots + n_r h_r$. Hence, the coefficients of p belong to F_2 . Let $g_i = \gcd(p, q_i)$ and $p_i = p/g_i$. By definition, the coefficients of p_i are in F_2 , so the coefficients of g_i are in F_2 . By definition, the coefficients of q_i/g_i are in F_2 , so the coefficients of q_i are in F_2 . Thus, $F_1 \subset F_2$.

To prove $F_2 \subset F_1$, note that the coefficients of the remainder and quotient of two polynomials belong to the field generated by the coefficients of the polynomials. Since the numerator and denominator of h_i are equal to $q_i/\gcd(p, q_i)$ and $p/\gcd(p, q_i)$, respectively, we have $F_2 \subset F_1$, so $F_1 = F_2$. ■

ACKNOWLEDGMENTS

We are grateful to the CCiS at CUNY Queens College for the computational resources and to Julio Banga, Joseph DiStefano, Marisa Eisenberg, Nikki Meshkat, Maria Pia Saccomani, Anne Shiu, Seth Sullivan, Alejandro Villaverde, and to referees for useful discussions and suggestions.

REFERENCES

- [1] L. Ljung and T. Glad, "On global identifiability for arbitrary model parametrizations," *Automatica*, vol. 30, no. 2, pp. 265–276, 1994. [Online]. Available: [https://doi.org/10.1016/0005-1098\(94\)90029-9](https://doi.org/10.1016/0005-1098(94)90029-9)
- [2] A. F. Villaverde, N. D. Evans, M. J. Chappell, and J. R. Banga, "Sufficiently exciting inputs for structurally identifiable systems biology models," *IFAC-PapersOnLine*, vol. 51, no. 19, pp. 16–19, 2018. [Online]. Available: <https://doi.org/10.1016/j.ifacol.2018.09.015>
- [3] X. Xia and C. Moog, "Identifiability of nonlinear systems with application to HIV/AIDS models," *IEEE Transactions on Automatic Control*, vol. 48, no. 2, pp. 330–336, Feb. 2003. [Online]. Available: <https://doi.org/10.1109/tac.2002.808494>
- [4] O.-T. Chis, J. R. Banga, and E. Balsa-Canto, "Structural identifiability of systems biology models: A critical comparison of methods," *PLoS ONE*, vol. 6, no. 11, pp. 1–16, 11 2011. [Online]. Available: <https://doi.org/10.1371/journal.pone.0027755>
- [5] H. Hong, A. Ovchinnikov, G. Pogudin, and C. Yap, "Global identifiability of differential models," *Communications on Pure and Applied Mathematics*, vol. 73, no. 9, pp. 1831–1879, 2020. [Online]. Available: <https://doi.org/10.1002/cpa.21921>
- [6] A. F. Villaverde, "Observability and structural identifiability of nonlinear biological systems," *Complexity*, vol. 2019, pp. 1–12, Jan. 2019. [Online]. Available: <https://doi.org/10.1155/2019/8497093>
- [7] J. DiStefano, *Dynamic Systems Biology Modeling and Simulation*. Academic Press, 2014.
- [8] E. Walter and L. Pronzato, *Identification of Parametric Models from Experimental Data*. Springer, 1997.
- [9] J. Karlsson, M. Anguelova, and M. Jirstrand, "An efficient method for structural identifiability analysis of large dynamic systems," *IFAC Proceedings Volumes*, vol. 45, no. 16, pp. 941 – 946, 2012, 16th IFAC Symposium on System Identification. [Online]. Available: <https://doi.org/10.3182/20120711-3-BE-2027.00381>
- [10] A. Sedoglavic, "A probabilistic algorithm to test local algebraic observability in polynomial time," *Journal of Symbolic Computation*, vol. 33, no. 5, pp. 735–755, 2002. [Online]. Available: <https://doi.org/10.1006/jsc.2002.0532>
- [11] A. F. Villaverde, A. Barreiro, and A. Papachristodoulou, "Structural identifiability of dynamic systems biology models," *PLOS Computational Biology*, vol. 12, no. 10, p. e1005153, 2016. [Online]. Available: <https://doi.org/10.1371/journal.pcbi.1005153>
- [12] J. Baaijens and J. Draisma, "On the existence of identifiable reparametrizations for linear compartment models," *SIAM Journal on Applied Mathematics*, vol. 76, no. 4, 2016. [Online]. Available: <https://doi.org/10.1137/15M1038013>
- [13] M. Saccomani, S. Audoly, and L. D'Angi , "Parameter identifiability of nonlinear systems: the role of initial conditions," *Automatica*, vol. 39, pp. 619–632, 2003. [Online]. Available: [https://doi.org/10.1016/S0005-1098\(02\)00302-3](https://doi.org/10.1016/S0005-1098(02)00302-3)
- [14] N. Meshkat, S. Sullivan, and M. Eisenberg, "Identifiability results for several classes of linear compartment models," *Bulletin of Mathematical Biology*, vol. 77, pp. 1620–1651, 10 2014. [Online]. Available: <https://doi.org/10.1007/s11538-015-0098-0>
- [15] E. Gross, H. A. Harrington, N. Meshkat, and A. Shiu, "Linear compartmental models: input-output equations and operations that preserve identifiability," *SIAM Journal on Applied Mathematics*, vol. 79, no. 4, pp. 1423–1447, 2019. [Online]. Available: <https://doi.org/10.1137/18M1204826>
- [16] D. Bearup, N. Evans, and M. Chappell, "The input-output relationship approach to structural identifiability analysis," *Computer Methods and Programs in Biomedicine*, vol. 109, no. 2, pp. 171–181, 2013. [Online]. Available: <https://doi.org/10.1016/j.cmpb.2012.10.012>
- [17] G. Bellu, M. P. Saccomani, S. Audoly, and L. D'Angi , "DAISY: A new software tool to test global identifiability of biological and physiological systems," *Computer Methods and Programs in Biomedicine*, vol. 88, no. 1, pp. 52–61, 2007. [Online]. Available: <https://doi.org/10.1016/j.cmpb.2007.07.002>
- [18] M. Saccomani and L. D'Angi , "Examples of testing global identifiability with the DAISY software," *IFAC Proceedings Volumes*, vol. 42, no. 10, pp. 48–53, 2009. [Online]. Available: <https://doi.org/10.3182/20090706-3-FR-2004.00007>
- [19] M. P. Saccomani and G. Bellu, "DAISY: An efficient tool to test global identifiability. Some case studies," in *2008 16th Mediterranean Conference on Control and Automation*, 2008, pp. 1723–1728. [Online]. Available: <https://doi.org/10.1109/MED.2008.4602152>
- [20] N. Meshkat, C. Kuo, and J. DiStefano, "On finding and using identifiable parameter combinations in nonlinear dynamic systems biology models and COMBOS: A novel web implementation," *PLoS ONE*, vol. 9, no. 10, p. e110261, 2014. [Online]. Available: <https://doi.org/10.1371/journal.pone.0110261>
- [21] A. Kalamı Yazdı, M. Nadjafkhan, and J. DiStefano, "COMBOS2: an algorithm to the input-output equations of dynamic biosystems via Gaussian elimination," *Journal of Taibah University for Science*, vol. 14, no. 1, pp. 896–907, 2020. [Online]. Available: <https://doi.org/10.1080/16583655.2020.1776466>
- [22] N. Meshkat, Z. Rosen, and S. Sullivan, "Algebraic tools for the analysis of state space models," in *The 50th Anniversary of Gr bner Bases, July 1–10, 2015, Osaka, Japan*, ser. Advances in Pure Mathematics, T. Hibi, Ed., vol. 77. Tokyo, Japan: Mathematical Society of Japan, 2018, pp. 171–205. [Online]. Available: <https://doi.org/10.2969/aspm/07710171>
- [23] T. Boukhobza, F. Hamelin, and C. Simon, "A graph theoretical approach to the parameters identifiability characterisation," *International Journal of Control*, vol. 87, no. 4, pp. 751–763, 2013. [Online]. Available: <https://doi.org/10.1080/00207179.2013.856519>
- [24] M. Komatsu, T. Yaguchi, and K. Nakajima, "Algebraic approach towards the exploitation of "softness": the input-output equation for morphological computation," *The International*

- Journal of Robotics Research*, 2020. [Online]. Available: <https://doi.org/10.1177/0278364920912298>
- [25] A. Ovchinnikov, A. Pillay, G. Pogudin, and T. Scanlon, "Computing all identifiable functions of parameters for ODE models," *Systems & Control Letters*, vol. 157, p. 105030, 2021. [Online]. Available: <https://doi.org/10.1016/j.sysconle.2021.105030>
- [26] H. Hong, A. Ovchinnikov, G. Pogudin, and C. Yap, "SIAN: software for structural identifiability analysis of ODE models," *Bioinformatics*, vol. 35, no. 16, pp. 2873–2874, 2019. [Online]. Available: <https://doi.org/10.1093/bioinformatics/bty1069>
- [27] A. Ovchinnikov, G. Pogudin, and P. Thompson, "Parameter identifiability and input-output equations," *Applicable Algebra in Engineering, Communication and Computing*, 2021. [Online]. Available: <https://doi.org/10.1007/s00200-021-00486-8>
- [28] A. Ben-Zvi, P. McLellan, and K. McAuley, "Identifiability of linear time-invariant differential-algebraic systems. 2. The differential-algebraic approach," *Industrial & Engineering Chemistry Research*, vol. 43, no. 5, pp. 1251–1259, 2004. [Online]. Available: <https://doi.org/10.1021/ie030534j>
- [29] N. Meshkat and S. Sullivant, "Identifiable reparametrizations of linear compartment models," *Journal of Symbolic Computation*, vol. 63, pp. 46–67, 2014. [Online]. Available: <https://doi.org/10.1016/j.jsc.2013.11.002>
- [30] L. Denis-Vidal, G. Joly-Blanchard, C. Noiret, and M. Petitot, "An algorithm to test identifiability of non-linear systems," *IFAC Proceedings Volumes*, vol. 34, no. 6, pp. 197–201, 2001. [Online]. Available: [https://doi.org/10.1016/S1474-6670\(17\)35173-X](https://doi.org/10.1016/S1474-6670(17)35173-X)
- [31] H. Miao, X. Xia, A. S. Perelson, and H. Wu, "On identifiability of nonlinear ODE models and applications in viral dynamics," *SIAM Review*, vol. 53, no. 1, pp. 3–39, 2011. [Online]. Available: <https://doi.org/10.1137/090757009>
- [32] F. Anstett-Collin, L. Denis-Vidal, and G. Millérioux, "A priori identifiability: An overview on definitions and approaches," *Annual Reviews in Control*, vol. 50, pp. 139–149, 2020. [Online]. Available: <https://doi.org/10.1016/j.arcontrol.2020.10.006>
- [33] S. Audoly, L. D'Angiò, M. Saccomani, and C. Cobelli, "Global identifiability of linear compartmental models – a computer algebra algorithm," *IEEE Transactions on Biomedical Engineering*, vol. 45, no. 1, pp. 36–47, 1998. [Online]. Available: <https://doi.org/10.1109/10.650350>
- [34] R. Bellman and K. Åström, "On structural identifiability," *Mathematical Biosciences*, vol. 7, no. 3–4, pp. 329–339, 1970. [Online]. Available: [https://doi.org/10.1016/0025-5564\(70\)90132-X](https://doi.org/10.1016/0025-5564(70)90132-X)
- [35] K. Godfrey and M. Chapman, "Identifiability and indistinguishability of linear compartmental models," *Mathematics and Computers in Simulation*, vol. 32, no. 3, pp. 273–295, 1990. [Online]. Available: [https://doi.org/10.1016/0378-4754\(90\)90185-L](https://doi.org/10.1016/0378-4754(90)90185-L)
- [36] K. Godfrey and J. DiStefano, "Identifiability of model parameters," *IFAC Proceedings Volumes*, vol. 18, no. 5, pp. 89–114, 1985. [Online]. Available: [https://doi.org/10.1016/S1474-6670\(17\)60544-5](https://doi.org/10.1016/S1474-6670(17)60544-5)
- [37] J. van den Hof, "Structural identifiability of linear compartmental systems," *IEEE Transactions on Automatic Control*, vol. 43, no. 6, pp. 800–818, 1998. [Online]. Available: <https://doi.org/10.1109/9.679020>
- [38] E. Walter and Y. Lecourtier, "Unidentifiable compartmental models: what to do?" *Mathematical Biosciences*, vol. 56, no. 1–2, pp. 1–25, 1981. [Online]. Available: [https://doi.org/10.1016/0025-5564\(81\)90025-0](https://doi.org/10.1016/0025-5564(81)90025-0)
- [39] A. Ovchinnikov, A. Pillay, G. Pogudin, and T. Scanlon, "Multi-experiment parameter identifiability of ODEs and model theory," 2021. [Online]. Available: <https://arxiv.org/abs/2011.10868>
- [40] J. J. DiStefano, *Dynamic systems biology modeling and simulation*. Elsevier, Academic Press, 2013.
- [41] E. Hille, *Ordinary Differential Equations in the Complex Domain*. Dover, 1997.
- [42] D. H. Anderson, *Compartmental Modeling and Tracer Kinetics*, ser. Lecture Notes in Bioinformatics. Springer, 1983. [Online]. Available: <https://doi.org/10.1007/978-3-642-51861-4>
- [43] E. Walter, *Identifiability of State Space Models*, ser. Lecture Notes in Biomathematics. Springer, 1982, vol. 46. [Online]. Available: <https://doi.org/10.1007/978-3-642-61823-9>
- [44] N. Meshkat, M. Eisenberg, and J. DiStefano, "An algorithm for finding globally identifiable parameter combinations of nonlinear ODE models using Gröbner bases," *Mathematical Biosciences*, vol. 222, no. 2, pp. 61–72, 2009. [Online]. Available: <https://doi.org/10.1016/j.mbs.2009.08.010>
- [45] K. Iima and Y. Yoshino, "Gröbner bases for the polynomial ring with infinite variables and their applications," *Communications in Algebra*, vol. 37, no. 10, pp. 3424–3437, 2009. [Online]. Available: <https://doi.org/10.1080/00927870802502878>
- [46] I. Kaplansky, *An Introduction to Differential Algebra*. Paris: Hermann, 1957.
- [47] T. H. Cormen, C. E. Leiserson, R. L. Rivest, and C. Stein, *Introduction to Algorithms, Third Edition*. The MIT Press, 2009.
- [48] A. Nowicki, *Polynomial derivations and their rings of constants*. Toruń: Uniwersytet Mikołaja Kopernika, 1994.
- [49] J. S. Milne, "Fields and Galois theory (v4.60)," 2018. [Online]. Available: <https://www.jmilne.org/math/CourseNotes/ft.html>
- [50] N. Meshkat and M. Eisenberg, "On reducibility of linear compartmental models and implications for structural identifiability analysis," 2021, in preparation.

This figure "lincomp.png" is available in "png" format from:

<http://arxiv.org/ps/1910.03960v5>