

MINIMAL ASYMPTOTIC TRANSLATION LENGTHS ON CURVE COMPLEXES AND HOMOLOGY OF MAPPING TORI

HYUNGRYUL BAIK, DONGRYUL M. KIM, AND CHENXI WU

ABSTRACT. Let S_g be a closed orientable surface of genus $g > 1$. Consider the minimal asymptotic translation length $L_{\mathcal{T}}(k, g)$ on the Teichmüller space of S_g , among pseudo-Anosov mapping classes of S_g acting trivially on a k -dimensional subspace of $H_1(S_g)$, $0 \leq k \leq 2g$. The asymptotics of $L_{\mathcal{T}}(k, g)$ for extreme cases $k = 0, 2g$ have been shown by several authors. Jordan Ellenberg asked whether there is a lower bound for $L_{\mathcal{T}}(k, g)$ interpolating the known results on $L_{\mathcal{T}}(0, g)$ and $L_{\mathcal{T}}(2g, g)$, which was affirmatively answered by Agol, Leininger, and Margalit.

In this paper, we study an analogue of Ellenberg's question, replacing Teichmüller spaces with curve complexes. We provide lower and upper bound on the minimal asymptotic translation length $L_{\mathcal{C}}(k, g)$ on the curve complex, whose lower bound interpolates the known results on $L_{\mathcal{C}}(0, g)$ and $L_{\mathcal{C}}(2g, g)$.

Finally, for each g , we construct a non-Torelli pseudo-Anosov $f_g \in \text{Mod}(S_g)$ which does not normally generates $\text{Mod}(S_g)$ and so that the asymptotic translation length of f_g on curve complexes decays more quickly than a constant multiple of $1/g$ as $g \rightarrow \infty$. From this, we provide a restriction on how small the asymptotic translation lengths on curve complexes should be if the similar phenomenon as in the work of Lanier and Margalit on Teichmüller spaces holds for curve complexes.

MSC 2020 classes: 37E30, 57K20

1. INTRODUCTION

Let S_g be a closed connected orientable surface of genus $g > 1$, $\text{Mod}(S_g)$ be its mapping class group, and $\mathcal{C}(S_g)$ be its curve complex. Then $\text{Mod}(S_g)$ isometrically acts on $\mathcal{C}(S_g)$, hence the *asymptotic translation length* $\ell_{\mathcal{C}}(f)$ of $f \in \text{Mod}(S_g)$ on $\mathcal{C}(S_g)$ is defined:

$$\ell_{\mathcal{C}}(f) := \liminf_{n \rightarrow \infty} \frac{d_{\mathcal{C}}(x, f^n(x))}{n}$$

for any $x \in \mathcal{C}(S_g)$ where $d_{\mathcal{C}}$ is the standard metric on $\mathcal{C}(S_g)$. The asymptotic translation length is also called *stable translation length*.

Note that $\text{Mod}(S_g)$ also acts on $H_1(S_g)$, the first homology group of S_g with real coefficients. For $f \in \text{Mod}(S_g)$, we denote the dimension of a maximal subspace of $H_1(S_g)$ on which f is trivial by $m(f)$. In particular,

$m(f) = 2g$ if and only if f is in the *Torelli group* $\mathcal{I}_g \leq \text{Mod}(S_g)$, the subgroup consisting of elements which act trivially on $H_1(S_g)$. As an application of Mayer-Vietoris sequence, one can observe that $m(f) + 1$ is same as the first betti number of the mapping torus of f , which is hyperbolic if and only if f is pseudo-Anosov by Thurston [Thu98].

In this paper, we mainly study the *minimal asymptotic translation lengths* among pseudo-Anosov mapping classes acting trivially on some subspaces of homology groups. Namely, for $0 \leq k \leq 2g$, we define

$$L_{\mathcal{C}}(k, g) := \inf \{ \ell_{\mathcal{C}}(f) : f \in \text{Mod}(S_g), f \text{ is pseudo-Anosov}, m(f) \geq k \}.$$

Then we investigate asymptotics of $L_{\mathcal{C}}(k, g)$ with varying k and g .

By replacing the curve complex $\mathcal{C}(S_g)$ with Teichmüller space $\mathcal{T}(S_g)$, one can also define $L_{\mathcal{T}}(k, g)$ analogously. Note that $\ell_{\mathcal{T}}(f)$ for a pseudo-Anosov element f is same as the logarithm of the stretch factor [B⁺78], hence coincides with the topological entropy of f [FLP12, Exposé Ten].

In each setting, there are two extreme cases: the first extreme is the case $k = 0$ that the minimal asymptotic translation length is considered in the *entire mapping class group* $\text{Mod}(S_g)$. The other extreme is $k = 2g$, which means that the minimal asymptotic translation length is considered in the *Torelli subgroup* $\mathcal{I}_g < \text{Mod}(S_g)$. These four cases have been resolved by various authors as below:¹

	Teichmüller spaces	Curve complexes
$\text{Mod}(S_g)$	(Penner [Pen91]) $L_{\mathcal{T}}(0, g) \asymp 1/g$	(Gadre-Tsai [GT11]) $L_{\mathcal{C}}(0, g) \asymp 1/g^2$
$\mathcal{I}_g$	(Farb-Leininger-Margalit [FLM08]) $L_{\mathcal{T}}(2g, g) \asymp 1$	(Baik-Shin [BS20]) $L_{\mathcal{C}}(2g, g) \asymp 1/g$

TABLE 1. Four extreme cases of $L_{\mathcal{T}}(k, g)$ and $L_{\mathcal{C}}(k, g)$.

Ellenberg [Ell] asked if $L_{\mathcal{T}}(k, g)$ interpolates $L_{\mathcal{T}}(0, g)$ and $L_{\mathcal{T}}(2g, g)$ in the sense that there exists $C > 0$ such that

$$(1.1) \quad L_{\mathcal{T}}(k, g) \geq C(k+1)/g$$

for all $g > 1$ and $0 \leq k \leq 2g$. This was answered affirmatively by Agol-Leininger-Margalit in [ALM16]. Indeed, they actually showed $L_{\mathcal{T}}(k, g) \asymp (k+1)/g$.

¹Throughout the paper, we write $A(x) \gtrsim B(x)$ if there exists a uniform constant $C > 0$ such that $A(x) \leq CB(x)$ for all x in the domain. We also write $A(x) \asymp B(x)$ if $A(x) \gtrsim B(x)$ and $B(x) \gtrsim A(x)$.

We ask an analogous question whether $L_C(k, g)$ interpolates $L_C(0, g)$ and $L_C(2g, g)$ in a similar sense as Ellenberg's question (1.1). We show that this is indeed the case and more concretely we obtain the following:

Theorem 1.1. *There exists $C, C' > 0$ such that*

$$\frac{C}{g(2g - k + 1)} \leq L_C(k, g) \leq C' \frac{k + 1}{g \log g}$$

for all $g > 1$ and $0 \leq k \leq 2g$.

From the statement, if k grows at least $2g - C'$ for some constant $C' > 0$, then $L_C(k, g) \gtrsim 1/g$ while $L_C(0, g) \asymp 1/g^2$. Observing this, we ask about minimal k with $L_C(k, g) \asymp 1/g$. For this discussion, see Section 4.

While the lower bound in Theorem 1.1 interpolates $L_C(0, g) \asymp 1/g^2$ and $L_C(2g, g) \asymp 1/g$, the upper bound in Theorem 1.1 does not interpolate these two values well. Indeed, we construct some values of k and g showing that $\frac{k+1}{g \log g}$ is larger than the actual asymptotic. We also show that k/g^2 works as an upper bound for some choices of (k, g) , which interpolates $L_C(0, g) \asymp 1/g^2$ and $L_C(2g, g) \asymp 1/g$.

Theorem 1.2. *There is a uniform constant $C > 0$ satisfying the following:*

For any integers $g, k \geq 0$, there exists a pseudo-Anosov $f : S_{g'} \rightarrow S_{g'}$ such that $g' > g$, $m(f) = k' > k$, and

$$\ell_C(f) \leq C \frac{k'}{g'^2}.$$

Applying Theorem 1.2 inductively, it follows that there is a diverging sequence $(k_j, g_j) \rightarrow \infty$ so that $L_C(k_j, g_j) \lesssim k_j/g_j^2$. See Corollary 3.1. Based on Table 1, we conjecture that the upper bound in Theorem 1.2 is actually the asymptotic for $L_C(k, g)$.

Conjecture 1.3. *We have*

$$L_C(k, g) \asymp \frac{k}{g^2}$$

for $g > 1$ and $0 \leq k \leq 2g$.

We focus on specific dimensions of maximal invariant subspaces. In [BS20], Torelli pseudo-Anosovs are constructed in a concrete way based on Penner's or Thurston's construction. In a similar line of thought, we utilize finite cyclic covers of S_2 so that we get pseudo-Anosovs $f \in \text{Mod}(S_g)$ with $m(f) = 2g - 1$ and satisfying the upper bound in Theorem 1.2. As a consequence, this yields the asymptotic of $L_C(2g - 1, g)$; only two extreme cases $\text{Mod}(S_g)$ and $\mathcal{I}_g$ were previously known. It is also interesting to figure out the asymptote $L_C(k, g)$ for other values (k, g) :

Question 1.4. *Can we give a sequence (k_j, g_j) , other than $(0, g)$ and $(2g, g)$, with explicit asymptote for $L_C(k_j, g_j)$ as $j \rightarrow \infty$?*

We give one such example in the following:

Theorem 1.5. *There exists a uniform constant $C > 0$ and pseudo-Anosovs $f_g \in \text{Mod}(S_g)$ such that*

$$m(f_g) = 2g - 1 \quad \text{and} \quad \ell_C(f_g) \leq \frac{C}{g}$$

for all $g > 1$. Moreover, the following asymptotic holds:

$$L_C(2g - 1, g) \asymp \frac{1}{g}.$$

The construction involved in Theorem 1.5 can be modified to deal with the Torelli case. Such a modification gives an asymptotic for $L_C(2g, g)$ which was already shown by [BS20] in a different way. See Remark 4.1. Further, only the last assertion can also be deduced from Theorem 1.1 and [BS20]. See Section 4 for details.

In [LM22], Lanier and Margalit showed that a pseudo-Anosov with small asymptotic translation length on the Teichmüller space has an entire mapping class group as its normal closure. The first and the third authors, Kim, and Shin, made an analogous question for asymptotic translation lengths on curve complexes in [BKSW23] (see [BKSW23, Question 1.2]). We later show that pseudo-Anosovs f_g constructed in Theorem 1.5 never normally generate the mapping class groups. As $\ell_C(f_g)$ is concretely estimated in Section 4, it provides how small the asymptotic translation length should be in order to observe the similar phenomenon as in [LM22]. In other words, we prove the following.

Theorem 1.6. *Suppose that there exists a universal constant C so that if a non-Torelli pseudo-Anosov $f \in \text{Mod}(S_g)$ has $\ell_C(f) < C/g$ then f normally generates $\text{Mod}(S_g)$ for large g . Then*

$$C \leq 1152.$$

Organization. In Section 2, we prove Theorem 1.1. Theorem 1.2 is proved in Section 3. In Section 4, explicit construction of pseudo-Anosovs realizing the asymptotic for $L_C(2g - 1, g)$ is provided, implying Theorem 1.5. The discussion on small asymptotic translation lengths on curve complexes and normal generation of mapping class groups is provided in Section 5.

Acknowledgements. The authors greatly appreciate Changsub Kim and Yair N. Minsky for helpful discussions. We also thank the anonymous referee for helpful comments. The first author was partially supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIT) (No. 2020R1C1C1A01006912).

2. PROOF OF THEOREM 1.1

In this section, we prove Theorem 1.1.

Lower bound. The main idea of showing the lower bound is similar to the one used in the proof in [BS20] of that $L_C(2g, g) \geq C/g$ for some constant. First note that for any $f : S_g \rightarrow S_g$, the Lefschetz number $L(f)$ is $2 - \text{Tr}(f_*)$ where $\text{Tr}(f_*)$ is the trace of the induced map $f_* : H_1(S_g) \rightarrow H_1(S_g)$.

Let us fix a pseudo-Anosov $f : S_g \rightarrow S_g$ whose restriction onto a k -dimensional subspace of $H_1(S_g)$ is the identity.

Pick a suitable basis for $H_1(S_g)$, the matrix for f_* can be written as

$$\begin{pmatrix} I_k & * \\ 0 & M \end{pmatrix}.$$

Suppose first that $k > 0$. When k is odd, let $m = 2g - k$ and when k is even, let $m = 2g - (k - 1)$. By taking the upper left block to be I_{k-1} in case k is even, one may assume M is a $m \times m$ square matrix with determinant 1 and m is odd (determinant 1 comes from the fact that f_* is actually a symplectic matrix).

Recall that there is a relation between trace and determinant as follows:

Lemma 2.1 ([KK92, Appendix B]). *For any $m \times m$ matrix A ,*

$$(-1)^m \det A = \sum_{\substack{c_1, \dots, c_m \geq 0, \\ c_1 + 2c_2 + \dots + mc_m = m}} \prod_{i=1}^m \frac{1}{c_i!} \left(-\frac{\text{Tr}(A^i)}{i} \right)^{c_i}$$

Observe that at least one of the matrices $M, M^2, \dots, M^m$ must have positive trace. Otherwise the right-hand side of the equality in Lemma 2.1 is always non-negative when we plug in M in the place of A in the lemma. On the other hand, since $\det(M) = 1$ and m is always odd by our choice, the left-hand side is -1 , a contradiction.

This implies that for some j satisfying $1 \leq j \leq m \leq 2g - k + 1$, $\text{Tr}(M^j)$ is positive, i.e., at least 1 since it is an integral matrix. $\text{Tr}(f_*^j)$ is the sum of $\text{Tr}(M^j)$ and the trace of the upper left block which is $2g - m \geq 1$. Therefore, $\text{Tr}(f_*^j)$ is at least 2 in general. But in fact $2g - m \geq 3$ as long as $k \geq 3$.

Assume $k \geq 3$. Now we have that $L(f^j) = 2 - \text{Tr}(f_*^j) < 0$ and we can apply a result of Tsai [Tsa09]. Then $\ell_C(f^j) \geq C/g$ for some constant $C > 0$ and consequently,

$$\ell_C \geq \frac{C}{gj} \geq \frac{C}{g(2g - k + 1)}.$$

For any fixed k , $\ell_C \geq L_C(0, g) \asymp 1/g^2$. Hence $\ell_C \geq \frac{C_k}{g(2g - k + 1)}$ for some constant C_k . Since the above argument works for any $k \geq 3$, replacing C by $\min\{C, C_0, C_1, C_2\}$, we obtain the lower bound in Theorem 1.1.

Upper bound. We now prove the upper bound provided in Theorem 1.1.

Recall that the Teichmüller space $\mathcal{T}(S_g)$ is a space of marked hyperbolic structures on S_g and vertices of the curve complex $\mathcal{C}(S_g)$ are essential simple closed curves on S_g . Hence, we can associate each point $x \in \mathcal{T}(S_g)$ with

systoles on S_g , the shortest simple closed geodesics, in the hyperbolic structure x . As systoles are within a uniformly bounded distance in the curve complex, it gives a coarsely well-defined map $\pi_g : \mathcal{T}(S_g) \rightarrow \mathcal{C}(S_g)$.

Masur-Misnky studied $\pi_g : \mathcal{T}(S_g) \rightarrow \mathcal{C}(S_g)$ in [MM99] and showed that π_g is coarsely Lipschitz.

Proposition 2.2 ((K_g, D_g) -coarsely Lipschitz, [MM99]). *There exist constants $K_g, D_g > 0$ such that for any $x, y \in \mathcal{T}(S_g)$ we have*

$$d_{\mathcal{C}}(\pi_g(x), \pi_g(y)) \leq K_g d_{\mathcal{T}}(x, y) + D_g$$

where $d_{\mathcal{T}}$ is the Teichmüller metric.

Furthermore, π_g is coarsely $\text{Mod}(S_g)$ -equivariant in the sense that there exists a constant A_g such that $d_{\mathcal{C}}((\pi_g \circ f)(x), (f \circ \pi_g)(x)) \leq A_g$ for any $x \in \mathcal{T}(S_g)$ and $f \in \text{Mod}(S_g)$. Then for $f \in \text{Mod}(S_g)$, $n > 0$, and $x \in \mathcal{T}(S_g)$, we have

$$\begin{aligned} d_{\mathcal{C}}(\pi_g(x), f^n(\pi_g(x))) &\leq d_{\mathcal{C}}(\pi_g(x), \pi_g(f^n(x))) + A_g \\ &\leq K_g d_{\mathcal{T}}(x, f^n(x)) + D_g + A_g \end{aligned}$$

Hence, we now have the comparison between asymptotic translation lengths of $f \in \text{Mod}(S_g)$ measured on $\mathcal{C}(S_g)$ and $\mathcal{T}(S_g)$:

$$\ell_{\mathcal{C}}(f) \leq K_g \ell_{\mathcal{T}}(f)$$

In particular, we have

$$(2.1) \quad L_{\mathcal{C}}(k, g) \leq K_g L_{\mathcal{T}}(k, g).$$

Due to the work [ALM16] of Agol-Leininger-Margalit, we already know the asymptotic of $L_{\mathcal{T}}(k, g)$. Hence, it remains to figure out the asymptotic of K_g .

In [GHKL13], Gadre-Hironaka-Kent-Leininger considered the minimal possible Lipschitz constant K_g which is defined as

$$\kappa_g := \inf \{K_g \geq 0 : \pi_g \text{ is } (K_g, D_g)\text{-coarsely Lipschitz for some } D_g > 0\}.$$

Then they showed that

$$\kappa_g \asymp \frac{1}{\log g}.$$

Combining this with [ALM16] and Inequality (2.1), we deduce the upper bound in Theorem 1.1.

3. UPPER BOUND INTERPOLATES $L_{\mathcal{C}}(0, g)$ AND $L_{\mathcal{C}}(2g, g)$

The upper bound provided in Theorem 1.1 does not interpolate $L_{\mathcal{C}}(0, g)$ and $L_{\mathcal{C}}(2g, g)$, and it is not sharp enough as one can see in Section 4. As stated in Theorem 1.2, the upper bound conjectured in Conjecture 1.3 can be observed along certain sequence $(k_j, g_j) \rightarrow \infty$. This section is devoted to proving Theorem 1.2.

Proof of Theorem 1.2. Let f_0 be a pseudo-Anosov map in the Torelli group of genus $g_0 > 1$. Let M be its mapping torus, $\alpha \in H^1(M)$ be the first cohomology class of M corresponding to f_0 , β be an element in $H^1(M)$ which is restricted to a cohomology class dual to a simple closed curve γ on S_{g_0} . For large enough $n > g + k$, let f_n be the pseudo-Anosov monodromy corresponding to $2^n \alpha + \beta$. Then f_n has the fiber of genus $O(2^n)$, and $\ell_C(f_n)$ is $O(2^{-2n})$. (cf. [BSW21])

A way to construct the surface S_n and map f_n corresponding to $2^n \alpha + \beta$ is as follows: Let $\widehat{S}$ be the $\mathbb{Z}$ -fold cover corresponding to β restricted to S_{g_0} , $\widehat{f}$ a lift of f_0 , h the deck transformation, then with a suitable choice of $\widehat{f}$ we have $S_n = \widehat{S}/(h^{2^n} \widehat{f})$ and f_n is lifted to h . Now consider a simple closed curve on a fundamental domain of $\widehat{S}$ which is not homologous to the boundary, such that the homology class c represented by this curve γ is preserved by $\widehat{f}$. The existence of such a homology class is due to the construction in Baik-Shin [BS20]. Then $\sum_{i=0}^{2^n-1} f_n^i c$ is invariant under f_n , and for $k < n$, let $c_k = \sum_{i=0}^{2^{n-k}-1} f_n^{i2^k} c$. Now $\text{Span}\{c_k, f_n c_k, \dots, f_n^{2^k-1} c_k\}$ is a 2^k dimensional invariant subspace of $f_n^{2^k}$. This proves Theorem 1.2.

Since the constant C in Theorem 1.2 does not depend on the choice of given g and k , we can apply the theorem inductively: At each j -th step with g_j and k_j , Theorem 1.2 applied to g_j and k_j gives $g' > g_j$, $k' > k_j$, and a pseudo-Anosov $f_{j+1} : S_{g'_j} \rightarrow S_{g'_j}$ with $\ell_C(f_{j+1}) \leq C k'_j / g_j'^2$. Then we set $g_{j+1} := g'_j$ and $k_{j+1} := k'_j$. As a consequence, we obtain the following corollary that interpolates $L_C(0, g)$ [GT11] and $L_C(2g, g)$ [BS20] in a partial way.

Corollary 3.1. *There is a constant C and a diverging sequence $(k_j, g_j) \rightarrow \infty$ as $j \rightarrow \infty$ such that*

$$L_C(k_j, g_j) \leq C \frac{k_j}{g_j^2}.$$

Corollary 3.1 can be regarded as an evidence for Conjecture 1.3 as it has a similar form to the desired asymptote. On the other hand, due to the inexplicit choice made in the proof of Theorem 1.2, it is hard to explicitly understand from which diverging sequence (k_j, g_j) we can deduce the desired asymptote. Hence it may require different approaches to make a concrete progress towards Conjecture 1.3.

However, pseudo-Anosov mapping classes we construct in the later section (Section 4) satisfy the asymptotes in Theorem 1.2 and Corollary 3.1.

4. PSEUDO-ANOSOVs WITH SPECIFIED INVARIANT HOMOLOGY DIMENSION

To the best of the authors' knowledge, asymptotics for $L_C(k, g)$ are known only when $k = 0$ (whole mapping class groups) and $k = 2g$ (Torelli groups). In this section, we construct pseudo-Anosovs $f_g \in \text{Mod}(S_g)$ with $m(f_g) = 2g - 1$ and realizing the asymptotic for $L_C(2g - 1, g)$.

From the definition of $L_C(k, g)$, $L_C(k, g) \leq L_C(k', g)$ if $k \leq k'$. Since $L_C(2g, g) \asymp 1/g$ from [BS20], the lower bound in Theorem 1.1, implies that $L_C(k, g) \asymp 1/g$ if k behaves like $2g$; for instance, $k \geq 2g - C$ for some constant $C > 0$. However, $L_C(0, g) \asymp 1/g^2$ by [GT11]. In this regard, we ask whether there is a sort of threshold for k that $L_C(k, g)$ becomes strictly smaller than $1/g$, such as $1/g^2$.

As a potential approach for this question, we think of constructing pseudo-Anosovs of specified maximal invariant homology dimensions on surfaces of large genera with small asymptotic translation lengths. In order to get pseudo-Anosov maps on surfaces of large genera, some previous results ([ALM16], [BSW21], [BKSW23]) employ a fixed hyperbolic mapping torus and consider its monodromy obtained from a fibered cone. Since the first betti number of a mapping torus of f is same as $m(f) + 1$, such monodromies in a fibered cone share the same m -value.

In contrast, we come up with a finite cyclic cover of a genus 2 surface to get the desired pseudo-Anosov maps on large genera surfaces as lifts of a fixed map. From the concrete estimation on how covering maps distort the distances on curve complexes [APT22], asymptotic translation length of the lift via degree g covering would be at least $1/g$, up to a constant multiple. We believe that constructing such lifts with specified maximal invariant homology dimensions would help to figure out the minimal $k = k(g)$ with $L_C(k, g) \asymp 1/g$.

We start with a non-separating curve α on the genus 2 surface S_2 , and take g copies of $S_2 \setminus \alpha$ for $g > 1$. Gluing two different copies of $S_2 \setminus \alpha$ along one boundary component in a cyclic way, we obtain the finite cyclic cover p_{g+1} of degree g as in Figure 1. Let us denote the resulting cover by S_{g+1} since it is of genus $g + 1$.




FIGURE 1. g -fold finite cyclic covering

This cover p_{g+1} corresponds to the kernel of the composed map

$$\pi_1(S_2) \xrightarrow{\hat{i}(\cdot, \alpha)} \mathbb{Z} \xrightarrow{\text{mod } g} \mathbb{Z}/g\mathbb{Z}$$

where $\hat{i}(\cdot, \cdot)$ stands for the algebraic intersection number. To see this, one can observe that an element of $\pi_1(S_2)$ can be lifted to $\pi_1(S_{g+1})$ via p_{g+1} if and only if its lift departs one copy of $S_2 \setminus \alpha$ and then returns to the same copy. If the lift departs and returns through the same boundary component

of $S_2 \setminus \alpha$, then the element of $\pi_1(S_2)$ has the algebraic intersection number 0 with α . Otherwise, if the lift departs and returns through different boundary components, then the algebraic intersection number is an integer multiple of g .

In [BS20], the first author and Shin directly constructed pseudo-Anosovs on S_g that are Torelli and of small asymptotic translation lengths on curve complexes. In the following, we construct pseudo-Anosovs with specific maximal invariant homology dimensions and satisfying the upper bound provided in Theorem 1.2 and Corollary 3.1. As a result, we obtain Theorem 1.5. Our strategy is to fix a suitable pseudo-Anosov on S_2 and then to lift it via p_{g+1} . Due to the symmetry of the covering, we can find a number of invariant homology classes proportional to the degree of the cover.

Proof of Theorem 1.5. The last assertion is a direct consequence of the first assertion and Theorem 1.1. By Theorem 1.1, there exists $C' > 0$ so that $L_{\mathcal{C}}(2g - 1, g) \geq \frac{C'}{g}$ for all $g > 1$. Hence, it remains to show the existence of the desired pseudo-Anosovs.

Fix $g > 1$ and in the rest of the proof, we simply denote by $p := p_{g+1}$. Let α be a non-separating curve on S_2 and let β be a separating curve on S_2 disjoint from α . Then each lift of β through p is also separating. For instance, see Figure 2.




FIGURE 2. A non-separating curve α and a separating curve β on S_2 with $\alpha \cap \beta = \emptyset$

Now let φ be a Torelli pseudo-Anosov mapping class in $\text{Mod}(S_2)$ such that $d_{\mathcal{C}}(\beta, \varphi\beta) \geq 3$. Note that this means that β and $\varphi\beta$ are separating simple closed curves on S_2 that fill the surface. Such a φ may be obtained as follows: let $\psi \in \text{Mod}(S_2)$ be a pseudo-Anosov mapping class. Since a pseudo-Anosov mapping class acts on $\mathcal{C}(S_2)$ as a loxodromic isometry by [MM99], $\lim_{n \rightarrow \infty} d_{\mathcal{C}}(\beta, \psi^n\beta)/n = \ell_{\mathcal{C}}(\psi) > 0$. Hence replacing ψ with ψ^n for some large n if necessary, we may assume that $d_{\mathcal{C}}(\beta, \psi\beta) \geq 3$. We then consider a mapping class $T_{\beta}T_{\psi\beta}^{-1}$ which is pseudo-Anosov by Thurston [Thu88] or Penner [Pen88]. Again, since $T_{\beta}T_{\psi\beta}^{-1}$ acts on $\mathcal{C}(S_2)$ as a loxodromic isometry, we can set φ to be some power of $T_{\beta}T_{\psi\beta}^{-1}$ so that φ is a pseudo-Anosov mapping class with the property that $d_{\mathcal{C}}(\beta, \varphi\beta) \geq 3$.

Now let $f = T_\beta T_{\varphi\beta}^{-1} T_{\varphi\alpha}^{-1}$. Since β , $\varphi\beta$, and $\varphi\alpha$ fill the surface and $\varphi\beta \cap \varphi\alpha = \emptyset$, f is pseudo-Anosov again by Thurston [Thu88] and Penner [Pen88]. Since β and $\varphi\beta$ are separating, T_β and $T_{\varphi\beta}^{-1}$ are Torelli, in particular, they preserve the homology class $[\alpha]$ of α . Furthermore, since φ is Torelli, $[\varphi\alpha] = [\alpha]$, which implies that $T_{\varphi\alpha}^{-1}$ also preserves $[\alpha]$. Hence, f preserves $[\alpha]$, and thus $\hat{i}(f(\cdot), \alpha) = \hat{i}(\cdot, \alpha)$. In particular, f preserves the kernel of $\pi_1(S_2) \xrightarrow{\hat{i}(\cdot, \alpha)} \mathbb{Z} \xrightarrow{\text{mod } g} \mathbb{Z}/g\mathbb{Z}$. Consequently, f can be lifted through p .

Let $\tilde{f} = T_{p^{-1}(\beta)} T_{p^{-1}(\varphi\beta)}^{-1} T_{p^{-1}(\varphi\alpha)}^{-1}$ be a lift of f via p . We now estimate $\ell_C(\tilde{f})$. Our strategy to obtain the desired upper bound for $\ell_C \tilde{F}$ is to find a simple closed curve $\tilde{\alpha}$ such that $\tilde{\alpha}$ and its image under a sufficiently high power of $\tilde{f}$ do not fill the surface, which means that they are within distance 2 in the curve complex. We do this by counting the number of intersections of images of $\tilde{\alpha}$ and lifts of subsurfaces.

Recall the construction of p : Take g copies $X_1, \dots, X_g$ of $S_2 \setminus \alpha$ and glue X_i and X_{i+1} along one of their boundary components. Throughout, we write each index i modulo g . Let $\tilde{\alpha} = \partial X_0 \cap \partial X_1$. That is, $\tilde{\alpha}$ be a boundary component of X_0 and X_1 where they are glued. Due to the construction, $\tilde{\alpha}$ is a lift of α .

Noting that $\hat{i}(\varphi\alpha, \alpha) = 0$ since φ is Torelli, we get

$$(4.1) \quad T_{p^{-1}(\varphi\alpha)}^{-1} \tilde{\alpha} \subseteq \bigcup_{j=-i(\varphi\alpha, \alpha)/2}^{i(\varphi\alpha, \alpha)/2} X_j$$

where $i(\cdot, \cdot)$ is the geometric intersection number. (cf. Figure 3) Similarly, $\hat{i}(\varphi\beta, \alpha) = 0$ and

$$T_{p^{-1}(\varphi\beta)}^{-1} T_{p^{-1}(\varphi\alpha)}^{-1} \tilde{\alpha} \subseteq \bigcup_{j=-\frac{i(\varphi\beta, \alpha) + i(\varphi\alpha, \alpha)}{2}}^{\frac{i(\varphi\beta, \alpha) + i(\varphi\alpha, \alpha)}{2}} X_j.$$

Since $T_{p^{-1}(\beta)}$ fixes each X_j , we have

$$\tilde{f} \tilde{\alpha} \subseteq \bigcup_{j=-\frac{i(\varphi\beta, \alpha) + i(\varphi\alpha, \alpha)}{2}}^{\frac{i(\varphi\beta, \alpha) + i(\varphi\alpha, \alpha)}{2}} X_j.$$

Conducting this procedure inductively, we finally get

$$\tilde{f}^n \tilde{\alpha} \subseteq \bigcup_{j=-n \cdot \frac{i(\varphi\beta, \alpha) + i(\varphi\alpha, \alpha)}{2}}^{n \cdot \frac{i(\varphi\beta, \alpha) + i(\varphi\alpha, \alpha)}{2}} X_j.$$

Hence, there exists $\tilde{j}$ such that $\tilde{f}^{\lfloor \frac{g-2}{i(\varphi\beta, \alpha) + i(\varphi\alpha, \alpha)} \rfloor} \tilde{\alpha} \cap X_{\tilde{j}} = \emptyset$. Since there exists an essential simple closed curve in $X_{\tilde{j}}$ which is a 2-holed torus, we have




FIGURE 3. A curve intersecting a lift of the new curve (non-indexed one) in some X_j spreads into $X_{j-1} \cup X_{j+1}$ by twisting along the lift. It describes how the image of $\tilde{\alpha}$ under multi-twists is trapped in the certain number of lifts of a subsurface, as in (4.1).

$d_C \left(\tilde{\alpha}, \tilde{f}^{\lfloor \frac{g-2}{i(\varphi\beta, \alpha) + i(\varphi\alpha, \alpha)} \rfloor} \tilde{\alpha} \right) \leq 2$ so $\ell_C \left(\tilde{f}^{\lfloor \frac{g-2}{i(\varphi\beta, \alpha) + i(\varphi\alpha, \alpha)} \rfloor} \right) \leq 2$. This implies the following estimate. Note that φ , $i(\varphi\beta, \alpha)$, and $i(\varphi\alpha, \alpha)$ are universal quantities independent on p and g .

$$(4.2) \quad \ell_C(\tilde{f}) \leq \frac{2}{\left\lfloor \frac{g-2}{i(\varphi\beta, \alpha) + i(\varphi\alpha, \alpha)} \right\rfloor}$$

We now show that $m(\tilde{f}) = 2g + 1$. Recall that $\beta \subseteq S_2$ is a separating curve and $\alpha \subseteq S_2$ is a non-separating curve disjoint from β . Temporarily let us denote Y be a component of $S_2 \setminus \beta$ that does not contain α . Let γ and δ be non-separating simple closed curves whose homology classes form a basis for $H_1(Y) \cong \mathbb{R}^2$. Let us also denote η be a non-separating curve on $S_2 \setminus \beta$ with $i(\eta, \alpha) = 1$. Then $p^{-1}(\gamma)$, $p^{-1}(\delta)$, $p^{-1}(\eta)$, and one component of $p^{-1}(\alpha)$ form a basis for $H_1(S_{g+1}) \cong \mathbb{R}^{2g+2}$. See Figure 4 for instance.




FIGURE 4. Basis for $H_1(S_{g+1})$

Let $\tilde{\gamma}_j = p^{-1}(\gamma) \cap X_j$, $\tilde{\delta}_j = p^{-1}(\delta) \cap X_j$, and $\tilde{\eta} = p^{-1}(\eta)$. Further, let $\tilde{\alpha} = \partial X_0 \cap \partial X_1$ which is a component of $p^{-1}(\alpha)$. Since φ is Torelli, it has a lift $\tilde{\varphi}$ through p . Hence, homology classes $\left\{ [\tilde{\varphi}\tilde{\gamma}_j], [\tilde{\varphi}\tilde{\delta}_j], [\tilde{\varphi}\tilde{\eta}], [\tilde{\varphi}\tilde{\alpha}] \right\}$ also form a basis for $H_1(S_{g+1})$.

Recall that $\tilde{f} = T_{p^{-1}(\beta)} T_{p^{-1}(\varphi\beta)}^{-1} T_{p^{-1}(\varphi\alpha)}^{-1}$. Since $\gamma \cap (\alpha \cup \beta) = \emptyset$, we have $\tilde{\varphi}\tilde{\gamma}_j \cap (p^{-1}(\varphi\alpha) \cap p^{-1}(\varphi\beta)) = \emptyset$. Here, note that $\tilde{\varphi}\tilde{\gamma}_j$ is a lift of $\varphi\gamma$ which is a component of $p^{-1}(\varphi\gamma)$. Hence it follows that $[\tilde{f}\tilde{\varphi}\tilde{\gamma}_j] = [T_{p^{-1}(\beta)}\tilde{\varphi}\tilde{\gamma}_j]$. Since each component of $p^{-1}(\beta)$, which is a lift of β , is separating, $T_{p^{-1}(\beta)}$ is Torelli. As a result, $[\tilde{f}\tilde{\varphi}\tilde{\gamma}_j] = [\tilde{\varphi}\tilde{\gamma}_j]$. Similarly, we have $[\tilde{f}\tilde{\varphi}\tilde{\delta}_j] = [\tilde{\varphi}\tilde{\delta}_j]$.

Now we consider $[\tilde{f}\tilde{\varphi}\tilde{\alpha}]$. As $\tilde{\varphi}\tilde{\alpha}$ is a lift of $\varphi\alpha$, $T_{p^{-1}(\varphi\alpha)}^{-1}\tilde{\varphi}\tilde{\alpha} = \tilde{\varphi}\tilde{\alpha}$. Furthermore, since $\alpha \cap \beta = \emptyset$, $\tilde{\varphi}\tilde{\alpha}$, a lift of $\varphi\alpha$, does not intersect $p^{-1}(\varphi\beta)$. It implies that $T_{p^{-1}(\varphi\beta)}^{-1}\tilde{\varphi}\tilde{\alpha} = \tilde{\varphi}\tilde{\alpha}$. Finally, since $T_{p^{-1}(\beta)}$ is Torelli again, we conclude $[\tilde{f}\tilde{\varphi}\tilde{\alpha}] = [\tilde{\varphi}\tilde{\alpha}]$.

So far, we have proved $m(\tilde{f}) \geq 2g + 1$. Suppose to the contrary that $m(\tilde{f}) = 2g + 2$, which means that $\tilde{f}$ is Torelli. Then $\tilde{f}\tilde{\varphi}\tilde{\eta}$ should be homologous to $\tilde{\varphi}\tilde{\eta}$. It implies $[T_{p^{-1}(\varphi\beta)}^{-1} T_{p^{-1}(\varphi\alpha)}^{-1} \tilde{\varphi}\tilde{\eta}] = [\tilde{\varphi}\tilde{\eta}]$ since $T_{p^{-1}(\beta)}$ is Torelli. As any two components of $p^{-1}(\varphi\alpha)$ bound a subsurface, they are homologous. In particular, as $\tilde{\varphi}\tilde{\alpha}$ is a component of $p^{-1}(\varphi\alpha)$, each of its components is homologous to $\tilde{\varphi}\tilde{\alpha}$. Hence, $[T_{p^{-1}(\varphi\beta)}^{-1} T_{p^{-1}(\varphi\alpha)}^{-1} \tilde{\varphi}\tilde{\eta}] = [T_{p^{-1}(\varphi\beta)}^{-1} T_{\tilde{\varphi}\tilde{\alpha}}^{-g} \tilde{\varphi}\tilde{\eta}]$. Noting that $T_{\tilde{\alpha}}^{-g}\tilde{\eta}$ can be isotoped into arbitrary neighborhood of $\tilde{\alpha} \cup \tilde{\eta}$, $T_{\tilde{\varphi}\tilde{\alpha}}^{-g}\tilde{\varphi}\tilde{\eta}$ can also be isotoped into arbitrary neighborhood of $\tilde{\varphi}\tilde{\alpha} \cup \tilde{\varphi}\tilde{\eta}$. Since $\tilde{\varphi}\tilde{\alpha} \cup \tilde{\varphi}\tilde{\eta}$ and $p^{-1}(\varphi\beta)$ are disjoint compact sets, we have $T_{p^{-1}(\varphi\beta)}^{-1} T_{\tilde{\varphi}\tilde{\alpha}}^{-g} \tilde{\varphi}\tilde{\eta} = T_{\tilde{\varphi}\tilde{\alpha}}^{-g} \tilde{\varphi}\tilde{\eta}$. Summing up the above argument, we obtain

$$[\tilde{\varphi}\tilde{\eta}] = [\tilde{f}\tilde{\varphi}\tilde{\eta}] = [T_{p^{-1}(\varphi\beta)}^{-1} T_{p^{-1}(\varphi\alpha)}^{-1} \tilde{\varphi}\tilde{\eta}] = [T_{p^{-1}(\varphi\beta)}^{-1} T_{\tilde{\varphi}\tilde{\alpha}}^{-g} \tilde{\varphi}\tilde{\eta}] = [T_{\tilde{\varphi}\tilde{\alpha}}^{-g} \tilde{\varphi}\tilde{\eta}]$$

where the first equality is the assumption. However,

$$[T_{\tilde{\varphi}\tilde{\alpha}}^{-g} \tilde{\varphi}\tilde{\eta}] = [\tilde{\varphi}\tilde{\eta}] - g \cdot \hat{i}(\tilde{\varphi}\tilde{\eta}, \tilde{\varphi}\tilde{\alpha}) [\tilde{\varphi}\tilde{\alpha}]$$

which implies that $\hat{i}(\tilde{\varphi}\tilde{\eta}, \tilde{\varphi}\tilde{\alpha}) = 0$. It contradicts our choice of η that $i(\tilde{\eta}, \tilde{\alpha}) = 1$. Therefore, $m(\tilde{f}) = 2g + 1$. Setting $f_{g+1} = \tilde{f}$ completes the proof of Theorem 1.5.

The lower bound on $\ell_{\mathcal{C}}(f_g)$ for f_g constructed in the proof can also be calculated in a concrete way by Aougab-Patel-Taylor [APT22] as follows.

$$\frac{\ell_{\mathcal{C}}(f)}{(g-1) \cdot 80 \cdot 2^{13} e^{54} \pi} \leq \ell_{\mathcal{C}}(f_g)$$

Remark 4.1. *In the proof, all figures describe one specific example. Any choice of α , β , γ , δ , and η works if it satisfies the condition we provide. That is,*

- α and β are non-separating and separating curves on S_2 respectively, and are disjoint;

- γ and δ are non-separating simple closed curves that form a basis for the first homology group of the component of $S_2 \setminus \beta$ disjoint from α ;
- η is a non-separating curve on $S_2 \setminus \beta$ with $i(\eta, \alpha) = 1$.

Furthermore, if we modify the map on S_2 to be $f = T_\beta T_{\varphi\beta}^{-1}$, then its lift via p_{g+1} is Torelli, which gives another proof of $L_C(2g, g) \asymp \frac{1}{g}$.

5. SMALL TRANSLATION LENGTH AND NORMAL GENERATION

In this section, we discuss pseudo-Anosov mapping classes with small asymptotic translation lengths and normal generation of mapping class groups. For a general group G and $g \in G$, the *normal closure* $\langle\langle g \rangle\rangle$ of g is the smallest normal subgroup of G containing g . The normal closure can be described in various way:

$$\langle\langle g \rangle\rangle = \bigcap_{g \in N \trianglelefteq G} N = \langle hgh^{-1} : h \in G \rangle$$

In a particular case that $\langle\langle g \rangle\rangle = G$, we say g *normally generates* G , and g is said to be a *normal generator* of G .

Normal generators of mapping class groups of surfaces have been studied by various authors. In [Lon86], Long asked whether there is a pseudo-Anosov that normally generates a mapping class group. This question recently answered affirmatively by Lanier and Margalit in [LM22]. Indeed, they showed that there is a universal constant so that pseudo-Anosovs with stretch factors less than the constant should be normal generators. Then the asymptotic $L_{\mathcal{T}}(0, g) \asymp 1/g$ by Penner [Pen91] deduces the answer. Precisely, Lanier and Margalit proved the following.

Theorem 5.1 (Lanier-Margalit [LM22]). *If a pseudo-Anosov $\varphi \in \text{Mod}(S_g)$ has the stretch factor less than $\sqrt{2}$, then φ normally generates $\text{Mod}(S_g)$.*

As the logarithm of stretch factor of a pseudo-Anosov equals to the translation length of the pseudo-Anosov on the Teichmüller space, Lanier and Margalit's result also means that the small translation length on the Teichmüller space implies the normal generation of the mapping class group. One natural question in this philosophy is whether the same holds in the circumstance of curve complexes. There are several ways to formalize this question:

- (1) Is there a universal constant $C > 0$ so that if a pseudo-Anosov $\varphi \in \text{Mod}(S_g)$ has $\ell_C(\varphi) < C/g$ then $\langle\langle \varphi \rangle\rangle = \text{Mod}(S_g)$?
- (2) Is there a universal constant $C > 0$ so that if a non-Torelli pseudo-Anosov $\varphi \in \text{Mod}(S_g)$ has $\ell_C(\varphi) < C/g$ then $\langle\langle \varphi \rangle\rangle = \text{Mod}(S_g)$?

Indeed, the first and the third authors of current paper, Kin and Shin stated (1) in [BKSW23, Question 1.2].

Remark 5.2. *In the above questions, the factor $1/g$ is necessary since $L_{\mathcal{C}}(2g, g) \asymp 1/g$ [BS20] and due to Proposition 1.6. Furthermore, we separately state above two questions in order to forbid the trivial (Torelli) case in (2) and deal with the same problem.*

Proof of Theorem 1.6. The family of pseudo-Anosovs constructed in Theorem 1.5 actually consists of non-normal generators, that is, $\langle\langle f_g \rangle\rangle \neq \text{Mod}(S_g)$. To see this, recall that $f_g = T_{p_g^{-1}(\beta)} T_{p_g^{-1}(\varphi\beta)}^{-1} T_{p_g^{-1}(\varphi\alpha)}^{-1}$. It can be rewritten as

$$f_g = T_{p_g^{-1}(\beta)} \left(\tilde{\varphi} T_{p_g^{-1}(\beta)}^{-1} \tilde{\varphi}^{-1} \right) \left(\tilde{\varphi} T_{p_g^{-1}(\alpha)}^{-1} \tilde{\varphi}^{-1} \right).$$

Hence, it follows that $\langle\langle f_g \rangle\rangle \leq \langle\langle T_{p_g^{-1}(\beta)}, T_{p_g^{-1}(\alpha)} \rangle\rangle$ where the right-hand-side means the smallest normal subgroup containing $T_{p_g^{-1}(\beta)}$ and $T_{p_g^{-1}(\alpha)}$.

Since each component of $p_g^{-1}(\beta)$ is separating, $T_{p_g^{-1}(\beta)}$ is Torelli, namely, contained in the kernel of the symplectic representation $\text{Mod}(S_g) \rightarrow \text{Sp}(2g, \mathbb{Z})$. Moreover, any two components of $p_g^{-1}(\alpha)$ bound an essential subsurface so they are homologous, which means that $T_{p_g^{-1}(\alpha)}$ acts samely as $T_{\tilde{\alpha}}^{g-1}$ on $H_1(S_g; \mathbb{Z})$. As such, $T_{p_g^{-1}(\alpha)}$ acts trivially on the mod $(g-1)$ homology $H_1(S_g, \mathbb{Z}/(g-1)\mathbb{Z})$. Hence, we have that the symplectic representation of $T_{p_g^{-1}(\alpha)}$ is contained in the kernel of $\text{Sp}(2g, \mathbb{Z}) \rightarrow \text{Sp}(2g, \mathbb{Z}/(g-1)\mathbb{Z})$. Consequently, the normal closure $\langle\langle T_{p_g^{-1}(\beta)}, T_{p_g^{-1}(\alpha)} \rangle\rangle$ is contained in the kernel of the composition

$$\text{Mod}(S_g) \rightarrow \text{Sp}(2g, \mathbb{Z}) \rightarrow \text{Sp}(2g, \mathbb{Z}/(g-1)\mathbb{Z})$$

which is surjective. It follows that

$$\langle\langle f_g \rangle\rangle \leq \langle\langle T_{p_g^{-1}(\beta)}, T_{p_g^{-1}(\alpha)} \rangle\rangle \neq \text{Mod}(S_g)$$

so f_g is not a normal generator as desired.

Note that we have a concrete upper bound for $\ell_{\mathcal{C}}(f_g)$ in (4.2):

$$\ell_{\mathcal{C}}(f_g) \leq \frac{2}{\left\lfloor \frac{g-3}{i(\varphi\beta, \alpha) + i(\varphi\alpha, \alpha)} \right\rfloor} \leq \frac{2(i(\varphi\beta, \alpha) + i(\varphi\alpha, \alpha))}{g-3 - (i(\varphi\beta, \alpha) + i(\varphi\alpha, \alpha))}$$

Hence, once we fix α , β , and φ , we get a quantitative restriction on the constant C in the above questions. For instance, we can consider the configuration as in Figure 5.




FIGURE 5. β and ξ fill the surface

Let $\lambda = T_\xi \beta$. As β and ξ fill the surface S_2 , β and $\lambda = T_\xi \beta$ also fill the surface. Since β is separating, $\lambda = T_\xi \beta$ is also separating. Hence, due to Penner [Pen88] or Thurston [Thu88], $\varphi = T_\lambda T_\beta^{-1}$ is a Torelli pseudo-Anosov. Furthermore, it follows that β and $\varphi\beta$ also fill the surface. Therefore, we can construct f_g as in Theorem 1.5 starting with α , β , and φ depicted above.

Since $i(\xi, \beta) = 6$, $i(\lambda, \beta) = i(T_\xi \beta, \beta) = i(\xi, \beta)^2 = 36$ by [FM11, Proposition 3.2]. Now from $\varphi\alpha = T_\lambda \alpha$ and $\varphi\beta = T_\lambda \beta$, we have:

$$\begin{aligned} i(\varphi\alpha, \alpha) &= i(T_\lambda \alpha, \alpha) = i(\lambda, \alpha)^2 &= 144 \\ i(\varphi\beta, \alpha) &= i(T_\lambda \beta, \alpha) = i(\lambda, \beta)i(\lambda, \alpha) &= 432 \end{aligned}$$

Hence, for the resulting f_g ,

$$\ell_C(f_g) \leq \frac{1152}{g - 579}$$

for $g > 579$. Consequently, we conclude Theorem 1.6.

REFERENCES

- [ALM16] Ian Agol, Christopher J Leininger, and Dan Margalit. Pseudo-anosov stretch factors and homology of mapping tori. *Journal of the London Mathematical Society*, 93(3):664–682, 2016.
- [APT22] Tarik Aougab, Priyam Patel, and Samuel J Taylor. Covers of surfaces, kleinian groups and the curve complex. *Journal of Topology*, 15(4):1833–1863, 2022.
- [B⁺78] Lipman Bers et al. An extremal problem for quasiconformal mappings and a theorem by thurston. *Acta Mathematica*, 141:73–98, 1978.
- [BKSW23] Hyungryul Baik, Eiko Kin, Hyunshik Shin, and Chenxi Wu. Asymptotic translation lengths and normal generations of pseudo-anosov monodromies for fibered 3-manifolds. *Algebraic & Geometric Topology*, 23(3):1363–1398, 2023.
- [BS20] Hyungryul Baik and Hyunshik Shin. Minimal asymptotic translation lengths of torelli groups and pure braid groups on the curve graph. *International Mathematics Research Notices*, 2020(24):9974–9987, 2020.
- [BSW21] Hyungyul Baik, Hyunshik Shin, and Chenxi Wu. An upper bound on the asymptotic translation lengths on the curve graph and fibered faces. *Indiana Univ. Math. J.*, 70(4):1625–1637, 2021.
- [Ell] Jordan Ellenberg. Pseudo-anosov puzzle 2: homology rank and dilatation. Quomodocumque, <http://quomodocumque.wordpress.com/2010/04/26>, April 26, 2010.
- [FLM08] Benson Farb, Christopher J Leininger, and Dan Margalit. The lower central series and pseudo-anosov dilatations. *American journal of mathematics*, 130(3):799–827, 2008.
- [FLP12] Albert Fathi, François Laudenbach, and Valentin Poénaru. *Thurston’s Work on Surfaces (MN-48)*, volume 48. Princeton University Press, 2012.
- [FM11] Benson Farb and Dan Margalit. *A primer on mapping class groups (pms-49)*. Princeton university press, 2011.
- [GHKL13] V. Gadre, E. Hironaka, A. E. Kent, and C. J. Leininger. Lipschitz constants to curve complexes. *Math. Res. Lett.*, 20(4):647–656, 2013.
- [GT11] Vaibhav Gadre and Chia-Yen Tsai. Minimal pseudo-Anosov translation lengths on the complex of curves. *Geom. Topol.*, 15(3):1297–1312, 2011.

- [KK92] LA Kondratyuk and MI Krivoruchenko. Superconducting quark matter in $su(2)$ colour group. *Zeitschrift für Physik A Hadrons and Nuclei*, 344(1):99–115, 1992.
- [LM22] Justin Lanier and Dan Margalit. Normal generators for mapping class groups are abundant. *Commentarii Mathematici Helvetici*, 97(1):1–59, 2022.
- [Lon86] DD Long. A note on the normal subgroups of mapping class groups. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 99, pages 79–87. Cambridge University Press, 1986.
- [MM99] Howard A. Masur and Yair N. Minsky. Geometry of the complex of curves I: Hyperbolicity. *Invent. Math.*, 138(1):103–149, 1999.
- [Pen88] Robert C Penner. A construction of pseudo-anosov homeomorphisms. *Transactions of the American Mathematical Society*, 310(1):179–197, 1988.
- [Pen91] Robert C Penner. Bounds on least dilatations. *Proceedings of the American Mathematical Society*, 113(2):443–450, 1991.
- [Thu88] William P Thurston. On the geometry and dynamics of diffeomorphisms of surfaces. *Bulletin (new series) of the american mathematical society*, 19(2):417–431, 1988.
- [Thu98] William P Thurston. Hyperbolic structures on 3-manifolds, ii: Surface groups and 3-manifolds which fiber over the circle. *arXiv preprint math/9801045*, 1998.
- [Tsa09] Chia-Yen Tsai. The asymptotic behavior of least pseudo-Anosov dilatations. *Geom. Topol.*, 13(4):2253–2278, 2009.

DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, 291 DAEHAK-RO YUSEONG-GU, DAEJEON, 34141, SOUTH KOREA

Email address: hrbaik@kaist.ac.kr

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, 219 PROSPECT STREET, NEW HAVEN, CT 06511, USA

Email address: dongryul.kim@yale.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN–MADISON, 480 LINCOLN DRIVE, MADISON, WI 53706, USA

Email address: cwu367@math.wisc.edu