

Ising Model Close to $d = 2$

Wenliang Li^{1,2,*}

¹*School of Physics, Sun Yat-Sen University, Guangzhou 510275, China*

²*Okinawa Institute of Science and Technology Graduate University, 1919-1 Tancha, Onna-son, Okinawa 904-0495, Japan*

The $d = 2$ critical Ising model is described by an exactly solvable Conformal Field Theory (CFT). The deformation to $d = 2 + \epsilon$ is a relatively simple system at strong coupling outside of even dimensions. Using novel numerical and analytical conformal bootstrap methods in Lorentzian signature, we show that the leading corrections to the Ising data are more singular than ϵ . There must be infinitely many new states due to the d -dependence of conformal symmetry. The linear independence of conformal blocks is central to this bootstrap approach, which can be extended to more rigorous studies of non-positive systems, such as non-unitary, defect/boundary and thermal CFTs.

INTRODUCTION

The d -dimensional Ising model is a fundamental model in statistical physics and condensed matter physics. Historically, it was proposed by Lenz to describe ferromagnetism and the case of $d = 1$ was solved by Ising. This simple model displays rich physics and captures some main traits of phase transitions and many-body problems. At criticality, it belongs to one of the simplest universality classes, characterized by the global \mathbb{Z}_2 symmetry. For $d > 4$, the critical behaviour of the Ising model is described by Landau's mean-field theory [1], in which fluctuations are neglected due to the averaging effects of many adjacent spins. At lower d , fluctuations play a more significant role. The mean-field description is not sufficient for $d < 4$ and the Ising critical exponents have non-trivial d -dependence [2]. As a natural continuation of Landau's theory, Wilson and Fisher calculated the critical exponents in $d = 4 - \epsilon$ dimensions using the perturbative ϵ expansion [4]. The ϵ expansion has proved to be a valuable tool in the studies of critical phenomena [5, 6].

At $d = 2$, it is well-known that the Ising model is solvable since Onsager's groundbreaking results [7]. The critical behaviour is described by the fixed point of renormalization group flows. In particular, scale invariance of the fixed point is promoted to conformal invariance. As another natural continuation, it would be interesting to deform the $2d$ exact solution to $d = 2 + \epsilon$ dimensions. The ϵ expansion usually concerns weakly coupled systems [8–11], but the case here remains strongly coupled [12], so the intriguing strong coupling physics becomes more manifest. More recently, the $\epsilon = d - 2$ expansion has also been used to study deconfined quantum criticality [13, 14]. (See [15] for a numerical conformal bootstrap study.) We notice a deceptively simple question:

Is the ϵ expansion of a strongly coupled system given by integer power series?

In the standard ϵ expansion, the corrections to the $d = 4$ data can be computed order by order in ϵ , given by asymptotic series [16]. It has been argued that they are integer power series based on the Renormalization Group (RG) analysis in the minimal subtraction scheme [17, 18]. Naively, one might think that the $\epsilon = d - 2$ expansion should also be the case. For instance, the scaling dimension of the lowest \mathbb{Z}_2 -even operator

was assumed to receive integer power corrections in the study of disorder effects in $2 + \epsilon$ dimensions [19]. However, the standard ϵ expansion is around a Gaussian theory. The weak coupling techniques and arguments do not easily extend to the strong coupling situation. For the $O(n)$ model, Cardy and Hamber performed an elegant analysis around $n = d = 2$ based on some analyticity assumptions on the RG equations [20], but these results do not apply for $d = 2 + \epsilon$ with $n < 2$.

In this letter, we will study the $d = 2 + \epsilon$ Ising model using the conformal bootstrap. The conformal bootstrap program aims to classify and solve CFTs by general principles and consistency conditions [21, 22], without resorting to the weak coupling expansion. For $d = 2$, conformal symmetry becomes infinite-dimensional and this program can be carried out rather successfully [23, 24]. The studies in $d > 2$ dimensions are more challenging as conformal symmetry is less constraining. Nevertheless, considerable progress has been achieved due to the seminal work [25], in which the unitarity assumption and the crossing equations are formulated as inequalities. This modern bootstrap approach has led to rigorous bounds on the space of unitary CFTs, such as the most precise determinations of the $d = 3$ Ising critical exponents [26–29]. We refer to [30–34] for useful reviews and lecture notes.

The critical Ising model can be viewed as a continuous family of \mathbb{Z}_2 -covariant CFTs parametrized by d . The case of non-integer d has also been studied by the unitary bootstrap methods in [35–37]. The bounds exhibit similar features as those at $d = 2, 3$ and the results are consistent with the $(4 - d)$ expansion. However, a subtlety is that the Wilson-Fisher fixed point is non-unitary in non-integer dimensions, because the spectrum contains descendant states of complex scaling dimensions [38]. It would be helpful to consider complementary approaches that are not based on unitarity, such as the flow method [39] and the truncation method [40]. The truncated bootstrap approach has been applied to the study of non-positive problems [41–52]. In the original formulation [40], the truncated problem is encoded in determinants. In [53], we proposed some new ingredients, which we believe are important to a more systematic formulation. We emphasized the essential role of linear independence and introduced the concept of norm to the truncation approach. (See [54, 55] for

the recent implementation using reinforcement-learning algorithms.) We will apply these notions to the numerical bootstrap study of the $d = 2 + \epsilon$ Ising CFT.

On the other hand, it was noticed in [37] that the tentative spectrum from the unitary numerical approach exhibits a transition at $d = 2 + \epsilon$ with $\epsilon \sim 0.2$ small but finite. Such a transition is expected since $d = 2$ is special. At $d = 2$, the spectrum is organized into Virasoro multiplets and the corresponding Regge trajectories have constant twists $\tau = \Delta - \ell$ with integer spacing. At $d = 3$, the twist spectrum of the Ising CFT is additive and the Regge trajectories have more interesting dependence on spin. Infinitely many high spin operators have twists asymptotic to the sum of two lower twists [56, 57]. For example, the Regge trajectories $[\sigma\sigma]_n$ are associated with the lowest \mathbb{Z}_2 -odd scalar σ and they have twist accumulation points at $2\Delta_\sigma + 2n$. We will discuss the location of the transition to the double-twist spectrum using analytic bootstrap techniques.

To address our question, we will study the 4-point function of the lowest \mathbb{Z}_2 -odd scalar operator σ . We focus on the leading corrections and assume:

1. The critical Ising model is conformally invariant in $d = 2 + \epsilon$ dimensions with $|\epsilon| \ll 1$.
2. The leading corrections to the $2d$ data are linear in ϵ :

$$\Delta_\sigma = \Delta_\sigma^{(0)} + \epsilon \Delta_\sigma^{(1)} + \dots, \quad (1)$$

$$\Delta_i = \Delta_i^{(0)} + \epsilon \Delta_i^{(1)} + \dots, \quad (2)$$

$$\lambda_i = \lambda_i^{(0)} + \epsilon \lambda_i^{(1)} + \dots, \quad (3)$$

where $\Delta_i = \Delta_{\mathcal{O}_i}$ and $\lambda_i = \lambda_{\sigma\sigma\mathcal{O}_i}$ are the scaling dimension and OPE coefficient of \mathcal{O}_i . The zeroth order values can be derived from the exact solution at $d = 2$. We will further assume that $\Delta_i^{(1)}$, $\lambda_i^{(1)}$ do not grow too rapidly with $\Delta_i^{(0)}$, so the conformal block summation is convergent [58].

In the first assumption, scale invariance of the Ising fixed point is enhanced to conformal invariance. There is ample evidence for conformal invariance in $d = 2, 3, 4 - \epsilon$ dimensions, so we expect that this property extends to $d = 2 + \epsilon$ [59]. In the second assumption, the leading corrections cannot be more singular since they have positive integer powers. They cannot start from second or higher orders in ϵ because the d -dependence of conformal blocks leads to first-order terms in the crossing equation.

Below we will examine if this is a consistent scenario. It turns out that the assumptions 1 and 2 are not consistent, so the corrections are expected to be more singular than ϵ^1 .

THE CROSSING EQUATION

We consider the 4-point function of the lowest \mathbb{Z}_2 -odd operator σ :

$$\langle \sigma(x_1) \sigma(x_2) \sigma(x_3) \sigma(x_4) \rangle = \frac{\mathcal{G}(z, \bar{z})}{x_{12}^{2\Delta_\sigma} x_{34}^{2\Delta_\sigma}}. \quad (4)$$

The conformally invariant cross-ratios are

$$u = z\bar{z} = \frac{x_{12}^2 x_{34}^2}{x_{13}^2 x_{24}^2}, \quad v = (1-z)(1-\bar{z}) = \frac{x_{14}^2 x_{23}^2}{x_{13}^2 x_{24}^2}. \quad (5)$$

The crossing equation for $\mathcal{G}(z, \bar{z})$ reads:

$$v^{\Delta_\sigma} \mathcal{G}(z, \bar{z}) = u^{\Delta_\sigma} \mathcal{G}(1 - \bar{z}, 1 - z). \quad (6)$$

In the $\epsilon = d - 2$ expansion, we have

$$\mathcal{G}(z, \bar{z}) = \mathcal{G}^{(0)}(z, \bar{z}) + \epsilon \mathcal{G}^{(1)}(z, \bar{z}) + \dots, \quad (7)$$

where the 2d solution reads

$$\mathcal{G}^{(0)}(z, \bar{z}) = \frac{\sqrt{1 + \sqrt{u} + \sqrt{v}}}{\sqrt{2} v^{1/8}}, \quad (8)$$

and $\mathcal{G}^{(1)}(z, \bar{z})$ can be written as convergent power series in z, \bar{z} in the regime $0 \leq z, \bar{z} < 1$. After the conformal block decomposition, the crossing equation becomes

$$\sum_i \lambda_i^2 F_i(z, \bar{z}) = 0, \quad (9)$$

where $F_i(z, \bar{z}) = v^{\Delta_\sigma} G_{\Delta_i, \ell_i}(z, \bar{z}) - (z \leftrightarrow 1 - \bar{z})$ and G_{Δ_i, ℓ_i} is the global conformal block for the conformal multiplet labelled by the primary operator \mathcal{O}_i . To first order in ϵ , the crossing equation reads

$$\begin{aligned} & \sum_i \lambda_i^{(0)} (\lambda_i^{(0)} \Delta_\sigma^{(1)} \partial_{\Delta_\sigma} + \lambda_i^{(0)} \Delta_i^{(1)} \partial_{\Delta_i} + 2 \lambda_i^{(1)}) F_i(z, \bar{z}) \\ & = (-1) \sum_i \lambda_i^2 \partial_d F_i(z, \bar{z}), \end{aligned} \quad (10)$$

which will be written more compactly in (13). Note that the derivative ∂_d extracts the d -dependence of $G_{\Delta, \ell}$. After taking the derivatives, we set $\{d, \Delta_i, \lambda_i\} \rightarrow \{2, \Delta_i^{(0)}, \lambda_i^{(0)}\}$. We do not make any assumptions about the signs of $\{\Delta_\sigma^{(1)}, \Delta_i^{(1)}, \lambda_i^{(1)}\}$.

Although the left hand side of (10) involves an infinite number of free parameters, the building blocks are the simple $2d$ conformal blocks [71]

$$G_{\Delta, \ell}^{d=2}(z, \bar{z}) = \frac{1}{1 + \delta_{\ell, 0}} \left(k_{\Delta + \ell}(z) k_{\Delta - \ell}(\bar{z}) + (z \leftrightarrow \bar{z}) \right), \quad (11)$$

where $k_\beta(x) = x^{\beta/2} {}_2F_1(\beta/2, \beta/2, \beta, x)$ is the $SL(2, \mathbb{R})$ block with identical external scaling dimensions. Since each term is multiplied by $\lambda_i^{(0)}$, the intermediate states are the same as those in 2d and their twists are given by

$$\{\tau_i^{(0)}\} = \{4n, 4n + 1\}, \quad (12)$$

where $n = 0, 1, 2, \dots$ but $\tau^{(0)} \neq 5$. Note that the twist-5 trajectory and the twist-1 spin-2 state are absent in the 2d intermediate spectrum of the $\sigma \times \sigma$ OPE. New states cannot contribute to the OPE at order ϵ^1 because their squared OPE coefficients are at least of order ϵ^2 . This applies to both primary and descendant states [78]. Since the 2d Ising model is unitary, we do not need to worry about potential cancellation of finite mixed contributions. On the contrary, the right hand side of (10) has no free parameter. We can compute the sum based on the 2d data using the general d formula of conformal blocks [79]. Then we take the d derivative and set $d \rightarrow 2$.

In the “bra-ket” notation, the crossing equation (10) reads

$$\Delta_\sigma^{(1)}|\Delta_\sigma\rangle + \sum_i \left(\Delta_i^{(1)}|\Delta_i\rangle + \lambda_i^{(1)}|\lambda_i\rangle \right) = -|d\rangle, \quad (13)$$

where $|a\rangle$ denotes the contribution generated by the change in a . Our question in the introduction becomes: *Do $|\Delta_\sigma\rangle, |\Delta_i\rangle, |\lambda_i\rangle$ form a complete set of basis for $|d\rangle$?* It turns out that the answer is negative [80]! The target $|d\rangle$ does not belong to the vector space spanned by $\{|\Delta_\sigma\rangle, |\Delta_i\rangle, |\lambda_i\rangle\}$.

Before analyzing the crossing equation (13), let us discuss the building blocks $|\Delta_\sigma\rangle, |\lambda_i\rangle, |\Delta_i\rangle, |d\rangle$. The first one can be easily derived from (8). Then a global conformal block takes a factorized form at $d = 2$, given in (11). (See [81] for a general d generalization.) According to the z dependence, we have

$$\begin{aligned} & \sum_i \left(\Delta_i^{(1)}|\Delta_i\rangle + \lambda_i^{(1)}|\lambda_i\rangle \right) \\ &= \sum_\beta v^{\frac{1}{8}} \left(A_\beta(\bar{z})k_\beta(z) + B_\beta(\bar{z})\partial_\beta k_\beta(z) \right) - (z \leftrightarrow 1 - \bar{z}), \end{aligned} \quad (14)$$

where $\beta \in \{\tau^{(0)}\}$ is defined in (12) and $A_\beta(\bar{z}), B_\beta(\bar{z})$ encode the dependence on \bar{z} . We can use the general d formula in [79] to compute numerically $|d\rangle$ order by order in z at any \bar{z} in $[0, 1)$ [82]. The analytic computation of $|d\rangle$ based on [83] is described in Supplemental Material.

NUMERICAL CONFORMAL BOOTSTRAP

Let us perform a numerical study of the crossing equation (13), which has no solution if $\{|d\rangle, |\Delta_\sigma\rangle, |\Delta_i\rangle, |\lambda_i\rangle\}$ are linearly independent. We can detect the linear independence by a norm:

$$\eta = \left\| |d\rangle + \Delta_\sigma^{(1)}|\Delta_\sigma\rangle + \sum_i \Delta_i^{(1)}|\Delta_i\rangle + \sum_i \lambda_i^{(1)}|\lambda_i\rangle \right\|, \quad (15)$$

which is the distance between the target point determined by $|d\rangle$ and a point in the space spanned by $|\Delta_\sigma\rangle, |\lambda_i\rangle, |\Delta_i\rangle$. If there exists at least one crossing solution, then we should find $\eta_{\min} = 0$. We define the norm in terms of sampling points [84]

$$\|H\| = \sqrt{\langle H|H\rangle} = \left(\frac{1}{N} \sum_{i=1}^N \mu(z_i, \bar{z}_i) |H(z_i, \bar{z}_i)|^2 \right)^{1/2}, \quad (16)$$

where the measure $\mu(z, \bar{z})$ will be specified later. The inner product $\langle H_1|H_2\rangle$ is defined as a weighted sum of the product $H_1^* H_2$. We consider the Lorentzian regime, so z, \bar{z} are

independent, real variables. We further concentrate on the region near the double-lightcone limit with $0 < z \ll 1$ and $0 \ll \bar{z} < 1$, which will also be studied analytically. We use sampling rather than derivative equations because it is easier to assign a proper measure $\mu(z, \bar{z})$.

In practice, we need to truncate the conformal block summation to a finite sum in the numerical studies. This is sometimes called OPE truncation [85]. Then we need to know if a finite η_{\min} is due to the OPE truncation or absence of crossing solution. Since we are sampling in a subregion, the prefactor of η_{\min} is scheme-dependent and the finite η_{\min} becomes smaller as we increase the truncation cutoff. To distinguish between the two origins, we examine the dependence of η_{\min} on the local sampling regions labelled by z_0 . If $\eta_{\min} > 0$ is mainly due to the OPE truncation, then the functional form of $\eta_{\min}(z_0)$ will change dramatically with the cutoff. Otherwise, $\eta_{\min}(z_0)$ will only get a smaller prefactor as more intermediate states are introduced. Near the lightcone, we can readily distinguish between them based on the scaling behaviour.




FIG. 1. Log-log plot of $\eta_{\min}(z_0)$ with a simple measure (17), where z_0 labels the sample region. The sampling points are at $z, 1 - \bar{z} = z_0 \times 10^{-k/10}$ with $z \neq 1 - \bar{z}$ and $k = 0, 1, 2, \dots, 10$. The scaling behaviour is not sensitive to the β truncation. A larger β cutoff reduces the prefactor, but does not modify the leading scaling behaviour. Therefore, a finite η_{\min} is mainly due to the absence of crossing solution, not the β truncation.

Near the lightcone $z = 0$, we can truncate the sum (14) to low β . But we will not truncate the spin sum, so $A_\beta(\bar{z}), B_\beta(\bar{z})$ in (14) remain arbitrary [89]. They will be evaluated near the other lightcone $\bar{z} = 1$. In the η minimization, $A_\beta(\bar{z})$ and $B_\beta(\bar{z})$ are approximated by truncated Taylor series about the center of the sampling region. We use high order Taylor polynomials to make the associated errors negligible. We can also view $A_\beta(\bar{z}), B_\beta(\bar{z})$ at different \bar{z} as independent parameters, but the results remain the same in the cases examined.

Now we discuss the choice of μ . A simple measure is

$$\mu_{\text{simple}}(z, \bar{z}) = 1. \quad (17)$$

In Fig. 1, we show the dependence of η_{\min} on the sampling region labelled by z_0 . One can notice the scaling behaviour

$$\eta_{\min}(z_0) \propto z_0^\alpha, \quad (18)$$

which becomes more precise at small z_0 . The exponent α is about 1.13(1) in the regime $10^{-6} < z_0 < 10^{-3}$, in which $|d\rangle$ can be computed by a direct summation over spin. For $z_0 < 10^{-6}$, we use the analytic expression of $|d\rangle$ in Supplemental Material to obtain a more precise value $\alpha \approx 1.125$. These results imply that $|d\rangle$ contains a vector that scales as $\lambda^{1.125}$ under $\{z, 1 - \bar{z}\} \rightarrow \{\lambda z, \lambda(1 - \bar{z})\}$. Furthermore, it does not belong to the space spanned by $\{|\Delta_\sigma\rangle, |\lambda_i\rangle, |\Delta_i\rangle\}$, so no crossing solution can be found. We will give an analytic understanding of the linear independence later.




FIG. 2. Log-log plot of $\eta_{\min}(z_0)$ with a refined measure (19). Here the measure depends on the cutoff $\beta_* = 1, 4, 8$. The sampling points are the same as those in Fig. 1. The scaling exponents decrease with the β cutoff and become negative, so the β truncation is not the main source and a finite η_{\min} is due to absence of crossing solution.

We can also consider a refined norm with a cutoff dependent measure. Near the lightcone, the lowest β contribution dominates the OPE truncation error, so we use

$$\mu_{\text{refined}}(z, \bar{z}) = |z^{\beta_*/2} - (1 - \bar{z})^{\beta_*/2}|^{-2}, \quad (19)$$

where β_* is the cutoff for the β summation in (14). If a crossing solution exists, the exponents should always be positive because the OPE truncation errors are of higher order in $z, 1 - \bar{z}$ than $\mu_*^{-1/2}$. In Fig. 2, we compare the results of different β_* . One can see that the exponent α decreases with the cutoff β_* and becomes negative, implying that the OPE truncation is not the main origin of $\eta_{\min} > 0$. The approximate values of the scaling exponents are 0.63(1), -0.87(1), -2.87(1), where the latter two are consistent with $\alpha_{\text{refined}} \approx \alpha_{\text{simple}} - \beta_*/2$. A negative exponent also implies a divergent η_{\min} in the double lightcone limit $z, 1 - \bar{z} \rightarrow 0$, providing a clear signature for absence of crossing solution.

The η minimization results have a geometric interpretation, as it induces a special vector $|N\rangle$ orthogonal to the basis vectors. The squared minimal distance η_{\min}^2 is precisely the inner product of $|N\rangle$ and $|d\rangle$. When $\eta_{\min} > 0$, there is no crossing solution due to a finite distance between the target point and the space spanned by the basis vectors.

ANALYTICAL CONFORMAL BOOTSTRAP

For a deeper understanding, let us study the crossing equation (13) in the analytic lightcone expansions. We will find obstructions from both regular and bi-singular terms.

First, we discuss the inconsistency from regular terms. Near the double lightcone limit, the target vector can be well approximated by

$$|d\rangle = \frac{z}{4\sqrt{2}} - \frac{3z(1 - \bar{z})^{\frac{1}{8}}}{16G} - (z \leftrightarrow 1 - \bar{z}) + \dots, \quad (20)$$

where $G = \Gamma(1/4)^2 (2\pi)^{-3/2}$ is Gauss's constant and \dots indicates higher order terms. The scaling behaviour $\eta_{\min}(z_0) \propto z_0^{1.125}$ in the numerical analysis is associated with the leading regular term in (20):

$$z(1 - \bar{z})^{\frac{1}{8}} - z^{\frac{1}{8}}(1 - \bar{z}). \quad (21)$$

In the lightcone limit $z \rightarrow 0$, the exponents of z are associated with the half twists of primary and descendant states in the direct-channel OPE. In the double lightcone expansion, we expect that the exponents of u, v are associated with intermediate twists [90]: $v^{\Delta_\sigma} \mathcal{G}(z, \bar{z}) = \sum_{i,j} c_{i,j} u^{\tau_i/2} v^{\tau_j/2}$, such as (8). Since the double-twist trajectories $[\sigma\sigma]_n$ are absent in 2d, the first term $z(1 - \bar{z})^{1/8}$ can only come from the direct-channel contribution. One can show that the structure of $k_\beta(z)$ in (14) is inconsistent with the explicit expression of $|d\rangle$, so the crossing equation (13) has no solution.

The $d = 2$ solution is very special. All the regular terms have vanishing coefficients, so the double-twist trajectories $[\sigma\sigma]_n$ can be absent. At first order in $\epsilon = d - 2$, the d -dependence of conformal symmetry requires the presence of double twist states in the $\sigma \times \sigma$ OPE. From this analytic perspective, the spectrum transition takes place at $d = 2 + 0^+$. Usually, the presence of double-twist states is based on the assumption of a twist gap [56, 57], but here we show that they are required by conformal symmetry even if the twist gap vanishes. Furthermore, we expect the existence of other double/multi-twist states, but the more complicated ones should be suppressed by higher powers of ϵ .

Second, we consider the inconsistency associated with bi-singular terms. The presence of double-twist trajectories is not sufficient. Another obstruction to solving (13) is the large spacing of the twist spectrum (12). We can simplify the analysis by focusing on the bi-singular terms. To match the power laws in $|d\rangle_{\text{b.s.}}$, the functions $A_\beta(\bar{z}), B_\beta(\bar{z})$ in (14) should take the form $\sum_{k=0}^{\infty} (a_{0,k} + a_{1,k} \log(1 - \bar{z})) (1 - \bar{z})^{k/2 - 1/8}$, with $a_{n,k}$ replaced by $b_{n,k}$ for $B_\beta(\bar{z})$. We introduce $\log(1 - \bar{z})$ because $\partial_\beta k_\beta(z)$ involves $\log z$. The exponents take the expected values and there is no double-twist exponents, so naively we may try to solve the crossing equation order by order. Let us count the total power of $z, 1 - \bar{z}$. For example, the first line of eq.(10) in Supplemental Material contains terms of order 1, 3/2. After solving the crossing equation to order 2, we substitute the solutions of $a_{n,k}, b_{n,k}$ into the 5/2 order

equation. We find that the sum below has a fixed coefficient

$$\begin{aligned} & \left(\Delta_\sigma^{(1)} |\Delta_\sigma\rangle + \sum_i \Delta_i^{(1)} |\Delta_i\rangle + \sum_i \lambda_i^{(1)} |\lambda_i\rangle + |d\rangle \right)_{\text{b.s.}} \\ &= -\frac{6\sqrt{2}}{539} \left(z(1-\bar{z})^{3/2} - z^{3/2}(1-\bar{z}) \right) + \dots, \end{aligned} \quad (22)$$

so the bi-singular part of the crossing equation (13) has no solution beyond order 2. To construct a crossing solution, one can reduce the spacing of the twist spectrum from 4 to 2, as in the standard case of generalized free theory.

The $d = 2$ solution is possible because of the special structure of 2d conformal blocks. As 2d global conformal blocks are invariant under $\ell \rightarrow -\ell$, the spectrum is symmetric in twist $\Delta - \ell$ and conformal spin $\Delta + \ell$. This explains the large spacing in intermediate twist spectrum, which is dual to that of 2ℓ [91]. As only even spin states appear in the $\sigma \times \sigma$ OPE, the 2d twist spacing is 4. This large spacing is inconsistent with the general d structure of conformal blocks.

DISCUSSION

We have investigated the $d = 2 + \epsilon$ critical Ising model using novel numerical and analytical conformal bootstrap methods. Our analyses of the crossing equation (13) disprove the naive expectation that the leading corrections are linear in ϵ . The d -dependence of global conformal symmetry implies the existence of new intermediate states, such as double twist trajectories. But the intermediate spectrum is the same as the $d = 2$ case at order ϵ if the naive expectation is correct. No solution to the crossing equation can be found due to the linear independence of conformal blocks. Since the obstructions are related to the speciality of $d = 2$, we expect them to appear also in other 2d CFTs.

A direct consequence is that the leading corrections to the 2d data should be more singular than ϵ^1 . The presence of ϵ^a corrections with $0 < a < 1$ will imply strong non-unitarity of the Ising CFT below $d = 2$. This is consistent with the observation of two kinks in the $d < 2$ unitary bootstrap bounds in [93]. The leading corrections may take the form of order $\epsilon^{1/k}$ with $k = 2, 3, \dots$. One can rule out the possibility that only scaling dimensions receive $\epsilon^{1/k}$ corrections by adding higher β derivatives of $k_\beta(z)$ to (14), then there should be infinitely many new states. Similar to the XY model results in [94], the simplest resolution could be that the leading corrections are of order $\epsilon^{1/2}$. For $d < 2$, the scaling dimensions can be complex conjugate pairs and the OPE coefficients of new states can be imaginary numbers. For perturbative RG fixed points, the $\epsilon^{1/2}$ behavior has also been found in the cases with two marginal operators, such as the $d = 4 - \epsilon$ random Ising model [95]. In general, the square root behaviour can appear around a bifurcation point at which two fixed points collide [96–99].

Although we show there are infinitely many new states, it is still unclear if the low-lying scaling dimensions receive singular corrections. It may be helpful to learn from other analytical insights [61–70, 100–105]. It would also be fascinating to

study other strongly coupled CFTs in $2 + \epsilon$ dimensions. For more complex problems, it could be useful to assume a hierarchical structure in operator product expansion [53, 106].

Many statistical physics models violate reflection positivity. Similarly, the boundary/defect bootstrap [107][42] [108–111] and thermal bootstrap [112–114] problems do not obey positivity constraints. In the usual numerical bootstrap, the positivity constraints are crucial to the derivation of rigorous bounds. Here we show that the inconsistent theory space can be ruled out without using positivity. We plan to revisit the non-positive bootstrap problems from the new perspective.

I would like to thank Hugh Osborn, Slava Rychkov, Ning Su and other organizers/participants of Bootstat 2021 for enlightening comments. I am particularly grateful to Slava Rychkov for valuable comments on the draft. The Bootstat 2021 program took place at Institut Pascal at Université Paris-Saclay with the support of the program “Investissements d’avenir” ANR-11-IDEX-0003-01. This work was partly supported by the 100 Talents Program of Sun Yat-Sen University, Okinawa Institute of Science and Technology Graduate University (OIST) and JSPS Grant-in-Aid for Early-Career Scientists (KAKENHI No. 19K14621).

* liwliang3@mail.sysu.edu.cn

- [1] L. D. Landau, “On the theory of phase transitions,” *Zh. Eksp. Teor. Fiz.* **7**, 19-32 (1937)
- [2] There are logarithmic corrections at $d = 4$ [3].
- [3] A. I. Larkin and D. E. Khmel’nitskii, “Phase transition in uniaxial ferroelectrics,” *Zh. Eksp. Teor. Fiz.* **56**, 2087-2098 (1969)
- [4] K. G. Wilson and M. E. Fisher, “Critical exponents in 3.99 dimensions,” *Phys. Rev. Lett.* **28**, 240-243 (1972) doi:10.1103/PhysRevLett.28.240
- [5] K. G. Wilson and J. B. Kogut, “The Renormalization group and the epsilon expansion,” *Phys. Rept.* **12**, 75-199 (1974) doi:10.1016/0370-1573(74)90023-4
- [6] A. Pelissetto and E. Vicari, “Critical phenomena and renormalization group theory,” *Phys. Rept.* **368**, 549-727 (2002) doi:10.1016/S0370-1573(02)00219-3 [arXiv:cond-mat/0012164 [cond-mat]].
- [7] L. Onsager, “Crystal statistics. 1. A Two-dimensional model with an order disorder transition,” *Phys. Rev.* **65**, 117-149 (1944) doi:10.1103/PhysRev.65.117
- [8] A. A. Migdal, “Phase transitions in gauge and spin-lattice systems,” *Sov. Phys. JETP* **42**, 743 (1975) PRINT-75-1042 (LANDAU-INST).
- [9] A. M. Polyakov, “Interaction of Goldstone Particles in Two-Dimensions. Applications to Ferromagnets and Massive Yang-Mills Fields,” *Phys. Lett. B* **59**, 79-81 (1975) doi:10.1016/0370-2693(75)90161-6
- [10] E. Brezin and J. Zinn-Justin, “Renormalization of the nonlinear sigma model in $2 + \epsilon$ dimensions. Application to the Heisenberg ferromagnets,” *Phys. Rev. Lett.* **36**, 691-694 (1976) doi:10.1103/PhysRevLett.36.691
- [11] E. Brezin and J. Zinn-Justin, “Spontaneous Breakdown of Continuous Symmetries Near Two-Dimensions,” *Phys. Rev. B* **14**, 3110 (1976) doi:10.1103/PhysRevB.14.3110

- [12] It is well known that the 2d Ising model can be mapped to a free fermion theory. It would be interesting to extend the fermionization to $2 + \epsilon$ dimensions.
- [13] R. Ma and C. Wang, “Theory of deconfined pseudo-criticality,” *Phys. Rev. B* **102**, no.2, 020407 (2020) doi:10.1103/PhysRevB.102.020407 [arXiv:1912.12315 [cond-mat.str-el]].
- [14] A. Nahum, “Note on Wess-Zumino-Witten models and quasiuniversality in 2+1 dimensions,” *Phys. Rev. B* **102**, no.20, 201116 (2020) doi:10.1103/PhysRevB.102.201116 [arXiv:1912.13468 [cond-mat.str-el]].
- [15] Y. C. He, J. Rong and N. Su, “Non-Wilson-Fisher kinks of $O(N)$ numerical bootstrap: from the deconfined phase transition to a putative new family of CFTs,” *SciPost Phys.* **10**, 115 (2021) doi:10.21468/SciPostPhys.10.5.115 [arXiv:2005.04250 [hep-th]].
- [16] E. Brezin, J. C. Le Guillou and J. Zinn-Justin, “Perturbation Theory at Large Order. I. The ϕ^{2N} Interaction,” *Phys. Rev. D* **15**, 1544-1557 (1977) doi:10.1103/PhysRevD.15.1544
- [17] C. Bagnuls and C. Bervillier, “Describing actual critical behaviour from field theory: a delicate matter,” *Phys. Lett. A* **195**, no.2, 163-170 (1994) doi:10.1016/0375-9601(94)90090-6
- [18] L. Schafer, “Sign of corrections to scaling amplitudes: Field-theoretic considerations and results for self-repelling walks,” *Phys. Rev. E* **50**, no.5, 3517 (1994) doi:10.1103/PhysRevE.50.3517
- [19] Z. Komargodski and D. Simmons-Duffin, “The Random-Bond Ising Model in 2.01 and 3 Dimensions,” *J. Phys. A* **50**, no.15, 154001 (2017) doi:10.1088/1751-8121/aa6087 [arXiv:1603.04444 [hep-th]].
- [20] J. L. Cardy and H. W. Hamber, “ $O(n)$ Heisenberg Model Close to $n = d = 2$,” *Phys. Rev. Lett.* **45**, 499 (1980) [erratum: *Phys. Rev. Lett.* **45**, 1217 (1980)] doi:10.1103/PhysRevLett.45.499
- [21] S. Ferrara, A. F. Grillo and R. Gatto, “Tensor representations of conformal algebra and conformally covariant operator product expansion,” *Annals Phys.* **76**, 161 (1973). doi:10.1016/0003-4916(73)90446-6
- [22] A. M. Polyakov, “Nonhamiltonian approach to conformal quantum field theory,” *Zh. Eksp. Teor. Fiz.* **66**, 23 (1974) [*Sov. Phys. JETP* **39**, 9 (1974)].
- [23] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, “Infinite Conformal Symmetry in Two-Dimensional Quantum Field Theory,” *Nucl. Phys. B* **241**, 333 (1984). doi:10.1016/0550-3213(84)90052-X
- [24] P. Di Francesco, P. Mathieu and D. Senechal, “Conformal Field Theory,” doi:10.1007/978-1-4612-2256-9
- [25] R. Rattazzi, V. S. Rychkov, E. Tonni and A. Vichi, “Bounding scalar operator dimensions in 4D CFT,” *JHEP* **0812**, 031 (2008) doi:10.1088/1126-6708/2008/12/031 [arXiv:0807.0004 [hep-th]].
- [26] S. El-Showk, M. F. Paulos, D. Poland, S. Rychkov, D. Simmons-Duffin and A. Vichi, “Solving the 3D Ising Model with the Conformal Bootstrap,” *Phys. Rev. D* **86**, 025022 (2012) doi:10.1103/PhysRevD.86.025022 [arXiv:1203.6064 [hep-th]].
- [27] S. El-Showk, M. F. Paulos, D. Poland, S. Rychkov, D. Simmons-Duffin and A. Vichi, “Solving the 3d Ising Model with the Conformal Bootstrap II. c-Minimization and Precise Critical Exponents,” *J. Stat. Phys.* **157**, 869 (2014) doi:10.1007/s10955-014-1042-7 [arXiv:1403.4545 [hep-th]].
- [28] F. Kos, D. Poland and D. Simmons-Duffin, “Bootstrapping Mixed Correlators in the 3D Ising Model,” *JHEP* **1411**, 109 (2014) doi:10.1007/JHEP11(2014)109 [arXiv:1406.4858 [hep-th]].
- [29] F. Kos, D. Poland, D. Simmons-Duffin and A. Vichi, “Precision Islands in the Ising and $O(N)$ Models,” *JHEP* **1608**, 036 (2016) doi:10.1007/JHEP08(2016)036 [arXiv:1603.04436 [hep-th]].
- [30] J. D. Qualls, “Lectures on Conformal Field Theory,” arXiv:1511.04074 [hep-th].
- [31] S. Rychkov, “EPFL Lectures on Conformal Field Theory in $D \geq 3$ Dimensions,” doi:10.1007/978-3-319-43626-5 arXiv:1601.05000 [hep-th].
- [32] D. Simmons-Duffin, “The Conformal Bootstrap,” doi:10.1142/9789813149441-0001 arXiv:1602.07982 [hep-th].
- [33] D. Poland, S. Rychkov and A. Vichi, “The Conformal Bootstrap: Theory, Numerical Techniques, and Applications,” *Rev. Mod. Phys.* **91**, no. 1, 15002 (2019) [*Rev. Mod. Phys.* **91**, 015002 (2019)] doi:10.1103/RevModPhys.91.015002 [arXiv:1805.04405 [hep-th]].
- [34] S. M. Chester, “Weizmann Lectures on the Numerical Conformal Bootstrap,” arXiv:1907.05147 [hep-th].
- [35] S. El-Showk, M. Paulos, D. Poland, S. Rychkov, D. Simmons-Duffin and A. Vichi, “Conformal Field Theories in Fractional Dimensions,” *Phys. Rev. Lett.* **112**, 141601 (2014) doi:10.1103/PhysRevLett.112.141601 [arXiv:1309.5089 [hep-th]].
- [36] C. Behan, “PyCFTBoot: A flexible interface for the conformal bootstrap,” *Commun. Comput. Phys.* **22**, no.1, 1-38 (2017) doi:10.4208/cicp.OA-2016-0107 [arXiv:1602.02810 [hep-th]].
- [37] A. Cappelli, L. Maffi and S. Okuda, “Critical Ising Model in Varying Dimension by Conformal Bootstrap,” *JHEP* **01**, 161 (2019) doi:10.1007/JHEP01(2019)161 [arXiv:1811.07751 [hep-th]].
- [38] M. Hogervorst, S. Rychkov and B. C. van Rees, “Unitarity violation at the Wilson-Fisher fixed point in $4 - \epsilon$ dimensions,” *Phys. Rev. D* **93**, no.12, 125025 (2016) doi:10.1103/PhysRevD.93.125025 [arXiv:1512.00013 [hep-th]].
- [39] S. El-Showk and M. F. Paulos, “Extremal bootstrapping: go with the flow,” *JHEP* **03**, 148 (2018) doi:10.1007/JHEP03(2018)148 [arXiv:1605.08087 [hep-th]].
- [40] F. Gliozzi, “More constraining conformal bootstrap,” *Phys. Rev. Lett.* **111**, 161602 (2013) doi:10.1103/PhysRevLett.111.161602 [arXiv:1307.3111 [hep-th]].
- [41] F. Gliozzi and A. Rago, “Critical exponents of the 3d Ising and related models from Conformal Bootstrap,” *JHEP* **1410**, 042 (2014) doi:10.1007/JHEP10(2014)042 [arXiv:1403.6003 [hep-th]].
- [42] F. Gliozzi, P. Liendo, M. Meineri and A. Rago, “Boundary and Interface CFTs from the Conformal Bootstrap,” *JHEP* **05**, 036 (2015) doi:10.1007/JHEP05(2015)036 [arXiv:1502.07217 [hep-th]].
- [43] Y. Nakayama, “Bootstrapping critical Ising model on three-dimensional real projective space,” *Phys. Rev. Lett.* **116**, no.14, 141602 (2016) doi:10.1103/PhysRevLett.116.141602 [arXiv:1601.06851 [hep-th]].
- [44] F. Gliozzi, “Truncatable bootstrap equations in algebraic form and critical surface exponents,” *JHEP* **10**, 037 (2016) doi:10.1007/JHEP10(2016)037 [arXiv:1605.04175 [hep-th]].
- [45] I. Esterlis, A. L. Fitzpatrick and D. Ramirez, “Closure of the Operator Product Expansion in the Non-Unitary Boot-

- strap,” JHEP **11**, 030 (2016) doi:10.1007/JHEP11(2016)030 [arXiv:1606.07458 [hep-th]].
- [46] S. Hikami, “Conformal bootstrap analysis for the Yang–Lee edge singularity,” PTEP **2018**, no.5, 053I01 (2018) doi:10.1093/ptep/pty054 [arXiv:1707.04813 [hep-th]].
- [47] S. Hikami, “Conformal Bootstrap Analysis for Single and Branched Polymers,” PTEP **2018**, no.12, 123I01 (2018) doi:10.1093/ptep/pty132 [arXiv:1708.03072 [hep-th]].
- [48] S. Hikami, “Dimensional Reduction by Conformal Bootstrap,” PTEP **2019**, no.8, 083A03 (2019) doi:10.1093/ptep/ptz081 [arXiv:1801.09052 [cond-mat.dis-nn]].
- [49] A. Leclair and J. Squires, “Conformal bootstrap for percolation and polymers,” J. Stat. Mech. **1812**, no.12, 123105 (2018) doi:10.1088/1742-5468/aaf10a [arXiv:1802.08911 [hep-th]].
- [50] S. Hikami, “Conformal Bootstrap Analysis for Localization: Symplectic Case,” [arXiv:1811.05918 [cond-mat.dis-nn]].
- [51] J. Rong and J. Zhu, “On the ϕ^3 theory above six dimensions,” JHEP **04**, 151 (2020) doi:10.1007/JHEP04(2020)151 [arXiv:2001.10864 [hep-th]].
- [52] Y. Nakayama, “Is there supersymmetric Lee-Yang fixed point in three dimensions?,” [arXiv:2104.13570 [hep-th]].
- [53] W. Li, “New method for the conformal bootstrap with OPE truncations,” arXiv:1711.09075 [hep-th].
- [54] G. Kántor, V. Niarchos and C. Papageorgakis, “Solving Conformal Field Theories with Artificial Intelligence,” Phys. Rev. Lett. **128**, no.4, 041601 (2022) doi:10.1103/PhysRevLett.128.041601 [arXiv:2108.08859 [hep-th]].
- [55] G. Kántor, V. Niarchos and C. Papageorgakis, “Conformal bootstrap with reinforcement learning,” Phys. Rev. D **105**, no.2, 025018 (2022) doi:10.1103/PhysRevD.105.025018 [arXiv:2108.09330 [hep-th]].
- [56] A. L. Fitzpatrick, J. Kaplan, D. Poland and D. Simmons-Duffin, “The Analytic Bootstrap and AdS Superhorizon Locality,” JHEP **1312**, 004 (2013) doi:10.1007/JHEP12(2013)004 [arXiv:1212.3616 [hep-th]].
- [57] Z. Komargodski and A. Zhiboedov, “Convexity and Liberation at Large Spin,” JHEP **1311**, 140 (2013) doi:10.1007/JHEP11(2013)140 [arXiv:1212.4103 [hep-th]].
- [58] We thank Slava Rychkov for emphasizing this point.
- [59] One needs to know if a virial current is present [60]. The case of $d = 4 - \epsilon$ has been revisited from various CFT perspectives [61–70].
- [60] J. Polchinski, “Scale and Conformal Invariance in Quantum Field Theory,” Nucl. Phys. B **303**, 226–236 (1988) doi:10.1016/0550-3213(88)90179-4
- [61] S. Rychkov and Z. M. Tan, “The ϵ -expansion from conformal field theory,” J. Phys. A **48**, no.29, 29FT01 (2015) doi:10.1088/1751-8113/48/29/29FT01 [arXiv:1505.00963 [hep-th]].
- [62] E. D. Skvortsov, “On (Un)Broken Higher-Spin Symmetry in Vector Models,” doi:10.1142/9789813144101_0008 [arXiv:1512.05994 [hep-th]].
- [63] S. Giombi and V. Kirilin, “Anomalous dimensions in CFT with weakly broken higher spin symmetry,” JHEP **11**, 068 (2016) doi:10.1007/JHEP11(2016)068 [arXiv:1601.01310 [hep-th]].
- [64] R. Gopakumar, A. Kaviraj, K. Sen and A. Sinha, “Conformal Bootstrap in Mellin Space,” Phys. Rev. Lett. **118**, no.8, 081601 (2017) doi:10.1103/PhysRevLett.118.081601 [arXiv:1609.00572 [hep-th]].
- [65] R. Gopakumar, A. Kaviraj, K. Sen and A. Sinha, “A Mellin space approach to the conformal bootstrap,” JHEP **05**, 027 (2017) doi:10.1007/JHEP05(2017)027 [arXiv:1611.08407 [hep-th]].
- [66] F. Gliozzi, A. Guerrieri, A. C. Petkou and C. Wen, “Generalized Wilson-Fisher Critical Points from the Conformal Operator Product Expansion,” Phys. Rev. Lett. **118**, no.6, 061601 (2017) doi:10.1103/PhysRevLett.118.061601 [arXiv:1611.10344 [hep-th]].
- [67] F. Gliozzi, A. L. Guerrieri, A. C. Petkou and C. Wen, “The analytic structure of conformal blocks and the generalized Wilson-Fisher fixed points,” JHEP **04**, 056 (2017) doi:10.1007/JHEP04(2017)056 [arXiv:1702.03938 [hep-th]].
- [68] L. F. Alday, “Solving CFTs with Weakly Broken Higher Spin Symmetry,” JHEP **10**, 161 (2017) doi:10.1007/JHEP10(2017)161 [arXiv:1612.00696 [hep-th]].
- [69] L. F. Alday, J. Henriksson and M. van Loon, “Taming the ϵ -expansion with large spin perturbation theory,” JHEP **07**, 131 (2018) doi:10.1007/JHEP07(2018)131 [arXiv:1712.02314 [hep-th]].
- [70] D. Carmi, J. Penedones, J. A. Silva and A. Zhiboedov, “Applications of dispersive sum rules: ϵ -expansion and holography,” SciPost Phys. **10**, 145 (2021) doi:10.21468/SciPostPhys.10.6.145 [arXiv:2009.13506 [hep-th]].
- [71] The study of conformal blocks dates back to the 1970’s [72–75]. For $d = 2, 4, 6$ the explicit expressions of 4-point scalar blocks were found in [76, 77].
- [72] S. Ferrara, A. F. Grillo, G. Parisi and R. Gatto, “Covariant expansion of the conformal four-point function,” Nucl. Phys. B **49**, 77 (1972) Erratum: [Nucl. Phys. B **53**, 643 (1973)]. doi:10.1016/0550-3213(72)90587-1, 10.1016/0550-3213(73)90467-7
- [73] S. Ferrara, A. F. Grillo, R. Gatto and G. Parisi, “Analyticity properties and asymptotic expansions of conformal covariant green’s functions,” Nuovo Cim. A **19**, 667 (1974). doi:10.1007/BF02813413
- [74] S. Ferrara, R. Gatto and A. F. Grillo, “Properties of Partial Wave Amplitudes in Conformal Invariant Field Theories,” Nuovo Cim. A **26**, 226 (1975). doi:10.1007/BF02769009
- [75] V. K. Dobrev, V. B. Petkova, S. G. Petrova and I. T. Todorov, “Dynamical Derivation of Vacuum Operator Product Expansion in Euclidean Conformal Quantum Field Theory,” Phys. Rev. D **13**, 887 (1976). doi:10.1103/PhysRevD.13.887
- [76] F. A. Dolan and H. Osborn, “Conformal four point functions and the operator product expansion,” Nucl. Phys. B **599**, 459 (2001) doi:10.1016/S0550-3213(01)00013-X [hep-th/0011040].
- [77] F. A. Dolan and H. Osborn, “Conformal partial waves and the operator product expansion,” Nucl. Phys. B **678**, 491 (2004) doi:10.1016/j.nuclphysb.2003.11.016 [hep-th/0309180].
- [78] For $d = 4 - \epsilon$, the absence of new states is an assumption in the conformal multiplet recombination in [61]. This can be deduced from the assumption of integer power series.
- [79] W. Li, “Lightcone expansions of conformal blocks in closed form,” JHEP **06**, 105 (2020) doi:10.1007/JHEP06(2020)105 [arXiv:1912.01168 [hep-th]].
- [80] A simple example is: $a_0 + a_2 t^2 = b_0 + b_1 t^1 + b_2 t^2$, which has no solution that is valid at generic t if $b_1 \neq 0$.
- [81] W. Li, “Factorized lightcone expansion of conformal blocks,” JHEP **05**, 128 (2021) doi:10.1007/JHEP05(2021)128 [arXiv:2012.09710 [hep-th]].
- [82] Near the lightcone $\bar{z} = 1$, a uniform precision evaluation requires summing the spin ℓ from 0 to $C(1 - \bar{z})^{-1/2}$, where $C \gg 1$ determines the precision.
- [83] D. Simmons-Duffin, “The Lightcone Bootstrap and the Spectrum of the 3d Ising CFT,” JHEP **1703**, 086 (2017)

- doi:10.1007/JHEP03(2017)086 [arXiv:1612.08471 [hep-th]].
- [84] M. Hogervorst and S. Rychkov, “Radial Coordinates for Conformal Blocks,” *Phys. Rev. D* **87**, 106004 (2013) doi:10.1103/PhysRevD.87.106004 [arXiv:1303.1111 [hep-th]].
- [85] For unitary CFTs, both the conformal block decompositions converge rapidly near $z = \bar{z} = 1/2$ [86]. Based on 2d examples, the non-unitary CFTs should share some features. See [87, 88] for recent discussions of the OPE convergence in the distributional sense and in Minkowski space.
- [86] D. Pappadopulo, S. Rychkov, J. Espin and R. Rattazzi, “OPE Convergence in Conformal Field Theory,” *Phys. Rev. D* **86**, 105043 (2012) doi:10.1103/PhysRevD.86.105043 [arXiv:1208.6449 [hep-th]].
- [87] P. Kravchuk, J. Qiao and S. Rychkov, “Distributions in CFT. Part I. Cross-ratio space,” *JHEP* **05**, 137 (2020) doi:10.1007/JHEP05(2020)137 [arXiv:2001.08778 [hep-th]].
- [88] P. Kravchuk, J. Qiao and S. Rychkov, “Distributions in CFT II. Minkowski Space,” [arXiv:2104.02090 [hep-th]].
- [89] In fact, $A_\beta(\bar{z}), B_\beta(\bar{z})$ should satisfy additional constraints, but the enlarged space has no solution as well.
- [90] L. F. Alday and A. Zhiboedov, “Conformal Bootstrap With Slightly Broken Higher Spin Symmetry,” *JHEP* **06**, 091 (2016) doi:10.1007/JHEP06(2016)091 [arXiv:1506.04659 [hep-th]].
- [91] Different Regge trajectories are connected by the analyticity in the $SL(2, \mathbb{R})$ conformal spin with negative ℓ and form a more rigid structure, i.e. Virasoro blocks. This can be generalized to $d > 2$. In [92], the closed results were inspired by the negative spin poles.
- [92] W. Li, “Closed-form expression for cross-channel conformal blocks near the lightcone,” *JHEP* **01**, 055 (2020) doi:10.1007/JHEP01(2020)055 [arXiv:1906.00707 [hep-th]].
- [93] J. Golden and M. F. Paulos, “No unitary bootstrap for the fractal Ising model,” *JHEP* **03**, 167 (2015) doi:10.1007/JHEP03(2015)167 [arXiv:1411.7932 [hep-th]].
- [94] D.R. Nelson and D. S. Fisher, “Dynamics of classical spins in one and two dimensions” *Phys. Rev. B* **16**, 4945-4955 (1977). doi:10.1103/PhysRevB.16.4945
- [95] D. Khmel'nitskii, “Second-order phase transition in inhomogeneous bodies,” *Zh. Eksp. Teor. Fiz* **41** (1975): 1960.
- [96] S. Gukov, “RG Flows and Bifurcations,” *Nucl. Phys. B* **919**, 583-638 (2017) doi:10.1016/j.nuclphysb.2017.03.025 [arXiv:1608.06638 [hep-th]].
- [97] H. Osborn and A. Stergiou, “Seeking fixed points in multiple coupling scalar theories in the ϵ expansion,” *JHEP* **05**, 051 (2018) doi:10.1007/JHEP05(2018)051 [arXiv:1707.06165 [hep-th]].
- [98] V. Gorbenko, S. Rychkov and B. Zan, “Walking, Weak first-order transitions, and Complex CFTs,” *JHEP* **10**, 108 (2018) doi:10.1007/JHEP10(2018)108 [arXiv:1807.11512 [hep-th]].
- [99] S. Rychkov and A. Stergiou, “General Properties of Multiscalar RG Flows in $d = 4 - \epsilon$,” *SciPost Phys.* **6**, no.1, 008 (2019) doi:10.21468/SciPostPhys.6.1.008 [arXiv:1810.10541 [hep-th]].
- [100] S. Caron-Huot, “Analyticity in Spin in Conformal Theories,” *JHEP* **09**, 078 (2017) doi:10.1007/JHEP09(2017)078 [arXiv:1703.00278 [hep-th]].
- [101] L. F. Alday, “Large Spin Perturbation Theory for Conformal Field Theories,” *Phys. Rev. Lett.* **119**, no.11, 111601 (2017) doi:10.1103/PhysRevLett.119.111601 [arXiv:1611.01500 [hep-th]].
- [102] D. Carmi and S. Caron-Huot, “A Conformal Dispersion Relation: Correlations from Absorption,” *JHEP* **09**, 009 (2020) doi:10.1007/JHEP09(2020)009 [arXiv:1910.12123 [hep-th]].
- [103] D. Mazáč, L. Rastelli and X. Zhou, “A Basis of Analytic Functionals for CFTs in General Dimension,” [arXiv:1910.12855 [hep-th]].
- [104] J. Penedones, J. A. Silva and A. Zhiboedov, “Nonperturbative Mellin Amplitudes: Existence, Properties, Applications,” *JHEP* **08**, 031 (2020) doi:10.1007/JHEP08(2020)031 [arXiv:1912.11100 [hep-th]].
- [105] S. Caron-Huot, D. Mazac, L. Rastelli and D. Simmons-Duffin, “Dispersive CFT Sum Rules,” *JHEP* **05**, 243 (2021) doi:10.1007/JHEP05(2021)243 [arXiv:2008.04931 [hep-th]].
- [106] W. Li, “Inverse Bootstrapping Conformal Field Theories,” *JHEP* **01**, 077 (2018) doi:10.1007/JHEP01(2018)077 [arXiv:1706.04054 [hep-th]].
- [107] P. Liendo, L. Rastelli and B. C. van Rees, “The Bootstrap Program for Boundary CFT_d,” *JHEP* **07**, 113 (2013) doi:10.1007/JHEP07(2013)113 [arXiv:1210.4258 [hep-th]].
- [108] M. Billo, V. Goncalves, E. Lauria and M. Meineri, “Defects in conformal field theory,” *JHEP* **04**, 091 (2016) doi:10.1007/JHEP04(2016)091 [arXiv:1601.02883 [hep-th]].
- [109] A. Gadde, “Conformal constraints on defects,” *JHEP* **01**, 038 (2020) doi:10.1007/JHEP01(2020)038 [arXiv:1602.06354 [hep-th]].
- [110] P. Liendo and C. Meneghelli, “Bootstrap equations for $\mathcal{N} = 4$ SYM with defects,” *JHEP* **01**, 122 (2017) doi:10.1007/JHEP01(2017)122 [arXiv:1608.05126 [hep-th]].
- [111] M. Lemos, P. Liendo, M. Meineri and S. Sarkar, “Universality at large transverse spin in defect CFT,” *JHEP* **09**, 091 (2018) doi:10.1007/JHEP09(2018)091 [arXiv:1712.08185 [hep-th]].
- [112] S. El-Showk and K. Papadodimas, “Emergent Spacetime and Holographic CFTs,” *JHEP* **10**, 106 (2012) doi:10.1007/JHEP10(2012)106 [arXiv:1101.4163 [hep-th]].
- [113] L. Iliesiu, M. Kologlu, R. Mahajan, E. Perlmutter and D. Simmons-Duffin, “The Conformal Bootstrap at Finite Temperature,” *JHEP* **10**, 070 (2018) doi:10.1007/JHEP10(2018)070 [arXiv:1802.10266 [hep-th]].
- [114] L. Iliesiu, M. Kologlu and D. Simmons-Duffin, “Bootstrapping the 3d Ising model at finite temperature,” *JHEP* **12**, 072 (2019) doi:10.1007/JHEP12(2019)072 [arXiv:1811.05451 [hep-th]].