# Logarithmic Quantum Time Crystal

Haipeng Xue (薛海鹏),<sup>1,\*</sup> Lingchii Kong (孔令琦),<sup>1,\*</sup> and Biao Wu (吴飙)<sup>1,2,3,†</sup>

<sup>1</sup>International Center for Quantum Materials, School of Physics, Peking University, Beijing 100871, China

<sup>2</sup>Wilczek Quantum Center, School of Physics and Astronomy,

Shanghai Jiao Tong University, Shanghai 200240, China

<sup>3</sup>Collaborative Innovation Center of Quantum Matter, Beijing 100871, China

(Dated: July 22, 2022)

We investigate a time-independent many-boson system, whose ground states are quasi-degenerate and become infinitely degenerate in the thermodynamic limit. Out of these quasi-degenerate ground states we construct a quantum state that evolves in time with a period that is logarithmically proportional to the number of particles, that is,  $T \sim \log N$ . This boson system in such a state is a quantum time crystal as it approaches the ground state in the thermodynamic limit. The logarithmic dependence of its period on the total particle number Nmakes it observable experimentally even for systems with very large number of particles. Possible experimental proposals are discussed.

#### I. INTRODUCTION

Spontaneous symmetry breaking is a well known physical phenomenon, where the observed ground state of a many-particle system does not possess the symmetries of its Hamiltonian[1], e.g., ferromagnetic materials break the rotational symmetry and crystals break the spatial translational symmetry. Wilczek suggested the possibility of spontaneous breaking of time translational symmetry [2, 3]: the observed ground state of a closed quantum system may oscillate periodically in time. This suggestion had drawn some quick criticism[4, 5]. In 2015, a no-go theorem was proved to exclude the possibility of spontaneous continuous time translational symmetry breaking in the ground state for a wide class of Hamiltonians with short-range interactions[6]. However, it was realized later that quantum time crystals can exist in a periodically driven system<sup>[7]</sup>. This is now known as discrete time crystal; its equilibrium state can vary with a period that is multiple of the driving period [8-10]. Two experiments with trapped ions and nitrogen-vacancy centers were performed. confirming the existence of discrete time crystals[11, 12]. Recently, the discussion about time crystals has expanded to the systems with long-range interaction[13].

When spontaneous symmetry breaking occurs in a time independent system, the observed ground state is not the true ground state but a superposition of many (quasi-)degenerate ground states[14]. Two energy states are quasi-degenerate if the energy gap between them approaches zero in the thermodynamic limit. There have some efforts to construct a time crystal with these (quasi-)degenerate ground states[15–17]. For such a time crystal, it typically oscillates with a period that grows polynomially with particle number, i.e.  $T \sim O(N^n)$ . It will be impossible in experiments to keep a many-particle state from decoherence for such a long time.

In this work we study a time-independent many-boson system. This boson system has many quasi-degenerate ground states, which become infinitely degenerate in the thermodynamic limit. With these ground states, we are able to construct states that oscillate with time with a period *T* that is logarithmically proportional to the number of particles *N*, that is,  $T \sim \log N$ . Due to this logarithmic dependence, the period is short enough for possible experimental observation even for very large *N*. We find that the observables that are usually used in time crystal experiments are not suitable for our systems. We suggest to observe the time crystal by measuring the square of overlap between the states of such a quantum time crystal at different times, i.e.,  $|\langle \Psi(0)|\Psi(t)\rangle|^2$  with a technique based on the Hong–Ou–Mandel interference[18].

## II. TWO-MODE INTERACTING BOSON SYSTEMS

The interacting system of *N* bosons that we are going to consider has only two modes. In a certain parameter range, this system has many quasi-ground states, which approach the true ground state in the thermodynamics limit  $N \rightarrow \infty$ . This is the critical feature of this system and a necessary condition for spontaneous symmetry breaking to occur.

## A. Theoretical model

The two-mode interacting boson system is described by the following Hamiltonian[14, 19, 20]

$$\hat{H} = -(\hat{a}_{1}^{\dagger}\hat{a}_{2} + \hat{a}_{2}^{\dagger}\hat{a}_{1}) + \frac{\gamma}{N}[\hat{n}_{1}(\hat{n}_{1} - 1) + \hat{n}_{2}(\hat{n}_{2} - 1)] + \frac{4\gamma}{N}\hat{n}_{1}\hat{n}_{2} + \frac{\gamma}{N}(\hat{a}_{1}^{\dagger}\hat{a}_{1}^{\dagger}\hat{a}_{2}\hat{a}_{2} + \hat{a}_{2}^{\dagger}\hat{a}_{2}^{\dagger}\hat{a}_{1}\hat{a}_{1}),$$
(1)

where  $\hat{a}_1^{\dagger}(\hat{a}_1)$  and  $\hat{a}_2^{\dagger}(\hat{a}_2)$  are the creation (annihilation) operators of two modes and *N* is the total number of bosons. This simple theoretical model can now be realized in experiments[21]. In the above, we have set the strength of single particle hopping as the unit of energy and therefore  $\gamma$  is a dimensionless parameter characterizing the interaction strength and the pair hopping. In addition, for simplicity, we set  $\hbar \equiv 1$ .

<sup>\*</sup> These two authors contributed equally.

<sup>&</sup>lt;sup>†</sup> wubiao@pku.edu.cn

In the thermodynamic limit  $N \to \infty$ , this quantum manybody system can be described by a mean-field model. Mathematically, one can simply replace the operators  $\hat{a}_{1,2}^{\dagger}$  and  $\hat{a}_{1,2}$  with their corresponding complex variables  $a_{1,2}^{*}$  and  $a_{1,2}$ in Eq. (1)[14]. We choose a different set of variables  $P = (|a_2|^2 - |a_1|^2)/2$  and  $Q = \arg(a_2) - \arg(a_1)$ . Physically, P and Q are the particle number difference and the phase difference between the two modes, respectively. In terms of P and Q, the mean-field model of the system is

$$H_m = -\sqrt{1 - 4P^2} \cos Q + \gamma (1 - 4P^2) \cos^2 Q.$$
 (2)

Note that *P* and *Q* are a pair of dynamical variables that are canonically conjugate to each other. Fig. 1 is the energy landscape in the phase space of this mean-field Hamiltonian. When  $\gamma < 1/2$ , the center of the phase space P = Q = 0 has the lowest energy. When  $\gamma > 1/2$ , all the points on the dashed line in Fig. 1(b) have the same lowest energy. That means that the system is infinitely degenerate when  $\gamma > 1/2$ . As we shall see in the next subsection, this infinite degeneracy corresponds to a set of quasi-degenerate ground states in the quantum model (1).



FIG. 1. The energy landscape in the phase space of the mean-field model  $H_m$ . (a) When  $\gamma < 1/2$ , the system has only one ground state at (Q, P) = (0, 0); (b) when  $\gamma > 1/2$ , the system has infinitely degenerate ground states, which are located on the dashed line.

#### B. Quantum energy levels

For mathematical simplicity, we transform this boson system into a mathematically equivalent spin system. With  $\hat{n}_{1,2} = \hat{a}^{\dagger}_{1,2}\hat{a}_{1,2}$ , we introduce  $\hat{S}_x = (\hat{a}^{\dagger}_1\hat{a}_2 + \hat{a}^{\dagger}_2\hat{a}_1)/2$ ,  $\hat{S}_y = (\hat{a}^{\dagger}_1\hat{a}_2 - \hat{a}^{\dagger}_2\hat{a}_1)/2i$ ,  $\hat{S}_z = (\hat{n}_1 - \hat{n}_2)/2$ . The quantum model (1) becomes

$$\hat{H}' = \hat{H}/2 = -\hat{S}_x + \frac{2\gamma}{N}\hat{S}_x^2.$$
 (3)

This Hamiltonian is also called Lipkin-Meshkov-Glick model and describes a spin system with S = N/2 [22]. In this spin formalism the energy eigenstates and energy levels are rather obvious. As  $\hat{S}_x$  commutes with  $\hat{H}$ , the eigenstates of the system are the eigenstates of  $\hat{S}_x$ ,

$$\hat{S}_{x}|m\rangle = m|m\rangle, \ m = -S, -S+1, \cdots, 0, \cdots, S.$$
(4)

For eigenstate  $|m\rangle$ , its corresponding energy level is

$$E_m = \frac{2\gamma}{N}m^2 - m.$$
 (5)

When  $\gamma < 1/2$ , the lowest energy level is at m = S = N/2 and the energy level increases monotonically as *m* decreases. And the energy gap  $\delta E_m$  between two neighboring levels  $|m-1\rangle$  and  $|m\rangle$  is

$$\delta E_m = 1 - \frac{2\gamma}{N}(2m-1) > 1 - 2\gamma. \tag{6}$$

This means that the energy gap remains finite even in the thermodynamic limit  $N \to \infty$ .



FIG. 2. The energy levels for  $\gamma > 1/2$  which are labeled by quantum number *m* and denoted by black bars. The ground state is at  $m_0$ . The energy gap  $\delta E$  between neighboring levels around  $m_0$  is  $\propto 1/N$ . All the energy levels marked by the gray area are quasi-ground states, whose energies become degenerate when  $N \rightarrow \infty$ . The width of the gray area is about  $\sim \sqrt{N}$ . The energy gaps outside the gray area remain finite even at the large *N* limit.

When  $\gamma > 1/2$ , the energy level is the lowest at  $m_0$ , which is the largest integer smaller than or equal to  $N/(4\gamma)$ . The highest energy level is at m = -S. Without loss of essential physics, we choose  $m_0 = N/(4\gamma)$  by choosing an appropriate value of  $\gamma$ . The energy gap between the ground state and the first excited state is  $2\gamma/N$ ; the gap between the highest energy level and the second highest is  $1 + 2\gamma - 2\gamma/N$ . The former approaches zero and the latter remains finite at the limit  $N \rightarrow \infty$ . This means that for  $\gamma > 1/2$  the energy gap  $\delta E_m$  between two neighboring levels has different dependence on N at different levels m. We find that for a set of energy levels near the ground state the gap approaches zero and for others it remains finite at  $N \rightarrow \infty$ . This is shown schematically in Fig. 2. To accurately to describe this behavior, we define the energy gap  $\Delta E_m$  between energy level  $|m\rangle$  and the ground state  $|m_0\rangle$  and we find that

$$\Delta E_m = \frac{2\gamma}{N} (m - m_0)^2 \,. \tag{7}$$

This means that any energy level  $|m\rangle$  will approach the ground state energy in the thermodynamic limit  $N \rightarrow \infty$  when *m* satisfies

$$|m-m_0| \sim O\left(N^{1/2-\delta}\right),\tag{8}$$

where  $0 < \delta < 1/2$ . These energy levels are located in the shadow area in Fig. 2. We call them quasi-ground states, which form a sub-Hilbert space *C*. At the limit  $N \rightarrow \infty$ , all these quasi-ground states become the mean-field ground states on the black dashed line in Fig. 1(b).



FIG. 3. Energy eigenstates  $|m\rangle$  in the quantum phase space for N = 440 and  $\gamma = 3/4$ . (a) The true ground state; (b) the 29th exited state. The black lines represent the mean-field infinite degenerate ground states.

#### C. Eigenstates in quantum phase space

As discussed above, in the case of  $\gamma > 1/2$  there are many quasi-ground states in the quantum model  $\hat{H}$ , which correspond to the infinitely many ground states in the mean-field model. Such a correspondence becomes more apparent and insightful when we plot these quasi-ground states in the quantum phase space (see Fig. 3).

The quantum phase space is obtained by dividing the meanfield phase space in Fig. 1 into Planck cells. The size of a Planck cell, which is denoted by  $(Q_i, P_j)$ , is 1/N. A Wannier function  $\{|Q_i, P_j\rangle\}$ , which is localized along both *P* and *Q* directions, is assigned for each Planck cell and all the Wannier functions form a complete set of orthonormal basis[23– 27]. An energy eigenstate  $|m\rangle$  is then projected to the quantum phase space as

$$|m\rangle = \sum_{i,j} |Q_i, P_j\rangle \langle Q_i, P_j | m \rangle .$$
(9)

What is plotted in Fig. 3 is  $|\langle Q, P|m \rangle|^2$ . Two examples are shown in Fig. 3: one is the true ground state  $|m_0\rangle$  and

the other is a quasi-ground state. It is clear from the figure that both eigenstates are concentrated around the mean-field ground states, which are marked by black solid lines in Fig. 3.

## III. LOGARITHMIC TIME CRYSTAL

The quantum time crystal can be constructed as follows. We choose an eigenstate  $|m_1\rangle$  with

$$m_1 = m_0 - \left\lfloor \sqrt{\frac{N}{\log N}} \right\rfloor,\tag{10}$$

where  $\lfloor x \rfloor$  is the largest integer  $\leq x$ . This state  $|m_1\rangle$  certainly belongs to the sub-Hilbert space *C* as  $\Delta E_{m_1} = 2\gamma/\log N$  approaches zero when  $N \to \infty$ . We construct a superposition state  $|\Phi_0\rangle = (|m_0\rangle + |m_1\rangle)/\sqrt{2}$ . It will evolve with time as

$$|\Phi(t)\rangle = \frac{e^{-iE_{m_0}t}}{\sqrt{2}} \left[ |m_0\rangle + \exp\left(-\frac{i2\pi t}{T_0 \log N}\right) |m_1\rangle \right], \quad (11)$$

where  $T_0 = \pi/\gamma$  is a constant with order O(1) and we have ignored the minor difference between *x* and  $\lfloor x \rfloor$ . It is clear that this quantum state will oscillate with a period of  $T_0 \log(N)$ .

In real experiments, it is hard to prepare a state that is a superposition of only two eigenstates. We consider a double-Gaussian superposition state  $|\Psi(0)\rangle = \sum_{m} c_{m} |m\rangle$ , where

$$c_m = \frac{1}{\sqrt{2\pi\sigma}} \left[ e^{-\frac{(m-m_0)^2}{4\sigma^2}} + e^{-\frac{(m-m_1)^2}{4\sigma^2}} \right],$$
 (12)

where  $\sigma$  is the width of the Gaussian distribution. Since we do not want the Gaussian distributions to spread out the whole sub-Hilbert space *C* of quasi-ground states, the width  $\sigma$  should be much smaller than  $|m_1 - m_0|$ . For simplicity, we choose  $\sigma \sim O(1)$ .

The overlap between  $|\Psi(t)\rangle$  and  $|\Psi(0)\rangle$  can be evaluated as

$$\langle \Psi(0)|\Psi(t)\rangle = \sum_{m=-N/2}^{N/2} |c_m|^2 \exp\left(-i\Delta E_m t\right).$$
(13)

When *N* is very large, we can replace the summation by integral,

$$\langle \Psi(0)|\Psi(t)\rangle \approx \int_{-\infty}^{\infty} dm \, |c_m|^2 \exp\left(-i\Delta E_m t\right)$$
$$=A(t) \left\{ 1 + \exp\left[-\frac{N^2}{4\Sigma^2(t)\log^2 N}\right] \exp\left(-i\frac{2\pi t}{T(t)}\right) \right\}, \quad (14)$$

where

$$T(t) = T_0 \log N \left( 1 + \frac{64\gamma^2 \sigma^4 t^2}{N^2} \right),$$
  

$$\Sigma(t) = \sqrt{\left(\frac{N}{8\gamma\sigma t}\right)^2 + \sigma^2},$$
  

$$A(t) = \frac{1}{2} \sqrt{\frac{N}{N + 8i\gamma\sigma^2 t}}.$$
(15)

It is clear from T(t) that when  $t \ll O(N)$  the system oscillates with a period of  $T_0 \log N$ . A(t) shows that the amplitude of the oscillation will gradually decrease with time. However, within the time interval  $0 < t \ll O(N)$  we should be able to observe many periods of oscillations since the period of oscillation is of order  $O(\log(N))$ . In addition, the oscillations are enveloped by a Gaussian function with width  $\Sigma(t)$  which increases with time. Instead of doing integral, one can also do the summation in Eq.(13) numerically. Both results are shown in Fig. 4 for comparison; they agree very well.



FIG. 4. The oscillations of logarithmic time crystal (12). The solid line is the analytical result in Eq.(14) and the squares are numerical results obtained by directly summing Eq.(13). N = 29240,  $m_1 = m_0 + 91$ ,  $\sigma = 1$ .

#### **IV. EXPERIMENTAL SCHEME**

In literature, the time crystal dynamics is usually illustrated with the correlation function of the polarization along the z direction,  $\langle \hat{S}_z(0)\hat{S}_z(t)\rangle$  [28]. For the quantum time crystal Eq.(11), we have

$$\langle \hat{S}_{z}(0)\hat{S}_{z}(t)\rangle = A_{0,+}e^{-i\delta E_{m_{0}+1}t} + A_{0,-}e^{-i\delta E_{m_{0}}t} + A_{1,+}e^{-i\delta E_{m_{1}+1}t} + A_{1,-}e^{-i\delta E_{m_{1}}t},$$
(16)

where  $A_{j,\pm} = (N/2+1)N/2 - m_j(m_j\pm 1), j = 0, 1$ . The four characteristic frequencies  $\delta E_{m_j\pm 1}$  are just the energy gaps between the neighboring levels at the two states  $|m_0\rangle$  and  $|m_1\rangle$ . Therefore, we have  $\delta E_{m_j\pm 1} \ll O(1/\log N)$  and the oscillation period is much longer than  $\log N$ . So the correlation function Eq.(16) can not be used to observe the dynamic behavior of our logarithmic quantum time crystal. We propose to use the experimental scheme shown in Fig. 5, where  $|\langle \Psi(0)|\Psi(t)\rangle|^2$  is measured.

In Fig. 5, there are two identical two-mode boson systems, which are denoted as a and b, created with atoms and optical devices. The tunneling between system a and b can be tuned: initially the optical potential barrier between them is high so that there is no tunneling; the barrier is lowered in the later stage of the experiment to allow Hong-Ou-Mandel interference. Initially when the tunneling is off, both system a and b have the same number of bosons and evolve with time under the Hamiltonian Eq. 1.

A swap operator  $\hat{V}$  between these two systems is defined as

$$\hat{V}|\psi\rangle_a \otimes |\phi\rangle_b = |\phi\rangle_a \otimes |\psi\rangle_b, \qquad (17)$$



FIG. 5. The experiment scheme. (i) Two identical boson systems *a* and *b*; each has two modes, denoted by 1,2. Initially, the two systems are independent and they are prepared in states,  $|\Psi(0)\rangle$  and  $|\Psi(t)\rangle$ , respectively. (ii) We allow the atom tunneling between the modes  $a_i$  and  $b_i$  (i = 1,2) to realize Hong-Ou-Mandel interference. (iii) We count particle number parity in system *b*. Repeat the above process for many times. This scheme effectively measures  $|\langle \Psi(0)|\Psi(t)\rangle|^2$ .

where  $|\cdot\rangle_a$  and  $|\cdot\rangle_b$  are states of systems *a* and *b*. It is easy to see that  $\langle \hat{V} \rangle = |\langle \psi | \phi \rangle|^2$ , i.e., the expectation value of the swap operator  $\hat{V}$  is the overlap between two quantum states  $|\psi\rangle$  and  $|\phi\rangle$ . As  $\hat{V}$  has only two eigenvalues  $\pm 1$ , measuring the expectation of operator  $\hat{V}$  requires statistically counting its eigenvalues within a large ensemble. In our system,  $\hat{V} =$  $\hat{V}_1 \otimes \hat{V}_2$ , where  $\hat{V}_i$  is the swapping on individual modes i = 1, 2. So measuring  $\hat{V}$  can be achieved by measuring  $\hat{V}_1$  and  $\hat{V}_2$ .

The symmetric eigenstate  $|S\rangle$  and asymmetric eigenstate  $|AS\rangle$  of  $\hat{V}_i$  can be expressed in terms of Fock states as follows

$$|S\rangle = \text{superpose}\left\{ (\hat{a}_i^{\dagger} - \hat{b}_i^{\dagger})^{2\alpha} (\hat{a}_i^{\dagger} + \hat{b}_i^{\dagger})^{\beta} |\text{vac}\rangle \right\}, \qquad (18)$$

$$|AS\rangle = \text{superpose}\left\{ (\hat{a}_i^{\dagger} - \hat{b}_i^{\dagger})^{2\alpha - 1} (\hat{a}_i^{\dagger} + \hat{b}_i^{\dagger})^{\beta} |\text{vac}\rangle \right\}, \quad (19)$$

where  $\hat{a}_i^{\dagger}(\hat{b}_i^{\dagger})$  refers to the creation operator on mode  $a_i(b_i)$ and  $\alpha$  and  $\beta$  are positive integers. We perform the following transformation,

$$\left(\hat{a}_{i}^{\dagger}+\hat{b}_{i}^{\dagger}\right)/\sqrt{2}\rightarrow\hat{a}_{i}^{\dagger},\quad \left(\hat{a}_{i}^{\dagger}-\hat{b}_{i}^{\dagger}\right)/\sqrt{2}\rightarrow\hat{b}_{i}^{\dagger}.$$
 (20)

This transformation can be realized experimentally with atoms tunneling between  $a_i$  and  $b_i$  under a Bose-Hubbard Hamiltonian [29]. This is essentially Hong-Ou-Mandel in-

$$|S'\rangle = \text{superpose}\left\{ (\hat{b}_{i}^{\dagger})^{2\alpha} (\hat{a}_{i}^{\dagger})^{\beta} |\text{vac}\rangle \right\},$$
$$|AS'\rangle = \text{superpose}\left\{ (\hat{b}_{i}^{\dagger})^{2\alpha-1} (\hat{a}_{i}^{\dagger})^{\beta} |\text{vac}\rangle \right\}.$$
(21)

This means that observing even(odd) particle numbers in mode  $b_i$  corresponds to eigenvalues +1(-1) of  $\hat{V}_i$ . Hence if the particle numbers in mode  $b_1$  and  $b_2$  have the same parity, we know the measured quantity of  $\hat{V}$  is +1; otherwise, it is -1.

The basic experimental steps are shown in Fig. 5. Initially, we prepare system *b* in state  $|\Psi(0)\rangle$  and let it evolve for a period of time *t* so the state becomes  $|\Psi(t)\rangle$ . At this same moment, we prepare system *a* in state  $|\Psi(0)\rangle$ . Then we turn off the tunneling between different modes and turn on the the tunneling between system *a* and *b*, realizing Hong-Ou-Mandel interference. Finally, we observe the number of particles in both modes of system *b*. By repeat this experiment many times for different *t*,  $\langle \hat{V} \rangle = |\langle \Psi(0) | \Psi(t) \rangle|^2$  is obtained.

# V. DISCUSSION AND CONCLUSION

Before concluding, we need to mention that performing perturbed external fields [16] actually fails to produce a logarithmic time crystal state. Because the field, like  $g\hat{S}_z$ , can only superpose the state  $|m_0\rangle$  with its nearest levels. To be honest, how to realize logarithmic time crystal is still an open question for us. But we do have some speculations. In solid

- A. J. Beekman, L. Rademaker, and J. van Wezel, An introduction to spontaneous symmetry breaking, SciPost Physics Lecture Notes, 11 (2019).
- [2] A. Shapere and F. Wilczek, Classical time crystals, Physical Review Letters 109, 160402 (2012).
- [3] F. Wilczek, Quantum time crystals, Physical Review Letters 109, 160401 (2012).
- [4] P. Nozières, Time crystals: Can diamagnetic currents drive a charge density wave into rotation?, EPL (Europhysics Letters) 103, 57008 (2013).
- [5] P. Bruno, Impossibility of spontaneously rotating time crystals: a no-go theorem, Physical Review Letters 111, 070402 (2013).
- [6] H. Watanabe and M. Oshikawa, Absence of quantum time crystals, Physical Review Letters 114, 251603 (2015).
- [7] K. Sacha, Modeling spontaneous breaking of time-translation symmetry, Physical Review A **91**, 033617 (2015).
- [8] D. V. Else, B. Bauer, and C. Nayak, Floquet time crystals, Physical Review Letters 117, 090402 (2016).
- [9] V. Khemani, A. Lazarides, R. Moessner, and S. L. Sondhi, Phase structure of driven quantum systems, Physical Review Letters 116, 250401 (2016).
- [10] N. Y. Yao, A. C. Potter, I.-D. Potirniche, and A. Vishwanath, Discrete time crystals: Rigidity, criticality, and realizations, Physical Review Letters **118**, 030401 (2017).
- [11] S. Choi, J. Choi, R. Landig, G. Kucsko, H. Zhou, J. Isoya,

physics, electrons can be excited from the band bottom to the band top by the phonons with specific momentum and energy, where the momentum of phonons is tuned by the structure of Brillouin-zone. In our model, the system can be excited by a laser from  $|m_0\rangle$  to  $|m_1\rangle$  when the frequency of laser is equal to  $\Delta E_{m_1} = E_{m_1} - E_{m_0}$ . But only controlling the frequency cannot assure that the intermediate states are not superposed. So, we need to speculate that there may be some extra characteristics (like Brillouin-zone with respect to electrons) about our logarithmic time crystal state to rule the momentum of laser so that it can be achieved.

In conclusion, we have studied a two-mode boson system of N particles. In a certain parameter regime, this system has many quasi-ground states, which forms a sub-Hilbert space C and become degenerate with the true ground state in the thermodynamic limit  $N \to \infty$ . We have been able to superpose these quasi-ground states and construct a quantum time crystal that oscillates with a period that is logarithmically proportional to the number of particles,  $T \sim O(\log(N))$ . Such a logarithmic dependence makes the period short enough for possible experimental observation even when the number of particle is large. A experimental scheme based on Hong-Ou-Mandel interference has been proposed.

# ACKNOWLEDGMENTS

We thank Yijie Wang for helpful discussion. This work is supported by the National Key R&D Program of China (Grants No. 2017YFA0303302, No. 2018YFA0305602), National Natural Science Foundation of China (Grant No. 11921005), and Shanghai Municipal Science and Technology Major Project (Grant No.2019SHZDZX01).

F. Jelezko, S. Onoda, H. Sumiya, V. Khemani, <u>et al.</u>, Observation of discrete time-crystalline order in a disordered dipolar many-body system, Nature **543**, 221 (2017).

- [12] J. Zhang, P. W. Hess, A. Kyprianidis, P. Becker, A. Lee, J. Smith, G. Pagano, I.-D. Potirniche, A. C. Potter, A. Vishwanath, et al., Observation of a discrete time crystal, Nature 543, 217 (2017).
- [13] V. K. Kozin and O. Kyriienko, Quantum time crystals from hamiltonians with long-range interactions, Physical Review Letters 123, 210602 (2019).
- [14] Q. Zhu, Q. Zhang, and B. Wu, Extended two-site bose-hubbard model with pair tunneling: spontaneous symmetry breaking, effective ground state and fragmentation, Journal of Physics B: Atomic, Molecular and Optical Physics 48, 045301 (2015).
- [15] A. D. Shapere and F. Wilczek, Regularizations of time-crystal dynamics, Proceedings of the National Academy of Sciences 116, 18772 (2019).
- [16] Y. Huang, T. Li, and Z. Yin, Symmetry-breaking dynamics of the finite-size lipkin-meshkov-glick model near ground state, Physical Review A 97, 012115 (2018).
- [17] A. Syrwid, A. Kosior, and K. Sacha, Can a bright soliton model reveal a genuine time crystal for a finite number of bosons?, EPL (Europhysics Letters) 134, 66001 (2021).
- [18] C.-K. Hong, Z.-Y. Ou, and L. Mandel, Measurement of subpicosecond time intervals between two photons by interference,

Physical Review Letters 59, 2044 (1987).

- [19] J.-Q. Liang, J.-L. Liu, W.-D. Li, and Z.-J. Li, Atom-pair tunneling and quantum phase transition in the strong-interaction regime, Physical Review A 79, 033617 (2009).
- [20] H. Cao and L. Fu, Quantum phase transition and dynamics induced by atom-pair tunnelling of bose-einstein condensates in a double-well potential, The European Physical Journal D 66, 1 (2012).
- [21] A. Hemmerich, Bosons condensed in two modes with flavorchanging interaction, Physical Review A 99, 013623 (2019).
- [22] H. J. Lipkin, N. Meshkov, and A. Glick, Validity of many-body approximation methods for a solvable model:(i). exact solutions and perturbation theory, Nuclear Physics 62, 188 (1965).
- [23] J. v. Neumann, Beweis des ergodensatzes und des h-theorems in der neuen mechanik, Zeitschrift für Physik 57, 30 (1929).
- [24] X. Han and B. Wu, Entropy for quantum pure states and quan-

tum h theorem, Physical Review E 91, 062106 (2015).

- [25] J. Jiang, Y. Chen, and B. Wu, Quantum ergodicity and mixing and their classical limits with quantum kicked rotor, arXiv preprint arXiv:1712.04533 (2017).
- [26] Y. Fang, F. Wu, and B. Wu, Quantum phase space with a basis of wannier functions, Journal of Statistical Mechanics: Theory and Experiment 2018, 023113 (2018).
- [27] Z. Wang, J. Feng, and B. Wu, Microscope for quantum dynamics with planck cell resolution, Physical Review Research 3, 033239 (2021).
- [28] K. Sacha and J. Zakrzewski, Time crystals: a review, Reports on Progress in Physics 81, 016401 (2017).
- [29] R. Islam, R. Ma, P. M. Preiss, M. Eric Tai, A. Lukin, M. Rispoli, and M. Greiner, Measuring entanglement entropy in a quantum many-body system, Nature 528, 77 (2015).