

Wetting and nonwetting near a tricritical point

Joseph O. Indekeu¹ and Kenichiro Koga²

¹*Institute for Theoretical Physics, KU Leuven, BE-3001 Leuven, Belgium*

²*Research Institute for Interdisciplinary Science, Okayama University, Okayama 700-8530, Japan*
(Dated: November 8, 2022)

The dihedral contact angles between interfaces in three-fluid-phase equilibria must be continuous functions of the bulk thermodynamic fields. This general argument, which we propose, predicts a nonwetting gap in the phase diagram, challenging the common belief in “critical-point wetting”, even for short-range forces. A demonstration is provided by exact solution of a mean-field two-density functional theory for three-phase equilibria near a tricritical point (TCP). Complete wetting is found in a tiny vicinity of the TCP. Away from it, nonwetting prevails and no wetting transition takes place, not even when a critical endpoint is approached. Far from the TCP, reentrant wetting may occur, with a different wetting phase. These findings shed light on hitherto unexplained experiments on ternary H₂O-oil-nonionic amphiphile mixtures in which nonwetting continues to exist as one approaches either one of the two critical endpoints.

Wetting phenomena and in particular wetting phase transitions have enjoyed much attention from physicists and other scientists or engineers. The first theoretical predictions and experimental observations of a phase transition from a state of partial wetting (“nonwet”), in which three phases are pairwise in contact at their mutual interfaces and meet at a common contact line, to a state of complete wetting (“wet”), in which one phase intrudes at the interface between the other two, date from the late 1970s [1–3]. Not only first-order but also critical wetting transitions (of second or higher order) may occur [4, 5]. Numerous reviews have since been dedicated to the development of this fascinating field, e.g., [6–11].

In 1977 Cahn predicted that complete wetting must be expected upon approach of a critical point where two phases become identical. This is termed “critical-point wetting” (CPW) [1]. Subsequently, insight was gained in which systems CPW does occur and in which it does not. For short-range forces, CPW must take place when two coexisting phases adsorbed at a wall with a nonzero “surface field” approach bulk criticality, as found in [5] and [12] (surface phase transition class C defined in [12]). For a vanishing surface field CPW is suppressed [13–15]. For long-range wall-fluid and/or fluid-fluid forces, nonwetting gaps exist in which CPW does not occur, as predicted in [16, 17] and recently demonstrated in [12] (classes A, B and D defined in [12]).

In this Letter we ask whether CPW should be expected when three coexisting fluid phases are treated on equal footing instead of replacing one of them by an *ad hoc* “wall” boundary condition. We consider molecular fluids governed by van der Waals forces, or fluids with forces of shorter range, such as plasmas and electrolytes [6] or colloid-polymer mixtures [18]. CPW can be tested approaching a critical endpoint (CEP) where two (out of three) phases become identical and the resulting critical phase coexists with the third, noncritical, phase. Especially relevant is the vicinity of a tricritical point (TCP) at which two CEP lines meet, in accord with Gibbs’ phase rule. Pioneering experiments in this arena were carried out by Widom *et al.* [19, 20] and [20] was judged to be

consistent with CPW [21]. However, in ternary H₂O-oil-nonionic amphiphile mixtures nonwetting was observed to persist [22].

To shed light on these issues, we consider a mixture with three components and study three-phase equilibria with coexisting phases α , β and γ in mean-field density-functional theory (DFT) with two spatially varying densities ρ_1 and ρ_2 . Two densities are necessary because near a TCP, in a single-density theory the “middle” phase always wets the interface between the other two phases [23, 24]. Gibbs’ phase rule dictates that two linear combinations of pressure p , temperature T and chemical potentials μ_i , $i = 1, 2, 3$, can be taken as independent field coordinates, say s and t , in the two-dimensional three-phase coexistence surface in thermodynamic space. This surface is bounded by two CEP lines that meet tangentially (with a power $3/2$) at the TCP, located at $s = t = 0$, along an asymptote that we take to be the t -axis. We can use $t \geq 0$ to measure the “temperature distance” to the TCP, in the (s, t) -plane, and use s to interpolate between two CEPs at fixed t [19, 24, 25].

Of chief interest here are fluids with short-range forces, for which the single-density approximation with a wall boundary condition always predicts CPW. The model we employ is a square-gradient DFT akin to model T in [27], in which the bulk free-energy density is a product of three potential wells, centered about different points in the (ρ_1, ρ_2) -plane, with bulk phase density pairs (ρ_1^ν, ρ_2^ν) , with $\nu = \alpha, \beta, \gamma$. In the theory of tricritical phenomena due to Griffiths an important constraint reflects the asymptotic scaling properties of the three-phase region close to the TCP [24, 25]. Using linear combinations of the densities, which we rename to be just our ρ_1 and ρ_2 , the constraint on the bulk densities, in each phase, takes the form

$$\rho_2^\nu = -(\rho_1^\nu)^2. \quad (1)$$

The (dimensionless) interfacial free-energy density Ψ is

taken to be

$$\Psi(\rho_1, \rho_2) = \frac{1}{2} \sum_{i=1,2} \left(\frac{d\rho_i}{dz} \right)^2 + \prod_{\nu=\alpha,\beta,\gamma} \sum_{i=1,2} (\rho_i - \rho_i^\nu)^2. \quad (2)$$

This simple symmetric form has been chosen in the interest of obtaining an exactly solvable DFT. Additional parameters may affect the (order and locus of the) wetting transitions, as they do in [26], but the presence of a nonwetting gap, for which we will give a general argument, ought to be robust.

The bulk free-energy density is minimal and takes equal values (chosen to be zero) in all three bulk phases. The densities $\rho_1(z)$ and $\rho_2(z)$, with z the coordinate perpendicular to the interface between any two phases, characterize the structure of that interface. The equilibrium (or “optimal”) densities minimize the excess free-energy functional associated with the interface,

$$\sigma[\rho_1, \rho_2] = \int_{-\infty}^{\infty} dz \Psi(\rho_1, \rho_2), \quad (3)$$

subject to the boundary conditions that the interface connects two (spatially homogeneous) bulk phases that are a macroscopic distance apart. Therefore we impose one of the bulk phases, e.g., α , at $z = -\infty$ and the other, e.g., β , at $z = \infty$. The equilibrium interfacial tension $\sigma_{\alpha\beta}$ is then the value of $\sigma[\rho_1, \rho_2]$ for the optimal density profiles.

The densities ρ_1 for the three coexisting phases, calculated to be the zeroes of the third-degree polynomial $P(\rho_1) = \rho_1^3 - 3t\rho_1 + 2s$, are ordered in the manner $\rho_1^\alpha(s, t) \leq \rho_1^\beta(s, t) \leq \rho_1^\gamma(s, t)$ [24]. At fixed t , s can interpolate between two CEPs, one at $\alpha\beta$ criticality ($st^{-3/2} = -1$) and one at $\beta\gamma$ criticality ($st^{-3/2} = 1$). The results for $s < 0$ can be obtained from those for $s > 0$ by interchanging phases α and γ , so we consider $s \geq 0$ and investigate the range $0 \leq st^{-3/2} \leq 1$.

There are now two possibilities. Either the β phase does not wet the $\alpha\gamma$ interface, in which case

$$\sigma_{\alpha\gamma} < \sigma_{\alpha\beta} + \sigma_{\beta\gamma}, \quad \text{nonwet}, \quad (4)$$

or the β phase wets the $\alpha\gamma$ interface, and then,

$$\sigma_{\alpha\gamma} = \sigma_{\alpha\beta} + \sigma_{\beta\gamma}, \quad \text{wet}, \quad (5)$$

the latter of which is sometimes referred to as “Antonov’s rule” [24]. When β does not wet the $\alpha\gamma$ interface, it is possible that γ wets the $\alpha\beta$ interface (or that it does not).

We take a twofold approach. High-precision numerical integration of the free-energy density is performed to obtain the interfacial tensions. Also, recently conjectured simple analytic forms of these interfacial tensions [28] are used to calculate them exactly. The two methods provide indistinguishable results, while the analytic calculation is by far the simplest.

The analytic forms allow one to obtain a geometrical representation of the wetting criterion. To elucidate this

we recall the analytic expression for, say, $\sigma_{\alpha\gamma}$ uncovered in [28]. It is applicable to a three-phase triangle of arbitrary geometry and valid for a non-wet interface,

$$\sigma_{\alpha\gamma} = \frac{\sqrt{2}}{6} p^3 \ell, \quad (6)$$

with p the Euclidean distance from α to γ and ℓ the Euclidean distance from β to the midpoint of the $\alpha\gamma$ line, in the (ρ_1, ρ_2) -plane. Note that p is the length of an edge, and ℓ that of the conjugate median, in the three-phase triangle.




Figure 1. Two representative three-phase triangles, after a coordinate transformation from (ρ_1, ρ_2) to (x, y) , together with the two curves $y(x)$ that correspond to the exact phase boundaries for wetting of the $\alpha\gamma$ interface by the β -phase. The curves feature $y(0)^2 = 2\sqrt{3} - 3$ and $|dy/dx|_{x=\pm 1} = \sqrt{3}$. A (non)wet state results if β is (outside) inside the domain bounded by the curves.

We now construct the geometrical wetting criterion (see Fig.1). We translate, rotate and uniformly rescale the three-phase triangle without affecting its shape, in the (ρ_1, ρ_2) -plane. After this, the α -phase point is fixed at $(x, y) = (-1, 0)$ and the γ -phase point at $(x, y) = (1, 0)$. Next we express the wetting condition (5) in x and y , using (6) for each interface, assuming a nonwet state (none of the interfaces between any two phases is wet by the third phase). We find that supposing Antonov’s rule is equivalent to drawing two curves $y(x)$, which trace the boundary of the domain of wet states. An involved algebraic calculation, invoking Apollonius’ geometrical theorem, shows that the curves satisfy

$$y^2 = x^2 - 3 + 2\sqrt{x^4 - 3x^2 + 3}. \quad (7)$$

A state in which the $\alpha\gamma$ interface is wet by β results if and only if the β -phase point lies inside the domain bounded by these curves. The boundary corresponds to the wetting transition. A similar construction applies for wetting by α or γ . Consequently, a *necessary* condition for wetting of an interface between two phases by a third

two interfacial tensions are equal. These may act as “neutral lines” with respect to wetting. For example, the line $\sigma_{\alpha\beta} = \sigma_{\alpha\gamma}$ excludes wetting by β or γ , since $\hat{\beta} = \hat{\gamma}$. Some exactly calculated points on this line are ($s = 0, t = 1$), ($s = (1/\sqrt{3})^{3/2}/\sqrt{2}, t = 1/\sqrt{3}$) and ($s = (1/2)^{3/2}, t = 1/2$). The latter is a common point of the neutral line and the CEP line. It is characterized by $\hat{\alpha} = 180^\circ$ and, asymptotically, $\hat{\beta} = \hat{\gamma} = 90^\circ$. Next, we know that in the wet regimes the asymptotic angle is $\hat{\beta} = 0$ or $\hat{\beta} = 180^\circ$. If there were no nonwetting gap, the asymptotic value of $\hat{\beta}$ on the CEP would have to jump discontinuously from 0 to 180° at the neutral point, where, by symmetry, it is 90° .

A discontinuity of this caliber in a dihedral angle is possible, and does exist, in the single-density theory with a wall, when the surface field is varied through zero, asymptotically close to the critical temperature T_c . This is conspicuous in the wetting phase diagram in [5]. However, there are no surface fields or other boundary effects that can cause discontinuities in observable dihedral angles, when all three phases are treated on equal footing and only bulk thermodynamic fields are varied within the two-dimensional subspace of three-phase equilibria. Consequently, all angles $0 < \hat{\beta} < 180^\circ$ must be encountered in a continuous manner also when moving along, and infinitesimally close to, the CEP line. This implies the existence of a nonwetting gap.

This general argument is supported in detail by the exact solution of the DFT. Upon approach of a CEP, the $\beta\gamma$ interface becomes diffuse as β and γ become one and the same critical phase, named $\beta\gamma$. The angle $\hat{\alpha}$ approaches 180° , and α and $\beta\gamma$ each fill a half-space. They are separated by a planar non-critical interface, denoted by $\alpha, \beta\gamma$. For $t_u^- < t < t_u^+$, in the $\beta\gamma$ half-space the diffuse $\beta\gamma$ interface, with vanishing interfacial tension, makes a non-zero contact angle $\hat{\beta}$ with the $\alpha, \beta\gamma$ interface. The diffuse interface is still bound, or localized, at the non-critical interface in the limit that the CEP is attained.

From the analytic expressions for the interfacial tensions we calculate, using series expansion to third order in the small distance $\epsilon = 1 - st^{-3/2}$ from the CEP line, the asymptotic contact angle $\hat{\beta}$ along the CEP line. The exact result, displayed in Fig.3, is

$$\cos \hat{\beta} = \frac{(2t-1)(1-31t+4t^2)}{((1+t)(1+4t))^{3/2}}, \text{ for } t_u^- < t < t_u^+. \quad (10)$$

The values t_u^- and t_u^+ , which span the nonwetting gap, are found by solving $\cos^2 \hat{\beta} = 1$ (which leads to $t(1-7t+4t^2)^2 = 0$). When $t \downarrow t_u^-$, $\cos \hat{\beta} \rightarrow 1$. When $t \uparrow t_u^+$, $\cos \hat{\beta} \rightarrow -1$ and reentrant CPW is found.

Starting in the nonwetting gap and moving along, and infinitesimally close to, either one of the two CEP lines, we may encounter an interfacial phase transition in which the diffuse interface unbinds from the planar non-critical interface. These “diffuse-interface unbinding” transitions, at u^\pm , are critical and of second order,




Figure 3. Infinitesimally close to the CEP line for $\beta\gamma$ criticality, the cosine of the asymptotic contact angle $\hat{\beta}$ is a continuous function of the bulk thermodynamic field t . Second-order diffuse-interface unbinding occurs at $t = t_u^- \equiv (7 - \sqrt{33})/8$ where $\hat{\beta} = 0$, $\hat{\gamma} = 180^\circ$ and wetting by β terminates at a CEP, and also at $t = t_u^+ \equiv (7 + \sqrt{33})/8$ where $\hat{\beta} = 180^\circ$, $\hat{\gamma} = 0$ and wetting by γ terminates at a CEP. The three-phase configuration for $t = 1/2$ where a neutral line (see Fig.2) terminates at a CEP is also shown.

with

$$\cos \hat{\beta} \pm 1 \propto (t - t_u^\pm)^2, \text{ for } st^{-3/2} = 1^-, \quad (11)$$

Our findings make it necessary to reinterpret the experimental observation of a hitherto unexplained nonwetting state, which continues to exist when either one of the two CEPs is approached in ternary H_2O -oil-nonionic amphiphile mixtures at three-phase coexistence [22]. A drop of the “middle” phase was observed not to spread but to form a lens, with a contact angle close to 90° . It was thought that this angle must approach exactly 90° close to the CEP based on a scaling argument. Other values, $0 < \hat{\beta} < 180^\circ$, were, unfortunately, not given attention because the authors excluded the very case ($x = y$ in [22]) which is realized in the nonwetting gap of our exactly solved two-density DFT.

In conclusion, for three-fluid-phase equilibria in systems with a TCP that are not approximated by two-phase equilibria at a wall, but described by a two-density DFT, the global wetting phase diagram largely contradicts the CPW scenario. A pronounced nonwetting gap is found in an exactly solved DFT paradigm and, we argue, must also generally be present by virtue of thermodynamic continuity of the dihedral angles as a function of bulk thermodynamic field variables, in three-phase equilibria of fluids.

We are grateful to Ben Widom for his insights and interest in this research. We thank Jonas Berx for discussions. We are indebted to the three reviewers for useful remarks. JOI thanks Okayama University for generous hospitality, the Japan Society for the Promotion of Science for a JSPS Invitational Fellowship for Research in

Japan with ID No. S18131 and FWO-Flanders for grant K204422N for a short stay. This work was supported

in part by JSPS KAKENHI (Grant Nos. 18KK0151, 20H02696).

-
- [1] J.W. Cahn, *Critical point wetting*, J. Chem. Phys. **66**, 3667 (1977)
 - [2] C. Ebner and W. F. Saam, *New phase-transition phenomena in thin argon films*, Phys. Rev. Lett. **38**, 1486 (1977)
 - [3] M.R. Moldover and J.W. Cahn, *An interface phase transition: complete to partial wetting*, Science **207**, 1073 (1980)
 - [4] D.E. Sullivan, *Surface tension and contact angle of a liquid-solid interface*, J. Chem. Phys. **74**, 2604 (1981)
 - [5] H. Nakanishi and M.E. Fisher, *Multicriticality of wetting, prewetting, and surface transitions*, Phys. Rev. Lett. **49**, 1565 (1982)
 - [6] P.-G. de Gennes, *Wetting: statics and dynamics*, Rev. Mod. Phys. **57**, 827 (1985)
 - [7] D.E. Sullivan and M.M. Telo da Gama, *Wetting transitions and multilayer adsorption at fluid interfaces*, in Fluid interfacial phenomena **10**, 45 (1986), ed. C.A. Croxton, Wiley, Chichester
 - [8] S. Dietrich, *Wetting phenomena*, in Phase Transitions and Critical Phenomena, vol. 12, p.1, eds. C. Domb and J.L. Lebowitz, Academic Press, London (1988)
 - [9] G. Forgacs, R. Lipowsky and T.M. Nieuwenhuizen, *The behavior of interfaces in ordered and disordered systems*, in Phase transitions and critical phenomena, vol. 14, p.135, eds. C. Domb and J.L. Lebowitz, Academic Press, London (1991)
 - [10] D. Bonn and D. Ross, *Wetting transitions*, Rep. Prog. Phys. **64**, 1085 (2001)
 - [11] D. Bonn, J. Eggers, J.O. Indekeu, J. Meunier and E. Rolley, *Wetting and spreading*, Rev. Mod. Phys. **81**, 739 (2009)
 - [12] R. Evans, M.C. Stewart and N.B. Wilding, *A unified description of hydrophilic and superhydrophobic surfaces in terms of the wetting and drying transitions of liquids*, PNAS **116**, 23901 (2019)
 - [13] J.O. Indekeu, *How universal is critical-point wetting?*, Physica A **177**, 428 (1991)
 - [14] A. Sevrin and J.O. Indekeu, *Critical-point wetting near grain boundaries and defect planes*, Phys. Rev. B **39**, 4516 (1989)
 - [15] F. Igloi and J.O. Indekeu, *Wetting near grain boundaries and defect planes, and its connection with wetting at walls and free surfaces*, Phys. Rev. B **41**, 6836 (1990)
 - [16] M.P. Nightingale and J.O. Indekeu, *Examination of the necessity of complete wetting near critical points in systems with long-range forces*, Phys. Rev. B **32**, 3364 (1985)
 - [17] C. Ebner and W. F. Saam, *Effect of long-range forces on wetting near bulk critical temperatures: An Ising-model study*, Phys. Rev. B **35**, 1822 (1987)
 - [18] H. N. W. Lekkerkerker, V. W. A. de Villeneuve, J. W. J. de Folter, M. Schmidt, Y. Hennequin, D. Bonn, J. O. Indekeu and D. G. A. L. Aarts, *Life at Ultralow Interfacial Tension: wetting, waves and droplets in demixed colloid-polymer mixtures*, Eur. Phys. J. B **64**, 341 (2008)
 - [19] J.C. Lang, B. Widom, *Equilibrium of three liquid phases and approach to the tricritical point in benzene-ethanol-water-ammonium sulfate mixtures*, Physica **81A**, 190 (1975)
 - [20] N. Nagarajan, W.W. Webb and B. Widom, *Surface tension of a two-component liquid mixture near its critical solution point*, J. Chem. Phys. **77**, 5771 (1982)
 - [21] S.-Y. Zinn and M.E. Fisher, *Interfacial tensions near critical endpoints: experimental checks of EdGF theory*, Mol. Phys. **103**, 2927 (2005)
 - [22] M. Kahlweit and G. Busse, *Wetting in mixtures of water, nonionic amphiphiles, and nonpolar solvents*, J. Chem. Phys. **91**, 1339 (1989)
 - [23] B. Widom, *Interfacial tensions of three fluid phases in equilibrium*, J. Chem. Phys. **62**, 1332 (1975).
 - [24] J.S. Rowlinson, B. Widom, *Molecular Theory of Capillarity* (Oxford: Clarendon) (1982)
 - [25] R.B. Griffiths, *Thermodynamic model for tricritical points in ternary and quaternary fluid mixtures*, J. Chem. Phys. **60**, 195 (1974)
 - [26] K. Koga, J.O. Indekeu, B. Widom, *Infinite-order transitions in density-functional models of wetting*, Phys. Rev. Lett. **104**, 036101 (2010)
 - [27] K. Koga, B. Widom, *Density functional models of the interfacial tensions near the critical endpoints and tricritical point of three-phase equilibria*, J. Phys.: Condens. Matter **28**, 244016 (2016)
 - [28] K. Koga and J.O. Indekeu, *Three-phase equilibria in density-functional theory: Interfacial tensions*, J. Chem. Phys. **150**, 164701 (2019)