

Simulated performance of the molecular mapping for young giant exoplanets with the Medium Resolution Spectrometer of JWST/MIRI

M. Mâlin¹, A. Boccaletti¹, B. Charnay¹, F. Kiefer¹, and B. Bézard¹

LESIA, Observatoire de Paris, Université PSL, CNRS, Sorbonne Université, Université Paris Cité, 5 place Jules Janssen, 92195 Meudon, France.

Received ; accepted

ABSTRACT

Context. Young giant planets are the best targets for characterization with direct imaging. The Medium Resolution Spectrometer (MRS) of the Mid-Infrared Instrument (MIRI) of the recently launched James Webb Space Telescope (JWST) will give access to the first spectroscopic data for direct imaging above 5 μm with unprecedented sensitivity at a spectral resolution up to 3700. This will provide a valuable complement to near-infrared data from ground-based instruments for characterizing these objects.

Aims. We aim to evaluate the performance of MIRI/MRS to detect molecules in the atmosphere of exoplanets and to constrain atmospheric parameters using Exo-REM atmospheric models.

Methods. The molecular mapping technique, based on cross-correlation with synthetic models, has been introduced recently. This promising detection and characterization method is tested on simulated MIRI/MRS data.

Results. Directly imaged planets can be detected with MIRI/MRS, and we are able to detect molecules (H_2O , CO , NH_3 , CH_4 , HCN , PH_3 , CO_2) at various angular separation depending on the strength of the molecular features and brightness of the target. We find that the stellar spectral type has a weak impact on the detection level. This method is globally most efficient for planets with temperatures below 1500 K, for bright targets and angular separation greater than 1". Our parametric study allows us to anticipate the ability to characterize planets that would be detected in the future.

Conclusions. The MIRI/MRS will give access to molecular species not yet detected in exoplanetary atmospheres. The detection of molecules as indicators of the temperature of the planets will make it possible to discriminate between the various hypotheses of the preceding studies, and the derived molecular abundance ratios should bring new constraints on planetary formation scenarios.

Key words. Infrared: planetary systems – Techniques: imaging spectroscopy – Planets and satellites: atmospheres - Planets and satellites: gaseous planets - Methods: data analysis Space vehicles: instruments

1. Introduction

An important outcome of exoplanet searches in the last decades is the diversity of their orbital and bulk properties. Understanding the mechanisms at play during their formation, or migration history is identified as a promising avenue to account for such a diversity (Madhusudhan et al. 2014; Mordasini et al. 2016). The characterization of exoplanet atmospheres has now become a priority in this field, one goal being precise to put some meaningful constraints on their formation. In particular, according to formation models (Oberg & Bergin 2021), measuring molecular abundances, like the ratios C/O and N/O, are relevant to link the atmospheric properties of giant planets to the locations of the snowlines in a planetary system.

A wide range of methods has been used to explore the properties of exoplanetary atmospheres. The first molecules were detected with transit spectroscopy (Charbonneau et al. 2002). This successful method enables to observe transmission spectra of the day-night terminator, the thermal emission spectra of the day side, and the phase curve on the orbit (e.g. Deming et al. 2013), but is limited to planetary systems whose semi-major axis is less than ~ 1 au as it suffers from the low probability that the planet is perfectly aligned with the observer. Phase-resolved high-resolution Doppler spectroscopy has proven to be a powerful means to detect molecules in the atmosphere of transiting

close-in giant planets. For instance, this led to the detection of CO in several hot jupiters like HD 209458 b by Snellen et al. (2010), or τ boo by Brogi et al. (2012) as well as H_2O by Birkby et al. (2013) in HD 189733 b. In the case of long-period planets, direct imaging with coronagraphy can also provide spectral information, although mostly at low to medium resolution. Owing to the inherent contrast limitation, post-processing methods to attenuate the starlight such as Angular Differential Imaging (Marois et al. 2006) or Spectral Differential Imaging (Racine et al. 1999), were decisive to perform observations of young giant planets being warm and bright in the infrared. A dozen systems have been characterized by spectroscopy with adaptive optics (AO), for example, 51 Eri b with SPHERE/VLT and GPI (Samland et al. 2017; Macintosh et al. 2015), and with GRAVITY/VLTI the planet β Pictoris b (Nowak et al. 2020b).

Considering the best of both worlds, Snellen et al. (2015) proposed to combine high contrast imaging with high resolution spectroscopy, while some similar concepts were already formulated earlier (e.g. Sparks & Ford 2002). An implementation of this original idea was introduced as the so-called "molecular mapping" technique by Hoeijmakers et al. (2018) yielding the detection of H_2O and CO in β Pic b, atmosphere, taking advantage of archival VLT/SINFONI data. Recently, with the same instrument, Petrus et al. (2021) used this method to characterize

the planet HIP 65426 b. In contrast, no molecular species was found in the planets of PDS 70 (Cugno et al. 2021), likely because the planet’s atmosphere or surroundings is dominated by dust. Similarly, Petit dit de la Roche et al. (2018) and Ruffio et al. (2019) took advantage of cross correlation with molecular templates to characterize the HR 8799 system with Keck/OSIRIS IFS data.

As of today, all direct observations were obtained in the near IR because of the reduced transmission of the Earth’s atmosphere in the mid-IR. The James-Webb Space Telescope (JWST) is expected to be a game-changer in the characterization of directly imaged exoplanet atmospheres (Hinkley et al. 2022), in allowing us to explore a relatively new spectral range for wavelengths longer than $5\mu\text{m}$ where planets emit most of their flux (implying smaller brightness ratio between the star and a planet), and exhibit clear molecular signatures from their atmosphere. Complementary to near IR data, a broader wavelength coverage will help to recover for example the temperature of the planet with higher accuracy, as already shown with the early results of JWST coronagraphy on HIP 65426 b (Carter et al. 2022).

Starting in 2022, MIRI, one of the science instruments of JWST, optimized for mid-infrared observations (Wright et al. 2015), is offering a unique opportunity for exoplanet science. MIRI has four observing modes: imaging, coronagraphy, low-resolution spectroscopy (LRS), and medium-resolution spectroscopy (MRS). The MRS provides integral field spectroscopy across the wavelength range 4.9 to $28.3\mu\text{m}$ (Wells et al. 2015), which constitute interesting features for exoplanet atmospheres characterization. As one of the early outcomes of JWST programs, Miles et al. (2022) illustrated the potential of MIRI MRS on the planetary-mass companion VHS 1256 b, for which several molecules were detected (CH_4 , CO , CO_2 , H_2O , K , Na).

Following the work by Patapis et al. (2022), who demonstrated the potential of molecular mapping for MIRI in two well-known systems, HR 8799 and GJ 504, we aim to further explore this concept with the self-consistent atmosphere model Exo-REM (Charnay et al. 2018) using a parametric study and extending to other known directly imaged planets.

The paper is organized as follows: Section 2 presents the data simulation and reduction for the MRS. Section 3 introduces the molecular mapping method we implemented. Section 4 provides a parametric study and section 5 the application to a few known directly imaged systems. Section 6 presents a more in-depth atmospheric study for the target GJ 504 b and Section 7 discusses the results.

2. Data simulation and reduction for the MRS

2.1. The Medium Resolution Spectrometer of MIRI

The MRS is one of the four observing modes of MIRI. It is an integral field spectrometer that provides diffraction-limited spectroscopy between $4.9\mu\text{m}$ and $28.3\mu\text{m}$, within a field-of-view (FoV) ranging from $3.7'' \times 3.7''$ at the shortest wavelengths to $7.74'' \times 7.95''$ at the longest wavelengths. The MRS includes four channels that have co-aligned FoV, observing the wavelength ranges simultaneously. Three observations using each time a different set of gratings are needed to observe the entire wavelength range (SHORT, MEDIUM, LONG, Wells et al. 2015). The spectral resolution decreases with increasing wavelength. The parameters of each subchannel (bands) are indicated in Tab. 1. The MRS is spatially undersampled at all wavelengths and mostly in the first channel, therefore dithering is necessary to improve this spatial sampling. Different dithering patterns are possible and

depend on each scientific case, mainly the 4-point dither pattern is preferred as it provides robust performance at all wavelengths and adequate point source separation in all channels.

The minimum integration time of the MRS detector in full frame is $t_{\text{fast}} = 2.775\text{ s}$. One integration is a ramp composed of several groups (N_{group}), and an exposure is made of several integrations (N_{int}). A reset is applied after each ramp (overhead $= t_{\text{fast}}$). Therefore, a series of multiple exposures (N_{exp}) corresponds to an actual integration time of $N_{\text{exp}} \times N_{\text{int}} \times N_{\text{group}} \times t_{\text{fast}}$, and an observation time (including overheads) of $N_{\text{exp}}(N_{\text{int}} \times N_{\text{group}} \times t_{\text{fast}} + (N_{\text{int}} - 1) \times t_{\text{fast}})$.

2.2. MIRISim

MIRISim is a software to simulate representative MIRI data (Klaassen et al. 2020) that incorporates the best knowledge of the instrument. The simulation takes into account effects due to the detectors, slicers, distortion, and noise sources. MIRISim outputs are the detector images in the uncalibrated data format that can be used directly in the JWST pipeline. In this work, we used version 2.4.1¹. The simulations are parameterized using three configuration files that define the astronomical scene, the setup of the instrument and the parameters of the simulator itself, as described in the following.

Scene. For this study, the scene is composed of a host star and one or several planetary companions. Each object is simulated by attributing a spectrum and its position in the FoV that are calculated based on the known astrometric positions. Low background emission is added.

Simulation parameters. The number of groups, integrations, and expositions are determined using the Exposure Time Calculator (ETC)² in order to avoid saturation on the detector and to obtain the signal-to-noise ratio desired. The PSF of the MRS is under-sampled by design, a well-sampled PSF requires that the object is observed in at least two dithered positions that include an offset as explained in Wells et al. (2015). We chose to do our simulations using the 4-point dithering pattern. Of the two possible detector read modes we selected the FAST mode (2.775 s per frame) which is more appropriate for bright targets. The grating position, as well as the observing channel, are also specified in this file.

Simulator. The last configuration file defines the various noise components. We apply Poisson noise (for each object in the scene including the background), bad pixels, dark current, hot pixels, flat-field, gain, and non-linearity. Moreover, we include the effect of fringes, detector drifts, and latency. The cosmic ray environment is set to define a minimum solar environment. We note that MIRISim is producing excess noise on the integration ramps using the FAST mode, so it is advised to turn off the read noise component.

Channel 4 suffers from a drop in sensitivity (Glasse et al. 2015). Therefore, we do not expect to achieve the planetary mass regime at such wavelengths, and we intentionally omit wavelengths larger than $18\mu\text{m}$.

2.3. JWST Pipeline

The steps of the JWST pipeline for the MRS are detailed in Labiano-Ortega et al. (2016). Starting with the detector images simulated with MIRISim, the pipeline is divided into three successive steps: *calwebb_detector1*, *calwebb_spec2*,

¹ <https://wiki.miricle.org/bin/view/Public/>

² <https://jwst.etc.stsci.edu>

Channel	Band	Wavelength (μm) ¹	Resolution (best estimate) ¹	FoV (arcsec) ²	Pixel size (arcsec) ²
1	SHORT (A)	4.885 – 5.751	3300-4000	3.70×3.70	0.196
	MEDIUM (B)	5.634 – 6.632	3420-3990		
	LONG (C)	6.408 – 7.524	3330-3840		
2	SHORT (A)	7.477 – 8.765	3190 - 3620	4.71×4.52	0.196
	MEDIUM (B)	8.711 – 10.228	3040 - 3530		
	LONG (C)	10.017 – 11.753	2890 - 3374		
3	SHORT (A)	11.481 – 13.441	2450 - 3010	6.19×6.14	0.245
	MEDIUM (B)	13.319 – 15.592	2300 - 2460		
	LONG (C)	15.400 – 18.072	2020 - 2790		
4	SHORT (A)	17.651 – 20.938	1400 - 1960	7.74×7.95	0.273
	MEDIUM (B)	20.417 – 24.220	1660 - 1730		
	LONG (C)	23.884 – 28.329	1340 - 1520		

Table 1. MRS instrumental parameters in each MRS band. ¹ Labiano et al. (2021), ²Wells et al. (2015)

calwebb_spec3, each including several intermediate steps that are listed below. In this work, we use version 1.4.0³ of the pipeline that is compliant with the version used for MIRISim. In appendix A, we provided the relevant steps to reduce the simulated data of MIRI/MRS.

2.4. Background treatment

The PSF of bright stars we are studying extends across almost all the field of view. It is therefore impossible to define a region where the pipeline could estimate and subtract the background directly from the science image. To overcome this issue, we simulated a scene with only the background emission, all other MIRISim configuration files (simulation and simulator parameters) being the same as those used for the astrophysical target simulation. This simulated background goes through stage 1 of the pipeline to correct for detector effects and is subtracted from target exposures using the step *background* in stage 2.

3. Molecular mapping method

3.1. Atmospheric models

The basic concept of molecular mapping relies on the correlation of spectro-imaging data with a model of the exoplanet atmospheres we are trying to detect. In the following, we will use Exo-REM, a self-consistent 1D radiative-equilibrium model. It has been first developed to simulate the atmospheres and spectra of young giant exoplanets (Baudino et al. 2015; Charnay et al. 2018) and more recently extended to irradiated planets (Blain et al. 2021). This model has been used to characterize some directly imaged planets at low and medium spectral resolution (e.g. Delorme et al. 2017; Bonnefoy et al. 2018; Petrus et al. 2021). The radiative-convective equilibrium is solved by assuming that the net flux (radiative and convective) is conservative. The conservation of the flux over the pressure grid (64 pressure levels) is solved iteratively using a constrained linear inversion method. The input parameters of the model are the effective temperature of the planet, the acceleration of gravity at 1 bar, and the elemental abundances. The model includes non-equilibrium chemistry comparing chemical reaction timescales and vertical mixing, using parametrizations from Zahnle & Marley (2014). The cloud scheme is detailed in Charnay et al. (2018); it takes into account micro-physics and simulates the formation of silicate, iron, sulfide, alkali salt, and water clouds. The cloud dis-

tribution is computed by taking into account sedimentation and vertical mixing with realistic eddy mixing coefficient K_{zz} profiles based on the mixing length theory. It takes into account Rayleigh scattering from H_2 , He, and H_2O , as well as absorption and scattering by clouds – calculated from extinction coefficient, single scattering albedo, and asymmetry factor interpolated from pre-computed tables for a set of wavelengths and particle radii. Sources of opacity include the H_2 – H_2 , H_2 –He, H_2O – H_2O and H_2O –air collision-induced absorption, ro-vibrational bands from molecules (H_2O , CH_4 , CO , CO_2 , NH_3 , PH_3 , TiO , VO , H_2S , HCN , and FeH), and resonant lines from Na and K. Lines lists used in Exo-REM are indicated in Blain et al. (2021).

In our simulations, the planetary spectra are modeled with Exo-REM. We built a grid of models using the ranges of parameters provided in Tab. 2. In particular, we considered clouds of iron and silicates (forsterite), and the particle radii are computed with simple microphysics in the cloud scheme. This method is based on the comparison of the timescales of the main physical processes involved in the formation and growth of cloud particles, which includes a supersaturation factor S , that we fixed at $S = 0.03$. This model reproduces the L-T transition, with the passage of clouds below the photosphere at the transition. Therefore, for the T-types, the clouds are forming below the photosphere and have a weak impact on spectra. The clouds are calculated in a self-consistent way depending on the condensation curves at each temperature (Visscher et al. 2010).

Parameters	Values	steps
Temperature (K)	400 - 2000	50
Logg	3.0 - 5.0	0.5
C/O	0.1 - 0.8	0.05
Metallicity	0.32 ; 1.0 ; 3.16 ; 10.0	

Table 2. Exo-REM grid models

As for the molecular templates, they are computed from the pressure-temperature profile at equilibrium and from the abundance profiles that were previously calculated. The radiative transfer is computed again with all chemical species removed, except the one considered. The clouds are also removed but the collision-induced absorption (H_2 – H_2 , H_2 –He, H_2O – H_2O) is still included.

Stellar spectra are taken from the BT-NextGen online libraries⁴. For stars cooler than 3000 K we used BT-Settl models.

³ <https://jwst-pipeline.readthedocs.io/en/latest/>

⁴ <http://svo2.cab.inta-csic.es/theory/newov2/index.php>

3.2. Subtracting stellar contribution and cross-correlation

The stellar contribution in high-angular resolution data is a mixture of the ideal diffraction pattern, and speckles due to optical aberrations, whose intensity scales with the star's spectrum, and the phase-induced chromaticity of speckles. Both scales radially with the wavelength, at first order. A planet buried in the diffracted halo and a star have very different spectral dependence, hence they can be disentangled (Sparks & Ford 2002). In the infrared, and in space conditions (no telluric lines), the atmospheric signature of a giant planet would appear as a high spectral frequency as opposed to the star, due to molecular absorptions. Therefore, as a prerequisite to apply the correlation with a model, the stellar contribution can be greatly attenuated by high-pass filtering while preserving the molecular signatures of the planet's spectrum almost intact (Ruffio et al. 2019). In our case, we used a Gaussian filter to suppress low frequencies on each spaxel (spectral pixel) of the cube. We adopt experimentally a filter parameter of $\sigma = 10$ which globally maximizes the detection of the simulated planets in our sample. Prior to applying the correlation, the Exo-REM models are degraded to the maximum resolution of the MRS (3700 in the first band 1A) and interpolated on the wavelength values of each MRS channel. The very same high-pass filter is applied to the Exo-REM models. Finally, we calculated the cross-correlation function (CCF) between the model and the data (high-pass filtered) for each velocity offset (δV) between the two spectra. Models and data spectra which are provided at a constant $\delta\lambda$ in MIRISim, are converted to velocity and re-interpolated to get a constant step in velocity. We used the python function `scipy.signal.correlate` to perform the correlation between two spectra. An example of the process in two different spaxels is shown in Fig. 1, one at the position of the planet in red, and the other one at an arbitrary position, in a noise-limited region, in pink. The cross-correlation function shows a peak of correlation at a radial velocity $\delta V = 0$. Looking at the spaxel away from the position of the planet, no peak of correlation is observed. We note that the MRS does not have a high enough spectral resolution to resolve the Doppler shift of the known imaged planets. Therefore, no Doppler shift is included in our simulation, and we focus only on the value at $\delta V = 0$ of the correlation function. The method is repeated independently on each spaxel to derive a correlation coefficient map at $\delta V = 0$ in which a planet would correspond to the highest correlation in the FoV (Fig. 1). The value of the correlation map in each position i, j is given by the equation 1, with M the model spectra and S the spectra from the data.

$$C_{i,j} = \frac{\sum_{\lambda} S(\lambda)_{i,j} \times M(\lambda)_{i,j}}{\sqrt{\sum_{\lambda} S(\lambda)_{i,j}^2 \times \sum_{\lambda} M(\lambda)_{i,j}^2}} \quad (1)$$

3.3. Signal to noise ratio calculations

To evaluate the signal-to-noise (S/N) ratio, first Hoeijmakers et al. (2018) measured the average standard deviation of the CCF in an annulus away from the peak of correlation, and away from the planet's position, to avoid systematic variations in the CCF at the location of the planet due to the autocorrelation. The autocorrelation function arises from all the harmonics in a molecule's spectrum which produce a non-zero correlation signal away from $\delta V = 0$. For instance, the CO generates secondary correlation peaks which can be almost as strong as the main correlation peak (Fig. 2). To account for the autocorrelation, Cugno et al. (2021) added a correction. The autocorrelation function of the model

spectrum is calculated and subtracted away from the peak of the CCF. However, the impact of the autocorrelation signal justifies taking into account the spatial dimension in estimating the noise. Petrus et al. (2021) measured the noise as the standard deviation of a Gaussian distribution derived from all the spaxels, excluding those containing the planet's signal (Fig. 3). The correlation signal of the planet is averaged in the velocity space around the correlation peak and spatially in a region centered on the planet. This method is also used by Patapis et al. (2022), who measured the signal as the mean value of the CCF in an aperture centered at the position of the planet. This measurement assumes that the noise follows a Gaussian distribution which is not always the case depending on the instrument and on the residuals left after stellar subtraction.

These methods are also conservative since the planet's signal can be integrated in principle over several pixels. Nevertheless, measuring the S/N of a planet in the footprint of its correlation pattern is not straightforward. In Appendix B.1, we derive how the size of the correlation pattern varies at first order. We find that it is a function of the wavelength-dependent PSF size, the template used for the correlation, and the noise level. However, the derived formula is too approximate to be used to measure the size of the correlation pattern in the data.

To obtain a robust measurement of the S/N , we defined a new method compliant with both a low level and a high level of detection, and also able to deal with the residual correlations of the stellar spectrum itself, particularly important if the planet's and star's temperatures are close, like for late-type stars. Our method also accounts for the spatial variation of the noise to avoid being limited by the autocorrelation signal as noticed in Petrus et al. (2021). Hiding the planets with a radius of 6 spaxels (maximum size of the correlation pattern estimated experimentally for a single planet in the image), we measured the noise as the standard deviation of all the other spaxels, ie. in the correlation map at $\delta V = 0$. Based on the parametric study (Sect. 4), we note that the strength of the correlation depends on the separation between the star and the planet. In addition, if the data are noisier, the correlation pattern is smaller. Finally, the width of the correlation pattern also scales with the wavelength as does the PSF. To complement the formalism in Appendix B.1, we present more figures in Appendix B.2 to demonstrate these effects on simulated data. Concerning the astrometry, the maximum of the correlation pattern does not necessarily correspond to the real position of the planet, and this effect is more important at small angular separations. Indeed, the stellar flux can contaminate the spaxels located at the planet's position, so the net effect is a higher correlation value further out in the planet's signature. Therefore, we stress that the astrometry of a companion based on the correlation map is unreliable at high noise levels and/or short angular separations. Finally, in all of our MRS simulations, we notice that the correlation pattern decreases at increasing wavelengths, especially because of a loss of sensitivity and higher background level impacting the longer wavelengths. We also note that molecular features tend to become shallower at longer wavelengths. Given these observed behaviors, we chose to measure the S/N in correlation maps only spatially, and we define the size of the planet's correlation pattern (containing N_S spaxels) experimentally based on its radial profile. We imposed a maximum size of 6 spaxels for all channels. As a first criterion, we selected the spaxels (red area Fig. 4 to display a case with higher noise level) whose correlation value is higher than 3 times the noise (measured in the blue area of Fig. 4). To ensure that we do not integrate noise in the signal since the correlation pattern is not circular, we imposed a second criterion by selecting the spax-




Fig. 1. Top: Simulation in channel 1A for a star ($T = 6000$ K) and a planet ($T = 1000$ K) separated by $1.8''$. From left to right: direct image resulting from four dithered positions, the star being offset in the bottom left corner (sum over the wavelengths of the channel); high frequencies residuals after subtracting the spectral Gaussian filter in each spaxel; correlation map with the very same template spectra as injected into the simulation for $\delta V = 0$. The red cross indicates the planet’s position, and the pink cross is the arbitrary position chosen away from the planet. Bottom: Illustration of the molecular mapping technique in two spaxels, one at the position of the planet (red) and the other one at a position away from the planet (pink), for channel 1A ($4.885 - 5.751 \mu\text{m}$). From left to right: we display the filtering process applied to the model, the combined spectra and the Gaussian filter (black) in the two spaxels (pink and red), the high-frequency component after subtraction, and the cross-correlation function for $\delta V = [-2000; +2000]$ km/s.




Fig. 2. Autocorrelation of each molecule spectrum (at $T = 550$ K) in the band of the spectral features.

els in which the correlation is larger than 50% of the maximum correlation. In addition, to account for the particular situation where the whole profile is above 50% of the maximum correlation, for instance, if the correlation with the star itself dominates the pattern (mainly the case of CO, or a hot planet around a cold star), we only use the central spaxel to measure the correlation peak. Finally, the S/N is calculated with Eq. 2, where σ is the standard deviation of the noise, and C_i the correlation values for the N_S spaxels.

$$S/N = \frac{\sum_i C_i}{\sqrt{N_S} \times \sigma} \quad (2)$$

4. Parametric analysis of MIRI/MRS detection capacity

To evaluate the detection limit of the MRS with molecular mapping, we run two sets of simulations, and we study the impact of the spectral type and the angular separation on a planet’s detection, and on the detection of each molecule included in Exo-REM. The first set of simulations (Sect. 4.1) allows us to restrain the parameters space to pursue this parametric study.




Fig. 3. Histogram of the pixels in the correlation map (grey area and red model), compared to the correlation coefficient at the planet’s location, as proposed in Petrus et al. (2021) and used in Patapis et al. (2022).

4.1. Impact of the spectral type

As the molecular mapping method relies on the fact that planetary and stellar spectra are different, we investigated the impact of the spectral type for both the star and the planet. We defined a set of 21 simulations (with a star and a planet in each simulation) by varying the planet temperature from 500 K to 2000 K by steps of 250 K, and we assumed 3 stellar temperatures of 3000 K, 6000 K and 9000 K, typically corresponding to M, G, and A type stars. In order to study only the impact of the spectral features, we considered a non-realistic situation in which the stellar flux, as well as the planet-to-star contrast, are kept constant for all 21 cases. The data simulation is done with $N_{group} = 26$ and $N_{int} = 13$ for a total exposure of 1 hour determined to achieve sufficient S/N on the planet (located at an angular separation of $1.4''$), in a reasonable computing time. The contrast at $5 \mu\text{m}$ is set to 10^3 . The model spectrum to calculate the correlation is identical to the input spectrum.




Fig. 4. Mean azimuthal profile of the correlation pattern (left) and typical correlation map to illustrate the S/N measurement (right), the same configuration as in Fig. 1. In blue the pixels used to evaluate the noise as the standard deviation of the distribution. In red, the pixels that are considered for the signal of the planet, as defined in Sect. 3.3. Top: Simulation with a low noise level (example with a star at $1.8''$ from the planet) Bottom: Simulation with a higher noise level (example with a star at $0.6''$ from the planet).

The S/N measured for the 3 first MRS channels are displayed in Fig. 5 (similar results are observed if we look only at a single band). Globally, we find that colder planets are easier to detect with molecular mapping, as a result of molecular lines being more pronounced in the planetary spectrum. On the contrary, the hottest planet in our sample ($T_p = 2000$ K) features a much higher correlation with the stellar spectrum and may become almost undetectable, a feature that is even more remarkable in channels 2 and 3. These results stand regardless of the stellar spectrum, but we note that, as expected, the detection is globally worse for a colder star, which has more spectral features ($T_s = 3000$ K). We also observe a general trend of a lower correlation signal with increasing wavelength, with channel 1 providing the highest detection.

4.2. Planets spectral type and angular separation

Guided by the former analysis, we chose a single stellar temperature ($T_s = 6000$ K) and considered more realistic simulations in which the system is located at 30 pc. Planet fluxes were calculated for the same temperature range as in Sect. 4.1 and for a radius of $1 R_{Jup}$. We tested the dependency of the molecular mapping efficiency to the planet's temperature and to its angular separation from the star. For convenience, the planet was positioned at the center of the FoV, while the position of the star was offset from $0.2''$ to $3.2''$. Compared to the simulations in Sect. 4.1, the planets here have potentially lower fluxes, so we generated 2 hours of observation with $N_{group} = 33$, and $N_{int} = 19$ (again with the goal to minimize computing time).

Fig. 6 displays the S/N in channels 1, 2, and 3, for each planet's temperature as a function of the angular separation from the star. In general, the S/N does not depend only on the planet's flux, since the continuum is filtered out while suppressing the star's contribution so the high frequency is the dominant factor

in the correlation. The highest S/N values are observed for temperatures ranging between 750 K and 1750 K, while lower performances are obtained for the coldest ($T_p = 500$ K) and the warmest ($T_p = 2000$ K) planets, because of a lower absolute flux, and respectively, a lower level of correlation due to fewer absorption lines. The S/N increases rapidly with the angular separation up to about $\sim 1.5''$, and then becomes asymptotic (at least in channels 1 and 3). Channel 2 shows a more gradual increase of the S/N with the separation.

The same parametric study is performed for each individual molecule, focusing on the channel or the band in which the detection is optimal (Fig. 7). The molecules are detected in the bands for which the absorption is the largest, and obviously departing the most from the stellar spectrum, as long as this absorption is not hidden by another molecule's absorption. Molecules with spectral features spanning a wider range of wavelengths will benefit from calculating the S/N in the cube built over the three sub-bands of one channel.

- H_2O is the prominent molecule in a planet's spectrum for any temperature. It is detected in every channel but mostly in the first one, except for cold planets where CH_4 will dominate in channel 2 and hide H_2O features. We observe the same trend (rapid and then asymptotic increase versus angular separation) as in the case of the full atmospheric model, although with a slightly lower S/N (120 at maximum).

- CO is well detected ($S/N = 10 \sim 30$) for the warmest planets, from 1250 K to 1750 K (but not in the hottest one at 2000 K) and as close as $0.6-0.8''$. For colder planets at 750 K and 1000 K, CO is detectable for separations larger than $0.8''$. Because the molecule's spectrum is featureless at wavelengths longer than $6 \mu m$ we present the result for the band 1A only, which globally presents the highest S/N . We found that the star itself produces a non-negligible correlation with CO , yielding some spatial residuals in the correlation map responsible for strong variations in the S/N curves.

- CH_4 is only detectable in cold planets (500 K and 750 K) in channel 2, and for separations larger than $1.4''$ as a result of fewer spectral features as compared for instance with H_2O .

- NH_3 is detected in channel 2 for planets with $T < 1000$ K, farther than $0.8''$. The detection of NH_3 will be a good tracer to discriminate between several assumptions of a planet's temperature such as 2M 1207 b (see Sect. 5.2).

- PH_3 and HCN have fewer features than the previous molecules, they are by nature more difficult to detect. According to the S/N analysis, we expect potential detection for the coldest objects ($T < 1000$ K) at rather large separations ($> 2''$). For PH_3 , we restrict the analysis to the individual band 2B and HCN in channel 3.

The abundances of each molecule as a function of the planet's temperature are indicated in Fig. 8, justifying the detection of NH_3 , CH_4 , HCN , and PH_3 only in cold planets. For hot planets, clouds are masking the absorption lines, explaining that fewer molecular features are detected.

To provide a reference S/N , we performed the very same simulations without any star, but just a planet in the center of the FoV, indicated as a cross in Fig. 7. This confirms that PH_3 and HCN should be detected for cold planets, in the case where the planet is not contaminated by stellar speckles. It also confirms that H_2S and CO_2 , even if quite abundant, are not accessible to the MRS for this range of planet temperature and brightness. As for H_2S , most spectral features are localized between ~ 5 and $8 \mu m$, where the signature of H_2O is dominating the spectrum hence masking H_2S features. This molecule might be detectable




Fig. 5. S/N as simulated for a range of planet and star temperatures in the first 3 MRS channels (built over the three sub-bands).




Fig. 6. S/N obtained as a function of angular separation from the star for a range of planet temperatures in the first 3 MRS channels.




Fig. 7. Signal-to-noise ratio obtained at each separation from the star for a range of planet's temperature for the following molecules H_2O , CO , CH_4 , HCN , NH_3 , PH_3 (in the band or channel where the spectral features are the strongest), the cross represents simulation without any star.

in the case of very bright targets. The detection of CO_2 is limited by shallow spectral features at $15\ \mu m$, low sensitivity of the instrument at such wavelengths, and stellar contamination. Therefore, CO_2 could be detected only in very bright objects, if we can manage to strongly attenuate the star's contamination. For

instance, an optimistic simulation, with a bright system at 22 pc, in which the star/planet contrast is favorable ($R_{star} = 0.8 R_{sun}$, $R_{planet} = 1.2 R_{jup}$, separation= $1.5''$) confirms this assumption: in this case CO_2 is clearly detected. Other molecules present in the models do not show any detection. TiO has spectral features in




Fig. 8. Molecular abundance in Exo-REM models of each molecule as a function of the planetary temperature.

channel 2 for the hot planets ($T > 1750$ K), however, it is too faint to be detected. FeH, K, VO, and Na have either no spectral features or signatures too faint to be detected.

We notice that for CO, CH₄, PH₃, at some temperatures the S/N of the simulation without the star is smaller than the S/N with the star. This is explained by the non-zeros correlation of the star's spectrum with these molecular features which results in a broader correlation pattern and tends to increase the S/N . This means that the S/N calculation method isn't perfect yet and could still be improved. More efficient attenuation of the star prior to applying molecular mapping could also help to reduce these effects.

5. Molecules detection based on known systems

5.1. Choice of targets based on observational limits

To complement the parametric analysis in Sect. 4, we explored the performance with known directly imaged planets in order to estimate the relevance of future programs with JWST/MIRI. The sample was defined to fulfill observational requirements: first, the angular resolution that JWST can achieve in the mid-IR imposes angular separations larger than $\sim 0.3''$ (which is about the angular resolution for the mean MRS wavelength), and second, the sensitivity of the MRS allows us to observe targets with flux larger than $30 \mu\text{Jy}$ (10σ signal in 10000 sec, Glasse et al. 2015). Therefore, we considered the following systems: GJ 504, HR 8799, β Pic, HD 95086, HIP 65426, 51 Eri HD 106906, 2M 1207 and the brown dwarf companion GJ 758, the characteristics of which are provided in Tab. 3 for the stars, and Tab. 4 for the planets. These systems cover a broad range of temperatures, angular separations, and stellar types (See Fig. 9), hence are meaningful to test the ability to characterize atmospheric parameters in the mid-IR, as compared with previous analysis in the near IR. Furthermore, all of these planets will be observed in the GTO programs with coronagraphs, either with MIRI or NIRCam.

Stellar spectra are defined with the parameters from Tab. 3 and normalized to the mean flux density values at $5.03 \mu\text{m}$ ⁵ as measured in the M band of the Johnson photometric band and tabulated at the SIMBAD astronomical database (Wenger et al. 2000). Since we are studying young systems, we can expect unresolved inner dust rings to contribute to the mid-IR flux,

⁵ we choose the shortest MRS wavelengths to be representative of the actual saturation level of the targets in the sample.




Fig. 9. Temperature of the planet as a function of its projected separation with its host star, for all of the planets from the sample. Stellar temperature is color-coded.

which needs to be taken into account in the global stellar flux. Planetary spectra are generated with the Exo-REM model using a set of temperature, $\log g$, $[M/H]$, C/O ratio, as listed in Tab. 4, and their flux density is scaled to the distance of the system and the planet's radius. For some systems, the flux level is adapted so that the models globally match the near IR data. These data are then converted to MIRISim input requirements: μJy for the flux density, and μm for the wavelengths. We used whereistheplanet.com (Wang et al. 2021) to infer the astrometry of the planet for an arbitrary date of June 2023 (likely the start of JWST cycle 2). For long-period planets, their projected positions do not vary significantly with the date (except for β Pictoris b).

The position and the spectrum of each object are used in the Exposure Time Calculator (ETC) to calculate the observational parameters (N_{group} , N_{int}). The number of groups per integration N_{group} was determined to avoid saturation while maximizing counts. Then, we chose the number of integrations to reach a S/N larger than 3 (and ideally above 5) on the detector for the planet's flux in each spectral band for the complete observation. The S/N is extracted on an aperture of $0.4''$ centered on the planet. These parameters are indicated for each simulation in Tab 5. The ETC is also convenient to check, based on the astrometry, that the planet is contained within the FoV. When needed, we adapted the telescope pointing, to position either the planet at a suitable location on the detector (especially if the angular separation is of the order of the size of the FoV) or to move away from a bright star which may cause saturation and latency. The S/N of the detection for each molecule of each system is summed up in Tab. 6.

5.2. Simulations and molecular mapping analysis of this planet sample

GJ 504 b is a T8-T9.5 object, discovered by (Kuzuhara et al. 2013). Bonnefoy et al. (2018) has analyzed the system in detail, aiming at constraining atmospheric parameters with near IR data (from 1 to $2.5 \mu\text{m}$). The uncertainty on the age (21 Myr to 4 Gyr) of this system gives two mass regimes ($1 M_{\text{jup}}$ or $23 M_{\text{jup}}$), making this object either a young exoplanet or an older brown dwarf. More measurements on the molecular abundances

Parameters	GJ 504	HR 8799	β Pictoris	HD 95086	HIP 65426	51 Eri	HD 106906	2M 1207	GJ 758
Spectral type	G0V	A5V	A6V	A8III	A2V	F0IV	F5V	M8	G9V
Temperature (K)	6200	7600	8000	7600	8800	7000	6700	2600	5500
R (R_{\odot})	1.35	1.5	1.8	1.6	1.77	1.5	1.4	0.25	0.88
Distance (pc)	17.56	39.4	19.45	86.2	111.4	29.4	102.8	52.4	15.5

Table 3. Stars parameters chosen for simulation, all have $[M/H] = 0$ and $\log g = 4.0$.

References for stellar distances [van Leeuwen \(2007\)](#) and [Gaia Collaboration et al. \(2016\)](#)

Planet's parameters	GJ 504 b ¹	HR 8799 b ²	HR 8799 cde ²	β Pictoris b ³	HD 95086 b ⁴	HD 95086 b ⁴
T(K)	550	1000	1100	1700	800	1400
logg	4.0	3.5	4.0	4.0	4.0	4.0
Separation (au)	43	68	43 - 27 - 16	9	62	62
Angular separation (arcsec)	2.5	1.72	0.94 - 0.7 - 0.38	0.55	0.63	0.63
Molecules						
H ₂ O	-3.27	-3.62	-3.62	-3.62	-3.53	-3.62
CO	-3.68	-3.3	-3.3	-3.3	-3.35	-3.3
CO ₂	-6.95	-6.96	-7.0	-7.01	-6.91	-7.0
CH ₄	-3.53	-5.76	-5.76	-7.52	-4.23	-7.17
HCN	-6.92	-7.3	-7.3	-7.78	-6.79	-7.67
NH ₃	-5.17	-6.27	-6.27	-6.94	-5.61	-6.75
H ₂ S	-4.66	-4.66	-4.66	-4.63	-4.66	-4.64
PH ₃	-6.34	-6.34	-6.34	-7.50	-6.34	-6.76

Planet's parameters	HIP 65426 b ⁵	51 Eri b ⁶	HD 106906 ⁷	2M 1207 ⁸	2M 1207 ⁹	GJ 758 ¹⁰
T(K)	1500	700	1800	1000	1600	600
logg	4.0	4.0	4.0	4.0	4.0	3.5
Separation (au)	110	11	850	125	125	226
Angular separation (arcsec)	0.81	0.34	7.11	0.78	0.78	1.36
Molecules						
H ₂ O	-3.62	-3.44	-3.62	-3.62	-3.62	-3.32
CO	-3.3	-3.42	-3.3	-3.3	-3.3	-3.58
CO ₂	-6.96	-6.89	-7.06	-6.98	-7.02	-6.91
CH ₄	-7.06	-3.9	-7.83	-5.30	-7.61	-3.61
HCN	-7.6	-6.78	-7.88	-7.06	-7.81	-6.84
NH ₃	-6.71	-5.46	-7.05	-5.97	-6.93	-5.22
H ₂ S	-4.65	-4.66	-4.63	-4.66	-4.63	-4.66
PH ₃	-6.47	-6.34	-7.83	-6.34	-7.36	-6.34

Table 4. Parameters of the simulated planets. All of them are simulated with $[M/H] = 1$ and $C/O = 0.5$. The volume mixing ratio (in log scale) of molecules present in the Exo-REM models are given, for each of the simulated planets, at the top of the atmosphere ($\log X_{mol}$ at 10^{-2} bar). Other molecules used in the model have a volume mixing ratio less than 10^{-20} .

References : ¹[Bonnefoy et al. \(2018\)](#), ²[Konopacky et al. \(2013\)](#), ³[Bonnefoy et al. \(2013\)](#), ⁴[Desgrange et al. \(2022\)](#), ⁵[Petrus et al. \(2021\)](#) and [Chauvin et al. \(2017\)](#), ⁶[Samland et al. \(2017\)](#), ⁷[Daemgen et al. \(2017\)](#), ⁸[Barman et al. \(2011\)](#), ⁹[Patience et al. \(2010\)](#), ¹⁰[Vigan et al. \(2016\)](#)

and metallicity are needed to put more robust constraints on the planet and thus on its formation. Methane has been detected in the atmosphere of this planet by [Janson et al. \(2013\)](#), but no other molecular feature has been detected yet.

The system is simulated by offsetting the star outside of the FoV at coordinates (2.0, -2.5)". As it is a nearby system, the star is too bright for the MRS and the detector would saturate in a few groups. Offsetting the star allows longer integrations. Processing the simulated data with the molecular mapping method, we obtained the correlation maps shown in Fig. 10, which displays the S/N for each detection. We were able to detect: H₂O, CO, CH₄, NH₃, HCN, and PH₃. Moreover, the correlation with the full model allows for the detection of wavelengths up to 18 μ m (Appendix Fig. C.11).

As a test case, we ran the simulation without the star to assess the impact of the stellar contribution. We found an improvement of the S/N by a factor of 3, respectively 2, for NH₃ and HCN, respectively, CH₄. Other molecules are also easier to detect,

and in particular, a $S/N = 6.8$ is achieved for CO₂. These results argue for better stellar removal to improve the detections and possibly detect CO₂ in this system and other similar systems.

HR 8799 system harbors 4 young giant planets with similar characteristics in terms of temperature and luminosity. They have been discovered by [Marois et al. \(2008, 2010\)](#). The presence of a planetesimal belt has been inferred from sub-millimetric observations with ALMA ([Booth et al. 2016](#)). Water and carbon monoxide have been clearly identified at high S/N in HR 8799 b, c, and d ([Petit dit de la Roche et al. 2018](#); [Barman et al. 2015](#); [Ruffio et al. 2021](#)). However, the presence of methane is still debated. In the case of HR 8799 b, it was claimed by [Barman et al. \(2015\)](#) using cross-correlation with a model spectrum on Osiris Keck data, but not confirmed using molecular mapping on the same data ([Petit dit de la Roche et al. 2018](#)) neither from complementary data ([Ruffio et al. 2021](#)). Broadband photometry of planets b, c, and d has




Fig. 10. Example of correlation maps for the simulated system GJ 504 with Exo-REM full atmospheric template and molecular template. Each molecule is shown in the channel/ band where the S/N is the highest.




Fig. 11. Exo-REM molecular template spectra of GJ 504 b. The parts of the spectra corresponding to the channels/ bands shown in Fig. 10 are highlighted.

provided evidence of significant atmospheric cloud coverage, while spectroscopy of planets b and c shows evidence for non-equilibrium CO/CH₄ chemistry (Janson et al. 2010; Hinz et al. 2010).

In the simulation, the star is offset to the coordinates (-0.4, 0.0)'' from the center of the FoV to be sure that all four planets are contained in a single dither position. The planet e cannot be detected, as it is too close to the star to be resolved with the MRS. The correlation maps of these simulated data are displayed in Fig. C.1. We detect H₂O and CO (for planets b, c,

and d). There is also a faint detection of NH₃ and CH₄ for planet b. No other molecule is detected.

β Pictoris system has two discovered planets Lagrange et al. (2009, 2010, 2019); Nowak et al. (2020a). Planet c is too close to the star and cannot be resolved with the MRS, therefore we focus on planet b. This planet has a dusty atmosphere (Bonnefoy et al. 2013). Previously, water and carbon monoxide have been detected in its atmosphere with SINFONI data using molecular mapping (Hoeijmakers et al. 2018).

Using the observation parameters of the GTO 1294 (PI: C. Chen) we did not manage to detect the planet, therefore we chose to increase the number of integrations. From the angular separation and temperature of the planet, we do not expect a strong detection. However, the brightness of the target still allows us to detect the planet with the full atmospheric model, while H₂O is the only detected molecule. The correlation maps corresponding to this system are presented in Fig. C.2.

HD 95086 b's detection has been presented in Rameau et al. (2013), they showed that it has a cool and dusty atmosphere, where the effects of possible non-equilibrium chemistry, reduced surface gravity, and methane bands in the near-infrared might be explored in the future. Chauvin et al. (2018) found that its near-infrared spectral energy distribution is well-fitted by spectral models of dusty and/or young L7-L9 dwarfs. Here, we aim at testing the two scenarios highlighted in Desgrange et al. (2022) using SPHERE observations combined with archival observations from VLT/NaCo and Gemini/GPI. These scenarios indicate that the color of the planet can be explained by the presence of a circumplanetary disk around planet b, with a range of high-temperature solutions (1400–1600 K) and significant extinction, or by a super-solar metallicity atmosphere but lower temperatures (800–1300 K), and small to medium amount of extinction.

We performed two simulations, one with a planet at the temperature of 800 K and a second with a planet of 1400 K. With the full spectrum, the planet is only detected in the coldest scenario.

In terms of molecules, we secured the detection of H_2O together with a suspicion of NH_3 . These results are presented in Fig. C.3 for the cold planet scenario and in Fig. C.4 for the warm planet scenario. The planet is close to its star (one of the closest in our sample) and this system is located at a large distance, which explains the globally faint detection of the planet and non-detection of other molecules.

HIP 65426 b has been discovered by Chauvin et al. (2017) with VLT/SPHERE, it is a young giant exoplanet. The Y- to H- band photometry and low-resolution spectrum indicate a $L6 \pm 1$ spectral type and a warm, dusty atmosphere. Petrus et al. (2021) studied this system with different methods including molecular mapping using VLT/SINFONI data and detected carbon monoxide and water vapor.

The planet is detected in channel 1, and only the molecule H_2O is detected. At this temperature, the detection of CO was expected from our parametric study. However, in the case of this system, which is almost 4 times more distant than the systems simulated in the parametric studies, the fainter flux of the planet explains why we do not have better detection of the planet and no detection of CO.

51 Eri b has been discovered with GPI in the near IR, Macintosh et al. (2015) indicates that it has a T4.5-T6 spectral type and the J-band spectroscopy shows methane absorption. From VLT/SPHERE data in the near IR, Samland et al. (2017) derived the presence of a vertically extended, optically thick cloud cover with small particles.

51 Eri is a bright star, the number of groups is small to avoid saturation so that the integration time is larger than for other sources to achieve the S/N required to detect the planet. The correlation maps are shown in Fig. C.6. Even though the planet is bright, it is not detected as it is too close to the star to be observable with this method, as expected from our parametric study in Sect. 4.

HD 106906 b is a young low-mass companion near the deuterium burning limit (Daemgen et al. 2017). It has been characterized spectrally in the near IR with VLT/SINFONI. This planet is the hottest and the most distant from its star in our sample.

Therefore, to simulate this system, we chose to have the planet in the center of the FoV and the star's PSF located outside the FoV. In principle, molecular mapping is not required to detect the planet as it is far from its host star and sufficiently bright. Applying molecular mapping, the planet is detected in all 3 channels, we can detect H_2O and CO. At high temperatures, we do not expect other molecules to be detected, as we can see in Fig. C.7.

2M 1207 b is the first planet ever detected with direct imaging by Chauvin et al. (2004) and will be one of the first exoplanets targeted with the MRS (GTO 1270, PI : Stephan Birkmann). The atmospheric properties of 2M 1207 b are not well constrained, and the MRS observations have the ability to break the degeneracy between two radically different models. On the one hand, Barman et al. (2011) proposed a temperature of 1000 K, $\log(g) = 4$ and $1.5 R_{Jup}$ which is in agreement with the first estimate of Chauvin et al. (2004), and on the other hand, Patience et al. (2010) found a best fit model at about 1600 K and $\log(g) = 4.5$ with a smaller radius of $0.5 R_{Jup}$.

The correlation maps are shown in the appendix Fig. C.8 and Fig C.9. Considering the model at 1000 K, we obtained a detection

System	N_{group}	N_{int}	Exposure time (s)
GJ 504	52	9	5649.98
HR 8799	21	44	10733.85
β Pictoris	5	100	6649.00
HD 95086	76	20	7083.15
HIP 65426	79	10	8869.0
51 Eri	10	200	24409.25
HD 106906	100	4	473.36
2M 1207	76	1	843.61
GJ 758	59	5	3374.45

Table 5. Simulation parameters for each target. Exposure times are indicated for one observation (three observations are required to obtain the full wavelength range). The settings of the simulation for 2M 1207 are the ones that will be used for the GTO program.

of H_2O , CO, and NH_3 . The planet is detected at a very high S/N with the full model. For the model at 1600 K, the detection of H_2O is much weaker and CO is undetected simply because the planet is smaller than in the former case. NH_3 is also undetected, as expected for such a high temperature based on the parametric study. The star being an M8 brown dwarf, the warmer planet scenario represents an extreme case for the molecular mapping method, but based on our simulations the MRS has the ability to provide a definitive answer about the planet's temperature. The detection of NH_3 can be a good indicator of the temperature of the planet.

GJ 758 B is a brown dwarf companion to a solar-type star. It has been discovered with Subaru/HiCIAO (Thalmann et al. 2009) and characterized with VLT/SPHERE in Vigan et al. (2016). No atmospheric model perfectly reproduces the measured fluxes of GJ 758 B in the near IR. As one of the coldest companions that have been directly imaged, it also appears to be an interesting target to apply molecular mapping.

The star is bright, therefore we offset it outside the FoV at coordinates (2.3, 1.3)'', and we used $N_{group} = 59$, and $N_{int} = 5$ for a total exposure time of 3374.45 s for one observation. The dithering pattern is modified from the default pattern to be optimized for the system. Correlation with the full spectrum yields a detection in the three channels, while both H_2O and NH_3 are clearly detected, and CH_4 can be suspected. These results are displayed in Fig. C.10.

6. Atmospheric characterization using grids of Exo-REM models for GJ 504 b

The former section suggests that GJ 504 b is the main interesting target of our sample for molecular mapping with the MRS. Here, we explore the potential of characterization on this specific system using two methods, one using the correlation maps and the other one with χ^2 minimization.

6.1. Correlation maps with grid of models

After subtracting the low frequencies on data and models (same method as in Sect. 3.2), the data are correlated with a grid of Exo-REM models, varying the temperature, the metallicity, the surface gravity, and the C/O ratio of the models. Models are high-pass filtered in the same way as the data. For each model, we calculate the correlation map with the correlation coefficients using Eq. 3, where $\sigma_S(\lambda)$ is the uncertainty on the flux at each wavelength, as extracted from the ERR extension of the cubes

Planets	Full model Channel 1	H ₂ O Channel 1	CO Band 1A	NH ₃ Channel 2	CH ₄ Channel 2	HCN Channel 3	PH ₃ Band 2B
GJ 504 b	72.9	59.0	9.1	27.4	15.8	7.9	7.1
HR 8799 b	66.0	64.6	10.4	7.7	4.1	—	—
HR 8799 c	34.8	34.4	9.5	—	—	—	—
HR 8799 d	34.7	36.6	7.5	—	—	—	—
β Pictoris b	6.6	7.2	—	—	—	—	—
HD 95086 b (800 K)	5.9	6.8	—	—	3.6	—	—
HD 95086 b (1400 K)	—	3.5	—	—	—	—	—
HIP 65426 b	13.6	13.7	—	—	—	—	—
51 Eri b	—	—	—	—	—	—	—
HD 106906 b	43.5	39.0	18.2	—	—	—	—
2M 1207 b (1000 K)	57.4	52.9	9.7	10.8	—	—	—
2M 1207 b (1600 K)	7.9	6.7	—	—	—	—	—
GJ 758 B	17.5	17.0	—	9.7	3.5	—	—

Table 6. S/N detection measured in correlation maps with the full atmospheric model and molecules templates for each planet of the sample. Only $S/N > 3$ are indicated.

(see Sect. 2.3). This value takes into account the photon noise and detector noise in each spaxel.

$$C = \frac{\sum_{\lambda} S(\lambda) \times M(\lambda) / \sigma_S(\lambda)^2}{\sqrt{\sum_{\lambda} S(\lambda)^2 / \sigma_S(\lambda)^2 \times \sum_{\lambda} M(\lambda)^2 / \sigma_S(\lambda)^2}} \quad (3)$$

Fig. 12 shows the grids of correlation with the correlation value at the position of the planet, that we obtained when exploring two parameters at once. In practice, for each coordinate in any of the grids, corresponding to a couple of parameter values, we took the maximum correlation coefficient obtained when varying the two other parameters. The real parameters of the planet in input to the simulation are indicated with black crosses.

In general, a significant range of models is producing high correlation values, resulting in a broad peak around the input parameter values, which implies a relatively low accuracy on the retrieved parameters. Still, the C/O vs. temperature correlation grid matches reasonably well the input parameters (bottom subplot in Fig. 12). On the contrary, we observed a tendency for higher metallicity and higher surface gravity in the two upper subplots in Fig. 12. This apparent mismatch will be discussed in the next section.

6.2. χ^2 minimization with grid of models

The former method based on the correlation with models is not sufficient to evaluate the reliability of the best model (maximum of correlation) with respect to the data. In addition, the noise estimation does not take into account the spatial noise induced by speckles. We now investigated the characterization capabilities using χ^2 minimization. Starting from the high-frequency spectrum extracted at the planet's position, we compared it to the same grid of Exo-REM models high-pass filtered. We use the correlation map to define where the planet is located, and then extract a high-frequency spectrum in the cubes after the high-pass spectral filtering on the spaxels. We extracted that spectrum for the planet's signal by co-adding the flux in the spaxels that are defined by the S/N analysis (section 3.3).

We note that this high-frequency spectrum does not contain any information on the total flux of the planet, due to the subtraction of low frequencies. We introduced a factor R , chosen to minimize the χ^2 for a given model, as given in Eq. 5. It corresponds to a global scaling parameter that does not influence the shape of the synthetic spectra, but allows us to take into account




Fig. 12. Grids of the correlation coefficient between the spectrum at the position of the planet and the Exo-REM models. Case of GJ 504 in channel 1. Similar results are observed on the other bands or reconstructed channels.

the planet's radius (as done in Baudino et al. 2015), and also captures some photometric calibration issues between the MIRISim simulated model and the actual model.

For each model, we determined the χ^2 using the equations 5 and 4, in which $\sigma_F(\lambda)$ is the uncertainty on the flux measured at each wavelength in an annulus at the same planet's separation from the star in the high frequencies cubes. The noise σ_S extracted from the JWST pipeline is now negligible, and this noise σ_F takes into account the spatial variation in the high-frequency cubes. Similarly to Fig. 12, we display the $\chi^2 - \chi^2_{min}$ values of the grid of models in Fig. 13.

$$\chi^2 = \sum_{\lambda} \left(\frac{S(\lambda) - R \times M(\lambda)}{\sigma_F(\lambda)} \right)^2 \quad (4)$$

$$R = \frac{\sum_{\lambda} (S(\lambda) \times M(\lambda)) / \sigma_F(\lambda)^2}{\sum_{\lambda} M(\lambda)^2 / \sigma_F(\lambda)^2} \quad (5)$$

As a sanity check, we obtained χ^2 values that are in agreement with the number of independent points in the spectrum. The χ^2

minimization gives similar results compared to the grid of correlation values. Again, several models yield χ^2 values that are close to the one of the input model, although the 1σ , 2σ , and 3σ contours point to a more restrained region in the C/O vs. temperature grid, when compared to Fig. 13. However, this is not the case for the metallicity and the surface gravity that are not well constrained, likely due to degeneracies between these atmospheric parameters. Indeed, the metallicity and the surface gravity information are mostly contained in the relative depth of the lines, which are affected by the filtering stage. Hence, it is more difficult to disentangle between two models that have different metallicity or surface gravity values. On the contrary, the abundance of molecules, such as H_2O , CO , and CH_4 that define the C/O ratio, depends on the temperature of the planet. Even though the continuum of the planetary spectrum is lost in the filtering, they still have a net effect in the high-frequency spectrum of the planet, explaining the rather good match obtained for the temperature and the C/O ratio. In the event that one of these parameters is well determined by other methods or other observations, the remaining parameters can be well constrained with high confidence as illustrated in Fig. 14 in which the surface gravity is fixed at the input values ($\log g = 4.0$) in each subplot.




Fig. 13. Values $\chi^2 - \chi^2_{\min}$ using the high-frequency spectrum and a grid of Exo-REM models. Green, blue and black lines are respectively 1σ , 2σ , and 3σ confidence regions (2.3, 6.18, and 11.8).

7. Discussion

7.1. Pros and cons of the mid IR for the molecular mapping

The Mid-IR spectral range offers advantages when combined with molecular mapping with respect to the near IR. For instance a planet at 700 K has its emission peak at $5.2 \mu\text{m}$ while the stellar contribution is getting less dominant. According to the parametric study (systems at 30 pc, planets of $1 R_{\text{jup}}$ around a star at 6000 K, 2 hours per observation), the MRS should allow us to detect planets at separations larger than $0.5''$, (except the hottest planet around the coldest star in our sample), as well as to allow the characterization of those that are further away than $1''$ from their star. We found that planets with temperatures above 500 K and below 1500 K will be easier to detect especially if they orbit G-type stars or younger. In addition, the mid-IR gives access to molecules that are not easily accessible at shorter wavelengths because of fainter absorption features (such as NH_3 , HCN).




Fig. 14. Values $\chi^2 - \chi^2_{\min}$ using the high-frequency spectrum and a grid of Exo-REM models. The log of the surface gravity is fixed at 4.0 to correspond to the input value in each subplot. Green, blue and black lines are respectively 1σ , 2σ , and 3σ confidence regions (2.3, 6.18, and 11.8).

On the contrary, the angular resolution is worse than in the near-IR, so stellar contamination becomes a more important issue. For this reason, we have chosen GJ 504 b to perform a detailed characterization study as it features favorable configurations in terms of angular separations and temperature. The comparison to a grid of models with a χ^2 approach provides reasonable results (section 6) but for other targets which are too close to their star, the confidence regions in the χ^2 maps are too large to reduce the accuracy on the atmospheric parameters. Only planets with S/N larger than 30 in our sample (HR 8799 planets, 2M 1207 b at 1000 K) are amenable to atmospheric characterization with the MRS. We note that the extracted spectra of the others planets (with $S/N < 30$) are too contaminated by stellar residuals. Although well detached, HD 106906 b is a special case because Exo-REM is not well adapted to model a planet at this temperature, therefore other atmospheric models would be needed to characterize this planet.

Moreover, having priors on the atmospheric parameters to correlate our data with models is necessary to reduce the explored parameter range. As a first approximation, one can use the high-frequency spectrum before applying the molecular mapping, to restrain this range using the χ^2 minimization. We will benefit from the use of both analyses: molecular mapping and χ^2 minimization, other methods of atmospheric retrievals would be relevant to further study of these planets.

Finally, Molecular mapping relies on the fact that the spectrum has lots of spectral features due to the molecular absorption and might be less efficient for young dusty planets since dust and clouds tend to flatten the spectrum, as noted for the PDS 70 planets in Cugno et al. (2021), (extinction due to dust has not been taken into account in the present work). Further works should be continued to test the impact of clouds and extinction on molecular mapping.

7.2. Possible improvements in the data reduction and molecular mapping

As a source of problems for the characterization, stellar subtraction should be tackled with more efficient methods, especially for planets located at separations below $1''$ which are more affected by the starlight. Subtracting a scaled template from each

spectrum in the data cube as did [Hoeijmakers et al. \(2018\)](#) gives a slightly higher S/N for the detection of the planets closer than $1''$ (such as β Pictoris b) but shows no improvement for more distant targets. As we noticed from the ideal simulations with no star in the scene and providing that the photon noise and the speckle noise are below the background noise, a significant improvement of the S/N is possible for the most impacted molecules (NH_3 , CH_4 , HCN , CO_2), which argues in favor of developing more efficient PSF subtraction algorithms. Consequently, more targets would be accessible for characterization with the MRS and molecular mapping. Other interesting targets might come from future ground-based surveys, such as The Young Suns Exoplanet Survey ([Bohn et al. 2020](#)), and SHINE with VLT/SPHERE ([Vigan et al. 2021](#)).

A second avenue for improving the performance of molecular mapping with MIRI is to take advantage of the full spectral range. Hitherto, because of current limitations in the pipeline, we have restrained the method to the MRS channels, but exploiting the full spectral range of MIRI should increase the detection of molecules that exhibit features at multiple wavelengths in different channels (such as NH_3). Two solutions are envisioned, either a full concatenation of the data cubes from each channel or the extraction of the planet spectrum channel per channel, prior to cross-correlation with the model. The first method is limited by interpolation artifacts while the second one requires the planet to be detected in each of the three channels.

7.3. Interpretation of the detection of the molecules in the targets sample

All planets in our sample, except 51 Eri b for which it is too challenging to disentangle the planet from the star's signal, are detected using the correlation with a full atmospheric model. Depending on the distance of the system, long integrations (more groups) are possible, effectively improving the ramp fitting. However, the planets will be fainter if distant, thus requiring longer exposure time (more than 4 hours per observation for HD 95086). In contrast, bright stars like β Pic and 51 Eri impose fewer groups per integration, which has a definitive impact on the planet detection, while at the same time the angular separation is smaller than $0.5''$, so that the star cannot be moved out of the FoV.

The planet GJ 504 b, although being the target with the largest contrast with its star, is the target with the largest S/N . On the contrary, 2M 1207 b is the one with the lowest contrast (in both temperature hypotheses) but its detection is much weaker. S/N is 8 times larger for GJ 504 b than for 2M 1207 b if the planet is at 1600 K.

We note that H_2O , the most abundant molecule in the atmosphere of giant planets and also the one with the most numerous spectral features, is always detected, even in planets that are close to their star and thus heavily contaminated by the starlight (β Pic b, HD 95086 b, HIP 65426 b). It is the only molecule unambiguously detected in these systems. This result is consistent with the parametric study. It is even detected with a slightly higher S/N than the correlation with the full atmospheric model, which can be explained by the fact that H_2O does not correlate with stellar residuals contrary to the full atmospheric model. CO is the second most abundant molecule in most planets of the sample. However, it is detected only for planets with separations larger than $0.8''$. Indeed, as confirmed by the parametric study, it is the molecule that suffers the most from stellar contamination, which also explains non-detection when the planet has a

low flux (such as HIP 65426, although it is at $0.8''$ and 2M 1207 b at 1600 K small radius hypothesis).

On the contrary, PH_3 and HCN are among the less abundant molecules, but their spectral features are such that they can be detected in cold planets, providing they are bright (nearby and well-separated planet like the case of GJ 504 b), in agreement with the parametric study.

The access to these molecules is a unique feature of JWST observations as it allows us to derive new constraints on planetary atmospheres as well as on planetary formation. In complement to NIRSpec data that can access the bright features of PH_3 around $4.3 \mu\text{m}$, we show that MIRI/MRS can also detect or confirm the presence of this molecule. [Mukherjee et al. \(2022\)](#) showed the importance of PH_3 in the emission spectrum of an atmosphere in thermochemical disequilibrium (based on MIRI-LRS simulations). Furthermore, measuring the phosphine abundance, which is the dominant phosphorus molecule, can provide an estimation of the P/O ratio. These abundances ratios are of interest to determine if one element is depleted compared to oxygen which provides valuable information on the planetary formation processes. In addition, [Zahnle & Marley \(2014\)](#) details the importance and implication of NH_3 , CH_4 , and HCN in the atmospheres of young giant planets and brown dwarfs. These molecules are keys to accurately deriving the C/O and N/O ratios, all being accessible to the MRS. They could also be indicators of chemical disequilibrium and provide constraints on the deep atmospheric temperatures and strength of vertical mixing that characterize the so-called "quench level".

NH_3 is also a powerful indicator of both planetary effective temperature and atmospheric chemical equilibrium state. Differences between two model spectra calculated at equilibrium and disequilibrium chemistry arise between $5\text{--}9 \mu\text{m}$. These differences mainly arise from different CH_4 , H_2O , and NH_3 abundances due to differences in the quench level ([Mukherjee et al. 2022](#)).

7.4. Influence of the atmospheric models

The molecules that are detected with molecular mapping are obviously strongly related to the atmospheric model that is used to generate the templates. This explains some differences in the case of HR 8799 and GJ 504 with ?, who used `petitRADTRANS` model. In their analysis, CO is not detected in GJ 504 b. Indeed, their molecular abundances computed with `petitRADTRANS` have significant differences with values computed with `Exo-REM` (for example lower CO abundance for a planet at 550 K). Similarly, in the case of the HR 8799 system, they obtained a higher detection of CO for planet c than for planet b, the former having a higher temperature and hence more CO according to `petitRADTRANS` as opposed to `Exo-REM`. In addition, CH_4 and NH_3 are less abundant in `Exo-REM`-based models with respect to models from `petitRADTRANS`, which explains why the detection of NH_3 and no CH_4 is lower in our analysis of the HR 8799 planets. These differences in the molecular compositions mostly come from a different treatment of the chemistry. In `Exo-REM`, chemical abundances are computed assuming chemical disequilibrium with quenching levels (generally between 1 and 10 bars) derived from a parametrization of the vertical mixing. In contrast, in [Patapis et al. \(2022\)](#), the quenching level is fixed at 10^{-2} bar.

Also, we can expect that a family of models that best fits the ground-based near-IR data might have large differences in the mid-IR regime. Depending on model assumptions, there is a large range of atmospheric parameter values that can repro-

duce the data. Therefore, the simulations presented in this paper should not be considered a perfect reproduction of each target, as long as we currently have little or no constraint in the mid-IR range for the known directly imaged planets.

7.5. Complementarity with MIRI coronagraphy

Although molecular mapping is powerful to identify the presence of molecules it comes with difficulties to put meaningful constraints on atmospheric parameters, as long as the planet continuum is significantly affected by the first step in which the stellar contribution is subtracted. In that respect, complementary observations would be useful, in particular, those with MIRI's coronagraphs, which are observing in the same spectral range as the MRS but with specific wavelengths and broader bandwidths. The 4QPM coronagraphs (Boccaletti et al. 2015) can provide photometric measurements of the planet's continuum flux (11.4 μm and 15.5 μm), as well as the NH_3 feature at 10.65 μm , which are certainly useful to provide priors for the MRS data analysis. The combination of observations with the MRS and the coronagraphs will be decisive in further constraining the atmospheres of known directly imaged planets as illustrated in Danielski et al. (2018).

7.6. Complementarity with future ground-based projects

MIRI/MRS has the advantage of not being impacted by telluric lines in comparison to the data taken with ground-based instruments. Still, future ELT instruments are highly complementary to the MRS in providing higher spectral resolution and different wavelength coverage.

In the near IR, HARMONI allows a range of spectral resolution settings (3000, 7000, 18000). Houllé et al. (2021) simulated molecular mapping observations combined with a matched-filter approach showing that planets with contrasts up to 16 mag and separations down to 75 mas can be detected at $> 5\sigma$.

In the mid-IR, METIS provides high-resolution spectro-imaging ($R \sim 100\,000$) at L/M band, including a mode with extended instantaneous wavelength coverage (assisted by a coronagraph). Snellen et al. (2015) showed that an Earth-like planet orbiting αCen could be detected at a S/N of 5 with an instrument like METIS. Note that the METIS M band overlaps with the shortest wavelength of the MRS.

8. Conclusions

In this paper, we showed that future JWST observations with MIRI-MRS are capable to provide clear detection of molecules in the atmospheres of young giant planets which is key in measuring abundance ratios and ultimately providing constraints on their formation and evolution. In the following, we sum up the main specific conclusions of our study.

- As already showcase in Patapis et al. (2022), we confirmed that the MIRI MRS mode has the potential to enhance planet detection owing to the molecular mapping technique which performs cross-correlation with atmospheric models.
- To determine the significance of the detection, we propose a data-driven experimental method, which takes into account the size and the shape of the correlation pattern, to avoid the impact of the autocorrelation, which arises for molecules that have harmonics in their spectra. While satisfactory for a majority of cases, the S/N may be over-estimated in

particular situations: with a CO template, or either with a hot planet ($T = 2000\text{ K}$) around a cold star.

- The correlation pattern differs from a PSF and scales with the wavelength at a high signal-to-noise ratio but also depends on the strength of the correlation, the atmospheric template, and the noise level. Importantly, the maximum of the correlation pattern does not necessarily correspond to the real position of the planet (stronger effect at small angular separations). As a result, astrometry is unreliable with correlation maps.
- From the parametric study, which samples a range of planetary temperatures and angular separations, we concluded that, while planets are detected as close as $0.5''$, a good level of characterization requires more angularly separated objects, typically larger than $1''$. The stellar spectral type has little impact on the performance, as opposed to the planetary temperature. The highest S/N values are achieved for planet temperatures ranging between 750 K and 1750 K, while lower performances are obtained for the coldest ($T_p = 500\text{ K}$), and the warmest ($T_p = 2000\text{ K}$) planets, respectively because of a lower absolute flux, and a lower level of correlation due to fewer spectral features.
- For planets typically colder than 1500 K, the following molecules are detectable: H_2O , CO, NH_3 , CH_4 , HCN, PH_3 , CO_2 . Some of these molecules have never been confirmed or even detected in the atmosphere of an exoplanet.
- We propose three directions for improving the performance of the molecular mapping method with MIRI. Firstly, exploring further the subtraction of the stellar contribution can allow the detection and characterization of planets closer than $1''$ as well as the detection of the molecules that have less spectral features or that are more hidden by other molecules. Secondly, a robust estimation of the S/N would require developing a more sophisticated analytical approximation to model the size and shape of the correlation pattern. And thirdly, we plan to perform Bayesian analysis, to better constrain the atmospheric parameters and to better evaluate the uncertainties of each parameter.
- Complementary data, such as coronagraphic mid-IR photometry with MIRI, for measuring a planet's continuum will provide temperature and surface gravity estimates, which in turn can be taken as constraints on the atmospheric parameters for characterization purposes using molecular mapping.
- The interpretation of the data processing with molecular mapping strongly depends on the assumptions of the models to generate synthetic spectra. A pilot program dedicated to a specific target will represent a benchmark for systematically comparing several models.

Acknowledgements. M.M. acknowledges the Centre National d'Études Spatiales (CNES, Toulouse, France) for supporting the PhD fellowship. B.C. and B.B. acknowledge financial support from the Programme National de Planétologie (PNP) of CNRS/INSU. This work was granted access to the HPC resources of MesoPSL financed by the Région Ile de France and the project Equip@Meso (reference ANR-10-EQPX-29-01) of the programme Investissements d'Avenir supervised by the Agence Nationale pour la Recherche. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (COBEX; grant agreement no 885593; EPIC, grant agreement no 819155). This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint

project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration. The results presented in this research paper were obtained using the python packages numpy (Harris et al. 2020), scipy (Virtanen et al. 2020), matplotlib (Hunter 2007) and astropy (Astropy Collaboration et al. 2013, 2018).

References

- Barman, T. S., Konopacky, Q. M., Macintosh, B., & Marois, C. 2015, *The Astrophysical Journal*, 804, 61
- Barman, T. S., Macintosh, B., Konopacky, Q. M., & Marois, C. 2011, *The Astrophysical Journal*, 735, L39
- Baudino, J.-L., Bézard, B., Boccaletti, A., et al. 2015, *Astronomy & Astrophysics*, 582, A83
- Birkby, J. L., de Kok, R. J., Brogi, M., et al. 2013, *Monthly Notices of the Royal Astronomical Society: Letters*, 436, L35
- Blain, D., Charnay, B., & Bézard, B. 2021, *Astronomy & Astrophysics*, 646, A15
- Boccaletti, A., Lagage, P.-O., Baudou, P., et al. 2015, *Publications of the Astronomical Society of the Pacific*, 127, 633
- Bohn, A. J., Kenworthy, M. A., Ginski, C., et al. 2020, *Monthly Notices of the Royal Astronomical Society*, 492, 431
- Bonnefoy, M., Boccaletti, A., Lagrange, A.-M., et al. 2013, *Astronomy & Astrophysics*, 555, A107
- Bonnefoy, M., Perraut, K., Lagrange, A.-M., et al. 2018, *Astronomy & Astrophysics*, 618, A63
- Booth, M., Jordán, A., Casassus, S., et al. 2016, *Monthly Notices of the Royal Astronomical Society: Letters*, 460, L10
- Brogi, M., Snellen, I. A. G., de Kok, R. J., et al. 2012, *Nature*, 486, 502
- Carter, A. L., Hinkley, S., Kammerer, J., et al. 2022, *The JWST Early Release Science Program for Direct Observations of Exoplanetary Systems I: High Contrast Imaging of the Exoplanet HIP 65426 b from 2-16 μm* , arXiv:2208.14990 [astro-ph]
- Charbonneau, D., Brown, T. M., Noyes, R. W., & Gilliland, R. L. 2002, *The Astrophysical Journal*, 568, 377
- Charnay, B., Bézard, B., Baudino, J.-L., et al. 2018, *The Astrophysical Journal*, 854, 172
- Chauvin, G., Desidera, S., Lagrange, A.-M., et al. 2017, *Astronomy & Astrophysics*, 605, L9
- Chauvin, G., Gratton, R., Bonnefoy, M., et al. 2018, *Astronomy & Astrophysics*, 617, A76
- Chauvin, G., Lagrange, A.-M., Dumas, C., et al. 2004, *Astronomy & Astrophysics*, 425, L29
- Cugno, G., Patapis, P., Stolker, T., et al. 2021, *Astronomy & Astrophysics*, 653, A12
- Daemgen, S., Todorov, K., Quanz, S. P., et al. 2017, *Astronomy & Astrophysics*, 608, A71
- Danielski, C., Baudino, J.-L., Lagage, P.-O., et al. 2018, *The Astronomical Journal*, 156, 276
- Delorme, P., Schmidt, T., Bonnefoy, M., et al. 2017, *Astronomy & Astrophysics*, 608, A79
- Deming, D., Wilkins, A., McCullough, P., et al. 2013, *The Astrophysical Journal*, 774, 95
- Desgrange, C., Chauvin, G., Christiaens, V., et al. 2022, *Astronomy & Astrophysics*, 664, A139
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2016, *Astronomy & Astrophysics*, 595, A2
- Glasse, A., Rieke, G. H., Bauwens, E., et al. 2015, *Publications of the Astronomical Society of the Pacific*, 127, 686
- Hinkley, S., Carter, A. L., Ray, S., et al. 2022, *Publications of the Astronomical Society of the Pacific*, 134, 095003
- Hinz, P. M., Rodigas, T. J., Kenworthy, M. A., et al. 2010, *The Astrophysical Journal*, 716, 417
- Hoeijmakers, H. J., Schwarz, H., Snellen, I. A. G., et al. 2018, *Astronomy & Astrophysics*, 617, A144
- Houllé, M., Vigan, A., Carlotti, A., et al. 2021, *Astronomy & Astrophysics*, 652, A67
- Janson, M., Bergfors, C., Goto, M., Brandner, W., & Lafrenière, D. 2010, *The Astrophysical Journal*, 710, L35
- Janson, M., Brandt, T. D., Kuzuhara, M., et al. 2013, *The Astrophysical Journal*, 778, L4
- Klaassen, P. D., Geers, V. C., Beard, S. M., et al. 2020, *Monthly Notices of the Royal Astronomical Society*, 500, 2813
- Konopacky, Q. M., Barman, T. S., Macintosh, B., & Marois, C. 2013, *Science*, 339, 1398
- Kuzuhara, M., Tamura, M., Kudo, T., et al. 2013, *The Astrophysical Journal*, 774, 11
- Labiano, A., Argyriou, I., Álvarez Márquez, J., et al. 2021, *Astronomy & Astrophysics*, 656, A57
- Labiano-Ortega, A., Dicken, D., Vandenbussche, B., et al. 2016, in *Observatory Operations: Strategies, Processes, and Systems VI*, ed. A. B. Peck, C. R. Benn, & R. L. Seaman (Edinburgh, United Kingdom: SPIE), 117
- Lagrange, A.-M., Boccaletti, A., Langlois, M., et al. 2019, *Astronomy & Astrophysics*, 621, L8
- Lagrange, A.-M., Bonnefoy, M., Chauvin, G., et al. 2010, *Science*, 329, 57
- Lagrange, A.-M., Gratadour, D., Chauvin, G., et al. 2009, *Astronomy & Astrophysics*, 493, L21
- Macintosh, B., Graham, J. R., Barman, T., et al. 2015, *Science*, 350, 64
- Madhusudhan, N., Amin, M. A., & Kennedy, G. M. 2014, *The Astrophysical Journal*, 794, L12
- Marois, C., Lafrenière, D., Doyon, R., Macintosh, B., & Nadeau, D. 2006, *The Astrophysical Journal*, 641, 556
- Marois, C., Macintosh, B., Barman, T., et al. 2008, *Science*, 322, 1348, arXiv:0811.2606
- Marois, C., Zuckerman, B., Konopacky, Q. M., Macintosh, B., & Barman, T. 2010, *Nature*, 468, 1080
- Miles, B. E., Biller, B. A., Patapis, P., et al. 2022, *The JWST Early Release Science Program for Direct Observations of Exoplanetary Systems II: A 1 to 20 Micron Spectrum of the Planetary-Mass Companion VHS 1256-1257 b*, arXiv:2209.00620 [astro-ph]
- Mordasini, C., van Boekel, R., Mollière, P., Henning, T., & Benneke, B. 2016, *The Astrophysical Journal*, 832, 41
- Mukherjee, S., Fortney, J. J., Batalha, N. E., et al. 2022, *The Astrophysical Journal*, 938, 107
- Nowak, M., Lacour, S., Lagrange, A.-M., et al. 2020a, *Astronomy & Astrophysics*, 642, L2
- Nowak, M., Lacour, S., Mollière, P., et al. 2020b, *Astronomy & Astrophysics*, 633, A110
- Oberg, K. I. & Bergin, E. A. 2021, *Physics Reports*, 893, 1, arXiv:2010.03529
- Patapis, P., Nasedkin, E., Cugno, G., et al. 2022, *Astronomy & Astrophysics*, 658, A72
- Patience, J., King, R. R., De Rosa, R. J., & Marois, C. 2010, *Astronomy and Astrophysics*, 517, A76
- Petit dit de la Roche, D. J. M., Hoeijmakers, H. J., & Snellen, I. A. G. 2018, *Astronomy & Astrophysics*, 616, A146
- Petrus, S., Bonnefoy, M., Chauvin, G., et al. 2021, *Astronomy & Astrophysics*, 648, A59
- Racine, R., Walker, G., Nadeau, D., Doyon, R., & Marois, C. 1999, *Publications of the Astronomical Society of the Pacific*, 111, 587
- Rameau, J., Chauvin, G., Lagrange, A.-M., et al. 2013, *The Astrophysical Journal*, 779, L26
- Ruffio, J.-B., Konopacky, Q. M., Barman, T., et al. 2021, *The Astronomical Journal*, 162, 290
- Ruffio, J.-B., Macintosh, B., Konopacky, Q. M., et al. 2019, *The Astronomical Journal*, 158, 200
- Samland, M., Mollière, P., Bonnefoy, M., et al. 2017, *Astronomy & Astrophysics*, 603, A57
- Snellen, I., de Kok, R., Birkby, J. L., et al. 2015, *Astronomy & Astrophysics*, 576, A59
- Snellen, I. A. G., de Kok, R. J., de Mooij, E. J. W., & Albrecht, S. 2010, *Nature*, 465, 1049
- Sparks, W. B. & Ford, H. C. 2002, *The Astrophysical Journal*, 578, 543
- Thalmann, C., Carson, J., Janson, M., et al. 2009, *The Astrophysical Journal*, 707, L123
- van Leeuwen, F. 2007, *Astronomy & Astrophysics*, 474, 653
- Vigan, A., Bonnefoy, M., Ginski, C., et al. 2016, *Astronomy & Astrophysics*, 587, A55
- Vigan, A., Fontanive, C., Meyer, M., et al. 2021, *Astronomy & Astrophysics*, 651, A72
- Visscher, C., Lodders, K., & Fegley, B. 2010, *The Astrophysical Journal*, 716, 1060
- Wang, J., Kulikauskas, M., & Blunt, S. 2021, *whereistheplanet: Predicting positions of directly imaged companions*, Wells, M., Pel, J.-W., Glasse, A., et al. 2015, *Publications of the Astronomical Society of the Pacific*, 127, 646
- Wenger, M., Ochsenbein, F., Egret, D., et al. 2000, *Astronomy and Astrophysics Supplement Series*, 143, 9
- Wright, G. S., Wright, D., Goodson, G. B., et al. 2015, *Publications of the Astronomical Society of the Pacific*, 127, 595
- Zahnle, K. J. & Marley, M. S. 2014, *The Astrophysical Journal*, 797, 41

Appendix A: JWST Pipeline

Stage 1. The first stage of the pipeline corrects for the detector effects. The input raw data are in the form of one or more ramps (integration) containing accumulating counts from the non-destructive detector readouts. The output is a corrected count-rate (slope) image. Corrections are applied group by group: first, the pipeline corrects the quality of the pixels, flagging those that will not be used. The step *dq_init* initializes the data quality using a reference file where known bad pixels are indicated. Saturated pixels are flagged with the step *saturation*. The first and the last groups of each integration are suppressed as they are the most affected by detector effects (*firstframe*, *lastframe*). The *linearity* step applies for each pixel a correction for the non-linear detector response, using the "classic" polynomial method where the coefficients of the polynomial are stored in a reference file. Dark current is corrected by subtracting dark current reference data from the input science data model (*dark*). On each integration ramp within an exposure, we perform cosmic rays/jumps detection (*jump*) by looking for outliers in the up-the-ramp signal in each pixel. Finally, ramp fitting determines the mean count rate, in units of counts per second, for each pixel by performing a linear fit to the data in the input file (*ramp_fitting*). This stage 1 takes 4D data in the shape of $(N_{int}, N_{group}, N_{pixel,x}, N_{pixel,y})$ to 2D images for the detector $(N_{pixel,x}, N_{pixel,y})$. Other steps of corrections are available but they will not be useful for simulated data, they will be considered for the real on-orbit data.

Stage 2. The second stage of the pipeline corresponds to the calibration. It includes additional corrections depending on the instrument and the observation mode to produce fully calibrated exposures. First, the pipeline associates a WCS object with each science exposure, which transforms the positions in the detector frame to positions using the International Celestial Reference System (ICRS) frame and wavelength (*assign_wcs*). The Source Type (*srctype*) step attempts to determine whether a spectroscopic source should be considered a point or extended object. Fringes in spectra are corrected (*fringe*). Finally, photometric calibrations allow converting count rates to surface brightness (in MJy/str) (*photom*). The outputs are 2D calibrated data of the detector.

Stage 3. The last stage is intended for combining all calibrated exposures. We can also subtract or equalize the sky background in science image (*mrs_imatch*). Outliers, bad pixels, and cosmic rays that would remain are flagged using the *outlier_detection* step. This stage takes into account the different dither positions to build a data cube with *cube_build* and extract a spatially averaged spectrum over the full field of view with *extract_1d*. We can choose to construct cubes for each band or combine bands and channels with a larger wavelength range. The outputs are 3D cubes with two extensions, the SCI image contains the surface brightness of the spaxels, and the ERR image is the uncertainty of the SCI values.

Appendix B: Size and shape of the correlation pattern in a simple case

Appendix B.1: Analytical first approximation

To have an idea of the parameters involved in the correlation that should help to understand the size of the correlation pattern, we derive a formula for the correlation depending on its parameters. We defined S as the observed spectrum and M as the model spectrum. The noise in the observed spectrum is $\xi \sim \mathcal{N}(0, \xi(\lambda))$, P is

the PSF.

$$S = PM + \xi \quad (\text{B.1})$$

The correlation can be written as :

$$C = \frac{M \otimes (PM + \xi)}{\|M\| \times \|(PM + \xi)\|} \quad (\text{B.2})$$

The numerator can be written as :

$$M \otimes (PM + \xi) = PM \otimes M + M \otimes \xi \quad (\text{B.3})$$

$$= \sum_{\lambda} P(\lambda) M^2 + \sum_{\lambda} M \xi \quad (\text{B.4})$$

$$\approx \sum_{\lambda} P(\lambda) M^2 \quad (\text{B.5})$$

In detail, the norm of the observed spectrum is:

$$\|(PM + \xi)\| = \sqrt{\sum_{\lambda} (PM)^2 + 2 \sum_{\lambda} PM \xi + \sum_{\lambda} \xi^2} \quad (\text{B.6})$$

$$\approx \sqrt{\sum_{\lambda} P(\lambda)^2 M(\lambda)^2 + \sum_{\lambda} \xi(\lambda)^2} \quad (\text{B.7})$$

Finally :

$$\|M\| = \sqrt{\sum_{\lambda} M^2} \quad (\text{B.8})$$

If the PSF function $P_x(\lambda)$ is a Gaussian $G_{\sigma}(x)$ with x the spaxel distance from the centroid and with a wavelength-dependent width $\sigma(\lambda) = \sigma_0 \lambda / \lambda_0$, then we may develop P at first order in $(\lambda - \lambda_0) / \lambda_0$ as:

$$P = G_{\sigma}(x) \quad (\text{B.9})$$

$$= \exp\left(-\frac{\mathbf{x}^2}{2\sigma_0^2 \lambda^2 / \lambda_0^2}\right) \quad (\text{B.10})$$

$$\approx \exp\left(-\frac{\mathbf{x}^2}{2\sigma_0^2}\right) \left(1 + \frac{x^2}{\sigma_0^2} \frac{\lambda - \lambda_0}{\lambda_0}\right) \quad (\text{B.11})$$

$$(\text{B.12})$$

In the following we will write $G_0(x) = \exp\left(-\frac{\mathbf{x}^2}{2\sigma_0^2}\right)$.

The different terms in the numerator and in the denominator involving $P_x(\lambda)$ can be developed according to the above equation at first order in $(\lambda - \lambda_0) / \lambda_0$. Linearising all first order terms leads to the following equation:

$$C \approx \frac{G_0(x)}{\sqrt{G_0(x)^2 + \delta^2}} + \frac{x^2}{\sigma_0^2} G_0(x) \frac{\delta^2}{(G_0(x)^2 + \delta^2)^{3/2}} \beta \quad (\text{B.13})$$

where we defined:

$$\beta = \frac{\sum_{\lambda} \frac{\lambda - \lambda_0}{\lambda_0} M(\lambda)^2}{\sum_{\lambda} M(\lambda)^2} \quad (\text{B.14})$$

$$\delta^2 = \frac{\sum_{\lambda} \xi(\lambda)^2}{\sum_{\lambda} M(\lambda)^2} \quad (\text{B.15})$$

$$(\text{B.16})$$

The parameter δ^2 can be considered as a positive constant that depends on the signal-to-noise level in the cube, while β can be bounded by the wavelength range of the considered channel. Indeed, since $\lambda_{\min} < \lambda < \lambda_{\max}$:

$$\frac{\lambda_{\min} - \lambda_0}{\lambda_0} < \beta < \frac{\lambda_{\max} - \lambda_0}{\lambda_0} \quad (\text{B.17})$$

Thus since the channels windows are on the order of $\lambda_0 \pm 14 - 18\%$, we expect $\beta < 0.18$. The zeroth-order term $C = G_0(x) / \sqrt{G_0(x)^2 + \delta^2}$ is thus a good approximation of the correlation pattern. It indicates that the correlation pattern will depend on the PSF, but also on the model spectrum and on the noise. However, this formula is insufficient to estimate a correlation radius, and it would result in an oversized correlation pattern. Indeed this is correct in the case of data dominated by Gaussian noise, which is not the case at long wavelengths and when we have more stellar residuals. It also assumes that the PSF is a Gaussian.

Appendix B.2: Example of simulations from the parametric study

Simulations show that we obtain different shapes and sizes of the correlation pattern depending on the noise level and the molecules studied. Fig. B.1 presents correlation maps for a planet at 500 K, it is a comparison between the direct image, and the correlation maps for three detected molecules (NH₃, CH₄, and PH₃) at the same wavelength (band 2B) and same star-planet separation. Fig. B.3 and Fig. B.4 is a comparison between CO and H₂O for four values of separation between a star and a planet. It corresponds to the simulation with a planet at 1750 K, in band 1A. Moreover, it is notable that astrometry is unreliable, mostly for short star-planet separation. The asymmetry of the correlation pattern is clearly visible in Fig. B.3. Concerning the mean azimuthal profile, we note that the profile of the PSF is unchanged whatever the noise level, whereas the profile of the correlation pattern depends on the noise level and on the template with which we correlate the data.




Fig. B.1. Simulations from the parametric study with the planet at 500 K and the star separated at 3.2". Correlations maps with different molecular templates: NH₃, CH₄, and PH₃ in the band 2B.




Fig. B.2. Corresponding mean azimuthal profiles for each correlation maps above Fig. B.1.




Fig. B.3. Simulations from the parametric study with the planet at 1750 K and the star at different separations. Correlation maps with CO in band 1A.




Fig. B.4. Simulations from the parametric study with the planet at 1750 K and the star at different separations. Correlation maps with H₂O in band 1A.




Fig. B.5. Corresponding radial profiles for each of the correlation maps of Fig. B.3 and Fig. B.4.

Appendix C: Simulations and correlation maps for each planet




Fig. C.1. MIRISim simulation and correlation maps of the simulated system HR 8799.




Fig. C.2. MIRISim simulation and correlation maps of the simulated system β Pictoris.




Fig. C.3. MIRISim simulation and correlation maps of the simulated system HD 95086 at 800 K.




Fig. C.4. MIRISim simulation and correlation maps of the simulated system HD 95086 at 1400 K.




Fig. C.5. MIRISim simulation and correlation maps of the simulated system HIP 65426.




Fig. C.6. MIRISim simulation and correlation maps of the simulated system 51 Eri.




Fig. C.7. MIRISim simulation and correlation maps of the simulated system HD 106906, the simulation plot shows directly the planet (in the center).




Fig. C.8. MIRISim simulation and correlation maps of the simulated system 2M 1207 at 1000 K.




Fig. C.9. MIRISim simulation and correlation maps of the simulated system 2M 1207 at 1600 K.




Fig. C.10. MIRISim simulation and correlation maps of the simulated system GJ 758.




Fig. C.11. MIRISim simulation and correlation maps of the simulated system GJ 504 for the 3 first channel. The scale is the same in each bands and channels.