

THE HOMOLOGY OF THE PARTITION ALGEBRAS

RACHAEL BOYD, RICHARD HEPWORTH, AND PETER PATZT

ABSTRACT. We show that the homology of the partition algebras, interpreted as appropriate Tor-groups, is isomorphic to that of the symmetric groups in a range of degrees that increases with the number of nodes. Furthermore, we show that when the defining parameter δ of the partition algebra is invertible, the homology of the partition algebra is in fact isomorphic to the homology of the symmetric group in all degrees. These results parallel those obtained for the Brauer algebras in the authors' earlier work, but with significant differences and difficulties in the *inductive resolution* and *high acyclicity* arguments required to prove them. Our results join the growing literature on homological stability for algebras, which now encompasses the Temperley-Lieb, Brauer and partition algebras, as well as the Iwahori-Hecke algebras of types A and B .

1. INTRODUCTION

In the last few years it has become increasingly apparent that the techniques of *homological stability*, which are most commonly applied to families of groups, can be successfully applied to families of *algebras*, where homology is interpreted as an appropriate Tor group. Indeed, the papers of Boyd–Hepworth [BH20], Boyd–Hepworth–Patz [BHP21], Hepworth [Hep22] and Moselle [Mos22] proved homological stability for Temperley-Lieb algebras, Brauer algebras, and Iwahori-Hecke algebras of types A and B respectively, and identified the stable homology in the first two cases. The Temperley-Lieb and Brauer algebras failed to satisfy a certain flatness condition that holds automatically for families of groups, necessitating the introduction of the new technique of *inductive resolutions*. Using related techniques, Sroka [Sro22] showed that the homology of the Temperley-Lieb algebra on an odd number of strands vanishes in positive degrees, in contrast to the known non-vanishing for an even number of strands. More recently, Boyde [Boy22] used a careful study of idempotents to unify and generalise the ‘invertible parameter’ results from [BH20, BHP21], together with Sroka’s vanishing result. In this paper, we prove homological stability for the partition algebras, and we identify their stable homology.

The *partition algebras* were introduced independently by Jones [Jon94] and Martin [Mar94] for their relevance in studying Potts models in statistical mechanics. They are also important in representation theory as a Schur–Weyl dual to the symmetric group, as in the work of Halverson–Ram [HR05, Theorems 5.4, 3.6] and


2020 *Mathematics Subject Classification.* 20J06, 16E40 (primary), 20B30 (secondary) .

Key words and phrases. Homology, homological stability, partition algebras.

For the purpose of open access, the authors have applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

Bowman–Doty–Martin [BDM22]. They contain a rich variety of subalgebras, including the planar partition, rook Brauer, rook, planar rook, Brauer, Motzkin and Temperley–Lieb algebras.

Given a commutative ring R , an element $\delta \in R$, and a non-negative integer n , the partition algebra $P_n(R, \delta)$ is defined to be the free module over R with basis given by the partitions of the set $\{-1, \dots, -n, 1, \dots, n\}$. These partitions can be drawn as diagrams with n nodes labelled $-1, \dots, -n$ on the left and n nodes labelled $1, \dots, n$ on the right. Nodes in the same block of a partition are then joined by edges. For ease of drawing, we do not include all edges but instead rely on transitivity. Disconnected nodes are allowed, corresponding to blocks of size one. For example, the following diagram shows the basis element $\{\{-1, -3\}, \{-2, -4, 4\}, \{2, 3\}, \{1\}\}$ of $P_4(R, \delta)$.



Multiplication is given by placing the diagrams side by side, identifying the middle nodes, and replacing any blocks not connected to the right or left by a factor of δ .

Diagrams in which every node on the left is connected to a single node on the right, and nothing else, are called *permutation diagrams*, and are in bijection with elements of the symmetric group $\mathfrak{S}_n$. This gives rise to inclusion and projection maps

$$R\mathfrak{S}_n \xrightarrow{\iota} P_n(R, \delta) \xrightarrow{\pi} R\mathfrak{S}_n$$

where ι sends permutations to permutation diagrams, and π does the reverse while sending all remaining diagrams to 0. In particular, $\pi \circ \iota$ is the identity map on $R\mathfrak{S}_n$.

We denote the trivial module of $R\mathfrak{S}_n$ by $\mathbb{1}$. Pulling back along π , we obtain the *trivial module* $\mathbb{1}$ of $P_n(R, \delta)$. This gives us the homology groups $H_*(\mathfrak{S}_n, \mathbb{1}) = \text{Tor}_*^{R\mathfrak{S}_n}(\mathbb{1}, \mathbb{1})$ of $\mathfrak{S}_n$ and $\text{Tor}_*^{P_n(R, \delta)}(\mathbb{1}, \mathbb{1})$ of $P_n(R, \delta)$. There are induced homomorphisms ι_* and π_* on homology groups for which $\pi_* \circ \iota_*$ is again the identity, so that the homology of $\mathfrak{S}_n$ appears as a direct summand of the homology of $P_n(R, \delta)$.

Theorem A. *Suppose that δ is invertible in R . Then the homology of the partition algebra is isomorphic to the homology of the symmetric group:*

$$\text{Tor}_*^{P_n(R, \delta)}(\mathbb{1}, \mathbb{1}) \cong H_*(\mathfrak{S}_n; \mathbb{1})$$

Indeed, the inclusion and projection maps

$$R\mathfrak{S}_n \xrightarrow{\iota} P_n(R, \delta) \xrightarrow{\pi} R\mathfrak{S}_n$$

induce inverse isomorphisms

$$\text{Tor}_*^{R\mathfrak{S}_n}(\mathbb{1}, \mathbb{1}) \xrightarrow[\cong]{\iota_*} \text{Tor}_*^{P_n(R, \delta)}(\mathbb{1}, \mathbb{1}) \xrightarrow[\cong]{\pi_*} \text{Tor}_*^{R\mathfrak{S}_n}(\mathbb{1}, \mathbb{1}).$$

Our second result holds without any assumptions on the value of δ .

Theorem B. *The inclusion map $\iota: R\mathfrak{S}_n \rightarrow P_n(R, \delta)$ induces a map in homology*

$$\iota_*: H_i(\mathfrak{S}_n; \mathbb{1}) \longrightarrow \text{Tor}_i^{P_n(R, \delta)}(\mathbb{1}, \mathbb{1})$$

that is an isomorphism in the range $n \geq 2i + 1$.

An immediate consequence of [Theorem B](#) is the following corollary.

Corollary C. *The partition algebras satisfy homological stability, that is, the inclusion $P_{n-1}(R, \delta) \hookrightarrow P_n(R, \delta)$ induces a map*

$$\mathrm{Tor}_i^{P_{n-1}(R, \delta)}(\mathbb{1}, \mathbb{1}) \longrightarrow \mathrm{Tor}_i^{P_n(R, \delta)}(\mathbb{1}, \mathbb{1})$$

that is an isomorphism in degrees $n \geq 2i + 1$, and this stable range is sharp. Furthermore, $P_n(R, \delta)$ and $\mathfrak{S}_n$ have the same stable homology:

$$\lim_{n \rightarrow \infty} H_*(\mathfrak{S}_n; \mathbb{1}) \cong \lim_{n \rightarrow \infty} \mathrm{Tor}_*^{P_n(R, \delta)}(\mathbb{1}, \mathbb{1}).$$

The first part of this corollary follows by combining [Theorem B](#) with Nakaoka's homological stability result for the symmetric groups, for which the stable range is sharp ([\[Nak60\]](#)). For the stable homology, the left hand side of this isomorphism is well known by the Barratt–Priddy–Quillen theorem [\[BP72, FM94\]](#). The above results exactly parallel the situation for the Brauer algebras, and as discussed in [\[BHP21\]](#) are reminiscent of the relationship between $\mathfrak{S}_n$ and the automorphism groups of free groups $\mathrm{Aut}(F_n)$ (see Galatius [\[Gal11\]](#)).

1.1. Outline, and comparison to previous work. In [Section 2](#) we introduce partition algebras and provide the necessary background needed for the rest of the paper. In [Section 3](#) we restate an abstract form of the principle that lies behind the technique of inductive resolutions that was introduced in [\[BH20\]](#), and was a crucial ingredient in [\[BH20\]](#) and [\[BHP21\]](#). In [Section 4](#) we establish the existence of inductive resolutions for the partition algebras. These are significantly more complicated than the Temperley–Lieb [\[BH20\]](#) and Brauer [\[BHP21\]](#) cases, and we find that we must consider several families of distinct modules in order to carry out our induction argument. In [Section 5](#) we follow the argument of [\[BHP21\]](#) to replace Shapiro's lemma in the setting of partition algebras. The high connectivity result required for any new proof of homological stability is found in [Section 6](#). Like our inductive resolutions argument, this is again more complicated than the analogous result in [\[BHP21\]](#), and heavily utilises the high connectivity of the complex of injective words with separators, introduced in that paper. We finish in [Section 7](#) by giving an account of the proof of the main theorem, which follows the same general argument as in [\[BHP21\]](#).

It is common, in homological stability for families of groups, to find that proofs of different results have a very similar overall structure, yet the proofs that the relevant complexes are highly acyclic can differ radically. What we can now see in homological stability for algebras, comparing the work of this paper to that of [\[BH20\]](#) and [\[BHP21\]](#), is an analogous situation where the overall technique is used in multiple situations, but the details of the acyclicity proofs — and now also of the inductive resolutions proofs — are where the important differences and difficulties lie.

1.2. Acknowledgements. The first author was supported by EPSRC Postdoctoral Fellowship EP/V043323/1 and EP/V043323/2. The third author was supported by a Simons Collaboration Grant. Our thanks to the anonymous referee for their detailed reading and helpful comments.

2. PARTITION ALGEBRAS

In this section we introduce the partition algebra, together with some specific elements and modules that will be important later in the paper.

Definition 2.1 (The partition algebra [Jon94, Mar94]). As explained in the introduction, if R is a commutative ring, δ is a chosen element in R , and n is a non-negative integer, then the partition algebra $P_n(R, \delta)$ is defined to be the free module over R with basis given by the partitions of the set $\{-n, \dots, -1, 1, \dots, n\}$. These are drawn as diagrams with nodes $-1, \dots, -n$ on the left and nodes $1, \dots, n$ on the right, with arcs indicating which nodes lie in the same block of the partition. (We allow ourselves to omit some arcs and instead use transitivity to determine the blocks.) An example is shown in Figure 1. Multiplication is given by placing the diagrams side by side, identifying the middle nodes, and replacing any blocks not connected to the right or left by a factor of δ , as in Figure 2.

We will use the terms ‘partition’ and ‘diagram’ interchangeably to mean a basis element of $P_n(R, \delta)$, and we will frequently abbreviate $P_n(R, \delta)$ as P_n .




FIGURE 1. Visualization of the partition $\{-5, -3\}, \{-4, -2, -1, 3, 4\}, \{1, 5\}, \{2\}$





FIGURE 2. Multiplication in the partition algebra

The partition algebra is generated by three types of diagrams [Mar96], corresponding to the following partitions:

- For $1 \leq i \leq n-1$, S_i is the diagram corresponding to the partition with blocks of pairs $\{-j, j\}$ for $j \neq i, i+1$, together with $\{-(i+1), i\}$ and $\{-i, (i+1)\}$. These generate the group ring of the symmetric group, $\mathfrak{S}_n$, as a subalgebra of P_n .
- For $1 \leq i \neq j \leq n-1$, V_{ij} is the diagram corresponding to the partition with blocks of pairs $\{-k, k\}$ for $k \neq i, j$ and one block of size four $\{-j, -i, i, j\}$.
- For $1 \leq i \leq n$, T_i is the diagram corresponding to the partition with blocks of pairs $\{-j, j\}$ for $j \neq i$ and two singleton blocks $\{-i\}$ and $\{i\}$.

See Figure 3 for depictions of some of these.

We now introduce the modules we will be working with.

FIGURE 3. The elements $S_2, V_{13}, T_3 \in P_4$

Recall that by a *permutation diagram* we mean a diagram in which each node on the left is joined to a single node on the right, and nothing else. Equivalently, permutation diagrams are ones that do not contain any singletons on the right *or* any blocks that contain ≥ 2 elements on the right.

Definition 2.2 (The trivial module $\mathbb{1}$). For any n , we define the *trivial* $R\mathfrak{S}_n$ -bimodule $\mathbb{1}$ to be the module given by the ring R , upon which the permutations act as the identity.

For any n , we define the *trivial* P_n -bimodule $\mathbb{1}$ to be the module given by the ring R , upon which the permutation diagrams act as the identity, and all other diagrams act as 0. This is the same as acting with P_n on R via the projection $\pi: P_n \rightarrow \mathfrak{S}_n$.

Definition 2.3. For $m \leq n$, we can view P_m as a subalgebra of P_n . Given a partition of $\{-m, \dots, -1, 1, \dots, m\}$, the map which sends $(\pm 1, \dots, \pm m)$ to $(\pm(n-m+1), \dots, \pm n)$ induces a partition on $\{-n, \dots, -(n-m+1), (n-m+1), \dots, n\}$. We add the blocks $\{-i, i\}$ for all $i \in \{1, \dots, (n-m)\}$, resulting in a partition in P_n . Pictorially, we are taking diagrams in P_m and extending them to ones in P_n by adding new nodes *below* the existing ones, with horizontal connections between the new nodes. Then, under the action of this subalgebra, P_n can be viewed as a left P_n -module and a right P_m -module, and we obtain the induced left P_n -module $P_n \otimes_{P_m} \mathbb{1}$.

The following proposition is taken from [Pat20].

Proposition 2.4 ([Pat20, Proposition 2.5]). *The induced module $P_n \otimes_{P_m} \mathbb{1}$ is a free R -module and a quotient of P_n .*

In terms of diagrams, a basis for this module is the set of diagrams in which the top m nodes on the right are placed under a box, satisfying the following condition:

- *The box is connected to exactly m distinct blocks.*

Under this description, the action of P_n on $P_n \otimes_{P_m} \mathbb{1}$ is given by pasting and simplifying the diagrams just as in the multiplication of P_n , and then identifying a diagram with 0 if it violates the condition above.

Thus there are two ways that a diagram could be identified with 0 after left multiplication by a diagram in P_n : One of the blocks attached to the box could, after pasting, consist only of nodes in the centre (visually, that block is free to be retracted into the box, and then disappears). Alternatively, two or more distinct blocks that were attached to the box can become fused into a single block (visually, there is now a path of arcs with both ends attached to the box). These two possibilities correspond to the two ways in which a diagram in P_m can fail to be a permutation

diagram, and therefore act as 0 on $\mathbb{1}$: It can have a singleton on the right, or it can have two nodes on the right belonging to the same block.

Example 2.5. Figure 4 depicts the module structure of $P_5 \otimes_{P_3} \mathbb{1}$. In the first example one of the blocks connected to the box consists entirely of nodes in the centre and therefore ‘vanishes’ or ‘retracts into the box’. In the second example the factor of δ arises due to a block that consists entirely of nodes in the centre and is not attached to the box.




FIGURE 4. The module structure of $P_5 \otimes_{P_3} \mathbb{1}$

3. THE PRINCIPLE OF INDUCTIVE RESOLUTIONS

In this brief section we state an abstract form of the principle that underlies the technique of inductive resolutions that appeared in [BH20] and [BHP21]. It allows us to identify modules that vanish under a fixed functor of the form $\text{Tor}_i^A(M, -)$ by resolving them using modules that already have this property, hence the name ‘inductive resolutions’. The theorem below is an abstraction of Section 3.3 of [BH20]. It can be regarded as an application of the general principle that a derived functor can be computed using resolutions by objects that are acyclic for that derived functor.

Theorem 3.1. *Let A be an algebra over a ring R , and let M be a right A -module. Suppose that N is a left A -module equipped with a resolution $Q_* \rightarrow N$ with the following two properties:*

- $\text{Tor}_*^A(M, Q_j)$ vanishes in positive degrees for all $j \geq 0$.
- $M \otimes_A Q_* \rightarrow M \otimes_A N$ is a resolution.

Then $\text{Tor}_^A(M, N)$ vanishes in positive degrees.*

Proof. Let $P_* \rightarrow M$ be a projective resolution, so that for any left A -module B , the groups $\text{Tor}_*^A(M, B)$ are computed by the complex $P_* \otimes_A B$. Consider the double complex $P_* \otimes_A Q_*$. There are two natural spectral sequences converging to the homology of the totalisation $\text{Tot}(P_* \otimes_A Q_*)$. For more on these spectral sequences, see Section 5.6 of [Wei94] and the summary in Section 3.2 of [BH20].

The first spectral sequence has E^1 -term given by

$${}^I E_{i,j}^1 = H_j(P_i \otimes_A Q_*) \cong P_i \otimes_A H_j(Q_*)$$

with d^1 induced by the differential of P . The isomorphism holds because each P_i is projective and therefore flat. It follows that the E^2 -term is

$${}^I E_{i,j}^2 = \mathrm{Tor}_i^A(M, H_j(Q_*)).$$

Since Q_* is a resolution of N , it follows that ${}^I E_{*,*}^2$ is simply $\mathrm{Tor}_*^A(M, N)$ concentrated on the x -axis, so that the same is true of ${}^I E_{*,*}^\infty$, and therefore we conclude that $H_*(\mathrm{Tot}(P_* \otimes_A Q_*)) \cong \mathrm{Tor}_*^A(M, N)$.

The second spectral sequence has E^1 -term given by ${}^{II} E_{i,j}^1 = H_j(P_* \otimes_A Q_i)$, *i.e.*

$${}^{II} E_{i,j}^1 = \mathrm{Tor}_j^A(M, Q_i)$$

with d^1 induced by the boundary maps of Q_* . Our first assumption now shows that ${}^{II} E_{*,*}^1$ is concentrated on the x -axis, where it is given by $\mathrm{Tor}_0^A(M, Q_*) = M \otimes_A Q_*$. Consequently ${}^{II} E_{*,*}^2$ is given by the homology of $M \otimes_A Q_*$, which by our second assumption is simply a copy of $M \otimes_A N$ at the origin. This shows that the homology of $\mathrm{Tot}(P_* \otimes_A Q_*)$ is simply a copy of $M \otimes_A N$ in degree 0.

Comparing the outcomes of the two spectral sequences, we see that $\mathrm{Tor}_*^A(M, N)$ vanishes in positive degrees, as required. $\square$

4. INDUCTIVE RESOLUTIONS

In this section, we will use the technique of *inductive resolutions*, which originated in [BH20] and was further used in [BHP21].

Definition 4.1. Suppose that X is a subset of the set $\{1, \dots, n\}$. Define J_X to be the left-ideal in P_n that is the R -span of all diagrams in which, among the nodes on the right labelled by elements of X , there is at least one singleton or one pair of nodes that are in the same block. For $m \leq n$, let $J_{\{n-m+1, \dots, n\}}$ be denoted by J_m .

Observe that $J_n = J_{\{1, \dots, n\}}$ is the span of precisely the diagrams that are *not* permutation diagrams. It is therefore the kernel of the projection map $\pi: P_n \rightarrow R\mathfrak{S}_n$.

Our aim in this section is to prove the following theorem, which will be used in the final section to understand the Tor groups $\mathrm{Tor}_*^{P_n}(\mathbb{1}, P_n \otimes_{P_m} \mathbb{1})$.

Theorem 4.2. *Let $X \subseteq \{1, \dots, n\}$ and suppose that one of the following conditions holds:*

- $|X| \leq n$ and δ is invertible in R .
- $|X| < n$.

Then the groups $\mathrm{Tor}_^{P_n}(\mathbb{1}, P_n/J_X)$ vanish in positive degrees.*

The proof of Theorem 4.2 will occupy the rest of the section. Aspects of the material are close to [BH20, Section 3] and [BHP21, Section 3], but overall the material here is significantly more complex.

Before we continue, we record the following lemma, which extends Theorem 4.2 to degree 0. We will need this lemma to prove the theorem.

Lemma 4.3. *Let J be a left ideal of P_n that is included in J_n . Then*

$$\mathbb{1} \otimes_{P_n} P_n / J \cong \mathbb{1}.$$

In particular,

$$\mathrm{Tor}_0^{P_n}(\mathbb{1}, P_n / J_X) \cong \mathbb{1}$$

for all $X \subset \{1, \dots, n\}$.

Proof. Due to the inclusions $0 \subset J \subset J_n$, we have the surjections

$$\mathbb{1} \otimes_{P_n} P_n \longrightarrow \mathbb{1} \otimes_{P_n} P_n / J \longrightarrow \mathbb{1} \otimes_{P_n} P_n / J_n.$$

Because

$$\mathbb{1} \otimes_{P_n} P_n \cong \mathbb{1}$$

and

$$\mathbb{1} \otimes_{P_n} P_n / J_n \cong \mathbb{1} \otimes_{P_n} R\mathfrak{S}_n \cong \mathbb{1} \otimes_{R\mathfrak{S}_n} R\mathfrak{S}_n \cong \mathbb{1},$$

the above composition is an isomorphism and the first map must be also injective. $\square$

4.1. Reducing to $\mathcal{A}_{X,x}$ and $\mathcal{B}_{X,x}$. Our proof of [Theorem 4.2](#) will be by induction on the cardinality of X . Ideally we would prove the inductive step by resolving P_n / J_X in terms of modules $P_n / J_{X'}$ with $|X'| < |X|$. However, we were not able to find a straightforward argument along these lines. To organise the argument, in this subsection we introduce some intermediate modules, and later on we will build our resolutions with these.

Definition 4.4. Let $Y \subseteq X \subseteq \{1, \dots, n\}$, and let $x \in X$ and $y \in Y$. We define three left P_n -submodules of P_n as follows:

- A_x is the span of all diagrams in which x is a singleton.
- $B_{X,x}$ is the span of all diagrams in which x lies in the same block as some other element of X .
- M_Y is the span of all diagrams in which the elements of Y lie in the same block.

We define quotients of these as follows:

$$\mathcal{A}_{X,x} = \frac{A_x}{A_x \cap J_{X-\{x\}}}, \quad \mathcal{B}_{X,x} = \frac{B_{X,x}}{B_{X,x} \cap J_{X-\{x\}}}, \quad \mathcal{M}_{X,Y} = \frac{M_Y}{M_Y \cap J_{X-Y}}$$

The following result will be useful when we come to verify the second condition of [Theorem 3.1](#).

Lemma 4.5. *The modules $\mathcal{A}_{X,x}$, $\mathcal{B}_{X,x}$, and $\mathcal{M}_{X,Y}$ behave as follows under tensor product with $\mathbb{1}$.*

- Let $x \in X \subseteq \{1, \dots, n\}$, and suppose that $n \geq 2$. Then $\mathbb{1} \otimes_{P_n} \mathcal{A}_{X,x} = 0$.
- Let $x \in X \subseteq \{1, \dots, n\}$. Then $\mathbb{1} \otimes_{P_n} \mathcal{B}_{X,x} = 0$.
- Let $Y \subseteq X \subseteq \{1, \dots, n\}$ with $|Y| \geq 2$. Then $\mathbb{1} \otimes_{P_n} \mathcal{M}_{X,Y} = 0$ and $\mathcal{M}_{X,Y}$ is a direct summand of P_n / J_{X-Y} .

Proof. We will show that, under the relevant conditions, each of A_x , $B_{X,x}$ and M_Y vanishes under $\mathbb{1} \otimes_{P_n} -$, and the same will then follow for $\mathcal{A}_{X,x}$, $\mathcal{B}_{X,x}$ and $\mathcal{M}_{X,Y}$.

To show that $\mathbb{1} \otimes_{P_n} A_x = 0$, we take a diagram α in A_x , so that x is a singleton in α . Let β denote a diagram obtained from α by placing x into the same block as some other element on the right. (This is possible by the assumption that $n \geq 2$.) Then $\alpha = \beta \cdot T_x$, and β acts as 0 on $\mathbb{1}$, so that

$$1 \otimes \alpha = 1 \otimes \beta \cdot T_x = 1 \cdot \beta \otimes T_x = 0 \otimes T_x = 0,$$

noting that $T_x \in A_x$. Since A_x is the span of such diagrams α , this completes the proof.

The argument for the other two modules is similar. For $\mathbb{1} \otimes_{P_n} B_{X,x}$, we take a diagram $\alpha \in B_{X,x}$, so that x is in the same block as some other element $w \in X$, and we use the factorisation $\alpha = \alpha \cdot V_{wx}$, noting that $V_{wx} \in B_{X,x}$.

For M_Y we take a diagram $\alpha \in M_Y$, so that all elements of Y lie in the same block, and factorise it as $\alpha = \alpha \cdot V_Y$ where $V_Y \in M_Y$ is the diagram with blocks $-Y \cup Y$ and $\{-p, p\}$ for $p \notin Y$; the assumption $|Y| \geq 2$ ensures that α acts as 0 on $\mathbb{1}$.

For the final claim about $\mathcal{M}_{X,Y}$, we use the fact that the element V_Y above is idempotent and sends J_{X-Y} into itself. $\square$

The following proposition breaks down the problem of resolving P_n/J_X into the analogous problem for $\mathcal{A}_{X,x}$ and $\mathcal{B}_{X,x}$.

Proposition 4.6. *Let $X \subseteq \{1, \dots, n\}$, let $x \in X$, and assume $n \geq 2$. The following sequence, in which all maps are induced by either an inclusion or an identity map, is a resolution of P_n/J_X .*

$$\begin{array}{ccccccc} \dots & \longrightarrow & 0 & \longrightarrow & \mathcal{A}_{X,x} \oplus \mathcal{B}_{X,x} & \longrightarrow & P_n/J_{X-\{x\}} \longrightarrow P_n/J_X \\ & & 2 & & 1 & & 0 & & -1 \end{array}$$

Moreover, applying $\mathbb{1} \otimes_{P_n} -$ to the sequence gives a resolution of $\mathbb{1} \otimes_{P_n} P_n/J_X$.

Proof. The map $P_n/J_{X-\{x\}} \rightarrow P_n/J_X$ is induced by the identity map on P_n and is well defined since $J_{X-\{x\}} \subset J_X$. The map $\mathcal{A}_{X,x} \rightarrow P_n/J_{X-\{x\}}$ is induced by the inclusion $A_x \subset P_n$ and is well defined since $(A_x \cap J_{X-\{x\}}) \subset J_{X-\{x\}}$, and a similar argument holds for the map $\mathcal{B}_{X,x} \rightarrow P_n/J_{X-\{x\}}$.

Surjectivity of the right hand map is immediate, giving exactness in degree -1 .

To show exactness in degree 0, observe that the ideals $J_{X-\{x\}} \subseteq J_X$ are both spanned by certain diagrams, so that the kernel of $P_n/J_{X-\{x\}} \rightarrow P_n/J_X$ is spanned by those diagrams that lie in J_X but not $J_{X-\{x\}}$. For a diagram to lie in J_X , some element of X must be a singleton, or two elements of X must lie in the same block. For it to not *also* be an element of $J_{X-\{x\}}$, there must only be one singleton, namely x , or only one pair of elements lying in the same block, of which one must be x . The diagrams with x a singleton are precisely the diagrams that span A_x , the diagrams in which x lies in the same block as some other element of X are precisely those that span $B_{X,x}$, and the proof follows.

To show exactness in degree 1, after unravelling the definitions of $\mathcal{A}_{X,x}$ and $\mathcal{B}_{X,x}$, it is sufficient to show that if we have $a \in A_x$ and $b \in B_{X,x}$ with $a + b \in J_{X-\{x\}}$,

then $a, b \in J_{X-\{x\}}$. This follows quickly from fact that A_x and $B_{X,x}$ have no basis elements in common.

To prove the second claim, we will show that after applying $\mathbb{1} \otimes_{P_n} -$, the resolution becomes

$$\begin{array}{ccccccc} \dots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \mathbb{1} \xrightarrow{\text{Id}} \mathbb{1} \\ & & 2 & & 1 & & 0 & & -1 \end{array}$$

so that the claim follows directly. The identification of the final two terms and the map between them follows from [Lemma 4.3](#). The terms in degree 1 vanish by [Lemma 4.5](#). (This is where we use the assumption $n \geq 2$.) $\square$

The last result, together with [Theorem 3.1](#), shows that, in order to prove vanishing of higher Tor's for P_n/J_X by induction, we must first do the same for $\mathcal{A}_{X,x}$ and $\mathcal{B}_{X,x}$. In the following two subsections we will construct resolutions of these.

4.2. Resolving $\mathcal{A}_{X,x}$. We now attempt to resolve $\mathcal{A}_{X,x}$. It will turn out that this requires different methods depending on which assumption from [Theorem 4.2](#) we use: that δ is invertible, or that $|X| < n$. Under the first assumption we have the following.

Proposition 4.7. *Suppose that $X \subseteq \{1, \dots, n\}$ and that δ is invertible in R . Then the module $\mathcal{A}_{X,x}$ is a direct summand of $P_n/J_{X-\{x\}}$.*

Proof. The element $\delta^{-1}T_x$ is an idempotent, thanks to the computation $T_x^2 = \delta T_x$. Right multiplication by $\delta^{-1}T_x$ sends $J_{X-\{x\}}$ into itself, and therefore induces an idempotent endomorphism of $P_n/J_{X-\{x\}}$. The image of this endomorphism consists of all left multiples of T_x , and this is precisely $\mathcal{A}_{X,x} = \frac{A_x}{A_x \cap J_{X-\{x\}}}$ as in the second paragraph of the proof of [Lemma 4.5](#). $\square$

The above result shows that, if $P_n/J_{X-\{x\}}$ has vanishing higher Tor's, then so does $\mathcal{A}_{X,x}$. When δ is not invertible, we need a more elaborate method using the following resolution.

Definition 4.8 (The resolution $C(X, x, y)$). Suppose that $X \subset \{1, \dots, n\}$ with $|X| < n$, let $x \in X$, and let $y \in \{1, \dots, n\} - X$. We define a resolution $C(X, x, y) \rightarrow \mathcal{A}_{X,x}$ as in [Figure 5](#). Thus $C(X, x, y)$ is given by $P_n/J_{X-\{x\}}$ in each degree. The maps are all given by right-multiplication by the indicated elements, so that the boundary maps alternate between $(1 - T_x V_{xy})$ and $T_x V_{xy}$, and the augmentation $P_n/J_{X-\{x\}} \rightarrow \mathcal{A}_{X,x}$ is given by T_x . The maps are well defined thanks to the fact that $T_x V_{xy}$ and T_x send $J_{X-\{x\}}$ into itself; in the former case this follows from the fact that $y \notin X$.

To check that consecutive maps compose to 0, one uses the computation $T_x V_{xy} T_x = T_x$ together with the resulting fact that $T_x V_{xy}$ is an idempotent. The fact that this really does define a resolution is given next.

Proposition 4.9. *Suppose that $X \subset \{1, \dots, n\}$ with $|X| < n$, let $x \in X$, and let $y \in \{1, \dots, n\} - X$. Assume that $n \geq 2$. Then*

$$C(X, x, y) \rightarrow \mathcal{A}_{X,x} \quad \text{and} \quad \mathbb{1} \otimes_{P_n} C(X, x, y) \rightarrow \mathbb{1} \otimes_{P_n} \mathcal{A}_{X,x}$$


are both resolutions.

$$\begin{array}{ccc}
& \vdots & \\
& \downarrow (1-T_x V_{xy}) & \\
P_n / J_{X-\{x\}} & & 2 \\
& \downarrow T_x V_{xy} & \\
P_n / J_{X-\{x\}} & & 1 \\
& \downarrow (1-T_x V_{xy}) & \\
P_n / J_{X-\{x\}} & & 0 \\
& \downarrow T_x & \\
\mathcal{A}_{X,x} & & -1
\end{array}$$

FIGURE 5. The resolution $C(X, x, y) \rightarrow \mathcal{A}_{X,x}$

Proof. First, we must show that $C(X, x, y) \rightarrow \mathcal{A}_{X,x}$ is acyclic. In degree -1 this is clear since A_x consists of all left multiples of T_x . In degrees 1 and above, this is an immediate consequence of the fact that $T_x V_{xy}$ is an idempotent. In degree 0 we require a more complex argument, as follows.

Suppose α is a diagram in which x is a singleton. If B is a block of α other than $\{x\}$, then we write α_B for the diagram obtained from α by incorporating x into B . And we write α_y for the diagram α_{B_y} , where B_y is the block containing y . For example, the following diagrams show α with B_y drawn in red and another block B drawn in blue, together with α_B and α_y .



Now observe that we have the relations

$$(4.1) \quad \alpha - \delta \alpha_y = \alpha(1 - T_x V_{xy})$$

and, for each block B in α ,

$$(4.2) \quad \alpha_B - \alpha_y = \alpha_B(1 - T_x V_{xy}).$$

Now consider an element $a \in P_n / J_{X-\{x\}}$ in degree 0 that lies in the kernel of the augmentation $T_x: P_n / J_{X-\{x\}} \rightarrow \mathcal{A}_{X,x}$. We wish to show that a is in the image of the differential, and we will do this by explaining how to adjust a by elements in the image of the differential (which does not change the fact that it lies in the kernel) in order to reduce it to 0. We can write a as a linear combination of diagrams in P_n that do not lie in $J_{X-\{x\}}$, and these can be divided into the following cases:

- (a) Diagrams α in which x is a singleton. Using elements of the form (4.1), we may adjust a by elements in the image of the differential in order to replace all such diagrams α with ones of the form α_y .
- (b) Diagrams in which x is connected to some element outwith $X - \{x\}$. These diagrams all have the form α_B , where α is the diagram obtained from the original by making x a singleton, and B is the block of α that originally contained x . Note that the assumption that the original diagram did not lie in $J_{X-\{x\}}$ means that α also does not lie in $J_{X-\{x\}}$. Using elements of the form (4.2), we may adjust a by elements in the image of the differential in order to replace all such diagrams α_B with ones of the form α_y .
- (c) Diagrams β in which x is connected to exactly one element, say w , in $X - \{x\}$. Then in $\beta T_x V_{xy}$ the element $w \in X - \{x\}$ is a singleton, so that $\beta T_x V_{xy} \in J_{X-\{x\}}$. Consequently $\beta = \beta(1 - T_x V_{xy})$ in $P_n / J_{X-\{x\}}$, and in particular β lies in the image of the differential. We may therefore adjust a by elements in the image of the differential to remove all diagrams of this form.

After modifying a as explained in each item above, we may now write it as a linear combination $a = \sum_{\alpha} \lambda_{\alpha} \alpha_y$ where α ranges over all diagrams that are not in $J_{X-\{x\}}$ and in which x is a singleton. We know that a lies in the kernel of the differential, so that $a \cdot T_x = 0$. However, we have $a \cdot T_x = \sum_{\alpha} \lambda_{\alpha} \alpha_y \cdot T_x = \sum_{\alpha} \lambda_{\alpha} \alpha$, and since the α are distinct diagrams not in $J_{X-\{x\}}$, we can conclude that $\lambda_{\alpha} = 0$ for all α , or in other words that $a = 0$. This completes the argument in degree 0, and so completes the proof that $C(X, x, y) \rightarrow \mathcal{A}_{X,x}$ is a resolution.

We now prove that $\mathbb{1} \otimes_{P_n} C(X, x, y) \rightarrow \mathbb{1} \otimes_{P_n} \mathcal{A}_{X,x}$ is a resolution. The target vanishes by Lemma 4.5, and $\mathbb{1} \otimes_{P_n} P_n / J_{X-\{x\}} = \mathbb{1}$ since $J_{X-\{x\}}$ acts as 0 on $\mathbb{1}$. Under the latter identification, the boundary maps, which used to be given by right-multiplication by the indicated elements, are now given by the action of those elements on $\mathbb{1}$, and therefore alternate between 0 and 1. The result follows. $\square$

4.3. Resolving $\mathcal{B}_{X,x}$. We now turn to the module $\mathcal{B}_{X,x}$, for which we build the following resolution.

Definition 4.10 (The resolution $D(X, x) \rightarrow \mathcal{B}_{X,x}$). Let $X \subseteq \{1, \dots, n\}$ and let $x \in X$. Define an augmented complex $D(X, x) \rightarrow \mathcal{B}_{X,x}$ as follows. In degree $i \geq 0$, $D(X, x)$ is given by

$$\bigoplus_{(x_0, \dots, x_i)} \mathcal{M}_{X, \{x, x_0, \dots, x_i\}}$$

where the sum is over all tuples $(x_0, \dots, x_i)$ of *distinct* elements of $X - \{x\}$. And on the summand corresponding to a tuple $(x_0, \dots, x_i)$, the map δ_i is given by the map

$$\mathcal{M}_{X, \{x, x_0, \dots, x_i\}} \longrightarrow \mathcal{M}_{X, \{x, x_0, \dots, x_{i-1}\}}$$

obtained from the inclusions

$$\begin{aligned} M_{\{x, x_0, \dots, x_i\}} &\hookrightarrow M_{\{x, x_0, \dots, x_{i-1}\}} \\ J_{X-\{x, x_0, \dots, x_i\}} &\hookrightarrow J_{X-\{x, x_0, \dots, x_{i-1}\}}. \end{aligned}$$

The map δ_i is simply the sum of these individual maps. To it put briefly, δ_i is the map that forgets that x_i had to be in the same block as $x, x_0, \dots, x_{i-1}$. The complex is illustrated in Figure 6. Each composite $\delta_{i-1} \circ \delta_i$ vanishes, because any element in its image is a sum of diagrams that each contain two elements $x_{i-1}, x_i \in$

$$\begin{array}{ccc}
& \vdots & \\
& \downarrow \delta_3 & \\
\bigoplus_{(x_0, x_1, x_2)} \mathcal{M}_{X, \{x, x_0, x_1, x_2\}} & & 2 \\
& \downarrow \delta_2 & \\
\bigoplus_{(x_0, x_1)} \mathcal{M}_{X, \{x, x_0, x_1\}} & & 1 \\
& \downarrow \delta_1 & \\
\bigoplus_{(x_0)} \mathcal{M}_{X, \{x, x_0\}} & & 0 \\
& \downarrow \delta_0 & \\
\mathcal{B}_{X, x} & & -1
\end{array}$$

FIGURE 6. The resolution $D(X, x) \rightarrow \mathcal{B}_{X, x}$. Summations are over tuples of distinct elements of $X - \{x\}$.

$X - \{x, x_1, \dots, x_{i-2}\}$ in the same block, and which therefore lie in $J_{X - \{x, x_1, \dots, x_{i-2}\}}$. We prove that this is indeed a resolution in [Proposition 4.11](#).

Proposition 4.11. *$D(X, x) \rightarrow \mathcal{B}_{X, x}$ is indeed a resolution, and the same is true for $\mathbb{1} \otimes_{P_n} D(X, x) \rightarrow \mathbb{1} \otimes_{P_n} \mathcal{B}_{X, x}$.*

Proof. We first prove that $D(X, x) \rightarrow \mathcal{B}_{X, x}$ is acyclic.

In degree -1 we must show that δ_0 is surjective. A diagram in $B_{X, x}$ has x in the same block as some other element x_0 of $X - \{x\}$, and therefore lies in the image of the inclusion $M_{\{x, x_0\}} \hookrightarrow B_{X, x}$, and surjectivity follows.

In degree 0 , we first observe that if we consider any two summands in degree 0 , then their images under δ_0 have trivial intersection. Indeed, this follows quickly from the fact that if x_0 and x'_0 are distinct elements of $X - \{x\}$, then $M_{\{x, x_0\}} \cap M_{\{x, x'_0\}} \subseteq J_{X - \{x\}}$, which itself holds because a diagram in $M_{\{x, x_0\}} \cap M_{\{x, x'_0\}}$ has x_0 and x'_0 in the same block as x , and therefore in the same block as one another. So to prove exactness in degree 0 we can look at just one x_0 -summand at a time:

$$\begin{array}{ccc}
\bigoplus_{x_1 \in X - \{x, x_0\}} \mathcal{M}_{X, \{x, x_0, x_1\}} & & 1 \\
& \downarrow \delta_1 & \\
\mathcal{M}_{X, \{x, x_0\}} & & 0 \\
& \downarrow \delta_0 & \\
\mathcal{B}_{X, x} & & -1
\end{array}$$

To prove that this sequence is exact at its middle term, observe that the kernel of δ_0 is spanned by diagrams in $M_{\{x,x_0\}}$ that lie in $J_{X-\{x\}} - J_{X-\{x,x_0\}}$. Pick such a diagram. For the diagram to lie in $J_{X-\{x\}}$, two elements of $X - \{x\}$ must be in the same block, or an element of $X - \{x\}$ must be a singleton. For the diagram to lie outwith $J_{X-\{x,x_0\}}$, since x_0 cannot be a singleton in $M_{\{x,x_0\}}$, we conclude that x_0 must be in the same block as some other element of $X - \{x\}$. So, x_0 lies in the same block as some element $x_1 \in X - \{x, x_0\}$, and since the diagram is in $M_{\{x,x_0\}}$ it follows that x, x_0, x_1 must all be in the same block. Thus the diagram is in $M_{\{x,x_0,x_1\}}$, and so lies in the image of δ_1 .

To prove exactness in degree $i \geq 1$ and above, one first observes that in degrees $i - 1, i, i + 1$ the complex splits as a direct sum over $(x_0, \dots, x_{i-1})$. It is therefore enough to concentrate on a single $(x_0, \dots, x_{i-1})$ -summand at a time. Having restricted to such a summand, one now proves exactness similarly to the degree 0 case, and we leave the details of this to the reader.

The fact that the resolution remains acyclic after applying $\mathbb{1} \otimes_{P_n} -$ follows immediately from [Lemma 4.5](#), which shows that in fact the resolution vanishes under this operation. $\square$

4.4. Proof of Theorem 4.2. We first tackle the cases $n = 0, 1$. When $n = 0$ we have $P_n = R$ and the claim follows immediately. When $n = 1$, we either have $X = \emptyset$, or we have $X = \{1\}$ and δ invertible. When $X = \emptyset$ we have $J_X = 0$, so that $P_n/J_X = P_n$ and the claim follows. Finally, when $X = \{1\}$ and δ is invertible, then J_X is the R -span, and indeed the P_n -span, of the idempotent $\delta^{-1}T_1$. Thus J_X and P_n/J_X are both direct summands of P_n , and in particular the latter is projective, so that the claim follows.

We now assume that $n \geq 2$, and prove the claim by strong induction on the cardinality of X . When $X = \emptyset$ we have $J_X = 0$, so that $P_n/J_X = P_n$ and the claim is immediate.

Suppose now that $|X| > 0$ and that the claim holds for all X' of a smaller cardinality. According to [Proposition 4.6](#) and [Theorem 3.1](#), it will be sufficient to show that the modules

$$P_n/J_{X-\{x\}} \quad \mathcal{A}_{X,x} \quad \mathcal{B}_{X,x}$$

all vanish under $\mathrm{Tor}_i^{P_n}(\mathbb{1}, -)$ for $i > 0$.

In the case of $P_n/J_{X-\{x\}}$ we have $\mathrm{Tor}_i^{P_n}(\mathbb{1}, P_n/J_{X-\{x\}}) = 0$ by the inductive hypothesis.

For $\mathcal{A}_{X,x}$, we divide into the case where δ is invertible, and the case where $|X| < n$. When δ is invertible, [Proposition 4.7](#) shows that $\mathcal{A}_{X,x}$ is a direct summand of $P_n/J_{X-\{x\}}$, which vanishes under $\mathrm{Tor}_i^{P_n}(\mathbb{1}, -)$ by the inductive hypothesis, so that $\mathcal{A}_{X,x}$ does as well. When $|X| < n$, [Proposition 4.9](#) gives us resolutions $C(X, x, y) \rightarrow \mathcal{A}_{X,x}$ and $\mathbb{1} \otimes_{P_n} C(X, x, y) \rightarrow \mathbb{1} \otimes_{P_n} \mathcal{A}_{X,x}$. The terms of $C(X, x, y)$ are all $P_n/J_{X-\{x\}}$, which vanish under $\mathrm{Tor}_i^{P_n}(\mathbb{1}, -)$ by the inductive hypothesis, so that [Theorem 3.1](#) applies to tell us that the same is true for $\mathcal{A}_{X,x}$ itself.

For $\mathcal{B}_{X,x}$, [Proposition 4.11](#) gives us the resolutions $D(X, x) \rightarrow \mathcal{B}_{X,x}$ and $\mathbb{1} \otimes_{P_n} D(X, x) \rightarrow \mathbb{1} \otimes_{P_n} \mathcal{B}_{X,x}$. The terms of $D(X, x)$ are direct sums of modules of the form $\mathcal{M}_{X,\{x,x_0,\dots,x_i\}}$. Each $\mathcal{M}_{X,\{x,x_0,\dots,x_i\}}$ is a direct summand of $P_n/J_{X-\{x,x_0,\dots,x_i\}}$

by Lemma 4.5, and since $\mathrm{Tor}_i^{\mathbf{P}_n}(\mathbb{1}, -)$ vanishes on the latter, it also vanishes on the former. (Note that this is the only place in our argument where we have used *strong* induction.) We can now apply Theorem 3.1 to $D(X, x)$ to find that $\mathcal{B}_{X, x}$ vanishes under $\mathrm{Tor}_i^{\mathbf{P}_n}(\mathbb{1}, -)$ as required. $\square$

5. REPLACING SHAPIRO'S LEMMA

This section closely follows Section 4 of [BHP21]. We include all statements, and proofs of the lemmas which slightly differ in the case of partition algebras. The proof of Theorem 5.1 is identical to that in [BHP21], with adapted inputs.

As in the case for the Brauer algebras, we have inclusion and projection maps

$$R\mathfrak{S}_m \xrightarrow{\iota} P_m \xrightarrow{\pi} R\mathfrak{S}_m.$$

These are compatible with the inclusions $P_m \rightarrow P_n$ and $R\mathfrak{S}_m \rightarrow R\mathfrak{S}_n$, and also respect the actions on the trivial module. They therefore induce the following maps of Tor-groups.

$$\mathrm{Tor}_*^{R\mathfrak{S}_n}(\mathbb{1}, R\mathfrak{S}_n \otimes_{R\mathfrak{S}_m} \mathbb{1}) \xrightarrow{\iota_*} \mathrm{Tor}_*^{\mathbf{P}_n}(\mathbb{1}, P_n \otimes_{P_m} \mathbb{1}) \xrightarrow{\pi_*} \mathrm{Tor}_*^{R\mathfrak{S}_n}(\mathbb{1}, R\mathfrak{S}_n \otimes_{R\mathfrak{S}_m} \mathbb{1})$$

Then the main result of this section is the following Theorem, which replaces Shapiro's Lemma in the Quillen style proof of homological stability for groups.

Theorem 5.1. *Let $n \geq m \geq 0$. Suppose that δ is invertible in R , or that $m < n$. Then the maps*

$$\iota_*: \mathrm{Tor}_*^{R\mathfrak{S}_n}(\mathbb{1}, R\mathfrak{S}_n \otimes_{R\mathfrak{S}_m} \mathbb{1}) \longrightarrow \mathrm{Tor}_*^{\mathbf{P}_n}(\mathbb{1}, P_n \otimes_{P_m} \mathbb{1})$$

and

$$\pi_*: \mathrm{Tor}_*^{\mathbf{P}_n}(\mathbb{1}, P_n \otimes_{P_m} \mathbb{1}) \longrightarrow \mathrm{Tor}_*^{R\mathfrak{S}_n}(\mathbb{1}, R\mathfrak{S}_n \otimes_{R\mathfrak{S}_m} \mathbb{1})$$

are mutually inverse isomorphisms.

Theorem A follows immediately from Theorem 5.1 by taking δ invertible and $m = n$, using the identifications $R\mathfrak{S}_n \otimes_{R\mathfrak{S}_m} \mathbb{1} \cong \mathbb{1}$ and $P_n \otimes_{P_m} \mathbb{1} \cong \mathbb{1}$.

The remainder of this section is devoted to proving Theorem 5.1, which follows in exactly the same way as Theorem 4.1 of [BHP21] after some preparatory definitions and lemmas.

Recall from Definition 4.1 that $J_m \subseteq P_n$ denotes the ideal consisting of all diagrams in which, among the nodes on the right labelled by $\{n - m + 1, \dots, n\}$, there is a least one singleton or one pair of nodes in the same block. Observe that P_n is a right $R\mathfrak{S}_m$ -module, via the inclusions $R\mathfrak{S}_m \subseteq P_m \subseteq P_n$, and that this module structure preserves J_m , since right multiplying by a diagram which permutes the nodes $\{n - m + 1, \dots, n\}$ does not change whether there exists a singleton or two nodes in the same block in this set. Therefore we have that P_n / J_m is a right $R\mathfrak{S}_m$ -module.

Lemma 5.2. *For $m \leq n$, P_n / J_m is free when regarded as a right $R\mathfrak{S}_m$ -module.*

Proof. P_n / J_m has basis consisting of the diagrams for which the nodes in $\{n - m + 1, \dots, n\}$ have no singleton, and no two nodes in the same block. This means that each node in $\{n - m + 1, \dots, n\}$ is attached to a distinct block in the diagram. Now, $\mathfrak{S}_m$ acts freely on this basis, since multiplying such a diagram with a permutation

in $\mathfrak{S}_m$ results again in a diagram where the nodes in $\{n-m+1, \dots, n\}$ are attached to distinct blocks. Under this action, the stabilizer of any such diagram is trivial. $\square$

Lemma 5.3. *For $m \leq n$, there is an isomorphism of left P_n -modules*

$$P_n / J_m \otimes_{R\mathfrak{S}_m} \mathbb{1} \cong P_n \otimes_{P_m} \mathbb{1}$$

under which $(b + J_m) \otimes r \in P_n / J_m \otimes_{R\mathfrak{S}_m} \mathbb{1}$ corresponds to $b \otimes r \in P_n \otimes_{P_m} \mathbb{1}$.

Proof. Throughout this proof we regard J_m as an ideal in P_n , and write $J_m \cap P_m$ for the corresponding ideal in P_m .

Let us show that the maps

$$(b + J_m) \otimes r \longmapsto b \otimes r$$

and

$$b \otimes r \longmapsto (b + J_m) \otimes r$$

are well-defined. It then immediately follows that they are inverses and thus isomorphisms.

For the first map, we need to show that $b\sigma \otimes r = b \otimes r$ for $\sigma \in \mathfrak{S}_m$ and that $j \otimes r \in P_n \otimes_{P_m} \mathbb{1}$ is zero if $j \in J_m$. The first equation follows immediately as $\sigma \in \mathfrak{S}_m \subset P_m$ acts as the identity on $\mathbb{1}$. The second condition holds because if $j \in J_m$, then we can write j as a sum of products of the form $b \cdot j'$ where $b \in P_n$ and $j' \in J_m \cap P_m$, and for each such summand we have $b \cdot j' \otimes r = b \otimes j' \cdot r = b \otimes 0 = 0$.

For the second map, we let $b \in P_n$, $b' \in P_m$, and $r \in \mathbb{1}$, and show that $(bb' + J_m) \otimes r = (b + J_m) \otimes (b' \cdot r)$. It is enough to prove this for $b' \in R\mathfrak{S}_m$ and $b' \in J_m \cap P_m$ as $P_m = R\mathfrak{S}_m \oplus (J_m \cap P_m)$. For $b' \in R\mathfrak{S}_m$, we get the equation directly from the definition of the tensor product. For $b' \in J_m \cap P_m$, we note that $bb' \in J_m = P_n \cdot J_m$ and thus $(bb' + J_m) \otimes r$ is zero, as is $(b + J_m) \otimes (b' \cdot r)$ since $b' \cdot r = 0$. $\square$

Now recall from [Theorem 4.2](#) that, under the hypotheses of [Theorem 5.1](#),

$$\mathrm{Tor}_*^{P_n}(\mathbb{1}, P_n / J_m) = \begin{cases} \mathbb{1} & \text{if } * = 0 \\ 0 & \text{if } * > 0 \end{cases}.$$

Proof of Theorem 5.1. The proof of [Theorem 5.1](#) now follows exactly as in [[BHP21](#), Proof of Theorem 4.1], replacing the occurrences of Br_n with P_n , and inputting [Lemma 5.3](#) and [Theorem 4.2](#) as appropriate. $\square$

6. HIGH CONNECTIVITY

We build a complex similar to the one in [[Hep22](#)] and [[BHP21](#)].

Definition 6.1. For n a non-negative integer, we define the chain complex $C_n = (C_n)_*$ of P_n -modules as follows. The degree p part $(C_n)_p$ is non-zero in degrees $-1 \leq p \leq n-1$, where it is given by

$$(C_n)_p = P_n \otimes_{P_{n-(p+1)}} \mathbb{1}.$$

So in degree -1 it follows that $(C_n)_{-1} = P_n \otimes_{P_n} \mathbb{1} \cong \mathbb{1}$. For $0 \leq p \leq n-1$ the degree p differential ∂^p is given by the alternating sum

$$\partial^p = \sum_{i=0}^p (-1)^i d_i^p : (C_n)_p \longrightarrow (C_n)_{p-1}.$$

Where, algebraically, the map d_i^p for $1 \leq i \leq p$ is given by

$$\begin{aligned} d_i^p : P_n \otimes_{P_{n-(p+1)}} \mathbb{1} &\longrightarrow P_n \otimes_{P_{n-p}} \mathbb{1} \\ x \otimes r &\longmapsto (x \cdot S_{n-p+i-1} \dots S_{n-p}) \otimes r \end{aligned}$$

and

$$\begin{aligned} d_0^p : P_n \otimes_{P_{n-(p+1)}} \mathbb{1} &\longrightarrow P_n \otimes_{P_{n-p}} \mathbb{1} \\ x \otimes r &\longmapsto x \otimes r. \end{aligned}$$

In other words, when $i = 0$ the product $S_{n-p+i-1} \dots S_{n-p}$ is taken to be the empty product, *i.e.* the identity element.

In terms of diagrams, elements in degree p can be described as diagrams with an $(n - (p + 1))$ -box at the top right, as in [Proposition 2.4](#) and the paragraph which follows it. If we label the nodes below the $(n - (p + 1))$ -box by $0, \dots, p$ from top to bottom, then d_i^p lifts up node i and plugs it into the box.

We now filter C_n . Note that in [\[BHP21\]](#) we first decomposed C_n based on the number of disjoint blocks on the left, and we could also do that here. However this is not necessary for the proof.

Definition 6.2. We define a filtration

$$F_0 C_n \subseteq F_1 C_n \subseteq \dots \subseteq F_{\lfloor \frac{n}{2} \rfloor} C_n = C_n$$

of C_n as follows. The j th level $F_j C_n$ is generated by diagrams with at most j blocks that have at least 2 positive (right hand) nodes and are not connected to the box. Note that this is indeed a filtration, since the boundary map can only decrease the number of blocks on the right not connected to the box.

We briefly recall the definition of the complex of injective words with separators.

Definition 6.3 (Injective words with separators). Let X be a finite set and let $k \geq 0$. An *injective word on X with k separators* is a word with letters taken from the set $X \sqcup \{|\}$ consisting of X and the *separator* $|$, where each letter from X appears at most once, and where the separator appears exactly k times. When $k = 0$, then these are simply the injective words on X .

Definition 6.4 (The complex of injective words with separators). Let X be a finite set, let $s \geq 0$, and let R be a commutative ring. The *complex of injective words with s separators* is the R -chain complex $W_X^{(s)}$ concentrated in degrees $-1 \leq p \leq |X| - 1$, and defined as follows. In degree p , $(W_X^{(s)})_p$ has basis given by the injective words on X with s separators, and with $(p + 1)$ letters from X . Thus such a word $\mathbf{a} \in (W_X^{(s)})_p$ has length $s + p + 1$. Let $r = s + p$ and $\mathbf{a} = a_0 a_1 \dots a_r$. The boundary operator $\partial^p : (W_X^{(s)})_p \rightarrow (W_X^{(s)})_{p-1}$ is defined by the rule

$$\partial^p(a_0 a_1 \dots a_r) = \sum_{i=0}^r (-1)^i a_0 \dots \widehat{a_i} \dots a_r$$

subject to the condition that if the omitted letter is a separator, then the corresponding term is omitted (or identified with 0). In other words, the boundary is the signed sum of the words obtained by deleting the letters that come from X and *not deleting any separators*, but with signs determined by the position of the deleted letter among all letters *including the separator*.

$$\partial^p(a_0 a_1 \dots a_r) = \sum_{a_i \in X} (-1)^i a_0 \dots \widehat{a_i} \dots a_r$$

We will aim to identify the filtration quotients $F_j C_n / F_{j-1} C_n$ with a sum of shifted copies of the complex of injective words with separators, as in [BHP21]. (Note that in [BHP21] the argument for the Brauer algebras is somewhat simpler, and so the reader may wish to look at the Brauer proof first.)

To complete this identification, we exhibit a one-to-one correspondence between diagrams and tuples of data. This correspondence is complicated, so we start with the simple example of the tuple corresponding to a diagram with no box, and no restriction on the right hand side blocks. Recall that a diagram is a pictorial way of representing a partition of the set $\{-n, \dots, -1, 1, \dots, n\}$.

A diagram D determines, and is determined by, a tuple (L, R, ϕ) consisting of:

- A partition L of $\{-1, \dots, -n\}$.
- A partition R of $\{1, \dots, n\}$.
- A labelling $\phi: R \rightarrow \{\emptyset\} \cup L$ with the property that $\phi(r) = \phi(r')$ only when $r = r'$ or $\phi(r) = \phi(r') = \emptyset$.

The correspondence sends a diagram D to the tuple (L, R, ϕ) for which:

- L is the induced partition on the left-hand nodes $-1, \dots, -n$.
- R is the induced partition on the right-hand nodes $1, \dots, n$.
- ϕ labels a block on the right by the (necessarily unique) block on the left to which it is attached, if any, and labels it by $\emptyset$ otherwise.




FIGURE 7. This figure shows the process of extracting from a diagram D the tuple (L, R, ϕ) . Blocks in L , and the labelings of R are indicated at their lowermost node.

An example is shown in Figure 7. Here, the process of restricting the partition to the left and right sides of the diagram amounts to discarding all the left-to-right

connections. Those left-to-right connections are instead recorded in the labelling ϕ . To see that ϕ satisfies the third property above because, observe that if it did not, then the diagram would have two distinct blocks r, r' on the right attached to the *same* block on the left. That would be a contradiction because then r and r' would in fact themselves be the same.

We now observe that the filtration quotient $F_j C_n / F_{j-1} C_n$ has a basis in degree p consisting of diagrams which have an $(n - (p + 1))$ -box on the right, and exactly j blocks with ≥ 2 nodes on the right that are not connected to the box. Here, the size of the box is determined by the degree as in [Definition 6.1](#), and the condition on the j blocks follows from the definition of the filtration given in [Definition 6.2](#). An example is given in [Figure 8](#). In the next definition, we explain how these basis




FIGURE 8. An example of a diagram D , when $n = 9$, $j = 1$ and $p = 6$.

diagrams determine a tuple of data, analogously to the discussion above. Once this data has been stripped from the diagram, we are left with the desired information of an injective word with separators. In this injective word, the letters encode left-to-right connections for which the block on the right has a single element; and the separators correspond to all other nodes below the box on the right. There are at least $2j$ of these separators, because there are precisely j blocks on the right that have 2 or more right-hand nodes and are not connected to the box. Later, in [Lemma 6.8](#) we show how to conversely start with an injective word with separators and our tuple of data and rebuild the diagram.

Definition 6.5. A diagram in the basis of $(F_j C_n / F_{j-1} C_n)_p$ determines a tuple (P, X, s, Y, f)

consisting of the following data:

- A partition P of $\{-1, \dots, -n\}$.
- A subset X of the blocks of P .
- A number $2j \leq s \leq n - |X|$.
- A partition Y of $\{1, \dots, s\}$, such that $\geq j$ blocks have size ≥ 2 .
- A labelling $f: Y \rightarrow (\{\emptyset\} \cup P \setminus X) \times \{\square, \neg\square\}$ (where the symbols $\square, \neg\square$ represent “box” and “not-box” respectively) satisfying the following properties:
 - singletons have first label $\emptyset$
 - no two blocks in Y can have the same first label in $P \setminus X$
 - exactly j blocks of size ≥ 2 have second label $\neg\square$

– exactly $n - s - |X|$ blocks have second label $\square$.

The diagram D determines the tuple as follows (an example is shown in Figure 9):

- P is the partition of $\{-1, \dots, -n\}$ given by restricting the blocks of D to the negative elements, *i.e.* to the nodes on the left hand side of the diagram.
- X is the set of blocks in P which, when viewed in D , are connected to exactly one thing on the right (this can be a connection to the box, or to a single node).
- The number s is equal to the number of nodes on the right of D not connected to a block in X . These nodes are precisely those which are singletons, or are connected to another element on the right, or to the box. Therefore, every node in one of the j blocks of D that have at least 2 positive (right hand) nodes and are not connected to the box (as in Definition 6.2) is included and so $s \geq 2j$. Also, none of the nodes that are connected to a block in X are included, so $s \leq n - |X|$. It follows that $n - |X| - s$ is the number of blocks connected to the box and to at least one node on the right.
- Y is the partition given by restricting D to the set of s nodes on the right that are not connected to the blocks of X (we re-label these $1, \dots, s$, maintaining the order).
- The first entry of the labelling f , in $\emptyset \cup P \setminus X$, indicates whether the blocks of Y are disconnected from the rest of D (in which case the label is $\emptyset$), or connected to the left hand side (in which case the label is the block in $P \setminus X$ that they are connected to). Singletons in Y cannot be connected to the left because otherwise they would be connected to a block in X on the left. Thus their first label must be $\emptyset$. Two blocks in Y cannot be connected to the same block in $P \setminus X$, so two first labels can only be the same if they are both $\emptyset$.

The second entry of the labelling f is $\square$ if the block in Y is connected to the box in D and $\neg\square$ if it is not. The condition that there are exactly j blocks of size ≥ 2 with second label $\neg\square$ accounts for the diagram being in the filtration quotient $F_j C_n / F_{j-1} C_n$. The condition that there are exactly $n - s - |X|$ blocks with second label $\square$ follows from the above observation that this is the number of blocks connected to the box, containing at least one node on the right.

The remaining data in the diagram determines an injective word with s separators $\mathbf{a}$, of length $p + 1 - s$, on the set X , obtained as follows: If the i th node (from the top) on the right is connected to a block in X , then the i th letter of $\mathbf{a}$ is the corresponding element of X . Otherwise the i th letter of $\mathbf{a}$ is a separator, and there are exactly s of these.

Definition 6.6. By the above discussion, we can define a map

$$\Phi_* : F_j C_n / F_{j-1} C_n \rightarrow \bigoplus_{P, X, s, Y, f} W_X^{(s)}[-s].$$

The direct sum is over all 5-tuples (P, X, s, Y, f) satisfying the properties listed at the start of Definition 6.5. A diagram D in $(F_j C_n / F_{j-1} C_n)_p$ is sent by Φ_p to the injective word with separators $\mathbf{a}$ in the degree p part of the summand $W_X^{(s)}[-s]$ corresponding to (P, X, s, Y, f) , where (P, X, s, Y, f) and $\mathbf{a}$ are obtained as in Definition 6.5.




FIGURE 9. This figure shows the process of extracting from the diagram D in Figure 8 the tuple (P, X, s, Y, f) and injective word $\mathbf{a}$ described in Definition 6.5. Blocks in P , and the labeling f of Y are indicated at their lowermost node.

We now prove that Φ_* is a chain map and isomorphism. This will allow us to leverage the high connectivity of the complex of injective words with separators ([BHP21, Proposition 5.14]) to a high connectivity result for C_n , via the filtration.

Lemma 6.7. Φ_* is a chain map.

Proof. First, we claim that the 5-tuple (P, X, s, Y, f) associated, via Φ_* , to a basis diagram D in $(F_j C_n / F_{j-1} C_n)_p$ is preserved in all diagrams appearing in the boundary of D . Recall from Definition 6.1 that the boundary map ∂^p sends a diagram to the alternating sum of the diagrams obtained as follows: work through the nodes on the right of the diagram, and in each case move the node into the box. This clearly does not change the left-hand end of the diagram, and therefore all of the diagrams in the boundary have the same X and P associated to them. If the node that is moved into the box is a singleton, or was part of a block that was connected to the box, then these nodes are included in the count for s , but after moving it into the box, the resulting diagram either has a singleton in the box or has a loop at the box, and therefore again vanishes. The other nodes counting towards s are those that are part of a block with ≥ 2 elements from the right, and are not connected to the box. There are exactly j such blocks, and so moving any of their nodes into the box gives zero in the quotient $(F_j C_n / F_{j-1} C_n)_p$. Therefore the only nodes we can move into the box without getting zero, are those that are not counted by s , i.e. s remains constant under the boundary map. It follows that Y and f remain constant, since Y partitions these s nodes and f labels them.

The above paragraph demonstrates that $F_j C_n / F_{j-1} C_n$ splits as a direct sum indexed by the 5-tuples (P, X, s, Y, f) . It now suffices to show that the assignment that sends a diagram with fixed (P, X, s, Y, f) to the corresponding injective word with separators $\mathbf{a}$ respects the boundary map. But this is clear: moving a node joined to a block in X into the box corresponds exactly to deleting one of the non-separator letters from $\mathbf{a}$. $\square$

Lemma 6.8. Φ_* is an isomorphism.

Proof. We will prove that Φ_* is an isomorphism by showing that it is obtained from a bijection between the basis of $(F_j C_n / F_{j-1} C_n)$, which is given by diagrams, and the basis of $\bigoplus_{X,P,s,Y,f} W_X^{(s)}[-s]$, which is given by injective words with separators. To do this, we will explain how to (re)build a diagram in $(F_j C_n / F_{j-1} C_n)$ from a tuple (P, X, s, Y, f) and an injective word with separators $\mathbf{a}$.

We work in degree $s + k - 1$ in the summand $W_X^{(s)}[-s]$ associated to a 5-tuple (P, X, s, Y, f) . We therefore take an injective word $\mathbf{a}$ of length k with s separators, and we will build a diagram in $(F_j C_n / F_{j-1} C_n)_{s+k-1}$. We begin with an empty diagram with $s + k$ nodes on the right hand side, and a box of size $n - s - k$; this is possible since $s + k \leq s + |X| \leq n$, where the latter inequality is one of the conditions imposed on the 5-tuple. Next, we build all the blocks on the left using P , and draw half-edges from the blocks in X to the right (don't connect these edges to anything yet). We place the injective word with separators vertically against the $s + k$ nodes on the right hand side, and the word indicates connections from k of the nodes to half-edges from X . We connect the remaining half edges from X to the box. The separators indicate the positioning of the s nodes $\{1, \dots, s\}$ which are then partitioned by Y , and labelled by f . The first labels of Y indicate which blocks are connected to blocks on the left hand side in $P \setminus X$. Finally, if the second label of a block in Y is $\square$ we connect the block to the box. Note that $|X| - k$ blocks of X are connected to the box, and $n - s - |X|$ blocks of Y are connected to the box, the latter property being another of our conditions on the 5-tuple. This means that exactly $n - s - k$ distinct blocks are connected to the box, and since this is the size of the box the diagram is non-zero in $(C_n)_{s+k}$. The diagram lies in $F_j C_n / F_{j-1} C_n$ since exactly j blocks in Y of size ≥ 2 have second label $\neg\square$ and are therefore not joined to the box, again by our conditions on the 5-tuple.

The last paragraph shows how to obtain, from a tuple (P, X, s, Y, f) and an injective word $\mathbf{a} \in W_X^{(s)}[-s]_{s+k-1}$, a diagram in the basis of $(F_j C_n / F_{j-1} C_n)_{s+k-1}$. It is now immediate to verify that this is inverse to the effect of Φ_* on bases, and this completes the proof. $\square$

Proposition 6.9. *For all $0 \leq j \leq \lfloor \frac{n}{2} \rfloor$, the filtration quotients $F_j C_n / F_{j-1} C_n$ satisfy $H_i(F_j C_n / F_{j-1} C_n) = 0$ for $i \leq \frac{n-3}{2}$.*

Proof. We first consider the case $n = 0$, where the only possibility is that $j = 0$ so that $F_0 C_0 = C_0$. The claim is then that $H_i(C_0) = 0$ for $i \leq -3/2$, but since C_0 consists of a single copy of $\mathbb{1}$ in degree -1 , this is immediate.

We now consider the case $n > 0$. Using [Lemma 6.8](#) this is equivalent to the homology of $W_X^{(s)}[-s]$ vanishing in the desired range, for each 5-tuple (P, X, s, Y, f) satisfying the conditions of [Definition 6.5](#). By [\[BHP21, Proposition 5.14.\]](#), $H_i(W_X^{(s)}) = 0$ for $i \leq |X| - 2$, so that $H_i(W_X^{(s)}[-s]) = 0$ for $i \leq |X| + s - 2$. It will therefore suffice to show that $\lfloor \frac{n-3}{2} \rfloor \leq |X| + s - 2$, or equivalently:

$$\begin{cases} n \leq 2|X| + 2s & \text{if } n \text{ even} \\ n \leq 2|X| + 2s - 1 & \text{if } n \text{ odd.} \end{cases}$$

Let us first prove that we always have $n \leq 2|X| + 2s$. Our conditions on the 5-tuple (P, X, s, Y, f) mean that $n - s - |X|$ is the number of blocks of Y with second

f -label $\square$, so that in particular $n - s - |X| \leq |Y|$. And since Y is a partition of $\{1, \dots, s\}$ we have $|Y| \leq s$. Combining the last two inequalities and rearranging gives us $n \leq |X| + 2s$. Because $|X| \geq 0$, we therefore have $n \leq 2|X| + 2s$. In particular, this proves the proposition if n is even. If n is odd, it certainly cannot be equal to $2|X| + 2s$ which is even. Therefore it can be at most one smaller: $n \leq 2|X| + 2s - 1$. $\square$

Theorem 6.10. $H_i(C_n) = 0$ for $i \leq \frac{n-3}{2}$.

Proof. By [Proposition 6.9](#), the homology of the filtration quotient $F_j C_n / F_{j-1} C_n$ vanishes in degrees $i \leq \frac{n-3}{2}$ for all j . The same then holds for C_n itself by considering the long exact sequences associated to the short exact sequences $0 \rightarrow F_{j-1} C_n \rightarrow F_j C_n \rightarrow F_j C_n / F_{j-1} C_n \rightarrow 0$. $\square$

7. PROOF OF [THEOREM B](#)

The proof of [Theorem B](#) directly mirrors the proof of [[BHP21](#), Theorem B], with the following substitutions:

- All instances of the Brauer algebra should be replaced with the partition algebra.
- The maps ι and π of [[BHP21](#)] should be replaced by the maps of the same name in the current paper. Similarly for the complex C_* .
- Theorem 5.4 of [[BHP21](#)] should be replaced with [Theorem 6.10](#).
- Theorem 4.1 of [[BHP21](#)] should be replaced with [Theorem 5.1](#).

We note that in the second paragraph of the proof of [[BHP21](#), Theorem 6.3], there is an error, and the words ‘odd’ and ‘even’ should be transposed.

REFERENCES

- [BDM22] Chris Bowman, Stephen Doty, and Stuart Martin. Integral Schur-Weyl duality for partition algebras. *Algebr. Comb.*, 5(2):371–399, 2022. [2](#)
- [BH20] Rachael Boyd and Richard Hepworth. Homological stability for Temperley-Lieb algebras, 2020. To appear in *Geometry & Topology*. [arXiv:2006.04256](#). [1](#), [3](#), [6](#), [7](#)
- [BHP21] Rachael Boyd, Richard Hepworth, and Peter Patzt. The homology of the Brauer algebras. *Selecta Math. (N.S.)*, 27(5):Paper No. 85, 31, 2021. [1](#), [3](#), [6](#), [7](#), [15](#), [16](#), [17](#), [18](#), [21](#), [22](#), [23](#)
- [Boy22] Guy Boyde. Idempotents and homology of diagram algebras, 2022. [arXiv:2212.01826](#). [1](#)
- [BP72] Michael Barratt and Stewart Priddy. On the homology of non-connected monoids and their associated groups. *Comment. Math. Helv.*, 47:1–14, 1972. [3](#)
- [FM94] Eric M. Friedlander and Barry Mazur. Filtrations on the homology of algebraic varieties. *Mem. Amer. Math. Soc.*, 110(529):x+110, 1994. With an appendix by Daniel Quillen. [3](#)
- [Gal11] Søren Galatius. Stable homology of automorphism groups of free groups. *Ann. of Math. (2)*, 173(2):705–768, 2011. [3](#)
- [Hep22] Richard Hepworth. Homological stability for Iwahori-Hecke algebras. *J. Topol.*, 15(4):2174–2215, 2022. [1](#), [16](#)
- [HR05] Tom Halverson and Arun Ram. Partition algebras. *European J. Combin.*, 26(6):869–921, 2005. [1](#)
- [Jon94] Vaughan F. R. Jones. The Potts model and the symmetric group. In *Subfactors (Kyuzeso, 1993)*, pages 259–267. World Sci. Publ., River Edge, NJ, 1994. [1](#), [4](#)
- [Mar94] Paul Martin. Temperley-Lieb algebras for nonplanar statistical mechanics—the partition algebra construction. *J. Knot Theory Ramifications*, 3(1):51–82, 1994. [1](#), [4](#)

- [Mar96] Paul Martin. The structure of the partition algebras. *J. Algebra*, 183(2):319–358, 1996. [4](#)
- [Mos22] Isaac Moselle. Homological stability for Iwahori-Hecke algebras of type B , 2022. [arXiv:2211.06230. 1](#)
- [Nak60] Minoru Nakaoka. Decomposition theorem for homology groups of symmetric groups. *Ann. of Math. (2)*, 71:16–42, 1960. [3](#)
- [Pat20] Peter Patzt. Representation stability for diagram algebras, 2020. [arXiv:2009.06346. 5](#)
- [Sro22] Robin Sroka. The homology of a Temperley-Lieb algebra on an odd number of strands, 2022. To appear in *Algebraic & Geometric Topology. 1*
- [Wei94] Charles A. Weibel. *An introduction to homological algebra*, volume 38 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1994. [6](#)

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF GLASGOW

Email address: `rachael.boyd@glasgow.ac.uk`

INSTITUTE OF MATHEMATICS, UNIVERSITY OF ABERDEEN

Email address: `r.hepworth@abdn.ac.uk`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA

Email address: `ppatz@ou.edu`