# Neutrino cooled disk in post-merger system studied via numerical GR MHD simulation with a composition-dependent equation of state \*

Agnieszka Janiuk

Center for Theoretical Physics of the Polish Academy of Sciences, Al. Lotników 32/46, 02-668 Warsaw, Poland

The code HARM\_COOL, a conservative scheme for relativistic magnetohydrodynamics, is being developed in our group and works with a tabulated equation of state of dense matter. This EOS can be chosen and used during dynamical simulation, instead of the simple ideal gas one. In this case, the inversion scheme between the conserved and primitive variables is not a trivial task. In principle, the code needs to solve numerically five coupled non-linear equations at every time-step. The 5-D recovery schemes were originally implemented in HARM and worked accurately for a simple polytropic EOS which has an analytic form. Our current simulations support the composition-dependent EOS, formulated in terms of rest-mass density, temperature and electron fraction. In this proceeding, I discuss and compare several recovery schemes that have been included in our code. I also present and discuss their convergence tests. Finally, I show set of preliminary results of a numerical simulation, addressed to the post-merger system formed after the binary neutron stars (BNS) coalescence.

# 1. Introduction

The first detection of a gravitational wave signal accompanied by an electromagnetic counterpart was announced on August 17, 2018, by the LIGO-Virgo team [1]. The signal originated from the coalescence of two neutron stars of masses in the range of 1.17-1.60  $M_{\odot}$  and the total mass of the system of 2.74  $M_{\odot}$ . It was followed by a short weak gamma ray burst, observed 1.7 seconds after the GW signal. Hence, the theoretical prediction that this class of short gamma ray bursts originate from compact binary mergers has been proven by the detection of the source GW 170817.

<sup>\*</sup> Presented at the 8<sup>th</sup> Conference of the Polish Society on Relativity, Warsaw, Poland, 19-23 September, 2022.

It was also suggested before that the radioactivities from dynamical ejecta after the first neutron star had been disrupted, can power an electromagnetic signal [8]. Subsequent accretion onto a newly formed black hole can provide bluer emission, if it is not absorbed by precedent ejecta [13, 7]. In this case, a day-timescale emission comes at optical wavelengths from lanthanide-free components of the ejecta, and is followed by a weeklong emission with a spectral peak in the near-infrared (NIR). This two component model fits well with observations of the kilonova, detected in coincidence with the source GRB-GW 170817.

In our studies, we investigate the very last stage of the system, namely the post-merger black hole accretion disk. Due to its high density, the accretion disk in post-merger system is opaque to photons. Neutrinos are produced via  $\beta$ -reactions, electron-positron pair annihilation, and plasmon decay and provide cooling mechanism. The plasma is composed of free nucleons, pairs, and Helium. In the outer regions, heavier nuclei can be also synthesized, under conditions of nuclear statistical equilibrium (NSE, see [5] for details). The kilonova signal is produced in the equatorial wind outflows, launched from the disk via magnetic instabilities. These ejecta are dominated by thermal energy of the dense plasma, and are accelerated to mildly relativistic velocities (of about 0.2-0.3 c, [6]). The matter is highly neutronized there, so due to the r-process nucleosynthesis, copious amounts of unstable heavy isotopes are formed in these winds, and power the Infrared/Optical emission through their radioactive decay.

### 2. Numerical modeling

Our study of the winds launched from the accretion disk, is done by evolving the general relativistic magnetohydrodynamic (GR MHD) equations in time. We use the HARM (High Accuracy Relativistic Magnetohydrodynamics) code [4] which is a conservative and shock capturing scheme. The numerical scheme advances the conserved quantities from one time step to the next by solving a set of non-linear hyperbolic equations for continuity, energy-momentum conservation and magnetic induction. In the GR MHD scheme they read:

$$\nabla_{\mu}(\rho u^{\mu}) = 0, \nabla_{\mu}(T^{\mu\nu}) = 0, \nabla_{\mu}(u^{\nu}b^{\mu} - u^{\mu}b^{\nu}) = 0$$
(1)

where

$$T^{\mu\nu} = T^{\mu\nu}_{\rm gas} + T^{\mu\nu}_{\rm EM} \tag{2}$$

is contributed by

$$T_{\rm gas}^{\mu\nu} = \rho h u^{\mu} u^{\nu} + p g^{\mu\nu} = (\rho + u + p) u^{\mu} u^{\nu} + p g^{\mu\nu}, \qquad (3)$$

Potor8'Ajaniuk'final printed on September 21, 2023

$$T_{\rm EM}^{\mu\nu} = b^2 u^{\mu} u^{\nu} + \frac{1}{2} b^2 g^{\mu\nu} - b^{\mu} b^{\nu}, \quad b^{\mu} = u_{\nu}^* F^{\mu\nu}.$$
 (4)

In the stress-energy tensor composed of the gas and electromagnetic terms,  $u^{\mu}$  is the four-velocity of the gas, u is the internal energy,  $\rho$  is the density, p is the pressure, and  $b^{\mu}$  is the magnetic four vector. F is the Faraday tensor and in a force-free approximation, we have  $E_{\nu} = u^{\nu}F^{\mu\nu} = 0$ . The unit convention is adopted such that G = c = M = 1.

Our initial conditions mimic the configuration after the transient (hypermassive-neutron star, HMNS) object has collapsed to a black hole, and a pressure equilibrium torus [3] has been formed. In this solution, the angular momentum along the radius of the disk is constant.

We parameterize our models with black hole Kerr parameter, a, and the inner radius and radius of pressure maximum,  $r_{in}$ , and  $r_{max}$ , respectively. The current simulations are run in an axisymmetric setup, i.e. 2D, with a resolution 256x256 points in the r, and  $\theta$  directions. The numerical code works in Kerr-Schild coordinates, which enables the matter to smoothly accrete under the horizon.

The torus is embedded in an initially poloidal magnetic field, prescribed with the vector potential of  $(0, 0, A_{\varphi})$ , with  $A_{\varphi} = (\frac{\bar{\rho}}{\rho_{\text{max}}} - \rho_0)$  where we use offset of  $\rho_0 = 0.2$ .

We parameterize the field strength with plasma  $\beta$  defined as the ratio of the gas-to-magnetic pressure,  $\beta = p_{\text{gas}}/p_{\text{mag}}$ . Here  $p_{\text{gas}} = (\gamma - 1)u_{\text{max}}$  and  $p_{\text{mag}} = b_{\text{max}}^2/2$ , where  $u_{\text{max}}$  is the internal energy of the gas at the radius of maximum pressure.

# 3. Chemical composition and structure of the disk

On eof significant challenges for numerical simulation is neutrino treatment. Neutrinos carry away energy and lepton number, so they alter electron fraction and composition of ejected material. Dynamical simulations must consider the realistic equation of state (EOS) and impact of neutrinos in the optically thin and thick regions. Lepton number conservation is expressed as follows:

$$\nabla_{\mu}(n_e u^{\mu}) = \mathcal{R}/m_b; \quad m_b = \rho/n_b; \quad Y_e = \frac{n_e}{n_b} = \frac{m_b n_e}{\rho} \tag{5}$$

where  $\mathcal{R}$  is the net rate of neutrino and antineutrino volume number densities in the fluid frame, and  $Y_e$  is the electron fraction.

Since the baryons dominate the rest-mass density, the baryon number conservation equation turns to the regular continuity equation. In the energy-momentum conservation equation, we must introduce an additional source term due to heating and cooling by neutrinos:

$$\nabla_{\mu}T^{\mu}_{\nu} = \mathcal{Q}u_{\nu},\tag{6}$$

where Q is the energy change per unit volume due to neutrino emission.

### 3.1. Conserved and primitive variables

Conservation equations solved by the GR MHD code can be expressed in a flux conservative form [14], and explicit choice of the conserved variables (which are analytic functions of primitive ones) is to some extent arbitrary. The recovery schemes for the primitive variables need numerical root finding, and can be broadly devided into two categories. In a 2D scheme, there are two independent variables, e.g,  $v^2$  and specific enthalpy W. Temperature is obtained from the EOS tables, solving  $h = h(\rho, T, Y_e)$ , and pressure and temperature are also obtained by Newton-Raphson method for W and  $v^2$ . Alternatively, the system of GR MHD equations is reduced to 3 equations, which have three unknowns. The chosen independent variables can be:  $\gamma$ , T, and  $W = h\rho\gamma^2$ . Pressure is interpolated from EOS tables, as  $P(\rho, T, Y_e)$ .

# 3.2. Recovery transformation

In the 2D recovery scheme of [9] the dimensionality of the recovery problem is reduced by making use of certain scalar quantities that can be computed from the conservatives. To avoid numerical pathologies o the  $1D_W$ scheme near the roots, this 2D scheme solves simultaneously the set of two equations:

$$f_1: \quad \tilde{Q}^2 = v^2 (\mathcal{B}^2 + W)^2 - \frac{(Q_\mu \mathcal{B}^\mu)^2 (\mathcal{B}^2 + 2W)}{W^2}$$
$$f_2: \quad Q_\mu n^\mu = -\frac{\mathcal{B}^2}{2} (1 + v^2) + \frac{(Q_\mu \mathcal{B}^\mu)^2}{2W^2} - W + p(u, \rho)$$

The independent variables used in this scheme are defined as:  $Q_{\mu} = -n_{\mu}T^{\nu}_{\mu} = \alpha T^{t}_{\mu}$ ; where  $\tilde{Q}^{\nu} = j^{\nu}_{\mu}Q^{\mu}$  is energy-momentum density in the normal obsever frame, and  $D = -\rho n_{\mu}u^{\mu} = \alpha \rho u^{t} = \gamma \rho$ ; is mass density in the observer's frame, and  $\mathcal{B}^{i} = \alpha B^{i} = \alpha^{*}F^{it}$  is magnetic 3-vector. Here,  $w = \rho + u + p$  with  $W = w\gamma^{2}$  is enthalpy. To solve this set of equations by means of Newton-Raphson method, the Jacobian matrix with  $\frac{\partial f_{1}}{\partial (v^{2})}$ ,  $\frac{\partial f_{2}}{\partial (v^{2})}$ ,  $\frac{\partial f_{1}}{\partial W}$ , and  $\frac{\partial f_{2}}{\partial W}$  is needed. Note that this scheme does not require an analytic EOS, and derivatives of pressure wtr. to  $\rho$ ,  $v^{2}$ , and u may be computed from tables using finite difference method.

In the 3D recovery scheme of [14], the system is extended to solve 3 equations, on W, z, and T, by adding a constraint on the internal energy given by EOS tables:

$$f_1: [\tau + D - z - B^2 + \frac{B^i S_i}{2z^2} + \rho]W^2 - \frac{B^2}{2} = 0$$

$$f_2: [(z+B^2)^2 - S^2 - \frac{2z+B^2}{z^2}(B^iS_i)^2]W^2 - (z+B^2)^2 = 0$$

$$f_3: \quad \epsilon - \epsilon(\rho, T, Y_e) = 0$$

. Here the temperature is employed directly as an unknown through  $\epsilon(W, zT) = h - 1 - \frac{P}{\rho} = \frac{z - DW - \rho W^2}{DW}$  and does not require inversion of the EOS. Notice here different notation:  $S_i$  is energy-momentum density, W is Lorentz factor, z is enthalpy, and  $\tau = -(n_{\mu}n_{\nu}T^{\mu\nu} + D)$ .

In method proposed in [12] the scheme is solving 1D equation, for the rescaled variable,  $\chi = \frac{\rho h \gamma^2}{\rho \gamma}$ . Other quantities are also rescaled, accordingly, to give Lorentz factor, and we give the brackets for  $\chi$ :  $2 - 2\frac{Q_{\mu}n^{\mu}+D}{D} - \frac{B^2}{D} < \chi < 1 - \frac{Q_{\mu}n^{\mu}+D}{D} - \frac{B^2}{D}$ . The equation

$$f(\chi) = \chi - \tilde{\gamma}(1 + \tilde{\epsilon} + \frac{\tilde{P}}{\tilde{\rho}}) = 0$$

is solved, with  $\tilde{P} = P(\tilde{\rho}, \tilde{\epsilon}, Y_e)$  found in tables.

The last method has proven to be most robust, for our wide parameter space. It works with smaller errors, while the performance spped is also slower. We performed the convergence tests to explore which recovery transformation works best for the wide parameter space. The parameters used for testing routines were:  $Y_e = 0.1$ ,  $\gamma = 2$ ,  $p_{\text{gas}}/p_{\text{mag}} = 10^5$ . We derived the conserved variables in Kerr metric, and then computed primitives perturbed by a factor of 1.05. The variables recovered through the 2D scheme were compared to the unperturbed, to calculate the total error, summed over all primitives  $\text{Err} = \Sigma_{k=0,NPR}(P_k - \bar{P}_k)^2$ . Figure 1 shows the results for all 3 methods probed.

# 3.3. Neutrino transport

We employ the neutrino leakage scheme that computes a gray optical depth estimate along radial rays for electron neutrinos, electron antineutrinos, and heavy-lepton neutrinos (nux), and then computes local energy and lepton number loss terms. The source code of the scheme has been



Fig. 1. Convergence test results for the 3D method (2 versions, with (1): we compute specific internal energy from state vector x and conservatives as in Eq. (25) in [2], and solve  $f_3$  or (2): we compute pressure from state vector x and conservatives and solve  $f_3$ ), shown in the top panels, the 2D method (bottom panel, left) and Palenzuela method (bottom panel, right).

publicly available and downloaded from https://stellarcollapse.org. Details are described e.g. in [11]. The initial tests were done within an optically thin regime for neutrinos.

$$\tau(r,\theta,\phi) = \int_{r}^{R} \sqrt{\gamma_{rr}} \bar{\kappa}_{\nu_{i}} dr' < 2/3.$$
(7)

# 4. Results

The structure of the flow in the evolved state is shown in Fig. 2. We present polar slices, taken at the evolved time of the simulation (t = 44 ms, which is equivalent to the 3000 gometric time units,  $t_g$ , for black hole mass of 3  $M_{\odot}$ ). Parameters of this model are black hole spin a = 0.6, initial torus radius  $r_{\text{in}} = 4r_g$  and pressure maximum radius of  $r_{\text{max}} = 11.8r_g$ . We also assume initial electron fraction in the torus of  $Y_{\text{e,disk}} = 0.1$ , and in the

#### Potor8'Ajaniuk' final printed on September 21, 2023



Fig. 2. Maps of density and magnetic field contours (left), electron fraction (middle) and neutrino emissivity (right panel). Model assumes optically thin plasma for neutrinos. Inversion scheme used the Palenzuela method. The model parameters are disk mass = 0.13  $M_{\odot}$ , gas-to-magnetic pressure ratio  $\beta = 50$  and black hole spin a = 0.6. The color maps are in logarithmic scale and are taken at t = 0.044 s.

atmosphere it is  $Y_{\rm e,atm} = 0.45$ . The specific entropy per baryon in the torus is assumed  $S = 10k_{\rm B}$ . This, with the enthalpy value resulting from FM torus sulution, gives the physical density scaling: units in the code are normalized such that  $\rho_{\rm max} = 1$ . The torus mass in cgs units is about  $0.03M_{\odot}$ .

As shown in the figure, the magnetic turbulence developed in the torus, and helped launching winds from its surface. The neutronized material is redistributed, and electron fraction in the winds became larger han in the torus, about  $Y_e = 0.3$ . The neutrino emissivity in the winds is couple of orders of magnitude less than in the torus. As can be seen in the 2D map, the accreting torus has a very high neutrino luminosity, but here neutrinos are partially trapped. A moderately high neutrino luminosity can be observed from the wind, where the neutrino-antineutrino pairs can contribute to the heating of the plasma.

Neutronisation of the plasma in the tours is still significant, and electron fraction values are below  $Y_e = 0.1 - 0.3$  in some densest regions (cf. [10]). The neutrinos which are created in weak interactions, and the electronpositron pair annihilation, nucleon bremsstrahlung and plasmon decay, provide an efficient cooling mechanism. In the currently developed new numerical scheme, we substituted a simplified description of the neutrino cooling rate given by the two-stream approximation previously used by [6] with a more advanced neutrino leakage scheme. The neutrino emissivity distribution image taken at an evolved state of the torus is presented in Fig. 2 for the optically thin case. For optically thick simulation (which assumed different initial parameters as for the specific entropy), the quantitative difference can be noticed. The neutrino and antineutrino luminosities, as a function of time, are shown in Figure 3.

The velocity of the wind outflows,  $v \sim (0.11 - 0.23) c$ , and mass loss via unbound outflows of 2- 17% of the initial disk mass were determined



Fig. 3. The electron neutrino and antineutrino (Lnu1 and Lnu2, respectively; Lnu3 denotes luminosity in other flavors) luminosity as a function of time, for our two models, optically thin and optically thick, calculated with the neutrino-leakage scheme.

previously by [6]. We showed that the details are sensitive to engine parameters: BH spin and magnetisation of the disk, namely the more magnetized disks produce faster outflows. We also found that the accretion disk ejecta produce heavy elements up to mass number  $A \sim 200$ , including platinium and gold isotopes (see details in [10]).

### 5. Conclusions

Numerical GR MHD simulations have been widely used to model postmerger systems and engines of short gamma ray bursts. We implemented a new scheme in our HARM-COOL code to the case study of post-merger system and kilonova source. The calculations are complex and need to cover physics of dense nuclear matter. Their performance is sensitive to the chosen recovery schemes. We tested several of them and chose the most robust for a lare parameter space in densities, temperatures, and electron fraction values, to work with the 3-parameter EOS implemented in hydrodynamical simulation. Proper source terms have been added in the system of equations for the neutrino losses, coupled with composition changes. We also use an advanced neutrino-leakage scheme and calculate emissivities of neutrino and antineutrino as a function of time.

The unbound outflows, i.e. winds, are powered by both neutrinos and magnetically driven acceleration. Therefore, winds may be more dense and powerful, if neutrino heating supports the magnetically-driven wind.

### Acknowledgment

This work was supported by grant 2019/35/B/ST9/04000 from the Polish National Science Center. We used computational resources of the ICM of the Warsaw University, and the PL-Grid, under grant *plggrb5*.

# REFERENCES

- B. P. Abbott and others. Gravitational Waves and Gamma-Rays from a Binary Neutron Star Merger: GW170817 and GRB 170817A. *ApJL*, 848(2):L13, 2017.
- [2] P. Cerdá-Durán, J. A. Font, L. Antón, and E. Müller. A new general relativistic magnetohydrodynamics code for dynamical spacetimes. A&A, 492(3):937–953, December 2008.
- [3] L. G. Fishbone and V. Moncrief. Relativistic fluid disks in orbit around Kerr black holes. ApJ, 207:962–976, 1976.
- [4] C. F. Gammie, J. C. McKinney, and G. Tóth. HARM: A Numerical Scheme for General Relativistic Magnetohydrodynamics. ApJ, 589(1):444–457, 2003.
- [5] A. Janiuk. Nucleosynthesis of elements in gamma-ray burst engines. A&A, 568:A105, 2014.
- [6] A. Janiuk. The r-process Nucleosynthesis in the Outflows from Short GRB Accretion Disks. ApJ, 882(2):163, 2019.
- [7] O. Korobkin, S. Rosswog, A. Arcones, and C. Winteler. On the astrophysical robustness of the neutron star merger r-process. *MNRAS*, 426(3):1940–1949, 2012.
- [8] L.-X. Li and B. Paczyński. Transient Events from Neutron Star Mergers. ApJL, 507(1):L59–L62, 1998.
- [9] S. C. Noble, C. F. Gammie, J. C. McKinney, and L. Del Zanna. Primitive Variable Solvers for Conservative General Relativistic Magnetohydrodynamics. ApJ, 641(1):626–637, 2006.
- [10] F. H. Nouri, A. Janiuk, and M. Przerwa. Studying Postmerger Outflows from Magnetized-neutrino-cooled Accretion Disks. ApJ, 944(2):220, 2023.
- [11] C. D. Ott, E. Abdikamalov, E. O'Connor, C. Reisswig, R. Haas, P. Kalmus, S. Drasco, A. Burrows, and E. Schnetter. Correlated gravitational wave and neutrino signals from general-relativistic rapidly rotating iron core collapse. *Phys.Rev.D*, 86(2):024026, 2012.

- [12] C. Palenzuela, S. L. Liebling, D. Neilsen, L. Lehner, O. L. Caballero, E. O'Connor, and M. Anderson. Effects of the microphysical equation of state in the mergers of magnetized neutron stars with neutrino cooling. *Phys.Rev.D*, 92(4):044045, 2015.
- [13] D. M. Siegel and B. D. Metzger. Three-Dimensional General-Relativistic Magnetohydrodynamic Simulations of Remnant Accretion Disks from Neutron Star Mergers: Outflows and r -Process Nucleosynthesis. *PRL*, 119(23):231102, 2017.
- [14] D. M. Siegel, P. Mösta, D. Desai, and S. Wu. Recovery Schemes for Primitive Variables in General-relativistic Magnetohydrodynamics. ApJ, 859(1):71, 2018.