

UNIVERSAL EQUATIONS FOR HIGHER GENUS GROMOV–WITTEN INVARIANTS FROM HODGE INTEGRALS

FELIX JANDA AND XIN WANG

ABSTRACT. We establish new universal equations for higher genus Gromov–Witten invariants of target manifolds, by studying both the Chern character and Chern classes of the Hodge bundle on the moduli space of curves. As a consequence, we find new push-forward relations on the moduli space of stable curves.

CONTENTS

1. Introduction	1
1.1. New universal equations	2
1.2. A new push-forward relation	3
1.3. Universal relations from Hodge class	4
1.4. Plan of the paper	5
2. Chern characters of the Hodge bundle	5
2.1. Review of Faber and Pandharipande’s formula	5
2.2. Pixton’s formula for double ramification cycles and the λ_g class	6
2.3. Universal equations from $\mathrm{ch}_{2g-1}(\mathbb{E})$	8
2.4. Proof of Theorem 1	10
2.5. Reformulation of Theorem 1	12
2.6. Proof of Corollary 2	13
3. Push-forward relations on $\overline{\mathcal{M}}_g$ for $(g \geq 2)$	14
3.1. Push-forward relations	14
3.2. Proof of Theorem 5	14
3.3. Connection to degree- g topological recursion relations	15
4. Chern class of Hodge bundle	16
4.1. Linear Hodge integrals	16
4.2. The Faber–Pandharipande relation and Givental quantization	16
4.3. Universal relations from the Chern classes of the Hodge bundle	18
4.4. Connection to Pixton’s 3-spin relations	19
References	21

1. INTRODUCTION

Finding universal partial differential equations for Gromov–Witten invariants of any target manifold is a long-standing problem in Gromov–Witten theory. Such equations capture fundamental properties for Gromov–Witten invariants. Usually, it is very difficult to find

2020 *Mathematics Subject Classification.* Primary 14N35; Secondary 14N10.

Key words and phrases. Gromov–Witten invariant, universal equation, push-forward relation.

explicit equations in all genera. So far, there have been mainly two approaches for finding such universal equations for Gromov–Witten invariants. One approach is to use tautological relations, in particular *topological recursion relations*, on the moduli space of stable curves. For example, the equality between the boundary divisors on $\overline{\mathcal{M}}_{0,4}$ leads to the well-known WDVV equation for genus-0 Gromov–Witten invariants. Another approach is to use tautological relations on the moduli space of stable maps. For example, in [FP00a], by using the Grothedieck–Riemann–Roch formula on the moduli space of stable maps, Faber and Pandharipande obtained a family of explicit universal equations for descendant Gromov–Witten invariants. In 2014, a remarkable formula for double ramification cycles was proposed by Pixton and later proved in [JPPZ17]. In this paper, we use this new ingredient to obtain new universal equations for descendant Gromov–Witten invariants.

1.1. New universal equations. Let X be a smooth projective variety and $\{\phi_\alpha : \alpha = 1, \dots, N\}$ be a basis of its cohomology ring $H^*(X; \mathbb{C})$ with $\phi_1 = \mathbf{1}$ the identity. Recall that the *big phase space* for Gromov–Witten invariants of X is defined to be $\prod_{n=0}^{\infty} H^*(X; \mathbb{C})$ with standard basis $\{\tau_n(\phi_\alpha) : \alpha = 1, \dots, N, n \geq 0\}$. Denote the coordinates on the big phase space with respect to the standard basis by $\{t_n^\alpha\}$. Let F_g be the genus- g generating function, which is a formal power series of $\mathbf{t} = (t_n^\alpha)$ with coefficients being the genus- g Gromov–Witten invariants. Denote $\langle\langle \tau_{n_1}(\phi_{\alpha_1}), \dots, \tau_{n_k}(\phi_{\alpha_k}) \rangle\rangle_g$ for the derivatives of F_g with respect to the variables $t_{n_1}^{\alpha_1}, \dots, t_{n_k}^{\alpha_k}$. Similarly, we use $\langle\langle \tau_{n_1}(\phi_{\alpha_1}), \dots, \tau_{n_k}(\phi_{\alpha_k}); c_{g,k} \rangle\rangle_g$ to denote the correlation function of Gromov–Witten theory twisted with a cohomology class $c_{g,k} \in H^*(\overline{\mathcal{M}}_{g,k}; \mathbb{C})$. For convenience, we identify $\tau_n(\phi_\alpha)$ with the coordinate vector field $\frac{\partial}{\partial t_n^\alpha}$ on $\prod_{n=0}^{\infty} H^*(X; \mathbb{C})$ for $n \geq 0$. If $n < 0$, $\tau_n(\phi_\alpha)$ is understood to be the 0 vector field. We also abbreviate $\tau_0(\phi_\alpha)$ by ϕ_α . Let $\phi^\alpha = \eta^{\alpha\beta} \phi_\beta$ with $(\eta^{\alpha\beta})$ representing the inverse matrix of the Poincaré intersection pairing on $H^*(X; \mathbb{C})$. As a convention, repeated Greek letter indices are summed over their entire range. Recall the following T operator which was studied in [Liu02]

$$T(W) := \tau_+(W) - \langle\langle W \phi^\alpha \rangle\rangle_0 \phi_\alpha \quad (1)$$

for any vector field W on the big phase space, where $\tau_+(W)$ is a linear operator defined by $\tau_+(\tau_n(\phi_\alpha)) = \tau_{n+1}(\phi_\alpha)$.

Let $A = (a_1, \dots, a_n)$ be a vector with integers $\sum_{i=1}^n a_i = 0$. Let $\mathbf{P}_g^A(A) \in H^{2g}(\overline{\mathcal{M}}_{g,n})$ be the cohomology class obtained from Pixton’s double ramification cycle formula [JPPZ17], which is roughly a linear combination of descendant stratum classes.

Our first result is:

Theorem 1. *For any $g \geq 2$, the following equation holds for pure descendant Gromov–Witten invariants of X*

$$\begin{aligned} & - \sum_{n,\alpha} \tilde{t}_n^\alpha \langle\langle \tau_{n+2g-1}(\phi_\alpha) \rangle\rangle_g + \frac{1}{2} \sum_{i=0}^{2g-2} (-1)^i \left\{ \langle\langle \tau_i(\phi_\alpha) \tau_{2g-2-i}(\phi^\alpha) \rangle\rangle_{g-1} + \sum_{h=0}^g \langle\langle \tau_i(\phi_\alpha) \rangle\rangle_h \langle\langle \tau_{2g-2-i}(\phi^\alpha) \rangle\rangle_{g-h} \right\} \\ &= \frac{(-1)^g g}{2^{2g-1}} \sum_{m=1}^{g-1} \frac{1}{m} \sum_{\substack{k_1+\dots+k_m=g \\ k_1,\dots,k_m \geq 1}} \sum_{\substack{g_1+\dots+g_m=g-1 \\ g_1,\dots,g_m \geq 1}} \sum_{\alpha_1,\dots,\alpha_m=1}^N \prod_{i=1}^m \frac{-1}{k_i!} \sum_{l_i=0}^{k_i-1} \binom{k_i-1}{l_i} \langle\langle T^{l_i}(\phi_{\alpha_{i-1}}) T^{k_i-1-l_i}(\phi^{\alpha_i}); \mathbf{P}_{g_i}^{g_i}(0,0) \rangle\rangle_{g_i} \end{aligned} \quad (2)$$

where $\tilde{t}_n^\alpha = t_n^\alpha - \delta_n^1 \delta_\alpha^1$ and $\alpha_0 = \alpha_m$.

Together with [LP11, Theorem 4], Theorem 1 implies the following identity:

Corollary 2. *For any $g \geq 2$, the following equation holds for the pure descendant Gromov-Witten invariants of X*

$$\begin{aligned} & \sum_{i=0}^{2g-2} (-1)^i \langle \langle T^i(\phi_\alpha) T^{2g-2-i}(\phi^\alpha) \rangle \rangle_{g-1} \\ &= \frac{(-1)^g}{2^{2g-2}} \sum_{m=1}^g \frac{1}{m} \sum_{\substack{k_1+\dots+k_m=g \\ k_1, \dots, k_m \geq 1}} \sum_{\substack{g_1+\dots+g_m=g-1 \\ g_1, \dots, g_m \geq 1}} \sum_{\alpha_1, \dots, \alpha_m=1}^N \prod_{i=1}^m \frac{-1}{k_i!} \sum_{l_i=0}^{k_i-1} \binom{k_i-1}{l_i} \langle \langle T^{l_i}(\phi_{\alpha_{i-1}}) T^{k_i-1-l_i}(\phi^{\alpha_i}) ; \mathbf{P}_{g_i}^{g_i}(0, 0) \rangle \rangle_{g_i}. \end{aligned} \quad (3)$$

Proposition 12 in Section 2.5 explains how to view the right hand side in (2) and (3) in terms of pure descendant Gromov-Witten invariants.

Remark 3. In [FP00a], it was pointed out that Faber–Pandharipande’s equations are related to the genus-0 $\tilde{L}_n$ -constraints proposed by Eguchi–Hori–Xiong in [EHX97]. However, the explicit expression of the higher genus $\tilde{L}_n$ -constraints remain unknown. We hope that (2) will give a hint at the explicit form of the higher genus corrections to the $\tilde{L}_n$ -constraints.


1.2. A new push-forward relation. It is very interesting to find nontrivial classes in the kernel of the boundary push-forward map

$$\iota: R^*(\overline{\mathcal{M}}_{g,2}) \rightarrow R^*(\overline{\mathcal{M}}_{g+1})$$

which can also be written in terms of ψ classes and boundary classes. Via the splitting principle in genus $g+1$ Gromov-Witten theory, this yields universal equations in genus g from linear combinations of descendant boundary classes in the kernel of ι_* . Such tautological relations were first constructed by Liu–Pandharipande in $R^{2g+r+1}(\overline{\mathcal{M}}_{g+1})$ for $r \geq 1$ (cf. [LP11, Theorem 2]). It is remarkable that the equation in Corollary 2 can be lifted to cycle level, which gives an extension of Liu–Pandharipande’s relation to the case of $r = 0$.

Definition 4. A stable graph Γ in the sense of [Pan18, §4.2] is a *stable circular graph* if its vertices are connected in a closed chain. We denote the set of circular graphs by $G_{g,n}^\circ$.

Any stable circular graph Γ has the same number of vertices and edges. For example, the following is a list of all stable circular graphs in $G_{4,0}^\circ$:



Theorem 5. *For $g \geq 1$, the following topological recursion relation holds in $R^{2g+1}(\overline{\mathcal{M}}_{g+1})$*

$$\begin{aligned} & \frac{2^{2g-1}}{(-1)^{g+1}(g+1)} \sum_{a+b=2g} (-1)^a \iota_*(\psi_1^a \psi_2^b) \\ &= \sum_{\Gamma \in G_{g,0}^\circ} \frac{1}{|\text{Aut}(\Gamma)|} \sum_{\substack{e \in E(\Gamma) \\ k(e) \geq 1, \forall e \in E(\Gamma)}} (\xi_\Gamma)_* \left(\prod_{e \in E(\Gamma)} \left(\frac{-1}{k(e)} \sum_{l(e)+m(e)=k(e)-1} \frac{\psi_{h(e)}^{l(e)}}{l(e)!} \frac{\psi_{h'(e)}^{m(e)}}{m(e)!} \right) \prod_{v \in V(\Gamma)} \mathbf{P}_{g(v)}^{g(v)}(0, 0) \right) \end{aligned} \quad (4)$$

where $\iota: \overline{\mathcal{M}}_{g,2} \rightarrow \overline{\mathcal{M}}_{g+1}$ and $\xi_\Gamma: \overline{\mathcal{M}}_\Gamma \rightarrow \overline{\mathcal{M}}_{g+1}$ are the canonical gluing maps. $h(e)$ and $h'(e)$ denote the two half edges of e .

Theorem 5 implies that Equation (3) of Corollary 2 holds for any cohomological field theory, as defined by Kontsevich and Manin in [KM97], in particular also for orbifold Gromov-Witten theory and quantum singularity theory (cf. [CR02], [FJR13]).

1.3. Universal relations from Hodge class. It was first mentioned in [FP00a] that a more sophisticated method of obtaining differential equations for pure descendant Gromov-Witten invariants from Faber–Pandharipande’s formula (cf. [FP00a, Proposition 2]) is to construct differential operators that correspond to multiplying by a Chern class of the Hodge bundle $\mathbb{E}$. The Chern classes of Hodge bundle $\mathbb{E}$ certainly vanish in degrees greater than g . One obtains from [FP00a, Proposition 2]) (Equation (37) in this paper) relations in degree greater than g (for each g). It would be interesting to understand these equations and their relation to topological recursion relations and the Virasoro constraints even when X is a point.

Let z be a formal variable. We consider the space $\mathbb{H}_+$ which is the space of polynomials in one variable z with coefficients in $H^*(X; \mathbb{C})$. It has a canonical linear basis $\{z^n \phi_\alpha : \alpha = 1, \dots, N, n \geq 0\}$. Denote the coordinates on $\mathbb{H}_+$ with respect to the canonical basis by $\{q_n^\alpha\}$. Then a general element in $\mathbb{H}_+$ can be written in the form $\mathbf{q}(z) = \sum_{n \geq 0, \alpha} q_n^\alpha \phi_\alpha z^n$. For convenience, we identify $\mathbb{H}_+$ with the big phase space $\prod_{n=0}^\infty H^*(X; \mathbb{C})$ via identification between basis $\{z^n \phi_\alpha\}$ and $\{\tau_n(\phi_\alpha)\}$, and dilaton shift about coordinates $q_n^\alpha = \tilde{t}_n^\alpha$. We will use the following convention

$$\tilde{\mathcal{P}}(\mathbf{q}) = \mathcal{P}(\mathbf{t})|_{q_n^\alpha = \tilde{t}_n^\alpha}$$

for any partition function $\mathcal{P}(\mathbf{t})$. Denote $\langle\langle \tau_{n_1}(\phi_{\alpha_1}), \dots, \tau_{n_k}(\phi_{\alpha_k}) \rangle\rangle_g(\mathbf{q}(z))$ for the derivatives of $\tilde{F}_g(\mathbf{q}) = F_g(\mathbf{t})|_{q_n^\alpha = \tilde{t}_n^\alpha}$ with respect to the variables $q_{n_1}^{\alpha_1}, \dots, q_{n_k}^{\alpha_k}$. Similarly, we use $\langle\langle \tau_{n_1}(\phi_{\alpha_1}), \dots, \tau_{n_k}(\phi_{\alpha_k}); c_{g,k} \rangle\rangle_g(\mathbf{q}(z))$ to denote the correlation function of Gromov–Witten theory twisted with a cohomology class $c_{g,k} \in H^*(\overline{\mathcal{M}}_{g,k}; \mathbb{C})$.

Theorem 6 (Theorem 17). *Let*

$$R(z) = \exp\left(\sum_{i=1}^{\infty} \frac{B_{2i}}{(2i)(2i-1)} u^{2i-1} z^{2i-1}\right).$$

Then, for $g \geq 2$, the following holds for descendant Gromov–Witten invariants

$$[u^i] \sum_{\Gamma \in G_g^{\text{Feyn}}} \frac{1}{|\text{Aut}(\Gamma)|} \text{Cont}_\Gamma = \begin{cases} 0, & i > g \\ \frac{(-1)^g}{2^g} \cdot \langle\langle \ ; \mathbf{P}_g^g(\emptyset) \rangle\rangle_g(\mathbf{q}(z)) & i = g \end{cases}$$

where $[u^i]$ denotes taking the coefficient of u^i and the contribution Cont_Γ of a Feynman graph Γ is defined as a contraction of tensors:

- Each vertex v is assigned a tensor bracket

$$\langle\langle \ \rangle\rangle_{g(v)}(R^{-1}\mathbf{q})$$

with insertion coming from the half edges.

- Each edge e is assigned a bivector with descendants $\frac{\sum_{\alpha, \beta} (\eta^{\alpha\beta} \phi_\alpha \otimes \phi_\beta - \eta^{\alpha\beta} R^{-1}(\psi_{h(e)})(\phi_\alpha) \otimes R^{-1}(\psi_{h'(e)})(\phi_\beta))}{\psi_{h(e)} + \psi_{h'(e)}}$

Proposition 7 (see Proposition 19). *Pixton’s 3-spin relations [Pix12] on the moduli space of stable curves imply the $i > g$ part of Theorem 17.*

1.4. Plan of the paper. This paper is organized as follows. In Section 2, we review Faber and Pandharipande's formula for Hodge classes, Pixton's double ramification cycle formula for the top Hodge class, and prove Theorem 1 and Corollary 2. In Section 3, we give prove the new push-forward relation of Theorem 5. In Section 4, we prove Theorem 6 and Proposition 7.

2. CHERN CHARACTERS OF THE HODGE BUNDLE

2.1. Review of Faber and Pandharipande's formula. Let $\overline{\mathcal{M}}_{g,n}(X, \beta)$ be the moduli space of stable maps $\{f : (C; p_1, \dots, p_n) \rightarrow X\}$, where $(C; p_1, \dots, p_n)$ is a genus- g nodal curve with n marked points and the image $f_*[C] = \beta \in H_2(X; \mathbb{Z})$ is the effective class β . The Hodge bundle $\mathbb{E}$ over $\overline{\mathcal{M}}_{g,n}(X, \beta)$ is the rank g vector bundle with fiber $H^0(C, \omega_C)$ over a moduli point $f : (C, p_1, \dots, p_n) \rightarrow X$, where ω_C is the dualizing sheaf of the domain curve C . In fact, the Hodge bundle over $\overline{\mathcal{M}}_{g,n}(X, \beta)$ is the pullback of the Hodge bundle from $\overline{\mathcal{M}}_{g,n}$ under the stabilization map $\text{St} : \overline{\mathcal{M}}_{g,n}(X, \beta) \rightarrow \overline{\mathcal{M}}_{g,n}$. Let $\text{ch}(\mathbb{E})$ be the Chern character of $\mathbb{E}$ and denote its degree k component by $\text{ch}_k(\mathbb{E})$. It is well-known that the positive even Chern characters $\text{ch}_{2k}(\mathbb{E})$ vanish (cf. [Mum83]). Let $\lambda_i := c_i(\mathbb{E})$ be the i -th Chern class of Hodge bundle.

In [FP00a], Faber and Pandharipande considered the following Hodge integrals over the moduli space of stable maps:

$$\left\langle \prod_{i=1}^n \tau_{k_i}(\phi_{\alpha_i}); \prod_{j=1}^m \text{ch}_{l_j}(\mathbb{E}) \right\rangle_{g, \beta} := \int_{[\overline{\mathcal{M}}_{g,n}(X, \beta)]^{\text{vir}}} \prod_{i=1}^n \psi_i^{k_i} \prod_{i=1}^n \text{ev}_i^* \phi_{\alpha_i} \prod_{j=1}^m \text{ch}_{l_j}(\mathbb{E})$$

where $[\overline{\mathcal{M}}_{g,n}(X, \beta)]^{\text{vir}}$ is the virtual fundamental class of $\overline{\mathcal{M}}_{g,n}(X, \beta)$, the descendant ψ -class ψ_i is the first Chern class of the tautological line bundle over $\overline{\mathcal{M}}_{g,n}(X, \beta)$ defined by the cotangent lines at the i -th marking of the domain curves, and $\text{ev}_i : \overline{\mathcal{M}}_{g,n}(X, \beta) \rightarrow X$ is the evaluation map defined by the image of the i -th marking.

The generating function of genus- g Hodge integrals is defined to be

$$F_g^{\mathbb{E}}(\mathbf{t}, \mathbf{s}) := \sum_{\beta \in H_2(X; \mathbb{Z})} Q^{\beta} \left\langle \exp \left(\sum_{n, \alpha} t_n^{\alpha} \tau_n(\phi_{\alpha}) \right); \exp \left(\sum_{m \geq 1} s_{2m-1} \text{ch}_{2m-1}(\mathbb{E}) \right) \right\rangle_g \quad (5)$$

where Q is the Novikov variable. In particular, after the restriction $\mathbf{s} = 0$, the generating function $F_g(\mathbf{t}) := F_g^{\mathbb{E}}|_{\mathbf{s}=0}$ becomes the usual generating function of genus- g pure descendant Gromov–Witten invariants. For convenience, we identify $\tau_n(\phi_{\alpha})$ with $\frac{\partial}{\partial t_n^{\alpha}}$ and $\text{ch}_n(\mathbb{E})$ with $\frac{\partial}{\partial s_n}$. Moreover we use the convention that $\tau_0(\phi_{\alpha}) = \phi_{\alpha}$ and $\tau_n(\phi_{\alpha}) = 0$ if $n < 0$. Double brackets denote differentiation of $F_g^{\mathbb{E}}(\mathbf{t}, \mathbf{s})$,

$$\left\langle \langle \tau_{k_1}(\phi_{\alpha_1}) \dots \tau_{k_n}(\phi_{\alpha_n}); \prod_{i=1}^m \text{ch}_{2l_i-1}(\mathbb{E}) \rangle \rangle_g^{\mathbb{E}} = \frac{\partial^{n+m}}{\partial t_{k_1}^{\alpha_1} \dots \partial t_{k_n}^{\alpha_n} \partial s_{2l_1-1} \dots \partial s_{2l_m-1}} F_g^{\mathbb{E}}(\mathbf{t}, \mathbf{s})$$

and

$$\left\langle \langle \tau_{k_1}(\phi_{\alpha_1}) \dots \tau_{k_n}(\phi_{\alpha_n}); \prod_{i=1}^m \text{ch}_{2l_i-1}(\mathbb{E}) \rangle \rangle_g = \left\langle \langle \tau_{k_1}(\phi_{\alpha_1}) \dots \tau_{k_n}(\phi_{\alpha_n}); \prod_{i=1}^m \text{ch}_{2l_i-1}(\mathbb{E}) \rangle \rangle_g^{\mathbb{E}}|_{\mathbf{s}=0}.$$

In [FP00a], the following universal equation for descendant Gromov–Witten invariants of any smooth projective variety was proved for any $l > g$:

$$-\tilde{t}_n^\alpha \langle \langle \tau_{n+2l-1}(\phi_\alpha) \rangle \rangle_g + \frac{1}{2} \sum_{i=0}^{2l-2} (-1)^i \left\{ \langle \langle \tau_i(\phi_\alpha) \tau_{2l-2-i}(\phi^\alpha) \rangle \rangle_{g-1} + \sum_{h=0}^g \langle \langle \tau_i(\phi_\alpha) \rangle \rangle_h \langle \langle \tau_{2l-2-i}(\phi^\alpha) \rangle \rangle_{g-h} \right\} = 0 \quad (6)$$

where $\tilde{t}_n^\alpha = t_n^\alpha - \delta_n^1 \delta_\alpha^1$. In fact, the Grothendieck–Riemann–Roch formula on the moduli space of stable maps (cf. [FP00a]) implies

$$\begin{aligned} & \frac{(2l)!}{B_{2l}} \langle \langle \text{ch}_{2l-1}(\mathbb{E}) \rangle \rangle_g^\mathbb{E} \\ &= -\tilde{t}_n^\alpha \langle \langle \tau_{n+2l-1}(\phi_\alpha) \rangle \rangle_g^\mathbb{E} + \frac{1}{2} \sum_{i=0}^{2l-2} (-1)^i \left\{ \langle \langle \tau_i(\phi_\alpha) \tau_{2l-2-i}(\phi^\alpha) \rangle \rangle_{g-1}^\mathbb{E} + \sum_{h=0}^g \langle \langle \tau_i(\phi_\alpha) \rangle \rangle_h^\mathbb{E} \langle \langle \tau_{2l-2-i}(\phi^\alpha) \rangle \rangle_{g-h}^\mathbb{E} \right\} \end{aligned} \quad (7)$$

where B_{2n} are the classical Bernoulli numbers defined by

$$\frac{x}{e^x - 1} = 1 + \frac{x}{2} + \sum_{n=1}^{\infty} B_{2n} \frac{x^{2n}}{(2n)!}.$$

Then (6) follows from (7), the vanishing of Chern character of Hodge bundle $\text{ch}_{2l-1}(\mathbb{E}) = 0$ for $l > g$, and setting $\mathbf{s} = 0$. This leads to the question how can these universal equations for pure descendant Gromov–Witten invariants be extended to the case $l \leq g$? In this paper, we give an answer to this question in the case $l = g$.

2.2. Pixton’s formula for double ramification cycles and the λ_g class.

2.2.1. Pixton’s formula for double ramification cycles. Denote $G_{g,n}$ to be the set of stable graphs of genus- g , n marked points [Pan18, §4.2]. Let $A = (a_1, \dots, a_n)$ be a vector of double ramification data, i.e. a vector of integers such that $\sum_{i=1}^n a_i = 0$. Let $\Gamma \in G_{g,n}$ be a stable graph of genus g with n legs and r be a positive integer. A weighting mod r of Γ is a function on the set of half-edges,

$$w: H(\Gamma) \rightarrow \{0, \dots, r-1\},$$

which satisfies the following three properties:

(i) $\forall h_i \in L(\Gamma)$, corresponding to the marking $i \in \{1, \dots, n\}$,

$$w(h_i) \equiv a_i \pmod{r},$$

(ii) $\forall e \in E(\Gamma)$, corresponding to two half-edges $h, h' \in H(\Gamma)$,

$$w(h) + w(h') \equiv 0 \pmod{r},$$

(iii) $\forall v \in V(\Gamma)$,

$$\sum_{v(h)=v} w(h) \equiv 0 \pmod{r},$$

where the sum is taken over all half-edges incident to v .

Let $\mathbf{W}_{\Gamma,r}$ be the set of weightings mod r of Γ . The set $\mathbf{W}_{\Gamma,r}$ is finite, with cardinality $r^{h^1(\Gamma)}$. For each stable graph $\Gamma \in G_{g,n}$, we associate a moduli space $\overline{\mathcal{M}}_{\Gamma} := \prod_{v \in V(\Gamma)} \overline{\mathcal{M}}_{g(v),n(v)}$ and define a natural map $\xi_{\Gamma} : \overline{\mathcal{M}}_{\Gamma} \rightarrow \overline{\mathcal{M}}_{g,n}$ to be the canonical gluing morphism.

We denote by $\mathbf{P}_g^{d,r}(A) \in R^d(\overline{\mathcal{M}}_{g,n})$ the degree d component of the tautological class

$$\sum_{\Gamma \in G_{g,n}} \sum_{w \in \mathbf{W}_{\Gamma,r}} \frac{1}{|\text{Aut}(\Gamma)|} \frac{1}{r^{h^1(\Gamma)}} \xi_{\Gamma*} \left[\prod_{i=1}^n \exp(a_i^2 \psi_{h_i}) \cdot \prod_{e=(h,h') \in E(\Gamma)} \frac{1 - \exp(-w(h)w(h')(\psi_h + \psi_{h'}))}{\psi_h + \psi_{h'}} \right]$$

in $R^*(\overline{\mathcal{M}}_{g,n})$.

Pixton proved for fixed g , A , and d , the class $\mathbf{P}_g^{d,r}(A) \in R^d(\overline{\mathcal{M}}_{g,n})$ is polynomial in r for r sufficiently large. We denote by $\mathbf{P}_g^d(A)$ the value at $r = 0$ of the polynomial associated to $\mathbf{P}_g^{d,r}(A)$. In other words, $\mathbf{P}_g^d(A)$ is the constant term of the associated polynomial in r .

The following formula for double ramification cycles was first conjectured by Pixton.

Theorem 8 ([JPPZ17]). *For $g \geq 0$ and double ramification data A , we have*

$$\text{DR}_g(A) = 2^{-g} \mathbf{P}_g^g(A) \in R^g(\overline{\mathcal{M}}_{g,n}).$$

2.2.2. Formula for λ_g . The top Chern class λ_g of the Hodge bundle $\mathbb{E}$ is a very special case of a double ramification cycle (see [JPPZ17, §0.5.3]):

$$\text{DR}_g(0, \dots, 0) = (-1)^g \lambda_g \in R^g(\overline{\mathcal{M}}_{g,n}). \quad (8)$$

Combining Theorem 8 and (8), it leads to a nice formula for the top Chern class of Hodge bundle λ_g , which is supported on the boundary divisor of curves with a non-separating node. That is, in $R^g(\overline{\mathcal{M}}_{g,n})$, we have

$$\lambda_g = \frac{(-1)^g}{2^g} \mathbf{P}_g^g(0, \dots, 0). \quad (9)$$

To illustrate this, we list some explicit formulae of λ_g on $\overline{\mathcal{M}}_g$ ($g \geq 2$) in low genus (cf. [JPPZ17, §3.1], which also lists a longer formula in genus 4):

Genus 1.

$$\lambda_1 = -\frac{1}{2} \mathbf{P}_1^1(0, 0) = \frac{1}{24} \xi_{\Gamma*}(1) \quad (10)$$

where $\Gamma \in G_{1,2}$ is the dual graph of the boundary divisor of singular stable curves with a non-separating node.

Genus 2.

$$\lambda_2 = \frac{1}{240} \text{graph}_1 + \frac{1}{1152} \text{graph}_2.$$

Genus 3.

$$\begin{aligned} \lambda_3 = & \frac{1}{2016} \text{graph}_3 + \frac{1}{2016} \text{graph}_4 - \frac{1}{672} \text{graph}_5 + \frac{1}{5760} \text{graph}_6 \\ & - \frac{13}{30240} \text{graph}_7 - \frac{1}{5760} \text{graph}_8 + \frac{1}{82944} \text{graph}_9. \end{aligned}$$

All these expressions are obtained by substituting $A = \emptyset$ in Pixton's formula for the double ramification cycle (except in the genus one case, where we should set $A = (0, 0)$).

2.3. Universal equations from $\text{ch}_{2g-1}(\mathbb{E})$. A well-known formula for $\text{ch}_{2g-1}(\mathbb{E})$ (cf. [FP00b, (46)]) is

$$\text{ch}_{2g-1}(\mathbb{E}) = \frac{(-1)^{g-1}}{(2g-1)!} \lambda_g \lambda_{g-1}. \quad (11)$$

In fact, this is a consequence from the following general relationship between the Chern characters and Chern classes of a vector bundle:

$$\sum_{k \geq 1} (-1)^{k-1} (k-1)! \text{ch}_k(\mathbb{E}) t^k = \log \left(\sum_{k \geq 0} c_k(\mathbb{E}) t^k \right).$$

Taking derivative $\frac{d}{dt}$, we get

$$\sum_{k \geq 1} (-1)^{k-1} k! \text{ch}_k(\mathbb{E}) t^{k-1} = \left(\sum_{k \geq 0} \lambda_k t^k \right)^{-1} \cdot \left(\sum_{k \geq 1} k \cdot \lambda_k t^{k-1} \right).$$

Since $c(\mathbb{E})^{-1} = c(\mathbb{E}^*)$, the above becomes

$$\sum_{k \geq 1} (-1)^{k-1} k! \text{ch}_k(\mathbb{E}) t^{k-1} = \left(\sum_{k=0}^g \lambda_k (-t)^k \right) \cdot \left(\sum_{k=1}^g k \cdot \lambda_k t^{k-1} \right). \quad (12)$$

Taking the coefficients of t^{2g-2} on both sides of (12) gives (11).

Recall the following important properties of the Hodge bundle:

- Pulling back via the gluing map $\iota_{g_1, g_2}: \overline{\mathcal{M}}_{g_1, n_1+1} \times \overline{\mathcal{M}}_{g_2, n_2+1} \rightarrow \overline{\mathcal{M}}_{g, n}$, we have

$$\iota_{g_1, g_2}^* \mathbb{E}_g \cong p_1^* \mathbb{E}_{g_1} \oplus p_2^* \mathbb{E}_{g_2} \quad (13)$$

where $p_i (i = 1, 2)$ denotes via the projection maps from $\overline{\mathcal{M}}_{g_1, n_1+1} \times \overline{\mathcal{M}}_{g_2, n_2+1}$ onto its factors.

- Pulling back via the gluing map: $\iota_{g-1}: \overline{\mathcal{M}}_{g-1, n+1} \rightarrow \overline{\mathcal{M}}_{g, n}$, we have

$$\iota_{g-1}^* \mathbb{E}_g \cong \mathbb{E}_{g-1} \oplus \mathcal{O} \quad (14)$$

where $\mathcal{O}$ is the structure sheaf.

Lemma 9. *For any $g \geq 2$ and a stable circular graph $\Gamma \in G_{g,0}^\circ$, the contribution of Γ to $\text{P}_g^g(\emptyset)$ is:*

$$\text{Cont}_\Gamma \text{P}_g^g(\emptyset) = \frac{B_{2g}}{|\text{Aut}(\Gamma)|} \sum_{\substack{\sum_{e \in E(\Gamma)=g} k(e)=g \\ k(e) \geq 1, \forall e \in E(\Gamma)}} (\xi_\Gamma)_* \prod_{e \in E(\Gamma)} \left(\frac{-1}{k(e)!} (\psi_{h(e)} + \psi_{h'(e)})^{k(e)-1} \right). \quad (15)$$

Proof. For any graph $\Gamma \in G_{g,0}^\circ$, the associated weighting modulo r is either identically equal to 0, or alternatingly a and $r-a$ for some $1 \leq a \leq r-1$. From [JPPZ17, Appendix], we know that

$$\left[\frac{1}{|\text{Aut}(\Gamma)|} \frac{1}{r} \sum_{a=1}^{r-1} (\xi_\Gamma)_* \prod_{e \in E(\Gamma)} \frac{1 - \exp(-a(r-a)(\psi_{h(e)} + \psi_{h'(e)}))}{\psi_{h(e)} + \psi_{h'(e)}} \right]_{\deg=g}$$

is polynomial in r for r sufficiently large. The contribution $\text{Cont}_\Gamma \mathbf{P}_g^g(\emptyset)$ of the graph Γ to $\mathbf{P}_g^g(\emptyset)$ equals the constant term of this polynomial. By a direct computation,

$$\begin{aligned} & \left[\frac{1}{|\text{Aut}(\Gamma)|} \frac{1}{r} \sum_{a=1}^{r-1} (\xi_\Gamma)_* \prod_{e \in E(\Gamma)} \frac{1 - \exp(-a(r-a)(\psi_{h(e)} + \psi_{h'(e)}))}{\psi_{h(e)} + \psi_{h'(e)}} \right]_{\deg=g} \\ &= \frac{1}{|\text{Aut}(\Gamma)|} \frac{1}{r} \sum_{a=1}^{r-1} \sum_{\sum_{e \in E(\Gamma)} k(e)=g} (\xi_\Gamma)_* \prod_{e \in E(\Gamma)} \frac{-1}{k(e)!} a^{k(e)} (a-r)^{k(e)} (\psi_{h(e)} + \psi_{h'(e)})^{k(e)-1} \\ &= \left(\frac{1}{r} \sum_{a=1}^{r-1} a^g (a-r)^g \right) \cdot \frac{1}{|\text{Aut}(\Gamma)|} \sum_{\sum_{e \in E(\Gamma)} k(e)=g} (\xi_\Gamma)_* \prod_{e \in E(\Gamma)} \frac{-1}{k(e)!} (\psi_{h(e)} + \psi_{h'(e)})^{k(e)-1}. \end{aligned}$$

Therefore,

$$\text{Cont}_\Gamma \mathbf{P}_g^g(\emptyset) = \frac{B_{2g}}{|\text{Aut}(\Gamma)|} \sum_{e \in E(\Gamma)=g} (\xi_\Gamma)_* \prod_{e \in E(\Gamma)} \left(\frac{-1}{k(e)!} (\psi_{h(e)} + \psi_{h'(e)})^{k(e)-1} \right),$$

where we have used Faulhaber's formula

$$\sum_{a=1}^{r-1} a^m = \frac{1}{m+1} \sum_{k=0}^m \binom{m+1}{k} B_k r^{m-k+1}$$

for any $m \geq 1$. □

Proposition 10. *For any $g \geq 2$, the following relation holds on $\overline{\mathcal{M}}_g$:*

$$\begin{aligned} & \lambda_g \lambda_{g-1} \\ &= \frac{-B_{2g}}{2^{2g-1}} \sum_{\Gamma \in G_{g,0}^\circ} \frac{1}{|\text{Aut}(\Gamma)|} \sum_{\substack{\sum_{e \in E(\Gamma)} k(e)=g \\ k(e) \geq 1, \forall e \in E(\Gamma)}} (\xi_\Gamma)_* \left(\prod_{e \in E(\Gamma)} \left(\frac{-1}{k(e)} \sum_{l(e)+m(e)=k(e)-1} \frac{\psi_{h(e)}^{l(e)} \psi_{h'(e)}^{m(e)}}{l(e)! m(e)!} \right) \prod_{v \in V(\Gamma)} \mathbf{P}_{g(v)}^{g(v)}(0,0) \right) \end{aligned} \quad (16)$$

Proof. We compute:

$$\begin{aligned} \lambda_{g-1} \cdot \mathbf{P}_g^g(\emptyset) &= \lambda_{g-1} \cdot \sum_{\Gamma \in G_{g,0}^\circ} \text{Cont}_\Gamma \mathbf{P}_g^g(\emptyset) = \sum_{\Gamma \in G_{g,0}^\circ} \lambda_{g-1} \cdot \text{Cont}_\Gamma \mathbf{P}_g^g(\emptyset) \\ &= \sum_{\Gamma \in G_{g,0}^\circ} \frac{B_{2g}}{|\text{Aut}(\Gamma)|} \sum_{\substack{\sum_{e \in E(\Gamma)} k(e)=g \\ k(e) \geq 1, \forall e \in E(\Gamma)}} (\xi_\Gamma)_* \left(\prod_{e \in E(\Gamma)} \frac{-1}{k(e)!} (\psi_{h(e)} + \psi_{h'(e)})^{k(e)-1} \cdot \prod_{v \in V(\Gamma)} \lambda_{g(v)} \right) \end{aligned}$$

In the second equality, we used both that, first, by definition, no stable graph with at least one separating edge contributes to $\mathbf{P}_g^g(\emptyset)$. Second, no graph with more than one loop contributes, since applying the properties of the Hodge bundle (13) and (14) to λ_{g-1} , the contribution of such a graph would involve a Hodge class λ_i for $i > g(v)$ on some vertex moduli space $\overline{\mathcal{M}}_{g(v),n(v)}$. Finally, in the third equality, we again used the properties of the Hodge bundle, combined with Lemma 9. We next apply (9) twice, and obtain

$$\lambda_g \lambda_{g-1} = \frac{(-1)^g}{2^g} \lambda_{g-1} \mathbf{P}_g^g(\emptyset)$$

$$\begin{aligned}
&= \frac{(-1)^g}{2^g} \sum_{\Gamma \in G_{g,0}^\circ} \frac{B_{2g}}{|\text{Aut}(\Gamma)|} \sum_{\substack{\sum_{e \in E(\Gamma)} k(e) = g \\ k(e) \geq 1 \forall e \in E(\Gamma)}} (\xi_\Gamma)^* \left(\prod_{e \in E(\Gamma)} \frac{-1}{k(e)!} (\psi_{h(e)} + \psi_{h'(e)})^{k(e)-1} \cdot \prod_{v \in V(\Gamma)} \lambda_{g(v)} \right) \\
&= \frac{(-1)^{2g-1}}{2^{2g-1}} \sum_{\Gamma \in G_{g,0}^\circ} \frac{B_{2g}}{|\text{Aut}(\Gamma)|} \sum_{\substack{\sum_{e \in E(\Gamma)} k(e) = g \\ k(e) \geq 1 \forall e \in E(\Gamma)}} (\xi_\Gamma)^* \left(\prod_{e \in E(\Gamma)} \frac{-1}{k(e)!} (\psi_{h(e)} + \psi_{h'(e)})^{k(e)-1} \cdot \prod_{v \in V(\Gamma)} P_{g(v)}^{g(v)}(0,0) \right).
\end{aligned}$$

After expanding each factor $(\psi_{h(e)} + \psi_{h'(e)})^{k(e)-1}$ via the binomial theorem, this yields (16). $\square$

To illustrate this, we list formulae for $\lambda_g \lambda_{g-1}$ on $\overline{\mathcal{M}}_g$ in low genus:
Genus 2.

$$\lambda_2 \lambda_1 = -\frac{1}{480} \begin{array}{c} \psi \\ \bullet \\ \text{P}_1^1(0,0) \end{array}.$$

Genus 3.

$$\lambda_3 \lambda_2 = \frac{1}{8064} \begin{array}{c} \psi^2 \\ \bullet \\ \text{P}_2^2(0,0) \end{array} + \frac{1}{8064} \begin{array}{c} \psi \psi \\ \bullet \\ \text{P}_2^2(0,0) \end{array} - \frac{1}{2688 \text{P}_1^1(0,0)} \begin{array}{c} \psi \\ \bullet \quad \bullet \\ \text{P}_1^1(0,0) \end{array}.$$

Genus 4.

$$\begin{aligned}
\lambda_4 \lambda_3 = & -\frac{1}{15360} \begin{array}{c} \psi^3 \\ \bullet \\ \text{P}_3^3(0,0) \end{array} - \frac{1}{5120} \begin{array}{c} \psi^2 \psi \\ \bullet \\ \text{P}_3^3(0,0) \end{array} + \frac{1}{23040 \text{P}_1^1(0,0)} \begin{array}{c} \psi^2 \\ \bullet \quad \bullet \\ \text{P}_2^2(0,0) \end{array} + \frac{1}{30720 \text{P}_1^1(0,0)} \begin{array}{c} \psi \\ \bullet \quad \bullet \\ \text{P}_2^2(0,0) \end{array} \\
& + \frac{1}{11520 \text{P}_1^1(0,0)} \begin{array}{c} \psi \psi \\ \bullet \quad \bullet \\ \text{P}_2^2(0,0) \end{array} + \frac{1}{15360 \text{P}_1^1(0,0)} \begin{array}{c} \psi \psi \\ \bullet \quad \bullet \\ \text{P}_2^2(0,0) \end{array} + \frac{1}{23040 \text{P}_1^1(0,0)} \begin{array}{c} \psi^2 \\ \bullet \quad \bullet \\ \text{P}_2^2(0,0) \end{array} \\
& + \frac{1}{30720 \text{P}_1^1(0,0)} \begin{array}{c} \psi \psi \\ \bullet \quad \bullet \\ \text{P}_2^2(0,0) \end{array} - \frac{1}{7680 \text{P}_1^1(0,0)} \begin{array}{c} \text{P}_1^1(0,0) \\ \bullet \quad \bullet \quad \bullet \\ \text{P}_1^1(0,0) \end{array}.
\end{aligned}$$

2.4. Proof of Theorem 1. We first recall some standard results about the correspondence between descendant and ancestor invariants (cf. [CG07, Appendix 2]). Consider the stabilization morphism

$$\text{St}: \overline{\mathcal{M}}_{g,n}(X, \beta) \rightarrow \overline{\mathcal{M}}_{g,n}$$

which forgets the map f in $(C; x_1, \dots, x_n; f) \in \overline{\mathcal{M}}_{g,n}(X, \beta)$ and stabilizes the curve $(C; x_1, \dots, x_n)$ by contracting unstable components to points. Let

$$\pi_m: \overline{\mathcal{M}}_{g,n+m}(X, \beta) \rightarrow \overline{\mathcal{M}}_{g,n}(X, \beta)$$

be the morphism, which forgets the last m markings, and stabilizes the curve. For $1 \leq i \leq n$, denote the cotangent line bundle along the i -th marked point over $\overline{\mathcal{M}}_{g,n}$ and $\overline{\mathcal{M}}_{g,n+m}(X, \beta)$ by L_i and $\mathcal{L}_i$, respectively. The bundles $\mathcal{L}_i$ and $\pi_m^* \text{St}^* L_i$ over $\overline{\mathcal{M}}_{g,n+m}(X, \beta)$ are identified outside the locus D consisting of maps such that the i -th marked point is situated on a component of the curve which gets collapsed by $\text{St} \circ \pi_m$.

This locus D is the image of the gluing map

$$\iota: \sqcup_{\substack{m_1+m_2=m \\ \beta_1+\beta_2=d}} \overline{\mathcal{M}}_{0,\{i\}+\bullet+m_1}(X, \beta_1) \times_X \overline{\mathcal{M}}_{g,[n]\setminus\{i\}+\circ+m_2}(X, \beta_2) \rightarrow \overline{\mathcal{M}}_{g,n+m}(X, \beta)$$

where the two markings $\bullet$ and $\circ$ are glued together under map ι . We denote the domain of this map by $Y_{n,m,\beta}^{(i)}$. The virtual normal bundle to D at a generic point is $\text{Hom}(\pi_m^* \text{St}^* c_1(L_i), \mathcal{L}_i)$,

and so D is “virtually Poincaré-dual” to $c_1(\mathcal{L}_i) - \pi_m^* \text{St}^* c_1(L_i)$ in the sense that

$$(c_1(\mathcal{L}_i) - \pi_m^* \text{St}^* c_1(L_i)) \cap [\overline{\mathcal{M}}_{n+m}(X, \beta)]^{\text{vir}} = \iota_* [Y_{n,m,\beta}^{(i)}]^{\text{vir}} \quad (17)$$

Define mixed type of ancestor and descendant twisted correlator as follows:

$$\begin{aligned} & \langle \langle \bar{\tau}_{j_1} \tau_{i_1}(\phi_1), \dots, \bar{\tau}_{j_n} \tau_{i_n}(\phi_n); c_{g,n} \rangle \rangle_g(\mathbf{t}(\psi)) \\ &:= \sum_{m, \beta \geq 0} \frac{Q^\beta}{m!} \int_{[\overline{\mathcal{M}}_{g,n+m}(X, \beta)]^{\text{vir}}} \prod_{k=1}^n \left(\bar{\psi}_k^{j_k} \psi_k^{i_k} \text{ev}_k^* \phi_k \right) \prod_{k=n+1}^{n+m} \text{ev}_k^* \mathbf{t}(\psi_k) \cdot \pi_m^* \text{St}^* c_{g,n}. \end{aligned} \quad (18)$$

for certain cohomology class $c_{g,n} \in H^*(\overline{\mathcal{M}}_{g,n})$, where $\psi_k := c_1(\mathcal{L}_k)$ and $\bar{\psi}_k := \pi_m^* \text{St}^* c_1(L_k)$. Below we only focus the cases of $c_{g,n} = 1$ or λ_g .

Lemma 11. *For any $2g - 2 + n > 0$, on the big phase space, we have*

$$\langle \langle \bar{\tau}_{j_1} \tau_{i_1}(\phi_{\alpha_1}), \dots, \bar{\tau}_{j_n} \tau_{i_n}(\phi_{\alpha_n}); \lambda_g \rangle \rangle_g = \langle \langle T^{j_1}(\tau_{i_1}(\phi_{\alpha_1})), \dots, T^{j_n}(\tau_{i_n}(\phi_{\alpha_n})); \lambda_g \rangle \rangle_g \quad (19)$$

for any $\{\phi_{\alpha_i} : i = 1, \dots, n\} \subset H^*(X; \mathbb{C})$ and T is the operator from (1). In particular,

$$\langle \langle \bar{\tau}_{j_1}(\phi_{\alpha_1}), \dots, \bar{\tau}_{j_n}(\phi_{\alpha_n}); \lambda_g \rangle \rangle_g = \langle \langle T^{j_1}(\phi_{\alpha_1}), \dots, T^{j_n}(\phi_{\alpha_n}); \lambda_g \rangle \rangle_g. \quad (20)$$

Note that by (9), these two equations also hold true if we replace λ_g by $\mathbf{P}_g^g(0, \dots, 0)$.

Proof. We will focus on the first marked point, and for simplicity, suppress the content of the other marked points from our notation. By (17) and (13), the following identity holds for mixed type twisted correlators on the big phase space

$$\langle \langle \bar{\tau}_{j_1-1} \tau_{i_1+1}(\phi), \dots; \lambda_g \rangle \rangle_g = \langle \langle \bar{\tau}_{j_1} \tau_{i_1}(\phi), \dots; \lambda_g \rangle \rangle_g + \langle \langle \tau_{i_1}(\phi), \phi_\beta \rangle \rangle_0 \langle \langle \bar{\tau}_{j_1-1}(\phi^\beta), \dots; \lambda_g \rangle \rangle_g$$

for any $\phi \in H^*(X; \mathbb{C})$.

This implies

$$\begin{aligned} & \langle \langle \bar{\tau}_{j_1} \tau_{i_1}(\phi_{\alpha_1}), \dots; \lambda_g \rangle \rangle_g = \langle \langle \bar{\tau}_{j_1-1} (\tau_{i_1+1}(\phi_{\alpha_1}) - \langle \langle \tau_{i_1}(\phi_{\alpha_1}), \phi_\beta \rangle \rangle_0 \phi^\beta), \dots; \lambda_g \rangle \rangle_g \\ &= \langle \langle \bar{\tau}_{j_1-1} T(\tau_{i_1}(\phi_{\alpha_1})), \dots; \lambda_g \rangle \rangle_g. \end{aligned}$$

Repeating this $j_1 - 1$ more times, we get

$$\langle \langle \bar{\tau}_{j_1} \tau_{i_1}(\phi_{\alpha_1}), \dots; \lambda_g \rangle \rangle_g = \langle \langle T^{j_1}(\tau_{i_1}(\phi_{\alpha_1})), \dots; \lambda_g \rangle \rangle_g$$

Similar analysis for the other markings, yields (19). $\square$

For any g, n such that $2g - 2 + n > 0$ and $\mathbf{t} = \sum_{n,\alpha} t_n^\alpha z^n \phi_\alpha \in H^*(X)[[\{t_n^\alpha\}]] [[z]]$, define a homomorphism $\Omega_{g,n}^{\mathbf{t}} : H^*(X)[[\psi]]^{\otimes n} \rightarrow H^*(\overline{\mathcal{M}}_{g,n})[[Q]][[\{t_n^\alpha\}]]$ via

$$\Omega_{g,n}^{\mathbf{t}}(\phi_{\alpha_1} \psi_1^{i_1}, \dots, \phi_{\alpha_n} \psi_n^{i_n}) = \sum_{m, \beta \geq 0} \frac{Q^\beta}{m!} \text{St}_* \pi_{m*} \left(\prod_{k=1}^n \text{ev}_k^* \phi_{\alpha_k} \psi_k^{i_k} \prod_{k=n+1}^{n+m} \text{ev}_k^* \mathbf{t}(\psi_k) \right).$$

From the “cutting edges” axiom of virtual cycles of moduli of stable maps to X , this system of homomorphisms satisfies the splitting axioms:

$$\iota_{g_1, g_2}^* \Omega_{g,n}^{\mathbf{t}}(\phi_{\alpha_1} \psi_1^{i_1}, \dots, \phi_{\alpha_n} \psi_n^{i_n}) = \sum_{\beta} \Omega_{g_1, n_1+1}^{\mathbf{t}}(\otimes_{k \in I_1} \phi_{\alpha_k} \psi_k^{i_k}, \phi_\beta) \otimes \Omega_{g_2, n_2+1}^{\mathbf{t}}(\otimes_{k \in I_2} \phi_{\alpha_k} \psi_k^{i_k} \otimes \phi^\beta) \quad (21)$$

and

$$\iota_{g-1}^* \Omega_{g,n}^{\mathbf{t}}(\phi_{\alpha_1} \psi_1^{i_1}, \dots, \phi_{\alpha_n} \psi_n^{i_n}) = \sum_{\beta} \Omega_{g-1, n+2}^{\mathbf{t}}(\phi_{\alpha_1} \psi_1^{i_1}, \dots, \phi_{\alpha_n} \psi_n^{i_n}, \phi_\beta, \phi^\beta) \quad (22)$$

where $\iota_{g_1, g_2} : \overline{\mathcal{M}}_{g_1, n_1+1} \times \overline{\mathcal{M}}_{g_2, n_2+1} \rightarrow \overline{\mathcal{M}}_{g, n}$ and $\iota_{g-1} : \overline{\mathcal{M}}_{g-1, n+2} \rightarrow \overline{\mathcal{M}}_{g, n}$ are the two canonical gluing maps. Note that, by definition, Ω is related to the double bracket of descendant potential via integration on the moduli space of curve, i.e.

$$\int_{\overline{\mathcal{M}}_{g, n}} \Omega_{g, n}^{\mathbf{t}}(v_1 \psi_1^{i_1}, \dots, v_n \psi_n^{i_n}) = \langle \langle \tau_{i_1}(v_1), \dots, \tau_{i_n}(v_n) \rangle \rangle_g(\mathbf{t})$$

Now let us prove Theorem 1. On the one hand, taking $l = g$ and $\mathbf{s} = 0$ in (7) yields

$$\begin{aligned} & \frac{(2g)!}{B_{2g}} \langle \langle \text{ch}_{2g-1}(\mathbb{E}) \rangle \rangle_g \\ &= -\tilde{t}_n^\alpha \langle \langle \tau_{n+2g-1}(\phi_\alpha) \rangle \rangle_g + \frac{1}{2} \sum_{i=0}^{2g-2} (-1)^i \left\{ \langle \langle \tau_i(\phi_\alpha) \tau_{2g-2-i}(\phi^\alpha) \rangle \rangle_{g-1} + \sum_{h=0}^g \langle \langle \tau_i(\phi_\alpha) \rangle \rangle_h \langle \langle \tau_{2g-2-i}(\phi^\alpha) \rangle \rangle_{g-h} \right\} \end{aligned} \quad (23)$$

On the other hand, by combining (11) and Proposition 10, we obtain

$$\begin{aligned} & \frac{(2g)!}{B_{2g}} \text{ch}_{2g-1}(\mathbb{E}) \\ &= \frac{(-1)^g g}{2^{2g-2}} \sum_{\Gamma \in G_{g,0}^\circ} \frac{1}{|\text{Aut}(\Gamma)|} \sum_{\substack{e \in E(\Gamma) \\ k(e)=g \\ k(e) \geq 1, \forall e \in E(\Gamma)}} (\xi_\Gamma)^* \left(\prod_{e \in E(\Gamma)} \left(\frac{-1}{k(e)} \sum_{l(e)+m(e)=k(e)-1} \frac{\psi_{h(e)}^{l(e)} \psi_{h'(e)}^{m(e)}}{l(e)! m(e)!} \right) \prod_{v \in V(\Gamma)} \mathbf{P}_{g(v)}^{g(v)}(0, 0) \right). \end{aligned}$$

Then combining with Lemma 11, the splitting axioms (21) and (22), we get

$$\begin{aligned} & \frac{(2g)!}{B_{2g}} \langle \langle \text{ch}_{2g-1}(\mathbb{E}) \rangle \rangle_g = \frac{(2g)!}{B_{2g}} \int_{\overline{\mathcal{M}}_g} \Omega_{g,0}^{\mathbf{t}} \cdot \text{ch}_{2g-1}(\mathbb{E}) \\ &= \frac{(-1)^g g}{2^{2g-2}} \sum_{\Gamma \in G_{g,0}^\circ} \frac{1}{|\text{Aut}(\Gamma)|} \sum_{\substack{e \in E(\Gamma) \\ k(e)=g \\ k(e) \geq 1, \forall e \in E(\Gamma)}} \int_{\overline{\mathcal{M}}_\Gamma} \xi_\Gamma^* \Omega_{g,0}^{\mathbf{t}} \cdot \left(\prod_{e \in E(\Gamma)} \left(\frac{-1}{k(e)} \sum_{l(e)+m(e)=k(e)-1} \frac{\psi_{h(e)}^{l(e)} \psi_{h'(e)}^{m(e)}}{l(e)! m(e)!} \right) \right. \\ & \quad \left. \cdot \prod_{v \in V(\Gamma)} \mathbf{P}_{g(v)}^{g(v)}(0, 0) \right) \\ &= \frac{(-1)^g g}{2^{2g-2}} \sum_{m=1}^{g-1} \frac{1}{2m} \sum_{\substack{k_1+\dots+k_m=g \\ k_1, \dots, k_m \geq 1}} \sum_{\substack{g_1+\dots+g_m=g-1 \\ g_1, \dots, g_m \geq 1}} \sum_{\alpha_1, \dots, \alpha_m=1}^N \prod_{i=1}^m \frac{-1}{k_i!} \sum_{l_i=0}^{k_i-1} \binom{k_i-1}{l_i} \langle \langle T^{l_i}(\phi_{\alpha_{i-1}}) T^{k_i-1-l_i}(\phi^{\alpha_i}); \mathbf{P}_{g_i}^{g_i}(0, 0) \rangle \rangle_{g_i} \end{aligned}$$

where $\alpha_0 = \alpha_m$. In the last equality, we took into account the symmetries from stable circular graphs Γ and indices $\{k_i\}_{i=1}^m, \{g_i\}_{i=1}^m$. Therefore, we prove Theorem 1.

2.5. Reformulation of Theorem 1. In this subsection, explain how to use Pixton's formula to rewrite each factor of the form $\langle \langle T^a(v_1) T^b(v_2); \mathbf{P}_g^g(0, 0) \rangle \rangle_g$ in the right hand side of (3) in terms of pure descendant Gromov-Witten invariants. For convenience, let $\mathbf{P}_\Gamma^g(0, 0)$ denote the constant term of the following polynomial in r for sufficiently large r

$$\mathbf{P}_\Gamma^g(0, 0) := \text{Coeff}_{r^0} \left[\sum_{w \in W_{\Gamma, r}} \frac{1}{|\text{Aut}(\Gamma)|} \frac{1}{r^{h^1(\Gamma)}} \prod_{e=(h, h') \in E(\Gamma)} \frac{1 - \exp(-w(h)w(h')(\psi_h + \psi_{h'}))}{\psi_h + \psi_{h'}} \right] \quad (24)$$

for any $\Gamma \in G_{g,2}$. We may view $\mathbf{P}_\Gamma^g(0, 0)$ as a polynomial in the ψ -classes $\{\psi_h : h \in H(\Gamma)\}$.

We next define a linear operator $P \mapsto \langle \langle T^a(v_1)T^b(v_2); P \rangle \rangle_\Gamma$ that turns any polynomial in the ψ -classes $\{\psi_h : h \in H(\Gamma)\}$ into a product of correlation functions. Given a monomial $P = c \cdot \prod_{h \in H(\Gamma)} \psi_h^{a_h}$, we let $\langle \langle T^a(v_1)T^b(v_2); P \rangle \rangle_\Gamma$ to be the contraction of the multilinear form

$$c \prod_{v \in V(\Gamma)} \langle \langle \cdot \cdot \cdot \rangle \rangle_{g(v)}$$

with arguments corresponding to half-edges $h \in H(\Gamma)$ via:

- for the argument corresponding to the first and second leg, use the vectors $T^a(v_1)$ and $T^b(v_2)$, respectively;
- for each edge $e = (h, h')$, use the bivector $T^{a_h}(\phi_\alpha) \otimes T^{a_{h'}}(\phi^\alpha)$ for the corresponding arguments.

Proposition 12.

$$\langle \langle T^a(v_1)T^b(v_2); P_g^g(0,0) \rangle \rangle_g = \sum_{\Gamma \in G_{g,2}} \langle \langle T^a(v_1)T^b(v_2); P_\Gamma^g(0,0) \rangle \rangle_\Gamma.$$

Proof. By (20) in Lemma 11 and (9), together with definition of Ω^t , we have

$$\begin{aligned} \langle \langle T^a(v_1)T^b(v_2); P_g^g(0,0) \rangle \rangle_g &= \langle \langle \bar{\tau}_a(v_1)\bar{\tau}_b(v_2); P_g^g(0,0) \rangle \rangle_g \\ &= \sum_{m,\beta \geq 0} \frac{Q^\beta}{m!} \int_{[\overline{\mathcal{M}}_{g,m+2}(X,\beta)]^{\text{vir}}} (\bar{\psi}_1^a \text{ev}_1^* v_1) (\bar{\psi}_2^b \text{ev}_2^* v_2) \prod_{k=3}^{m+2} \text{ev}_k^* \mathbf{t}(\psi_k) \cdot \pi_m^* \text{St}^* P_g^g(0,0) \\ &= \int_{\overline{\mathcal{M}}_{g,2}} \psi_1^a \psi_2^b \cdot \Omega_{g,2}^t(v_1, v_2) \cdot P_g^g(0,0). \end{aligned}$$

By the projection formula, this equals

$$\sum_{\Gamma \in G_{g,2}} \int_{\overline{\mathcal{M}}_{g,2}} \psi_1^a \psi_2^b \cdot \Omega_{g,2}^t(v_1, v_2) \cdot \xi_{\Gamma*} P_\Gamma^g(0,0) = \sum_{\Gamma \in G_{g,2}} \int_{\overline{\mathcal{M}}_\Gamma} \psi_1^a \psi_2^b \cdot \xi_\Gamma^* \Omega_{g,2}^t(v_1, v_2) \cdot P_\Gamma^g(0,0).$$

Then by the splitting axioms (21), (22) and equation (24), we can express each summand as a product of descendant correlation functions according to the operator $P \mapsto \langle \langle T^a(v_1)T^b(v_2); P \rangle \rangle_\Gamma$. In the process, the two marked points are assigned the vectors v_1, v_2 , and the cotangent line class ψ is translated to descendants using the operator T . Each node is translated into a sum of pairs of primary vector fields ϕ_α and ϕ^α . $\square$

2.6. Proof of Corollary 2. The following vanishing identity for the Gromov–Witten invariants of any smooth projective variety was first conjectured by K. Liu and H. Xu in [LX09], and was later proven in [LP11].

$$0 = -\tilde{t}_n^\alpha \langle \langle \tau_{n+2g-1}(\phi_\alpha) \rangle \rangle_g + \frac{1}{2} \sum_{i=0}^{2g-2} (-1)^i \left\{ \sum_{h=0}^g \langle \langle \tau_i(\phi_\alpha) \rangle \rangle_h \langle \langle \tau_{2g-2-i}(\phi^\alpha) \rangle \rangle_{g-h} \right\} \quad (25)$$

for $n \geq 0$.

Combining (2) and (25), we obtain

$$\sum_{i=0}^{2g-2} (-1)^i \langle \langle \tau_i(\phi_\alpha) \tau_{2g-2-i}(\phi^\alpha) \rangle \rangle_{g-1}$$

$$= \frac{(-1)^g g}{2^{2g-2}} \sum_{m=1}^{g-1} \frac{1}{m} \sum_{\substack{k_1+\dots+k_m=g \\ k_1, \dots, k_m \geq 1}} \sum_{\substack{g_1+\dots+g_m=g-1 \\ g_1, \dots, g_m \geq 1}} \sum_{\alpha_1, \dots, \alpha_m=1}^N \prod_{i=1}^m \frac{-1}{k_i!} \sum_{l_i=0}^{k_i-1} \binom{k_i-1}{l_i} \langle\langle T^{l_i}(\phi_{\alpha_{i-1}}) T^{k_i-1-l_i}(\phi^{\alpha_i}); \mathbf{P}_{g_i}^{g_i}(0,0) \rangle\rangle_{g_i}. \quad (26)$$

Recall that for any arbitrary contravariant tensors P and Q on the big phase space, the following formula was proved in [Liu11, Proposition 3.2]:

$$\sum_{i=0}^m (-1)^i P(\tau_i(\phi_\alpha)) Q(\tau_{m-i}(\phi^\alpha)) = \sum_{i=0}^m (-1)^i P(T^i(\phi_\alpha)) Q(T^{m-i}(\phi^\alpha))$$

for $m \geq 0$. In particular, if we take $P(W) = Q(W) = W$, the left hand side of (26) becomes

$$\sum_{i=0}^{2g-2} (-1)^i \langle\langle T^i(\phi_\alpha) T^{2g-2-i}(\phi^\alpha) \rangle\rangle_{g-1}.$$

This completes the proof of Corollary 2.

3. PUSH-FORWARD RELATIONS ON $\overline{\mathcal{M}}_g$ FOR $(g \geq 2)$

3.1. Push-forward relations. First, we introduce our definition of push-forward relations, which is slightly different from the notion proposed in [Pan18, Section 4.9.2]. Consider the gluing map,

$$\iota: \overline{\mathcal{M}}_{g,2} \rightarrow \overline{\mathcal{M}}_{g+1}$$

with image equal to the boundary divisor $\Delta_0 \subset \overline{\mathcal{M}}_{g+1}$ of curves with a nonseparating node. A push-forward relation can be understood as an element of $\ker(\iota_*)$, where $\iota_*: R^*(\overline{\mathcal{M}}_{g,2}) \rightarrow R^*(\overline{\mathcal{M}}_{g+1})$. Push-forward relations among descendant stratum classes yield universal equations for genus less than or equal to g Gromov–Witten invariants, despite the fact that the relations are embedded in genus $g+1$ moduli space of curves. A simple family of push-forward relations in $R^{2g+r+1}(\overline{\mathcal{M}}_{g+1})$ for $(g \geq 1, r \geq 1)$ were first found in [LP11]

$$\sum_{a+b=2g+r} (-1)^a \iota_* (\psi_1^a \psi_2^b) = 0. \quad (27)$$

Comparing equations (4) and (27), we see (4) gives the correction terms explicitly for the left hand side of equation (27) for $r = 0$.

3.2. Proof of Theorem 5. Let $j: \Delta(g_1, g_2) \cong \overline{\mathcal{M}}_{g_1,2} \times \overline{\mathcal{M}}_{g_2,2} \rightarrow \overline{\mathcal{M}}_{g+1,2}$ denote the boundary divisor parametrizing nodal curves with two components C_1 and C_2 with genus g_1 and g_2 , where $g_1 + g_2 = g + 1$. Let ψ_{*i} be the cotangent line class at the node along the curve C_i for $i = 1, 2$. By the relation [LP11, Proposition 2] in genus $g + 1$, we have

$$-\psi_1^{2g+3} - \psi_2^{2g+3} + \sum_{g_1+g_2=g+1: g_i > 0} \sum_{a+b=2g+2} (-1)^a j_* (\psi_{*1}^a \psi_{*2}^b \cap [\Delta(g_1, g_2)]) = 0 \quad (28)$$

in $R^{2g+3}(\overline{\mathcal{M}}_{g+1,2})$. Pushing forward equation (28) to $\overline{\mathcal{M}}_{g+1}$ by forgetting the two marked points, via string equation, we have the following relation on $\overline{\mathcal{M}}_{g+1}$

$$\kappa_{2g+1} + \frac{1}{2} \sum_{g_1+g_2=g: g_i > 0} \sum_{a+b=2g} (-1)^a j_* (\psi_{*1}^a \psi_{*2}^b \cap [\Delta(g_1, g_2)]) = 0. \quad (29)$$

Meanwhile, by Mumford's Grothendieck–Riemann–Roch calculation (cf. [Mum83]), the following holds on $\overline{\mathcal{M}}_{g+1}$:

$$\begin{aligned} & \frac{(2g+2)!}{B_{2g+2}} \text{ch}_{2g+1}(\mathbb{E}) \\ &= \kappa_{2g+1} + \frac{1}{2} \iota_* \left(\sum_{a+b=2g} (-1)^a \psi_1^a \psi_2^b \right) + \frac{1}{2} \sum_{g_1+g_2=g, g_i > 0} \sum_{a+b=2g} (-1)^a j_* (\psi_{*1}^a \psi_{*2}^b \cap [\Delta(g_1, g_2)]) \end{aligned}$$

Thus we get the following relation on $\overline{\mathcal{M}}_{g+1}$,

$$\frac{(2g+2)!}{B_{2g+2}} \text{ch}_{2g+1}(\mathbb{E}) = \frac{1}{2} \iota_* \left(\sum_{a+b=2g} (-1)^a \psi_1^a \psi_2^b \right)$$

Together with equation (11) and Proposition 10, we obtain Theorem 5.

3.3. Connection to degree- g topological recursion relations. We propose the following connection between the relations (3), and a class of degree- g topological recursion relations on the moduli space of stable curves:

Conjecture 13. *The universal equation (3) is a consequence of the degree- g topological recursion relations proposed in [CJWZ23].*

We verify the conjecture for $g = 2$ case in the example below.

Example 14. *For genus $g = 2$, the equation (3) specializes to*

$$\begin{aligned} & \sum_{i=0}^2 (-1)^i \langle \langle T^i(\phi_\alpha) T^{2-i}(\phi^\alpha) \rangle \rangle_1 \\ &= \frac{2}{2^2} \sum_{m=1}^1 \frac{1}{m} \sum_{\substack{k_1+\dots+k_m=2 \\ k_1, \dots, k_m \geq 1}} \sum_{\substack{g_1+\dots+g_m=1 \\ g_1, \dots, g_m \geq 1}} \prod_{i=1}^m \frac{-1}{k_i!} \sum_{l_i=0}^{k_i-1} \binom{k_i-1}{l_i} \langle \langle T^{l_i}(\phi_{\alpha_{i-1}}) T^{k_i-1-l_i}(\phi^{\alpha_i}); \mathbf{P}_{g_i}^{g_i}(0, 0) \rangle \rangle_{g_i} \end{aligned} \quad (30)$$

We will check that (30) is a consequence of the genus-1 topological recursion relation:

$$\langle \langle T(W) \rangle \rangle_1 = \frac{1}{24} \sum_{\alpha} \langle \langle W \phi_\alpha \phi^\alpha \rangle \rangle_0 \quad (31)$$

for any vector field W on the big phase space. We start by rewriting the left hand side of (30) as

$$\begin{aligned} & \sum_{\alpha} \sum_{i=0}^2 (-1)^i \langle \langle T^i(\phi_\alpha) T^{2-i}(\phi^\alpha) \rangle \rangle_1 = 2 \sum_{\alpha} \langle \langle \phi_\alpha T^2(\phi^\alpha) \rangle \rangle_1 - \sum_{\alpha} \langle \langle T(\phi_\alpha) T(\phi^\alpha) \rangle \rangle_1 \\ &= \frac{1}{24} \sum_{\alpha, \beta} \langle \langle \phi_\alpha T(\phi^\alpha) \phi_\beta \phi^\beta \rangle \rangle_0, \end{aligned}$$

where we have used (31) and its derivatives in the last equality. The right hand side of (30) equals

$$- \frac{1}{4} \sum_{\alpha} \sum_{l_1=0}^1 \binom{1}{l_1} \langle \langle T^{l_1}(\phi_\alpha) T^{1-l_1}(\phi^\alpha); \mathbf{P}_1^1(0, 0) \rangle \rangle_1$$

$$\begin{aligned}
&= -\frac{1}{4} \sum_{\alpha} \langle \langle T(\phi_{\alpha}) \phi^{\alpha}; P_1^1(0, 0) \rangle \rangle_1 - \frac{1}{4} \sum_{\alpha} \langle \langle \phi_{\alpha} T(\phi^{\alpha}); P_1^1(0, 0) \rangle \rangle_1 \\
&= -\frac{1}{2} \sum_{\alpha} \langle \langle \phi_{\alpha} T(\phi^{\alpha}); P_1^1(0, 0) \rangle \rangle_1 = \frac{1}{24} \sum_{\alpha, \beta} \langle \langle \phi_{\alpha} T(\phi^{\alpha}) \phi_{\beta} \phi^{\beta} \rangle \rangle_0
\end{aligned}$$

where we have used the identity

$$\sum_{\alpha} \langle \langle T(\phi_{\alpha}) \phi^{\alpha}; P_1^1(0, 0) \rangle \rangle_1 = \sum_{\alpha, \beta} \eta^{\alpha\beta} \langle \langle T(\phi_{\alpha}) \phi_{\beta}; P_1^1(0, 0) \rangle \rangle_1 = \sum_{\beta} \langle \langle \phi_{\beta} T(\phi^{\beta}); P_1^1(0, 0) \rangle \rangle_1.$$

in the second equality, and the formula (10) for $P_1^1(0, 0)$ in the last equality.

4. CHERN CLASS OF HODGE BUNDLE

Unlike the Chern characters, the Chern classes $\lambda_i = c_i(\mathbb{E})$ already vanish in degrees $i > g$. Thus for degree reasons, we believe the vanishing of the λ_i implies much stronger relations than the relations from the Chern characters we have studied in Section 2, and this makes it very interesting to study what kind of relations the vanishing of the λ_i for $i > g$ leads to.

4.1. Linear Hodge integrals. Define the linear Hodge integral

$$\langle \tau_{k_1}(\phi_{\alpha_1}), \dots, \tau_{k_n}(\phi_{\alpha_n}); \lambda_j \rangle_{g, n, \beta} := \int_{[\overline{\mathcal{M}}_{g, n}(X, \beta)]^{\text{vir}}} \prod_{i=1}^n \psi_i^{k_i} \text{ev}_i^* \phi_{\alpha_i} \cdot \lambda_j \quad (32)$$

where $\lambda_j = c_j(\mathbb{E})$ is the j -th Chern class of the Hodge bundle $\mathbb{E}$.

The generating function of genus- g linear Hodge integrals is given by

$$F_g^{\mathbb{E}}(\mathbf{t}, u) := \left\langle \exp \left(\sum_{n, \alpha} t_n^{\alpha} \tau_n(\phi_{\alpha}) \right); \lambda(u) \right\rangle_g \quad (33)$$

where $\lambda(u) = \sum_{i=0}^g \lambda_i u^i$ is the Chern polynomial of the Hodge bundle $\mathbb{E}$, and u is a formal parameter. Via a standard identity relating the Chern classes and Chern characters of a vector bundle, we have

$$\lambda(u) = \exp \left(\sum_{l \geq 1} (2l-2)! u^{2l-1} \text{ch}_{2l-1}(\mathbb{E}) \right). \quad (34)$$

The generating function of genus- g linear Hodge integrals $F_g^{\mathbb{E}}(\mathbf{t}, u)$ can be obtained from $F_g^{\mathbb{E}}(\mathbf{t}, \mathbf{s})$ via the substitution

$$s_{2i-1} = (2i-2)! \cdot u^{2i-1}. \quad (35)$$

4.2. The Faber–Pandharipande relation and Givental quantization.

4.2.1. The Faber–Pandharipande relation. Define the differential operator for $l \geq 1$

$$D_{2l-1} := - \sum_{n=0}^{\infty} \sum_{\alpha} \tilde{t}_n^{\alpha} \frac{\partial}{\partial t_{n+2l-1}^{\alpha}} + \frac{\hbar}{2} \sum_{n=0}^{2l-2} (-1)^n \eta^{\alpha\beta} \frac{\partial}{\partial t_n^{\alpha}} \frac{\partial}{\partial t_{2l-2-n}^{\beta}} \quad (36)$$

Let $\mathcal{D}^{\mathbb{E}}(\mathbf{t}, \mathbf{s}) = \exp\left(\sum_{g \geq 0} \hbar^{g-1} F_g^{\mathbb{E}}(\mathbf{t}, \mathbf{s})\right)$ be the total descendant potential of Hodge integrals. The following differential equation was proved in [FP00a, Proposition 2]

$$\frac{\partial}{\partial s_{2l-1}} \mathcal{D}^{\mathbb{E}}(\mathbf{t}, \mathbf{s}) = \frac{B_{2l}}{(2l)!} D_{2l-1} \cdot \mathcal{D}^{\mathbb{E}}(\mathbf{t}, \mathbf{s}) \quad (37)$$

for any $l \geq 1$. The identity (37) may be used to derive the following formula (cf. [Giv01])

$$\mathcal{D}^{\mathbb{E}}(\mathbf{t}, \mathbf{s}) = \exp\left(\sum_{l \geq 1} \frac{B_{2l}}{(2l)!} s_{2l-1} D_{2l-1}\right) \cdot \mathcal{D}^X(\mathbf{t}) \quad (38)$$

for $\mathcal{D}^{\mathbb{E}}(\mathbf{t}, \mathbf{s})$ in terms of $\mathcal{D}^X(\mathbf{t}) = \exp\left(\sum_{g \geq 0} \hbar^{g-1} F_g(\mathbf{t})\right)$, the total descendant potential of Gromov–Witten invariants of X .

4.2.2. Quantization of quadratic Hamiltonians. In this section, we give a brief review of Givental's quantization of quadratic Hamiltonians (see [Giv01] for more details).

Let z be a formal variable. We consider the space $\mathbb{H}$ which is the space of Laurent polynomials in one variable z with coefficients in $H^*(X; \mathbb{C})$. We define a symplectic form Ω on $\mathbb{H}$ via

$$\Omega(f, g) = \text{Res}_{z=0} \eta(f(-z), g(z)) dz$$

for any $f, g \in \mathbb{H}$. Note that we have $\Omega(f, g) = -\Omega(g, f)$. There is a natural polarization $\mathbb{H} = \mathbb{H}_+ \oplus \mathbb{H}_-$ corresponding to the decomposition $f(z, z^{-1}) = f_+(z) + f_-(z^{-1})z^{-1}$ of Laurent polynomials into polynomial and polar parts. It is easy to see that $\mathbb{H}_+$ and $\mathbb{H}_-$ are both Lagrangian subspaces of $\mathbb{H}$ with respect to Ω .

Introduce a Darboux coordinate system $\{p_m^\beta, q_n^\alpha\}$ on $\mathbb{H}$ with respect to the above polarization. A general element $f \in \mathbb{H}$ can be written in the form

$$\sum_{m \geq 0, \beta} p_m^\beta \phi^\beta(-z)^{-m-1} + \sum_{n \geq 0, \alpha} q_n^\alpha \phi_\alpha z^n,$$

where $\{\phi^\alpha\}$ is the dual basis to $\{\phi_\alpha\}$. Denote

$$\mathbf{p}(z) := \sum_{m \geq 0, \beta} p_m^\beta \phi^\beta(-z)^{-m-1}, \quad \mathbf{q}(z) := \sum_{n \geq 0, \alpha} q_n^\alpha \phi_\alpha z^n.$$

For convenience, we identify $\mathbb{H}_+$ with the big phase space via identification between basis $\{z^n \phi_\alpha\}$ and $\{\tau_n(\phi_\alpha)\}$, and dilaton shift about coordinates $q_n^\alpha = t_n^\alpha - \delta_n^1 \delta_\alpha^1$.

Let $A: \mathbb{H} \rightarrow \mathbb{H}$ be a linear infinitesimal symplectic transformation, i.e. $\Omega(Af, g) + \Omega(f, Ag) = 0$ for any $f, g \in \mathbb{H}$. In Darboux coordinates, the quadratic Hamiltonian

$$f \rightarrow \frac{1}{2} \Omega(Af, f)$$

is a series of homogeneous degree two monomials in $\{p_m^\beta, q_n^\alpha\}$.

We define the quantization of quadratic monomials as

$$\widehat{q_m^\beta q_n^\alpha} = \frac{q_m^\beta q_n^\alpha}{\hbar}, \quad \widehat{q_m^\beta p_n^\alpha} = q_m^\beta \frac{\partial}{\partial q_n^\alpha}, \quad \widehat{p_m^\beta p_n^\alpha} = \hbar \frac{\partial}{\partial q_m^\beta} \frac{\partial}{\partial q_n^\alpha},$$

where $\hbar$ is a formal variable. We define the quantization $\widehat{A}$ of a general A by extending the above equalities linearly. The differential operators $\widehat{q_m^\beta q_n^\alpha}, \widehat{q_m^\beta p_n^\alpha}, \widehat{p_m^\beta p_n^\alpha}$ act on the *Fock space*, which is the space of formal functions in $\mathbf{q}(z) \in \mathbb{H}_+$. For example, the descendant

potential $\mathcal{D}(\mathbf{t})$ may be regarded as an element in the Fock space via the dilaton shift $\mathbf{q}(z) = \mathbf{t}(z) - \mathbf{1}z$. The quantization of a symplectic transform of the form $\exp(A)$, with A infinitesimal symplectic, is defined as $\exp(\widehat{A}) = \sum_{n \geq 0} \frac{\widehat{A}^n}{n!}$.

4.3. Universal relations from the Chern classes of the Hodge bundle. It is easy to see that multiplication by $\widehat{z^{2k-1}}$ defines an infinitesimal symplectic transformation on $(\mathbb{H}, \Omega)$, and we denote by $\widehat{z^{2k-1}}$ the corresponding quantization. It may be checked that $\widehat{z^{2k-1}}$ coincides with the operator D_{2k-1} from (36). For convenience, we will use the following convention

$$\widetilde{\mathcal{P}}(\mathbf{q}) = \mathcal{P}(\mathbf{t})|_{\mathbf{t}(z)=\mathbf{q}(z)+\mathbf{1}z}$$

for any partition function $\mathcal{P}(\mathbf{t})$. This allows us to restate (38) as

$$\widetilde{\mathcal{D}}^{\mathbb{E}}(\mathbf{q}, \mathbf{s}) = \widehat{R(z)} \cdot \widetilde{\mathcal{D}}^X(\mathbf{q}) \quad (39)$$

for the R -matrix

$$R(z) = \exp\left(\sum_{i=1}^{\infty} \frac{B_{2i}}{(2i)!} s_{2i-1} z^{2i-1}\right).$$

So under the substitution (35), taking the logarithm of equation (39), we obtain

$$\sum_{g \geq 0} \hbar^{g-1} \widetilde{F}_g^{\mathbb{E}}(\mathbf{q}, u) = \log \left(\exp \left(\sum_{l \geq 1} \frac{B_{2l}}{(2l)(2l-1)} u^{2l-1} \widehat{z^{2l-1}} \right) \widetilde{\mathcal{D}}^X(\mathbf{q}) \right).$$

Denote

$$\langle \langle \tau_{i_1}(\phi_1), \dots, \tau_{i_n}(\phi_n); c_{g,n} \rangle \rangle_g^{\sim}(\mathbf{q}(z)) = \langle \langle \tau_{i_1}(\phi_1), \dots, \tau_{i_n}(\phi_n); c_{g,n} \rangle \rangle_g(\mathbf{t}(z))|_{\mathbf{t}(z)=\mathbf{q}(z)+\mathbf{1}z} \quad (40)$$

for certain cohomology class $c_{g,n} \in H^*(\overline{\mathcal{M}}_{g,n})$.

Proposition 15. *Let $g > 0$ and $i \geq 0$. Then, the following holds for descendant Gromov–Witten invariants*

$$[u^i \hbar^{g-1}] \log \left(\exp \left(\sum_{l \geq 1} \frac{B_{2l}}{(2l)(2l-1)} u^{2l-1} \widehat{z^{2l-1}} \right) \widetilde{\mathcal{D}}^X(\mathbf{q}) \right) = \begin{cases} 0 & , \text{ if } i > g, \\ \frac{(-1)^g}{2^g} \cdot \langle \langle ; \mathbf{P}_g^g(\emptyset) \rangle \rangle_g^{\sim}(\mathbf{q}(z)) & , \text{ if } i = g. \end{cases}$$

Here $[u^i \hbar^{g-1}]$ denotes taking the coefficient of $u^i \hbar^{g-1}$.

Proof. This follows from the vanishing $\lambda_i = 0$ if $i > g$, and equation (9). $\square$

Theorem 16 ([Giv04]). *Let $\mathbf{q}(z) = \sum_{m, \alpha} q_m^\alpha \phi_\alpha z^m$ be an element of $\mathbb{H}_+$. Given a partition function $\mathcal{D}(\mathbf{q})$ on $\mathbb{H}_+$, the quantized operators act as follows. The quantization of an R -matrix R acts on $\mathcal{D}$ via*

$$\widehat{R}\mathcal{D}(\mathbf{q}) = \left[e^{\frac{\hbar}{2} V \left(\frac{\partial}{\partial \mathbf{q}}, \frac{\partial}{\partial \mathbf{q}} \right)} \mathcal{D} \right] (R^{-1} \mathbf{q})$$

where $R^{-1} \mathbf{q}$ is the power series $R^{-1}(z) \mathbf{q}(z)$, and the quadratic form $V = \sum_{k,l} (p_k, V_{kl} p_l)$ is defined via

$$V(w, z) = \sum_{k,l \geq 0} V_{kl} w^k z^l = \frac{1 - R(-w)^* R(-z)}{w + z},$$

where $*$ denotes the adjoint with respect to the intersection pairing on $H^*(X; \mathbb{C})$.

The action of the exponential of the quadratic differential operator in Theorem 16 has a Feynman graph expansion, and its logarithm only involves connected graphs (c.f. [CPS18]). Let Γ denote a connected dual graph consisting of vertices V , edges E , and legs L , with each vertex v labeled by a genus $g(v)$. We define the genus of the graph by $g(\Gamma) = h^1(\Gamma) + \sum_{v \in V} g(v)$ where h^1 denotes the first Betti number of the Feynman graph Γ . Denote G_g^{Feyn} to be the set of connected Feynman graphs of genus g . Feynman graphs differ from the stable graphs from Section 2.2.1 in that they allow genus zero vertices of valence 1 or 2.

Theorem 17. *Let*

$$R(z) = \exp\left(\sum_{i=1}^{\infty} \frac{B_{2i}}{(2i)(2i-1)} u^{2i-1} z^{2i-1}\right).$$

Then, for $g \geq 2$, the following holds for descendant Gromov–Witten invariants

$$[u^i] \sum_{\Gamma \in G_g^{\text{Feyn}}} \frac{1}{|\text{Aut}(\Gamma)|} \text{Cont}_{\Gamma} = \begin{cases} 0, & i > g \\ \frac{(-1)^g}{2^g} \cdot \langle\langle \ ; \mathbf{P}_g^g(\emptyset) \rangle\rangle_g^{\sim}(\mathbf{q}(z)) & i = g \end{cases} \quad (41)$$

where $[u^i]$ denotes taking the coefficient of u^i and the contribution Cont_{Γ} of a Feynman graph Γ is defined as a contraction of tensors:

- Each vertex v is assigned a tensor bracket

$$\langle\langle \ \rangle\rangle_{g(v)}^{\sim}(R^{-1}\mathbf{q})$$

with insertion coming from the half edges.

- Each edge e is assigned a bivector with descendants

$$\frac{\sum_{\alpha, \beta} (\eta^{\alpha\beta} \phi_{\alpha} \otimes \phi_{\beta} - \eta^{\alpha\beta} R^{-1}(\psi_{h(e)})(\phi_{\alpha}) \otimes R^{-1}(\psi_{h'(e)})(\phi_{\beta}))}{\psi_{h(e)} + \psi_{h'(e)}}$$

4.4. Connection to Pixton’s 3-spin relations. In 2011, Pixton [Pix12] proposed a large class of conjectural relations in the tautological ring of $\overline{\mathcal{M}}_{g,n}$, which extend the Faber–Zagier relations on the moduli space of smooth curves $\mathcal{M}_g$. In [PPZ15], these relations were proven to hold in cohomology by studying the cohomological field theory of Witten’s 3-spin class, and so these relations are now usually called *Pixton’s 3-spin relations*. In [Jan17], the relations were established in Chow via a study of the equivariant Gromov–Witten theory of $\mathbb{P}^1$. Pulling back to the moduli space of stable maps, Pixton’s 3-spin relations lead to a large class of differential equations for ancestor Gromov–Witten invariants of any target manifold.

This work leads to a natural question:

Question 18. Is there any connection between the Hodge class relations (41) and Pixton’s 3-spin relations?

To answer this question, we start by making a connection between Pixton’s 3-spin relations, and the Chern classes of the Hodge bundle. There is a canonical way of writing the λ_i in terms of boundary classes, ψ -classes and κ -classes, i.e. as elements of the strata algebra. As explained in [Pan18, §1.3], the Chern polynomial $\lambda(u)$ forms a one-dimensional semisimple CohFT $\Omega^{\lambda(u)}$. The Givental–Teleman classification for $\Omega^{\lambda(u)}$ yields the required canonical expression for $\lambda(u)$.

Proposition 19. *Pixton’s 3-spin relations on the moduli space of stable curves imply the vanishing $\lambda_i = 0$ for $i > g$. More precisely, as elements of the strata algebra, we may write λ_i as a linear combination of Pixton’s 3-spin relations.*

Proof. We use the results of [Jan18], which show that a large class of tautological relations from cohomological field theories are linear combinations of the 3-spin relations. In particular, the discussion of [Jan18, Section 3.8] applies to the CohFT $\Omega^{\mathbb{P}^1}$ of the degree zero $\mathbb{C}^*$ -equivariant Gromov–Witten theory of $\mathbb{P}^1$. We first analyze this CohFT, then relate it to the Hodge CohFT, and finally study their tautological relations.

The state space of $\Omega^{\mathbb{P}^1}$ is the equivariant cohomology

$$H_{\mathbb{C}^*}^*(\mathbb{P}^1) \cong \mathbb{C}[\mu][H]/(H(H - \mu)) \cong \mathbb{C}[\mu]\langle [0], [\infty] \rangle$$

viewed as a 2-dimensional free module over the base ring $\mathbb{C}[\mu]$. Here, $[0] = H$ and $[\infty] = H - \mu$ are the cohomology classes of the points 0 and ∞ in $\mathbb{P}^1$, respectively. Then $e_0 = [0]/\mu_0$ and $e_\infty = [\infty]/\mu_\infty$ for $\mu_0 = \mu$ and $\mu_\infty = -\mu$ form a basis of orthogonal idempotents. In particular, this CohFT is semisimple. In the basis e_0, e_∞ , the pairing is given by the matrix

$$\begin{pmatrix} 1/\mu_0 & 0 \\ 0 & 1/\mu_\infty \end{pmatrix}.$$

Note that there is the formula

$$\Omega_{g,n}^{\mathbb{P}^1}(\phi_1, \dots, \phi_n) = \sum_{i=0}^g \lambda_i \cdot \int_{\mathbb{P}^1} \prod_j \phi_j \cdot ([0] + [\infty])^{g-i}$$

for $\phi_1, \dots, \phi_n \in H_{\mathbb{C}^*}^*(\mathbb{P}^1)$. If we decompose $\phi_i = \phi_{i0}e_0 + \phi_{i\infty}e_\infty$, we may rewrite this as

$$\Omega_{g,n}^{\mathbb{P}^1}(\phi_1, \dots, \phi_n) = \sum_{\alpha \in \{0, \infty\}} \mu_\alpha^{g-1} \prod_j \phi_{j\alpha} \cdot \sum_{i=0}^g \lambda_i \mu_\alpha^{-i} = \sum_{\alpha \in \{0, \infty\}} \mu_\alpha^{g-1} \Omega_{g,n}^{\lambda(\mu_\alpha^{-1})}(\phi_{1\alpha}, \dots, \phi_{n\alpha}).$$

Thus, $\Omega^{\mathbb{P}^1}$ is a direct sum of two Hodge CohFTs.

There is a Givental–Teleman reconstruction

$$\Omega^{\lambda(u)} = R.\omega, \tag{42}$$

where

$$R(z) = \exp \left(\sum_{i=1}^{\infty} \frac{B_{2i}}{(2i)(2i-1)} u^{2i-1} z^{2i-1} \right)$$

and $\omega_{g,n}(a_1, \dots, a_n) = a_1 \cdots a_n$ for $a_1, \dots, a_n \in \mathbb{C}$. For $\Omega^{\mathbb{P}^1}$, the Givental–Teleman reconstruction

$$\Omega_{g,n}^{\mathbb{P}^1}(\phi_1, \dots, \phi_n) = R^{\mathbb{P}^1}.\omega_{g,n}^{\mathbb{P}^1}(\phi_1, \dots, \phi_n) \tag{43}$$

respects the direct sum structure, with R -matrix

$$R^{\mathbb{P}^1}(z) = \begin{pmatrix} R(z)|_{u=\mu_0^{-1}} & 0 \\ 0 & R(z)|_{u=\mu_\infty^{-1}} \end{pmatrix}$$

if written in the basis $\{e_0, e_\infty\}$, and topological field theory defined via

$$\omega_{g,n}^{\mathbb{P}^1}(\phi_1, \dots, \phi_n) = \sum_{\alpha \in \{0, \infty\}} \mu_\alpha^{g-1} \omega_{g,n}(\phi_{1\alpha}, \dots, \phi_{n\alpha})$$

The results of [Jan18, Section 3.8] imply that for any choice of $\phi_1, \dots, \phi_n$ (well-defined in the non-equivariant limit $\mu \rightarrow 0$), the polar terms in μ on the right hand side of (43) are a

linear combination of Pixton's 3-spin relations. If $n > 0$, note that because of the diagonal form of the pairing and $R^{\mathbb{P}^1}$, we have

$$\Omega_{g,n}^{\mathbb{P}^1}([0], 1, 1, \dots, 1) = \mu^g ((R.\omega)(1, \dots, 1))|_{u=\mu^{-1}},$$

and thus the polar terms correspond exactly to the terms in (42) of degree $i > g$. This establishes the proposition in the case $n > 0$.

The case $n = 0$ follows from the case $n = 1$ via the dilaton equation, and the fact that Pixton's relations are stable under pushing-forward via forgetful morphisms. $\square$

Remark 20. The result of Proposition 19 immediately implies that the differential equations for ancestor Gromov–Witten invariants from the vanishing $\lambda_i = 0$ for $i > g$ are linear combinations of those for Pixton's 3-spin relations. We expect a compatibility between the Givental–Teleman reconstruction, quantization of quadratic Hamiltonian, and the descendant/ancestor correspondence, however we were not able to find a reference for this fact. This compatibility would imply that in a strong sense, the $i > g$ part of (41) is a consequence of Pixton's 3-spin relations.

Remark 21. Given that Pixton's 3-spin relations are expected to be a complete set of relations in the tautological ring, we also expect the $i = g$ part of (41) to also be a consequence of Pixton's 3-spin relations. However, we do not know how to prove this. One difficulty is that the two sides of (41) have very different origins. On a basic level, the left hand side of (41) comes from the Grothendieck–Riemann–Roch computation of the λ_g -class, while the right hand side comes from relative Gromov–Witten theory.

Acknowledgements. The authors would like to thank the anonymous referees for careful reading of the manuscript and for the many helpful suggestions and comments. The first author was partially supported by NSF grants DMS-2054830 and DMS-2239320. The second author was partially supported by National Key R&D Program of China (2022YFA1006200), National Science Foundation of China (Grant No. 12071255) and Shandong Provincial Natural Science Foundation (Grant No. ZR2021MA101).

Declarations

Conflict of interest The authors declared that they have no conflicts of interest to this work.

REFERENCES

- [CG07] Tom Coates and Alexander Givental. Quantum Riemann–Roch, Lefschetz and Serre. *Ann. of Math. (2)*, 165(1):15–53, 2007.
- [CGT15] Tom Coates, Alexander Givental, and Hsian-Hua Tseng. Virasoro constraints for toric bundles. *arXiv preprint arXiv:1508.06282*, 2015.
- [CJWZ23] Emily Clader, Felix Janda, Xin Wang, and Dmitry Zakharov. Topological recursion relations from Pixton's formula. *Michigan Math. J.*, 73(2):227–241, 2023.
- [CPS18] Emily Clader, Nathan Priddis, and Mark Shoemaker. Geometric quantization with applications to Gromov–Witten theory. In *B-model Gromov–Witten theory*, Trends Math., pages 399–462. Birkhäuser/Springer, Cham, 2018.
- [CR02] Weimin Chen and Yongbin Ruan. Orbifold Gromov–Witten theory. In *Orbifolds in mathematics and physics (Madison, WI, 2001)*, volume 310 of *Contemp. Math.*, pages 25–85. Amer. Math. Soc., Providence, RI, 2002.

- [EHX97] Tohru Eguchi, Kentaro Hori, and Chuan-Sheng Xiong. Quantum cohomology and Virasoro algebra. *Phys. Lett. B*, 402(1-2):71–80, 1997.
- [FJR13] Huijun Fan, Tyler Jarvis, and Yongbin Ruan. The Witten equation, mirror symmetry, and quantum singularity theory. *Ann. of Math. (2)*, 178(1):1–106, 2013.
- [FP00a] Carel Faber and Rahul Pandharipande. Hodge integrals and Gromov-Witten theory. *Invent. Math.*, 139(1):173–199, 2000.
- [FP00b] Carel Faber and Rahul Pandharipande. Logarithmic series and Hodge integrals in the tautological ring. *Michigan Math. J.*, 48:215–252, 2000. With an appendix by Don Zagier, Dedicated to William Fulton on the occasion of his 60th birthday.
- [Giv01] Alexander Givental. Gromov-Witten invariants and quantization of quadratic Hamiltonians. *Mosc. Math. J.*, 1(4):551–568, 645, 2001. Dedicated to the memory of I. G. Petrovskii on the occasion of his 100th anniversary.
- [Giv04] Alexander Givental. Symplectic geometry of Frobenius structures. In *Frobenius manifolds*, volume E36 of *Aspects Math.*, pages 91–112. Friedr. Vieweg, Wiesbaden, 2004.
- [Jan17] Felix Janda. Relations on $\overline{M}_{g,n}$ via equivariant Gromov-Witten theory of $\mathbb{P}^1$. *Algebr. Geom.*, 4(3):311–336, 2017.
- [Jan18] Felix Janda. Frobenius manifolds near the discriminant and relations in the tautological ring. *Letters in Mathematical Physics*, 108(7):1649–1675, 2018.
- [JPPZ17] Felix Janda, Rahul Pandharipande, Aaron Pixton, and Dimitri Zvonkine. Double ramification cycles on the moduli spaces of curves. *Publications mathématiques de l’IHÉS*, 125(1):221–266, 2017.
- [KM97] Maxim Kontsevich and Yuri Manin. Gromov-Witten classes, quantum cohomology, and enumerative geometry. In *Mirror symmetry, II*, volume 1 of *AMS/IP Stud. Adv. Math.*, pages 607–653. Amer. Math. Soc., Providence, RI, 1997.
- [Liu02] Xiaobo Liu. Quantum product on the big phase space and the Virasoro conjecture. *Adv. Math.*, 169(2):313–375, 2002.
- [Liu11] Xiaobo Liu. On certain vanishing identities for Gromov-Witten invariants. *Trans. Amer. Math. Soc.*, 363(6):2939–2953, 2011.
- [LP11] Xiaobo Liu and Rahul Pandharipande. New topological recursion relations. *Journal of Algebraic Geometry*, 20(3):479–494, 2011.
- [LX09] Kefeng Liu and Hao Xu. A proof of the Faber intersection number conjecture. *J. Differential Geom.*, 83(2):313–335, 2009.
- [Mum83] David Mumford. Towards an enumerative geometry of the moduli space of curves. In *Arithmetic and geometry, Vol. II*, volume 36 of *Progr. Math.*, pages 271–328. Birkhäuser Boston, Boston, MA, 1983.
- [Pan18] Rahul Pandharipande. A calculus for the moduli space of curves. In *Algebraic geometry: Salt Lake City 2015*, volume 97.1 of *Proc. Sympos. Pure Math.*, pages 459–487. Amer. Math. Soc., Providence, RI, 2018.
- [Pix12] Aaron Pixton. Conjectural relations in the tautological ring of $\overline{M}_{g,n}$. *arXiv preprint arXiv:1207.1918*, 2012.
- [PPZ15] Rahul Pandharipande, Aaron Pixton, and Dimitri Zvonkine. Relations on $\overline{M}_{g,n}$ via 3-spin structures. *Journal of the American Mathematical Society*, 28(1):279–309, 2015.

UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN, URBANA, IL 61801, USA
Email address: fjanda@illinois.edu

SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY, JINAN, CHINA
Email address: wangxin2015@sdu.edu.cn