

Hybridization induced triplet superconductivity with $S^z = 0$

Edine Silva¹, R. C. Bento Ribeiro¹, Heron Caldas², and Mucio A. Continentino^{1*}

¹ *Centro Brasileiro de Pesquisas Físicas, Rua Dr. Xavier Sigaud,
150, Urca 22290-180, Rio de Janeiro, RJ, Brazil and*

² *Departamento de Ciências Naturais, Universidade Federal de São João Del Rei,
Praça Dom Helvécio 74, 36301-160, São João Del Rei, MG, Brazil*

The Kitaev superconducting chain is a model of spinless fermions with triplet-like superconductivity. It has raised interest since for some values of its parameters it presents a non-trivial topological phase that host Majorana fermions. The physical realization of a Kitaev chain is complicated by the scarcity of triplet superconductivity in real physical systems. Many proposals have been put forward to overcome this difficulty and fabricate artificial triplet superconducting chains. In this work we study a superconducting chain of spinful fermions forming Cooper pairs, in a triplet $S = 1$ state, but with $S^z = 0$. The motivation is that such pairing can be induced in chains that couple through an antisymmetric hybridization to an s-wave superconducting substrate. We study the nature of edge states and the topological properties of these chains. In the presence of a magnetic field the chain can sustain gapless superconductivity with pairs of Fermi points. The momentum space topology of these Fermi points is non-trivial, in the sense that they can only disappear by annihilating each other. For small magnetic fields, we find well defined degenerate edge modes with finite Zeemann energy. These modes are not symmetry protected and decay abruptly in the bulk as their energy merges with the continuum of excitations.

I. INTRODUCTION

The Kitaev chain, a prototypical one-dimensional toy model of spinless fermions and p -wave superconductivity is well known to host Majorana zero modes (MZMs) at the ends of the chain. These spinless fermions form Cooper pairs in a triplet state with $S = 1$ and a finite S^z component of the total spin. The MZMs have, in principle, a prospect of being applied to fault-tolerant topological quantum computation¹⁻³. However, p -wave superconductivity is very difficult to find in real materials^{4,5}. One realistic model, proposed to experimentally search for MZMs is the one-dimensional (1D) semiconductor-superconductor (SM-SC) nanowire⁶⁻¹³, where these modes are predicted to appear at the ends of the wire under appropriate, specific conditions. The Majorana nanowire consists of a semiconductor with a large Rashba-type SOC in proximity to a conventional s -wave superconductor under a magnetic (Zeeman spin splitting) field to achieve an effective p -wave superconductor⁵.

In this work we consider a different model of a p -wave superconducting chain. It has spinful fermions¹⁴⁻¹⁶ that form Cooper pairs in a triplet state, $S = 1$, but with $S^z = 0$. This type of p -wave pairing opens the possibility for obtaining new chiral superconductors with topological properties^{17,18}.

The main motivation for studying these chains is the following. When atoms are deposited on a superconducting substrate, in general its orbitals hybridize with those of the substrate¹⁹ providing a mechanism for inducing superconductivity in the chain. If the substrate is a BCS superconductor and the hybridization V_{ij} between the orbitals at site i of the substrate and at site j of the chain is antisymmetric, i.e., $V_{ji} = -V_{ij}$, or in momentum space $V_{-k} = -V_k$, the *induced superconductivity* in the chain is of the type $S = 1$, $S^z = 0$. The energy scale involved in this process can be significant¹⁹.

The mechanism that induces this type of superconductivity in the wire is similar to that occurring in multi-band superconductors²⁰⁻²⁶. Consider a two-band (a and b) superconductor with an attractive interaction in the a band that gives rise to a BCS pairing of the electrons in this band (substrate). An anti-symmetric hybridization V_k between these electrons and those in the non-interacting b -band induces a pairing gap with p -wave symmetry, of the type $S = 1$, $S^z = 0$ in the b -band (the chain)^{20,22,23}. The induced p -wave pairing $\Delta_{ind}^p(k)$, with $S = 1$, $S^z = 0$, is given by²⁰⁻²³, (see also note²⁷),

$$\Delta_{ind}^p(k) = \frac{V_k}{\sqrt{(\epsilon_k^b - \epsilon_k^a)^2 + 4|V_k|^2}} \Delta_a^s, \quad (1.1)$$

where the quasi-particles in the interacting a -band, with a dispersion relation ϵ_k^a in the normal state, condense in an s -wave BCS singlet superconducting state with an s -wave gap Δ_a^s . The dispersion of the non-interacting band is given by, ϵ_k^b . The anti-symmetric hybridization V_k between the orbitals of different parities in the interacting and non-interacting bands is responsible for an induced p -wave pairing $\Delta_{ind}^p(k)$, of the type $S = 1$, $S^z = 0$, as given by Eq. 1.1(see note²⁷). Notice that V_k in Eq. 1.1 is a *one-body term*, or simply, an inter-band hopping that transfer electrons between orbitals of different parities between the bands. This mechanism for induced superconductivity is different from the usual proximity effect that involves Cooper pair tunneling or Andreev reflections at the non-superconductor-superconductor boundary^{28,29}.

In this paper we consider a model that describes a BCS superconducting substrate on top of which is deposited a chain with non-interacting electrons. This kind of setup has already been implemented experimentally^{26,31}. The electrons in the chain, the non-interacting b -band, hybridize with those in the a -band of the superconducting substrate where the Cooper pairs are formed. If this hybridization is anti-symmetric, the induced superconductivity in the chain is unconventional and corresponds to the pairing Δ_{ind}^p in Eq. 1.1. In this equation, Δ_a^s is the BCS s -wave pairing of the substrate²⁷. The model is valid whenever the main coupling between the chain and substrate is through the hybridization between their orbitals. Notice that the hybridization, contrary to the spin-orbit interaction, has no spin-flip terms and the Cooper pairs in the chains preserve the anti-parallel coupling of the spins inherited from the BCS Cooper pairs of the substrate. The antisymmetric character of the total wave-function of the induced Cooper pair is conferred by the antisymmetric hybridization V_k that plays the role of the spatial dependent wave function of the pair and allows for a symmetric state of the spins²⁷. Then, the superconductivity induced in the chain is triplet $S = 1$, but with $S^z = 0$. Notice that the antisymmetric character of the hybridization arises when it mixes orbitals with angular momenta that differ by an odd number^{20,22,23}. This includes the important cases that the orbitals in the chain-substrate system have $s-p$ or $p-d$ character. For completeness the model also includes the spin-orbit coupling (SOC) in the wire and the effect of a magnetic field. In the case of SOC, this limits the validity of our results to the case that the spin-orbit interaction vanishes or is smaller than the induced superconducting parameter in the wire.

With the motivation of a physical mechanism to obtain an $S = 1$, $S^z = 0$, p -wave-superconducting chain, and its possible applications³⁰, we present here a study of the topological properties and excitations of such a chain. In the absence of a magnetic field and the conditions stated above for the SOC, the $S = 1$, $S^z = 0$ chain is topological, for a range of parameters. It presents four Majorana modes, two in each extremity of a finite chain. As the magnetic field

is turned on, these modes acquire a finite Zeeman energy. They are not symmetry protected and disappear as they merge with the continuum of Bogoliubov excitations for a sufficiently high magnetic field.

The $S = 1$, $S^z = 0$ chain, in the presence of a large magnetic field¹², presents gapless superconductivity due to the presence of pairs of Fermi points^{32,33}. The non-trivial momentum space topology of these Fermi points implies their stability, since they can only be destroyed by annihilating in pairs^{34,35}. The dispersion at the Fermi points is linear, like in Dirac points. This is in contrast with the Kitaev chain that is a gapped superconductor with Majorana edge modes.

The paper is organized as follows. In Section II we present the Hamiltonian of the model. In Section III we obtain the energy spectra of an infinite chain. In Section IV we study a finite chain described by the Hamiltonian written in terms of Majorana operators. The numerical results are shown in Section V. Section VI is devoted to the calculation of the topological invariant and topological indexes of the several phases of the model. We conclude in Section VII.

II. HAMILTONIAN

The Hamiltonian describing the chain of N sites and spinful fermions with a $S = 1$, $S^z = 0$, p -wave induced superconducting interaction is given by $\mathcal{H} = \mathcal{H}_N + \mathcal{H}_{\mathcal{HS}}$, where

$$\begin{aligned}\mathcal{H}_N &= -\mu \sum_{j=1,\sigma}^N c_{j,\sigma}^\dagger c_{j,\sigma} - h \sum_{j=1,\sigma}^N \sigma c_{j,\sigma}^\dagger c_{j,\sigma} - \sum_{j=1,\sigma}^{N-1} \left(t c_{j,\sigma}^\dagger c_{j+1,\sigma} + h.c. \right) + i\lambda \sum_{j=1,\sigma,\bar{\sigma}}^{N-1} \left(c_{j,\sigma}^\dagger (\sigma^y)_{\sigma\bar{\sigma}} c_{j+1,\bar{\sigma}} + h.c. \right) \\ \mathcal{H}_{\mathcal{HS}} &= -\frac{1}{2} \sum_{j=1,\sigma}^{N-1} \left(\Delta (c_{j,\sigma}^\dagger c_{j+1,-\sigma}^\dagger - c_{j+1,\sigma}^\dagger c_{j,-\sigma}^\dagger) + h.c. \right).\end{aligned}\quad (2.1)$$

The first equation describes the normal chain in the presence of a uniform external magnetic field h parallel to the wire^{7,8,13}. The quantity μ is the chemical potential, t a nearest neighbor hopping. In our strict one-dimensional model, the Rashba-like term is essentially an antisymmetric spin-flip hopping λ due to the spin-orbit interaction^{5,6,11,36,37}.

The Hamiltonian $\mathcal{H}_{\mathcal{HS}}$ represents the induced superconductivity in the chain due to its hybridization with the BCS superconducting substrate (see Eq. 1.1). Since this hybridization is antisymmetric, Δ is the induced antisymmetric superconducting pairing ($\Delta_{ij} = -\Delta_{ji} = \Delta$) between fermions with antiparallel spins in neighboring sites of the wire. In momentum space it is defined by Eq. 1.1, where we removed all indices for simplicity. N is the number of sites in the chain, and $\sigma = \pm$, corresponds to spin up and spin down, respectively. The total antisymmetry of the order parameter with antiparallel spins is guaranteed by the spatial antisymmetric wave function of the Cooper pairs. Then the induced superconductivity is a triplet state with $S = 1$, but with the z -component of the total spin of the Cooper pair, $S^z = 0$. The term $h.c.$ stands for Hermitian conjugate.

III. INFINITE CHAIN

For an infinite chain with periodic boundary conditions we can Fourier transform Hamiltonian Eq. (1.1) in momentum space. We choose the basis $\Psi_{\mathbf{k}} = (c_{\mathbf{k},\sigma}^\dagger, c_{\mathbf{k},-\sigma}^\dagger, c_{-\mathbf{k},\sigma}, c_{-\mathbf{k},-\sigma})^T$, with $\sigma = \uparrow$ and $-\sigma = \downarrow$ to obtain,

$$H = \frac{1}{2} \sum_{\mathbf{k}} \Psi_{\mathbf{k}}^\dagger \mathcal{H}(\mathbf{k}) \Psi_{\mathbf{k}} + \frac{1}{2} \sum_{\mathbf{k}} [\varepsilon_{\mathbf{k}\uparrow} + \varepsilon_{-\mathbf{k}\downarrow}], \quad (3.1)$$

where $\varepsilon_{\mathbf{k}\uparrow\downarrow} = -(2t \cos k + \mu \pm h)$, $\lambda_k = 2i\lambda \sin(k)$ and $\Delta_k = 2i\Delta \sin k$,

$$\mathcal{H}(k) = \begin{pmatrix} \varepsilon_{\mathbf{k}\sigma} & \lambda_k & 0 & \Delta_k^* \\ \lambda_k^* & \varepsilon_{\mathbf{k}-\sigma} & \Delta_k^* & 0 \\ 0 & \Delta_k & -\varepsilon_{-\mathbf{k}\sigma} & -\lambda_{-k}^* \\ \Delta_k & 0 & -\lambda_{-k} & -\varepsilon_{-\mathbf{k}-\sigma} \end{pmatrix}. \quad (3.2)$$

Notice that $\lambda_{-k}^* = \lambda_k$, with a similar relation for Δ_k . We remark that in the present Hamiltonian, the gap parameter is k -dependent and has p -wave symmetry, distinctively from other previous approaches, as for instance, in Refs.^{8,9,33}.

The Hamiltonian, Eq. 3.2, can be written as $\mathcal{H} = (\epsilon_k - \mu)\sigma_z \otimes \tau_0 - h\sigma_z \otimes \tau_z + i\lambda_k\sigma_z \otimes \tau_y - i\Delta_k\sigma_y \otimes \tau_x$, where σ_i and τ_i are Pauli matrices ($i = x, y, z$) and $\tau_0 = \sigma_0$ are the 2×2 identity matrix and $\epsilon_k = -2t \cos k$. It has particle-hole




FIG. 1. (Color online) Gap closing lines. At the blue lines, $h_{\pm} \pm 2t \pm \mu$ the gap closes at the time-reversal invariant momenta $k = 0$ and $k = \pm\pi$. Along the red (dashed) lines, the gap closes at $k = \pm\pi/2$ (we consider only the case, $\lambda < \Delta$). In the figure we chose t as the unity of energy ($t = 1$). The red dashed lines correspond to $\lambda = 0.1$ and $\Delta = 0.5$.

symmetry since $U_p \mathcal{H}^*(k) U_p^\dagger = -\mathcal{H}(-k)$, where $U_p = i\sigma_x \otimes \sigma_0$. On the other hand, we have, $U_t \mathcal{H}^*(k) U_t^\dagger \neq \mathcal{H}(-k)$, with $U_t = i\sigma_0 \otimes \sigma_y$, implying that the system is not time-reversal invariant. Time-reversal invariance is broken by the magnetic field, but also by the triplet pairing of the quasi-particles. The Hamiltonian, Eq. 3.2, has an additional chiral symmetry that will be discussed further on.

The Hamiltonian of the infinite chain can be diagonalized and the dispersion relations of the quasi-particles are given by,

$$\begin{aligned}\omega_1(k) &= \sqrt{\epsilon_k^2 + h^2 + \lambda_k^2 + \Delta_k^2 + 2\sqrt{(h^2 + \lambda_k^2)(\epsilon_k^2 + \Delta_k^2)}} \\ \omega_2(k) &= \sqrt{\epsilon_k^2 + h^2 + \lambda_k^2 + \Delta_k^2 - 2\sqrt{(h^2 + \lambda_k^2)(\epsilon_k^2 + \Delta_k^2)}},\end{aligned}\quad (3.3)$$

$\omega_3(k) = -\omega_1(k)$ and $\omega_4(k) = -\omega_2(k)$. These dispersions have several *gap closing lines*, as shown in Fig.1. Closing of the gaps occur at the time-reversal invariant points of the Brillouin zone, $k = 0$ and $k = \pm\pi$, but also for $k = \pm\pi/2$. In the former case the critical fields at which the gap closes are given by, $h_c = \pm 2t \pm \mu$ and are independent of the other parameters of the model.

The critical fields for gap closing at $k = \pm\pi/2$ are given by, $h_c = \pm\sqrt{4(\Delta^2 - \lambda^2) + \mu^2}$ and depend on Δ and λ , besides the chemical potential μ . We can distinguish two cases, $\lambda > \Delta$ and $\lambda < \Delta$, but here we consider only the case of small spin orbit coupling, $\lambda < \Delta$. Notice that for $\mu = 0$, and assuming $\lambda = 0$, for simplicity, the first gap closing with increasing field occurs for $k_{\pm} = \pm\pi/2$ and $h_c = \pm 2\Delta$ ($\Delta < 2t$). This vanishing of the gap at a wavevector that is not time-reversal invariant is particularly interesting and we will explore this case numerically further on. We will also discuss the topological nature of the different phases of the model.

IV. FINITE SIZE CHAINS

For the purpose of studying the edge modes in the system, we consider a finite size chain with N sites and open boundary conditions. We neglect spin-orbit coupling ($\lambda = 0$), for simplicity, in this section. We rewrite Eq. 1.1 in terms of new real operators

$$\begin{aligned}c_{j,\sigma} &= \frac{1}{2}(\gamma_{j,\sigma}^B + i\gamma_{j,\sigma}^A) \\ c_{j,\sigma}^\dagger &= \frac{1}{2}(\gamma_{j,\sigma}^B - i\gamma_{j,\sigma}^A)\end{aligned}\quad (4.1)$$

where the Majorana operators satisfy, $\gamma_{j,\sigma}^\beta = \gamma_{j,\sigma}^{\beta\dagger}$, $\{\gamma_{i,\sigma}^\beta, \gamma_{j,\sigma'}^\beta\} = \delta_{ij,\sigma\sigma'}$ and $\gamma_{j,\sigma}^\beta \gamma_{j,\sigma}^\beta = 1$ ($\beta = A, B$). The symbol σ in $\gamma_{j,\sigma}^B$ does not mean a Majorana of spin σ . It is only a label to distinguish the various operators, since we need two Majoranas to represent an electron of spin up and two for an electron of spin down.

In terms of these new operators the Hamiltonian can be rewritten as,

$$\begin{aligned} \mathcal{H} = & -\frac{1}{2} \sum_{j=1,\sigma}^N (\mu + \sigma h) (1 + i\gamma_{j,\sigma}^B \gamma_{j,\sigma}^A) - \frac{it}{4} \sum_{j=1,\sigma}^{N-1} (\gamma_{j,\sigma}^B \gamma_{j+1,\sigma}^A - \gamma_{j,\sigma}^A \gamma_{j+1,\sigma}^B) - \\ & \frac{i\Delta}{4} \sum_{j=1,\sigma}^{N-1} (\gamma_{j,\sigma}^B \gamma_{j+1,-\sigma}^A + \gamma_{j,\sigma}^A \gamma_{j+1,-\sigma}^B). \end{aligned} \quad (4.2)$$

Next, we introduce more four Majorana operators given by,

$$\begin{aligned} \alpha_{j,\sigma}^{A\pm} &= \gamma_{j,\sigma}^A \pm \gamma_{j,-\sigma}^A \\ \alpha_{j,\sigma}^{B\pm} &= \gamma_{j,\sigma}^B \pm \gamma_{j,-\sigma}^B. \end{aligned} \quad (4.3)$$

Notice that

$$\begin{aligned} \alpha_{j,\sigma}^{A\pm} &= \pm \alpha_{j,-\sigma}^{A\pm} \\ \alpha_{j,\sigma}^{B\pm} &= \pm \alpha_{j,-\sigma}^{B\pm}. \end{aligned} \quad (4.4)$$

Finally, in term of these new operators, the Hamiltonian of the superconducting chain can be written as,

$$\begin{aligned} \mathcal{H} = & -\frac{i\mu}{4} \sum_{j=1}^N (\alpha_{j,\uparrow}^{B+} \alpha_{j,\uparrow}^{A+} + \alpha_{j,\uparrow}^{B-} \alpha_{j,\uparrow}^{A-}) - \frac{i}{8} \sum_{j=1}^{N-1} (t + \Delta) (\alpha_{j,\uparrow}^{B+} \alpha_{j+1,\uparrow}^{A+} - \alpha_{j,\uparrow}^{A-} \alpha_{j+1,\uparrow}^{B-}) \\ & - \frac{i}{8} \sum_{j=1}^{N-1} (t - \Delta) (\alpha_{j,\uparrow}^{B-} \alpha_{j+1,\uparrow}^{A-} - \alpha_{j,\uparrow}^{A+} \alpha_{j+1,\uparrow}^{B+}) - \frac{ih}{4} \sum_{j=1}^N (\alpha_{j,\uparrow}^{B+} \alpha_{j,\uparrow}^{A-} + \alpha_{j,\uparrow}^{B-} \alpha_{j,\uparrow}^{A+}), \end{aligned} \quad (4.5)$$

where the sum over σ has been performed and expressed in terms of $\sigma = \uparrow$. This index $\sigma = \uparrow$ now becomes redundant but we keep it anyway. The α -operators are such that,

$$\begin{aligned} \alpha_{\ell,\uparrow}^{A\pm} &= \gamma_{\ell,\uparrow}^A \pm \gamma_{\ell,\downarrow}^A \\ \alpha_{\ell,\uparrow}^{B\pm} &= \gamma_{\ell,\uparrow}^B \pm \gamma_{\ell,\downarrow}^B. \end{aligned} \quad (4.6)$$

The Hamiltonian Eq. 4.5 describes two independent (\pm) sub-chains that are coupled by the magnetic field term, the last term in Eq. 4.5 (see Fig 2). Notice from Eqs. 4.4 and Eq. 4.5 that the magnetic field term breaks time-reversal symmetry, as it is not invariant under the change $\sigma \rightarrow -\sigma$.




FIG. 2. (Color online) Eq. 4.7 describes two independent sub-chains (\pm) in terms of the operators α^{\pm} . The figure represents a topological phase, with Majoranas at the edges of each sub-chain.

For $t = \Delta = 0$, the Hamiltonian, Eq. 4.5, describes trivial chains coupled by the magnetic field. On the other hand, if we take $\mu = h = 0$ and $t = \Delta$, we obtain,

$$\mathcal{H} = \frac{-i\Delta}{4} \sum_{j=1}^{N-1} (\alpha_{j,\uparrow}^{B+} \alpha_{j+1,\uparrow}^{A+} - \alpha_{j,\uparrow}^{A-} \alpha_{j+1,\uparrow}^{B-}). \quad (4.7)$$

Now the system consists of two decoupled \pm chains, as shown schematically in Fig. 2. The Majoranas $\alpha_{1,\uparrow}^{B-}$ and $\alpha_{1,\uparrow}^{A+}$ at the beginning of these chains do not enter the Hamiltonian, as also the Majoranas $\alpha_{N,\uparrow}^{B+}$ and $\alpha_{N,\uparrow}^{A-}$ at the ends of the chains. These are the zero energy edge modes that signal the existence of a non-trivial topological phase in the superconducting chain. Notice that the \pm chains are *not* associated with a given spin direction. As we show below,

these zero modes persist for $|\mu/2t| < 1$, which characterizes the topological phase of the system in the absence of a magnetic field ($h = 0$).

Then, for $h = \mu = 0$ and $t = \Delta$, the system is formed of two independent Kitaev-like chains, each with two Majoranas. We can combine the Majoranas at the edges of each chain to obtain,

$$\begin{aligned} g^- &= \alpha_{1,\uparrow}^{B-} + i\alpha_{N,\uparrow}^{A-} \\ g^+ &= \alpha_{N,\uparrow}^{B+} + i\alpha_{1,\uparrow}^{A+}, \end{aligned} \quad (4.8)$$

to form a pair of non-local fermions, one in each sub-chain. Then, in the absence of a magnetic field, which couples the \pm sub-chains, the latter are the zero energy edge modes of the system. In terms of the original fermions operators, we have

$$g^- = (c_{1,\uparrow} - c_{N,\uparrow}^\dagger) - (c_{1,\downarrow}^\dagger + c_{N,\downarrow}) - \left((c_{1,\downarrow} - c_{N,\downarrow}^\dagger) - (c_{1,\uparrow}^\dagger + c_{N,\uparrow}) \right). \quad (4.9)$$

We recall that in the Kitaev spinless p-wave superconducting chain, with $S = 1$, $S^z = 1$, the non-local fermion is given by³⁸,

$$f_K = (c_1 - c_N^\dagger) - (c_1^\dagger + c_N). \quad (4.10)$$

In the presence of a magnetic field, we can combine two Majoranas, either in the same sites or on different edges, to form the fermions. In the former case we have,

$$\begin{aligned} g_1 &= \alpha_{1,\uparrow}^{B-} + i\alpha_{1,\uparrow}^{A+} \\ g_N &= \alpha_{N,\uparrow}^{B+} + i\alpha_{N,\uparrow}^{A-}, \end{aligned} \quad (4.11)$$

or in terms of fermion operators,

$$\begin{aligned} g_1 &= 2(c_{1,\uparrow} - c_{1,\downarrow}^\dagger) \\ g_N &= 2(c_{N,\uparrow} + c_{N,\downarrow}^\dagger). \end{aligned} \quad (4.12)$$

These modes are present for $|\mu/2t| < 1$, for $0 < h < h^*$, i.e., for finite fields but below a critical field h_c , as we show below. They are localized at the edges of the chain and have a finite Zeemann energy. They are not symmetry protected, as they exist in a region of the phase diagram that is topologically trivial, as evidenced by the topological invariant calculated further on.

V. NUMERICAL RESULTS

In this section, we discuss the dispersion relations for an infinite chain and obtain numerical results on finite chains of size N with open boundary conditions, in the presence of SOC and magnetic field. We consider two different situations that are distinguished by the points of the Brillouin zone at which the gap closes. This may occur either at the time-reversal invariant momenta, $k = 0$, $k = \pm\pi$, or at the points $k = \pm\pi/2$.

A. Gap closing at $k = \pm\pi/2$

First, for the infinite chain, with $\mu = \lambda = 0$, the superconducting gap decreases with increasing magnetic field and finally closes at the non-time-reversal wave-vectors, $k = \pm\pi/2$ at a critical magnetic field $h_c = 2\Delta$, as shown in the dispersion relations of Fig. 3. For magnetic fields $2\Delta < h < 2t$, the system enters a topological phase characterized by the presence of pairs of monopoles^{32,33,39} at the field-dependent wave-vectors $k = \pm k_1$ and $k = \pm k_2$, as shown in Fig. 3b. Finally, for $h = 2t$ the monopoles annihilate each other at the time-reversal k-points of the Brillouin zone, Fig. 3c. These results are not qualitatively affected by the presence of the spin-orbit interaction, as can be seen in Fig. 3, where it is taken finite. In the next section we will give a detailed analysis of the topological nature of this phase. Notice that we are neglecting the possibility of any finite momentum pairing, since the superconductivity in the wire is induced by the s-wave substrate.

For finite chains at $h = 0$, we confirm numerically, the presence of four zero energy Majorana modes, two at each edge of the chain, for $|\mu/2t| \leq 1$. Their wave functions are localized at the ends of the chain. This is expected since




FIG. 3. (Color online) a) Dispersion relations for $t = 1$, $\mu = 0$, $\lambda = 0.15$ and $\Delta = 0.5$ a) gap closing at $k = \pi/2$ for the critical field $h_c = \sqrt{4(\Delta^2 - \lambda^2)}$. b) Two pairs of monopoles in the non-trivial topological phase for $h_c < h < 2t$ ($h = 1.2$). c) For $h = 2t$ the two pairs of monopoles annihilate at $k = 0$, and $k = \pm\pi$. The color code indicates the degree of superposition between particles and hole states.

in this limit of $h = 0$, the system consists of two decoupled Kitaev chains, as we obtained before. Then, for $h = 0$ we find a topological phase with zero energy edge modes for $|\mu/2t| < 1$.

As we turn on the magnetic field in the finite chain, the four edge modes persist in the presence of a small field, but now they are separated in two double degenerate modes with finite energies, as shown in Fig. 4. The lower energy states correspond to the Zeemann energy of an electron with spin parallel to the field and the excited state, with positive energy, to that of an electron with spin antiparallel to the field. The wave functions of these edge modes decay in the bulk with a penetration length which is nearly field independent and corresponds to that of the system in the absence of the field. Further increasing the magnetic field ($\mu = \lambda = 0$), as it reaches the value, $h^* = \Delta$, i.e., *before the gap closes*, the energy of the local modes merge with the continuum of Bogoliubon excitations and their wave functions become abruptly delocalized. The closest analogy of this phenomenon is the Chandrasekhar-Clogston limit in conventional superconductors⁴⁰ where the magnetic field is screened up to a critical field beyond which it penetrates abruptly in the bulk.

Further increasing the field, the gap decreases and closes for a critical field $h_c = \pm\sqrt{4(\Delta^2 - \lambda^2) + \mu^2}$, in the presence of SOC ($\lambda < \Delta$). The effect of SOC in this case is to compete with the superconducting gap, reducing it. Finally, for $h > h_c$, Fermi points appear in the system. These Fermi points are topologically protected as we discuss further on. Notice in Fig. 4a that the spectra are always particle-hole symmetric and this holds for any values of the parameters of the Hamiltonian, in agreement with what was obtained previously. Further on, we will discuss the existence of a hidden chiral symmetry of the Hamiltonian.

It is interesting to investigate how the edge modes give rise to a full electronic spin. They can combine either on




FIG. 4. (Color online) Spectrum of eigenvalues for a chain of 300 sites with $\mu = 0$, $\Delta = 0.6$, $\lambda = 0.1$ and $h = 0.2$, in units of the hopping t ($t = 1$). In b) the energy of the 4 edge modes, with two degenerate, is emphasized. In c) the wave functions of the modes shown in b (see also Fig. 5).

the same edge or on different edges as a delocalized quasi-particle. The numerical results in Fig. 5 show that the wave functions of the modes with negative energy, i.e., with spins aligned parallel to the field are localized on *two opposite edges of the chain*. Consequently, the wave function of the *electron* with spin up, parallel to the magnetic field, is delocalized with equal weight on different edges of the chain, as shown in Fig. 5. The same holds for the excited state with spin down. Notice also in Fig. 5 that the energy of two edge modes are required to give the Zeeman energy of a single electron.

Finally it is shown in Fig. 6, that for large enough fields, local modes reappear when their Zeemann energy falls in a gap of the spectra of bogoliubons.

B. Gap closing at time-reversal wave vectors

The gap closing phenomena that occurs at the time-reversal invariant wave-vectors in the present study are associated with the annihilation of monopoles, shown in Fig. 3c. Monopoles of different charges annihilate each other exactly at these points, $k = 0$ and $k = \pm\pi$. These monopoles, or Fermi points are non-trivial topological objects in momentum space³². They are characterized by a topological charge, given by a winding number³².

For $h = 0$ we observe gap closing as a function of the chemical potential for $\mu = \pm 2t$ at the time-reversal point $k = 0$. Also for $h = 0$ the spin-orbit interaction can promote gap closing for $\lambda = \Delta$. However, we restrict our study here for the case $\lambda < \Delta$.

VI. TOPOLOGICAL INVARIANT

In order to characterize the different phases and transitions at the gap closing points, we calculate topological indices for our problem. First we seek for a chiral symmetry of the Hamiltonian, Eq. 3.2. We want to find an operator K , i.e., a matrix K that anticommutes with \mathcal{H} and, such that it satisfies $K.K = 1$. Imposing these conditions, $\{K, \mathcal{H}\} = 0$ and $K^2 = 1$, we find they can be satisfied with $K = \sigma_x \otimes \sigma_0$. Applying the same unitary transformation UKU^T that




FIG. 5. (Color online) The energy of the edge modes aligned with the magnetic field and their wave functions, for $\mu = 0$, $\Delta = 0.6$, $\lambda = 0.1$ and $h = 0.4$. These modes combine to form a single quasi-particle with the correct total Zeemann energy. This quasi-particle is delocalized since its wave function resides in the two opposite edges of the chain.




FIG. 6. (Color online) As a new gap opens and the Zeemann energy falls in a gap, the edge modes reappear. In this case at the trivial topological phase with $\mathcal{M} = 0$, for $t = 1$, $h = 2.3$, $\Delta = 0.5$, $\lambda = 0.1$ and $\mu = 0$ (see Fig. 7).

diagonalizes the matrix K , to the Hamiltonian \mathcal{H} , we obtain, $\mathcal{H}' = U\mathcal{H}U^T$ as,

$$\mathcal{H}' = \begin{bmatrix} 0 & A \\ A^* & 0 \end{bmatrix}$$

where,

$$A = \begin{bmatrix} -2(\epsilon_k + h - \mu) & -2(\Delta_k + \lambda_k) \\ -2(\Delta_k - \lambda_k) & -2(\epsilon_k - h - \mu) \end{bmatrix}$$

with $\epsilon_k = -2t \cos k$, $\Delta_k = 2i\Delta \sin k$ and $\lambda_k = 2i\lambda \sin k$. This chiral symmetry of the Hamiltonian is due to the particle-hole symmetry of the spectrum that we observe for all values of the parameters, even in the presence of a magnetic field.

In order to calculate the topological invariant we first obtain,

$$G(k) = \frac{\partial \ln \det[A(k)]}{\partial k}, \quad (6.1)$$

which is given by,

$$G(k) = \frac{-4t\mu \sin k - 4(t^2 - (\Delta^2 - \lambda^2)) \sin 2k}{-h^2 + \mu^2 + 4t \cos k(\mu + t \cos k) + 4(\Delta^2 - \lambda^2) \sin^2 k}. \quad (6.2)$$

Introducing,

$$M(k) = \frac{1}{2\pi i} \int G(k) dk \quad (6.3)$$

and performing the integration, we get

$$M(k) = \frac{1}{2\pi} \Im m \left[\ln(-h^2 + \mu^2 + 4t \cos k(\mu + t \cos k) + 4(\Delta^2 - \lambda^2) \sin^2 k) \right]. \quad (6.4)$$

The topological invariant is obtained from,

$$\mathcal{M} = 2[M(\pi) - M(0)]. \quad (6.5)$$

We get,

- $\mathcal{M} = 1$, for $(2t - \mu)^2 < h^2$ and $(2t + \mu)^2 > h^2$.
- $\mathcal{M} = -1$, for $(2t + \mu)^2 < h^2$ and $(2t - \mu)^2 > h^2$
- $\mathcal{M} = 2$, for $h = 0$ and $\mu < 2t$.
- $\mathcal{M} = 0$, for $h = 0$ and $\mu > 2t$.

For $\lambda = h = 0$ and $|\mu/2t| < 1$ we find $\mathcal{M} = 2$, due to the two pairs of Majoranas in the uncoupled chains, as we obtained previously, and in agreement with simulations.




FIG. 7. (Color online) Topological indexes \mathcal{M} of the different regions of the phase diagram. The red line, corresponding to $h = 0$, $|\mu/2t| < 1$, has $\mathcal{M} = 2$. To obtain the gap-closing lines at $(\pi/2)$, we used $\Delta = 0.5$ and $\lambda = 0.1$, in units of t . Phases with topological indexed (1,-1) and (2,-2) are gapless phases with one and two pairs of Fermi points, respectively.

Since the gap may also close at $k = \pi/2$, we have additionally^{36,37},

$$\mathcal{M} = 2M(\pi/2) - M(\pi) - M(0) \quad (6.6)$$

that yields,

- $\mathcal{M} = 2$, for $(2t \pm \mu)^2 > h^2$ and $h^2 > 4(\Delta^2 - \lambda^2) + \mu^2$.
- $\mathcal{M} = 1$, for $h^2 > \mu^2 + 4(\Delta^2 - \lambda^2)$ and

$$(2t + \mu)^2 > h^2 > (2t - \mu)^2 \text{ or}$$

$$(2t - \mu)^2 > h^2 > (2t + \mu)^2$$
- $\mathcal{M} = 0$, for $h^2 > \mu^2 + 4(\Delta^2 - \lambda^2)$ and $h^2 < (\mu \pm 2t)^2$.

In these cases the indexes are not universal in the sense that the phase boundaries depend on the parameters Δ and λ . The latter competes with superconductivity and it's role is to renormalize (reduce) the superconducting gap. The topological indexes of the different phases are shown in Fig. 7.

It is interesting to look at the phase diagram in Fig. 7 together with the dispersion relations shown in Fig. 3b. These relations were obtained for $\mu = 0$ and $2t > h > \sqrt{4(\Delta^2 - \lambda^2)}$. This region of the phase diagram is associated with a topological index $\mathcal{M} = 2$ that in this case counts the number of pairs of topological Fermi points in this phase. The topological charge of a Fermi point is characterized by a winding number, similar to the one we calculated. It is integrated in a closed contour in momentum space that embraces the Fermi point^{32,33}. In a one-dimensional system, this involves integrating all along the Brillouin zone, as in the calculation of the winding number, Eq. 6.3.




FIG. 8. (Color online) Region of the phase diagram, in color, where protected Majorana edge modes are observed. It is characterized by a topological index $\mathcal{M} = 2$, and requires that $h = 0$ and $|\mu/2t| < 1$. The uncolored region, $\lambda > \Delta$, has no Majorana edge modes.

It is also relevant that the region of the phase diagram with $\mu = 0$ and $h < \Delta$, where we observed magnetically polarized edge modes, is topologically trivial, with $\mathcal{M} = 0$. This implies that these modes are not symmetry protected.

Finally, Fig.8 shows the region of the phase diagram where protected Majorana modes are observed. They occur only at $h = 0$, for $|\mu/2t| < 1$ and $\lambda < \Delta$.

VII. CONCLUSIONS

In this work we presented a study of the topological properties of a superconducting chain with electrons pairing in a spin triplet state $S = 1$, but with the z -component $S^z = 0$. The motivation for this study is that this type of pairing can be induced in a chain whose orbitals have an anti-symmetric hybridization with those of a BCS s -wave superconducting substrate. Alternative pairings to the simple spinless case studied by Kitaev have been considered to obtain topological superconductors. In particular, equal spin pairing⁴¹ and d -wave pairing^{36,37} were proposed. Here we studied the case $S = 1$, $S^z = 0$ with a concrete physical motivation. We have obtained the symmetry properties and calculated the topological indexes of the model and obtained a rich phase diagram with trivial and topological phases distinguished by these indexes.

We pointed out the existence of a topological phase, in the absence of a magnetic field, with four Majorana modes, two in each edge of the finite chain. In small finite fields $h < \Delta$, these modes become spin polarized. Each pair of edge modes gives rise to a full electronic spin that can align parallel or antiparallel to the magnetic field. A full polarized electronic spin has a wave-function on both edges of the chain and corresponds to a delocalized quasi-particle. These edge modes are not symmetry protected, as they occur in a region of the phase diagram characterized by a trivial

topological index, $\mathcal{M} = 0$. For magnetic fields $h > \Delta$ these edge modes disappear as they merge with the spectrum of bogoliubons. This occurs before the gap closes at $h = 2\Delta$.

For $h > 2\sqrt{\Delta^2 - \lambda^2}$ the topological excitations are in momentum space and correspond to pairs of Fermi points. At these points the dispersion relations are linear in momentum. These Fermi points occur in a large region of parameters of the phase diagram. They appear in pairs (Weyl fermions) and can only be destroyed by annihilating each other^{32,33} at the time-reversal wave-vectors of the Brillouin zone. We have found nontrivial phases⁴² with a single pair, $\mathcal{M} = 1$ and two pairs of Fermi points, $\mathcal{M} = 2$.

ACKNOWLEDGMENTS

E. Silva would like to thank CAPES for financial support. R.C.B.R would like to thank FAPERJ and CAPES, for financial support. H.C. would like to thank FAPEMIG for partial financial support. H.C. also thanks the Centro Brasileiro de Pesquisas Físicas-CBPF, where part of this work was done, for the kind hospitality. M.A.C would like to thank CNPq and FAPERJ for partial financial support and Tobias Micklitz and Wei Chen for useful discussions.

-
- * corresponding author: muciocontinentino@gmail.com
- ¹ C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, Non-Abelian anyons and topological quantum computation, *Reviews of Modern Physics* **80**, 1083 (2008).
 - ² S. Das Sarma, M. Freedman, and C. Nayak, Majorana zero modes and topological quantum computation, *npj Quantum Information* **1**, 15001 (2015).
 - ³ J. Alicea, Y. Oreg, G. Refael, F. von Oppen, and M. P. A. Fisher, Non-Abelian statistics and topological quantum information processing in 1D wire networks, *Nature Physics* **7**, 412 (2011).
 - ⁴ P.-H. Chou, C.-H. Chen, S.-W. Liu, C.-H. Chung, and C.-Y. Mou, Geometry-induced topological superconductivity, *Phys. Rev. B* **103**, 014508 (2021).
 - ⁵ H. Pan, S. Das Sarma, Majorana nanowires, Kitaev chains, and spin models, *Phys. Rev. B* **107**, 035440 (2023).
 - ⁶ M. Aghaee et al., InAs-Al Hybrid Devices Passing the Topological Gap Protocol (2022).
 - ⁷ R. M. Lutchyn, J. D. Sau, and S. Das Sarma, Majorana Fermions and a Topological Phase Transition in Semiconductor-Superconductor Heterostructures, *Phys. Rev. Lett.* **105**, 077001 (2010).
 - ⁸ Y. Oreg, G. Refael, and F. von Oppen, Helical Liquids and Majorana Bound States in Quantum Wires, *Phys. Rev. Lett.* **105**, 177002 (2010).
 - ⁹ J. D. Sau, R. M. Lutchyn, S. Tewari, and S. Das Sarma, Generic New Platform for Topological Quantum Computation Using Semiconductor Heterostructures, *Phys. Rev. Lett.* **104**, 040502 (2010).
 - ¹⁰ J. D. Sau, S. Tewari, R. M. Lutchyn, T. D. Stanescu, and S. Das Sarma, Non-Abelian quantum order in spin-orbit-coupled semiconductors: Search for topological Majorana particles in solid-state systems, *Phys. Rev. B* **82**, 214509 (2010).
 - ¹¹ S. Tewari, T. D. Stanescu, J. D. Sau, and S. Das Sarma, Topological minigap in quasi-one-dimensional spin-orbit-coupled semiconductor Majorana wires, *Phys. Rev. B* **86**, 024504 (2012).
 - ¹² Panagiotis Kotetes, Classification of engineered topological superconductors, *New Journal of Physics* **15**, 105027 (2013).
 - ¹³ T. D. Stanescu and S. Tewari, Majorana fermions in semiconductor nanowires: fundamentals, modeling, and experiment, *J. Phys.: Condens. Matter* **25**, 233201 (2013).
 - ¹⁴ Maria Teresa Mercaldo, Mario Cuoco, and Panagiotis Kotetes, Magnetic-field-induced topological reorganization of a p-wave superconductor, *Phys. Rev. B* **94**, 140503(R) (2016).
 - ¹⁵ Timo Hyart, Anthony R. Wright, and Bernd Rosenow, Zeeman-field-induced topological phase transitions in triplet superconductors, *Phys. Rev. B* **90**, 064507 (2014).
 - ¹⁶ R. C. Bento Ribeiro, J. H. Correa, L. S. Ricco, A. C. Seridonio and M. S. Figueira, Spin-polarized Majorana zero modes in double zigzag honeycomb nanoribbons, *Phys. Rev. B* **105**, 205115 (2022).
 - ¹⁷ J. Linder and J. W. A. Robinson, Superconducting Spintronics, *Nat. Phys.* **11**, 307 (2015).
 - ¹⁸ A. Bhattacharyya, D. Adroja, Y. Feng, D. Das, P. K. Biswas, T. Das, J. Zhao, μ SR Study of Unconventional Pairing Symmetry in the Quasi-1D $\text{Na}_2\text{Cr}_3\text{As}_3$ Superconductor. *Magnetochemistry* **9**, 70 (2023).
 - ¹⁹ J. P. Muscat and D. M. Newns, Chemisorption in Metals, *Progress in Surface Science.* , **9**, 1 (1978); D. M. Newns, Self-Consistent Model of Hydrogen Chemisorption, *Phys. Rev.* **178**, 1123 (1969);
 - ²⁰ A. M. Black-Schaffer and A. V. Balatsky, Odd-frequency superconducting pairing in multiband superconductors, *Phys. Rev. B* **88**, 104514 (2013).
 - ²¹ A. M. Black-Schaffer and A. V. Balatsky, Proximity-induced unconventional superconductivity in topological insulators, *Phys. Rev. B* **87**, 220506(R) (2013).
 - ²² H. Caldas, F. S. Batista, M. A. Continentino, F. Deus and D. Nozadze, A two-band model for p-wave superconductivity, *Ann. Phys.* **384**, 211 (2017).
 - ²³ F. Deus, M. A. Continentino and H. Caldas, Induced p-wave superconductivity without spin-orbit interactions, *Ann. Phys.* **362**, 208 (2015).

- ²⁴ Annette Bussmann-Holder, Hugo Keller, Arndt Simon and Antonio Bianconi, Multi-Band Superconductivity and the Steep Band/Flat Band Scenario, *Condens. Matter*, **4**, 91 (2019).
- ²⁵ Wei Tao, Zheng Jue Tong, Anirban Das, Duc-Quan Ho, Yudai Sato, Masahiro Haze, Junxiang Jia, Yande Que, Fabio Bussolotti, K. E. Johnson Goh, BaoKai Wang, Hsin Lin, Arun Bansil, Shantanu Mukherjee, Yukio Hasegawa, and Bent Weber, Multiband superconductivity in strongly hybridized 1T'-WTe₂/NbSe₂ heterostructures, *Phys. Rev. B* **105**, 094512 (2022).
- ²⁶ E. Liebhaber, L. M. Rutten, G. Reece, et al. Quantum spins and hybridization in artificially-constructed chains of magnetic adatoms on a superconductor. *Nat Commun* **13**, 2160 (2022).
- ²⁷ Let us write Eq. 1.1 as, $\Delta_{ind}^p = g(k)\Delta_a^s$, where $\Delta_a^s = (\Delta_k^{\uparrow\downarrow} + \Delta_k^{\downarrow\uparrow})/2$ is the symmetrized BCS pairing of the a -band (substrate), with $\Delta_k^{\sigma-\sigma} = \langle a_{k\sigma}^\dagger a_{-k-\sigma}^\dagger \rangle$. We then obtain for the induced pairing that, if $g(k)$ is symmetric, $\Delta_{ind}^p = 0$, and if $g(k)$ is antisymmetric, i.e., $g(-k) = -g(k)$, then, $\Delta_{ind}^p = g(k)\langle a_{k\uparrow}^\dagger a_{-k\downarrow}^\dagger \rangle$.
- ²⁸ T. D. Stanescu and S. Das Sarma, Proximity-induced superconductivity generated by thin films: Effects of fermi surface mismatch and disorder in the superconductor, *Phys. Rev. B* **106**, 085429 (2022).
- ²⁹ N. van Loo, G. P. Mazur, T. Dvir, et al. Electrostatic control of the proximity effect in the bulk of semiconductor-superconductor hybrids. *Nat Commun* **14**, 3325 (2023).
- ³⁰ Jacob Linder and Jason W. A. Robinson, Superconducting spintronics, *Nature Physics*, **11**, 307 (2015).
- ³¹ L. Schneider, K. T. Ton, I. Ioannidis, et al., *Nature* **621**, 60 (2023).
- ³² G. E. Volovik, *The Universe in a Helium Droplet* (Clarendon Press, Oxford, 2003); G. Volovik, Quantum Phase Transitions from Topology in Momentum Space. In: Unruh, W.G., SchÄEtzhold, R. (eds) *Quantum Analogues: From Phase Transitions to Black Holes and Cosmology*. Lecture Notes in Physics, vol 718, 31 (Springer, Berlin, Heidelberg, 2007).
- ³³ M. Gong, S. Tewari, and C. Zhang, BCS-BEC Crossover and Topological Phase Transition in 3D Spin-Orbit Coupled Degenerate Fermi Gases, *Phys. Rev. Lett.* **107**, 195303 (2011).
- ³⁴ Petr Horava, Stability of Fermi Surfaces and K Theory, *Phys. Rev. Letts.* **95**, 016405 (2005).
- ³⁵ J. L. Manes, F. Guinea, and Maria A. H. Vozmediano, Existence and topological stability of Fermi points in multilayered graphene, *Phys. Rev. B* **75**, 155424 (2007).
- ³⁶ L. M. Wong, K. T. Law, arXiv:1110.4575v2 [cond-mat.supr-con], <https://doi.org/10.48550/arXiv.1110.4575>
- ³⁷ C. L. M. Wong and K. T. Law, Majorana Kramers doublets in $d_{x^2-y^2}$ -wave superconductors with Rashba spin-orbit coupling, *Phys. Rev. B* **86**, 184516 (2012).
- ³⁸ Jason Alicea, New directions in the pursuit of Majorana fermions in solid state systems, *Rep. Prog. Phys.* **75** 076501 (2012).
- ³⁹ R. Okugawa and S. Murakami, Dispersion of Fermi arcs in Weyl semimetals and their evolutions to Dirac cones, *Phys. Rev. B* **89**, 235315 (2014).
- ⁴⁰ B. S. Chandrasekhar, A note on the maximum critical field of high-field superconductors, *Appl. Phys. Lett.* **1**, 7 (1962); A. M. Clogston, Upper Limit for the Critical Field in Hard Superconductors, *Phys. Rev. Lett.* **9**, 266 (1962).
- ⁴¹ E. Dumitrescu and S. Tewari, Topological properties of the time-reversal-symmetric Kitaev chain and applications to organic superconductors, *Phys. Rev. B* **88**, 220505(R) (2013).
- ⁴² see, Bohm-Jung Yang and Naoto Nagaosa, Classification of stable three-dimensional Dirac semimetals with nontrivial topology, *Nature Communications*, **5**, 4898 (2014), for a classification of Dirac points in topological insulators.