Strict area law implies commuting parent Hamiltonian

Isaac H. Kim¹, Ting-Chun Lin², Daniel Ranard³, and Bowen Shi²

¹Department of Computer Science, University of California, Davis, CA 95616, USA ²Department of Physics, University of California San Diego, La Jolla, CA 92093, USA ³Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139

Abstract

We show that in two spatial dimensions, when a quantum state has entanglement entropy obeying a strict area law, meaning $S(A) = \alpha |\partial A| - \gamma$ for constants α, γ independent of lattice region A, then it admits a commuting parent Hamiltonian. More generally, we prove that the entanglement bootstrap axioms in 2D imply the existence of a commuting, local parent Hamiltonian with a stable spectral gap. We also extend our proof to states that describe gapped domain walls. Physically, these results imply that the states studied in the entanglement bootstrap program correspond to ground states of some local Hamiltonian, describing a stable phase of matter. Our result also suggests that systems with chiral gapless edge modes cannot obey a strict area law provided they have finite local Hilbert space.

1 Introduction

A fundamental question in the study of quantum many-body systems is the classification of phases. Formally, one can define gapped phases of matter by defining an equivalence relation in the space of Hamiltonians. If two ground states are adiabatically connected to each other, the two systems are said to be in the same phase.

The classification of gapped quantum many-body systems has been completed in one spatial dimensions [1, 2, 3]. However, in higher dimensions, much less is known. In two spatial dimensions, a common expectation is that each gapped phase of matter is uniquely labeled by the combination of some unitary modular tensor category and a rational number known as the chiral central charge [4]. Though this expectation seems plausible, establishing this fact from a microscopic point of view has been challenging. In three dimensions and higher, there is not even an exhaustive proposed classification, at least if one attempts to accommodate exotic models such as fractors [5].

Recently, a new paradigm to tackle this problem was developed: the entanglement bootstrap [6]. This is an approach to derive the universal properties of gapped quantum phases from ground state entanglement, and ideally to classify them. In entanglement bootstrap, one begins with a set of reasonable axioms about the ground state entanglement, which are believed to be approximately satisfied for gapped ground states. One sufficient axiom is to assume a "strict area law" on the entanglement entropy S(A) of region A, [7, 8]

$$S(A) = \alpha |\partial A| - \gamma \tag{1}$$

for constants α , γ independent of A. While this assumption is particularly compact, weaker axioms involving linear combinations of entropies are sufficient and perhaps more natural, described in Section 2.2.

While the entanglement bootstrap axioms are not *precisely* satisfied in most natural systems, they provide a simple framework in which otherwise difficult ideas can be explored rigorously. Early developments of the entanglement bootstrap program focused on proving familiar ideas about gapped systems in two spatial dimensions, such as the anyon fusion rules [6]. More recent progress in this direction led to novel superselection sectors and their fusion rules in gapped domain walls [9], exotic topological charges in three

spatial dimensions [10, 11], as well as an expression for chiral central charge [12, 13]. There is even a possible extension of this approach to critical systems [14].

While these past works suggest strongly that the entanglement bootstrap approach can lead to a novel predictions about the universal properties of gapped ground states, there is a potential concern. While the axioms of the entanglement bootstrap are known to hold for a large class of gapped ground states, one may wonder if the converse is true. Namely, given some state that satisfies the axioms, one might ask if this state can be viewed as a ground state of some local Hamiltonian. If such a parent Hamiltonian does not exist, not all the findings made in the entanglement bootstrap program may be physically relevant, because then the universal properties revealed by the purported "ground state" might not belong to any local Hamiltonian.

To that end, we prove rigorously that any state that satisfies the axioms of the entanglement bootstrap exactly has a gapped parent Hamiltonian with a stable spectral gap. More specifically, we show that the axioms in Ref. [6] — the ones that are applicable to the "bulk" of gapped ground states in 2D without any domain wall — imply that there is a parent Hamiltonian with a spectral gap. In fact, this parent Hamiltonian is a sum of local terms each of which are commuting projectors. Moreover, we also show that this Hamiltonian satisfies a set of conditions that guarantees the stability of the gap [15]. Therefore, any state that satisfies the axioms of the entanglement bootstrap can be viewed as a ground state of some local Hamiltonian, which is a representative of a stable phase of matter that is robust against sufficiently weak (but nonvanishing) perturbation.

The same argument also works in the presence of gapped domain walls [9]. This is a setup in which some of the axioms are relaxed along a codimension-1 defect, known as a *gapped domain wall*. The axioms for domain walls in Ref. [9] were motivated by considering entanglement properties (such as the strict area law), independent of other inputs such as topological quantum field theory. Therefore, the existence of the gapped parent Hamiltonian in this context was not a priori obvious.

The main insight behind our work is an observation about the property of a quantum Markov chain [16], which may be of independent interest. A tripartite quantum state ρ_{ABC} is said to be a quantum Markov chain if it satisfies $S(\rho_{AB}) + S(\rho_{BC}) = S(\rho_B) + S(\rho_{ABC})$, where $S(\rho) = -\text{Tr}(\rho \log \rho)$ is the von Neumann entropy of a density matrix ρ . We prove that the projector onto the support of ρ_{AB} and ρ_{BC} must necessarily commute. This fact, together with the known properties of quantum Markov chains used in the entanglement bootstrap program [6], lets us construct a local parent Hamiltonian consisting of commuting projectors.

Our result has two immediate physical implications. First, it implies that the states that satisfy the axioms of the entanglement bootstrap ought to be a representative ground state wavefunction of some stable phase of matter. This is because there is a local parent Hamiltonian with a stable spectral gap which has this wavefunction as a ground state. Second, our result implies that there is an obstruction to satisfying the axioms of the entanglement bootstrap *exactly* for systems with a nonzero chiral central charge. It is well-known that such systems must host a nonzero energy current along the edge at finite temperature [4]. If the axioms are satisfied exactly, by our result there must exist a local commuting parent Hamiltonian. For such Hamiltonian, any energy current is identically zero at all temperature, a direct contradiction with the fact that nonzero chiral central charge implies a nonzero energy current at finite temperature.

The rest of the paper is structured as follows. In Section 2 we discuss the setup and provide an executive summary of the main results. In Section 2.3 we discuss well-known facts that are relevant to this paper. In Section 3, we review the argument for extending the axioms of the entanglement bootstrap [6], focusing on the setup we advocated in Section 2. In Section 4, we introduce a class of parent Hamiltonians and show that they all satisfy a property known as the local topological quantum order condition [15]. In Section 5, we prove the aforementioned property of quantum Markov chain: that projectors onto the support over the reduced density matrices commute with each other. In Section 6, we construct the parent Hamiltonian that has a stable spectral gap. In Section 7, we extend the construction to the case in which a gapped domain wall is present. We note that the Hamiltonians constructed in these sections are quite elaborate, consisting of local terms acting on more than hundred sites. In Section 8, we introduce a technique to reduce this

number to three, though we no longer have a guarantee that the local terms commute with each other. We conclude with a discussion and open problems in Section 9.

2 Results and strategy

2.1 Informal overview

We begin with an informal sketch. Suppose a 2D state ρ satisfies

$$I(A:C|B)_{\rho} = 0 \tag{2}$$

for all regions ABC topologically equivalent to those of [Fig. 1]. Here, I(A : C|B) := S(AB) + S(BC) - S(B) - S(ABC) denotes the conditional mutual information. Note this property is true for states with strict area law, as in Eq. (1), or more generally for states that satisfy the entanglement bootstrap axioms [Section 2.2]. Our main result is the construction of a commuting parent Hamiltonian H for which ρ is a gapped and frustration-free ground state, with a local uniqueness property described below.

Figure 1: Sub-regions A, B and C in the 2D plane arranged like a chain.

The construction is easily summarized as follows. For any region X, let

$$P_X = \operatorname{proj}_{\operatorname{supp}(\rho_X)} \tag{3}$$

be the projector onto the support of ρ_X . For states ρ_{ABC} on a finite-dimensional Hilbert space satisfying $I(A:C|B)_{\rho} = 0$, it turns out [Proposition 3]

$$[P_{AB}, P_{BC}] = 0. (4)$$

We will construct commuting parent Hamiltonian for ρ by taking

$$H = \sum_{X \in \mathcal{S}} (1 - P_X) \tag{5}$$

where the sum is taken over some set S of specially chosen O(1)-size regions X. We guarantee the terms P_X commute by ensuring that whenever a pair of overlapping regions $X_1, X_2 \in S$ appear in the sum, they are topologically equivalent to the pair of regions AB, BC [Fig. 1], so that Eq. (4) guarantees $[P_{X_1}, P_{X_2}] = 0$.

We also want H to have a unique ground state in some sense – for instance, taking H = 0 gives a "parent Hamiltonian," but it is not useful. Still, we do not want to require H has unique ground state: models such as the toric code have degeneracies depending on the global topology of the manifold. A natural condition is that the ground state is "locally unique," in the sense that all zero-energy states on a disk look identical on the interior of the disk. One codification of this is the condition of *local topological quantum* order (LTQO) [15]. To guarantee LTQO, we require the regions in S form a sort of "open cover" of the plane, i.e., their interiors must cover the plane. The LTQO property then follows from basic entanglement bootstrap techniques.

To find a set of regions satisfying these geometric requirements, we make use of a cellular decomposition of the plane. In particular, referring to Fig. 5, each region X is given by the union of a single green region with its neighboring blue and red regions. One can check such regions X satisfy the above conditions.

Perhaps one can see that the basic ideas are topological and not too intricate. However, some care is required in the choice of regions X. Much of the technical detail is devoted to making a rigorous argument while assuming the entanglement bootstrap axioms only for a small number of explicitly chosen, O(1)-size lattice regions and their translations.

2.2 Summary of results

Throughout this paper, we shall consider a two-dimensional (2D) plane, covered by a set of faces, each being a closed hexagon-shaped disk [Fig. 2]. We assign a finite-dimensional Hilbert space to each face, and we informally view the global Hilbert space as a tensor product of the local Hilbert spaces. We choose a quantum state in the global Hilbert space, which we call the *reference state* and denoted as σ throughout the paper.¹



Figure 2: In this paper, we consider a partition of \mathbb{R}^2 into hexagonal cells, each representing a finitedimensional quantum system.

Denoting the set of faces as Λ , we define several notions that shall be useful. For a set $A \subset \Lambda$, we denote the *neighborhood* of A as N(A), which is a set of faces in $\Lambda \setminus A$ that are adjacent to at least one of the faces in A. We shall sometimes focus on the boundary of a single face $f \in \Lambda$. In those cases, we shall abuse the notation and denote the neighborhood as N(f), instead of $N(\{f\})$. We say that a set $A \subset \Lambda$ is *connected* if the union of the faces, viewed as a subset of \mathbb{R}^2 , is connected. Similarly, $A \subset \Lambda$ is *simply connected* if the union of the faces is, viewed as a subset of \mathbb{R}^2 , is simply connected. Examples are shown in Fig. 3. Sometimes we shall refer to a simply connected set as a *disk*. Lastly, if a disk is $f \cup N(f)$ for some $f \in \Lambda$, we refer to this disk as an *elementary disk*. We emphasize that elementary disk is not necessarily the smallest disk. Nevertheless, we advocate this convention due to the repeated appearance of elementary disk in this paper.



Figure 3: (Left) The faces colored in green is an example of a simply connected subsystem. (Right) The faces colored in blue is an example of a connected subsystem which is not simply connected.

Our first main result concerns a reference state that satisfies the axioms A0 and A1 in the vicinity of a face. More formally, consider a disk $A \subset \Lambda$.

Definition 1. The state σ satisfies A0 in the vicinity of $A \subset V$ if

$$(S(A) + S(A \cup N(A)) - S(N(A)))_{\sigma} = 0.$$
(6)

¹We treat the infinite lattice informally. More formally, one could adopt the algebraic framework [17], ultimately viewing the reference state as a positive linear functional on the algebra of quasi-local observables. Rather than assuming this "global state" as an input, one could first view the reference state as a collection of consistent density matrices on O(1)-size regions covering the plane, satisfying the entanglement bootstrap axioms locally. Then the techniques we discuss allow one to define a consistent "global state," e.g. a state on the algebra of quasi-local observables, consistent with the local data.

Definition 2. The state σ satisfies A1 in the vicinity of $A \subset V$ if for any partition of N(A) into disjoint simply connected subsets $N(A)_1$ and $N(A)_2$,

$$(S(A \cup N(A)_1) - S(N(A)_1) + S(A \cup N(A)_2) - S(N(A)_2))_{\sigma} = 0.$$
⁽⁷⁾

We say that a reference state is *valid* if for every face $f \in \Lambda$, A0 and A1 in the vicinity of f is satisfied. To be more precise, we demand

$$(S(C) + S(C|B))_{\sigma} = 0 \quad \text{for} \quad \overrightarrow{BC}$$
(8)

where $S(X|Y) := S_{XY} - S_Y$ and

$$(S(C|B) + S(C|D))_{\sigma} = 0 \quad \text{for} \quad \overrightarrow{BCD}, \overrightarrow{BCD}, \overrightarrow{BCD}, (9)$$

as well as rotated choices of subsystems B and D (while keeping C intact).

Our first main result establishes, for the valid reference states, the existence of a local commuting projector parent Hamiltonian. These are Hamiltonians that can be expressed as a sum of local projectors that commute with each other. Here is our first main theorem.

Theorem 1. For any valid reference state, there exists a local commuting projector parent Hamiltonian satisfying the local topological quantum order condition [15, Definition 4].

In Ref. [15], the local topological quantum order condition, together with a condition they refer to as the local gap condition, guarantees the stability of the gap under any perturbation which has a finite but sufficiently small strength.² It should be noted that the local gap condition is trivially satisfied by any local commuting projector models. Therefore, an immediate corollary of Theorem 1 is the existence of a parent Hamiltonian for any valid reference state, which has a stable spectral gap.

Our second main result establishes the existence of a local commuting projector parent Hamiltonian for a larger class of states. In Ref. [9], certain axioms of the entanglement bootstrap were relaxed along a one-dimensional line. As an example, we can consider a lattice shown in Fig. 4. The reference state, in this case, will continue to satisfy A0 everywhere, but A1 may be violated along the disks that touch the red line in Fig. 4. More precisely, while A1 continues to be satisfied for the following choice of B, C, and D,

it may not necessarily hold for the following choices:

$$\begin{array}{c} B \\ \hline C \\ \hline C \\ \hline \end{array}, \begin{array}{c} B \\ \hline C \\ \hline \end{array}.$$
(11)

In words, A1 may not hold if the boundaries between B and D are both away from the domain wall and lie on the same side with respect to the domain wall. Otherwise, the A1 continues to hold. Again, we say that the reference state is a *valid* reference state (for gapped domain walls) if it satisfies these constraints.

Theorem 2. For any valid reference state for gapped domain walls, there exists a local commuting projector parent Hamiltonian satisfying the local topological order condition.

We now turn to a brief discussion on basic facts about quantum entropies [Section 2.3]. The rest of the paper will focus on the proof of Theorem 1.

²The condition we use is actually TQO-2 in Ref. [18], though we use the local TQO condition for convenience.



Figure 4: A 2D plane with a gapped domain wall (red). For all the elementary disks that do not touch the red line, **A1** continues to hold. However, for the elementary disks that touch the red line, **A1** may not necessarily hold. The axiom **A0** continues to hold everywhere.

2.3 Preliminary: Entropy inequalities

Throughout this paper, density matrices are denoted by Greek letters, such as $\rho, \sigma, \tau, \ldots$ The subscripts refer to the Hilbert spaces the density matrices act on. For instance, ρ_{ABC} is a density matrix acting on a Hilbert space $\mathcal{H}_{ABC} = \mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$. The set of density matrices over subsystem X is denoted as \mathcal{D}_X . The von Neumann entropy is defined as $S(\rho) = -\text{Tr}(\rho \log \rho)$. Entanglement entropy is defined as $S(A)_{\rho} = S(\rho_A)$. Conditional mutual information over a tripartite state ρ_{ABC} is defined as

$$I(A:C|B)_{\rho} := S(AB)_{\rho} + S(BC)_{\rho} - S(B)_{\rho} - S(ABC)_{\rho}.$$
(12)

We remark that the strong subadditivity of entropy implies the following inequalities, which we shall often use:

$$I(AA': CC'|B)_{\rho} \ge I(A:C|B)_{\rho},$$

$$I(AA':C|B)_{\rho} \ge I(A:C|A'B)_{\rho}.$$
(13)

We also remark that, for any tripartite quantum state ρ_{ABC} , the following inequality always holds.

$$S(A|B)_{\rho} + S(A|C)_{\rho} \ge 0.$$
 (14)

Eq. (14) is known as the *weak monotonicity*, a fact that we shall often use as well.

3 Extensions of axioms

Recall our central assumption: that the reference state σ satisfies **A0** and **A1** in the vicinity of every face $f \in \Lambda$ [Eq. (8), (9)]. One of the foundational facts in Ref. [6] is that the axioms **A0** and **A1** imply the same axioms at a larger scale. Brought to our current setup, this would imply, for instance, that

$$(S(C) + S(C|B))_{\sigma} = 0 \quad \text{for} \qquad BC \tag{15}$$

and

$$(S(C|B) + S(C|D))_{\sigma} = 0 \quad \text{for} \qquad \overrightarrow{BC + D}$$
(16)

An important similarity between Eq. (8) and Eq. (15) (and similarly, between Eq. (9) and Eq. (16)) is that a disk (C) is surrounded by the union of two disks (B and D). Moreover, the union BCD is again a disk. While the precise shapes of the subsystems labeled in B, D, BC, and CD differ between Eq. (9) and Eq. (16), topologically they are the same. Eq. (15) and (16) are examples of what we mean by an extension of axioms A0 and A1. While the argument for such extensions has been discussed in Ref. [6] already, we revisit this argument in this paper for the sake of concreteness.

3.1 Extending A0

Let us first consider the simpler case: proving extensions of A0. While the argument we present below is general, we aid the readers with a concrete example. Suppose, for instance, that our goal is to prove Eq. (15). How would we proceed? The main idea can be schematically expressed as follows:

$$\xrightarrow{BC} \xrightarrow{A1} \xrightarrow{BC} \xrightarrow{BC} (17)$$

This diagram means that we begin with $(S(C) + S(BC) - S(B))_{\sigma} = 0$ for the subsystems B and C on the left hand side and then gradually change the subsystems via A1 to arrive at $(S(C) + S(BC) - S(B))_{\sigma} = 0$ for the subsystems shown on the right hand side.

As a first step, we can first argue

$$(S(C) + S(C|B))_{\sigma} = 0 \quad \text{for} \qquad BC \qquad (18)$$

This is a simple consequence of SSA:

$$(S(C) + S(C|B))_{\sigma} \ge \left(S(C) + S(C|BB')\right)_{\sigma} \tag{19}$$

for any subsystem B'. Now we can evaluate the difference of $(S(C) + S(C|B))_{\sigma}$ appearing in Eq. (15) and Eq. (18), the result of which is

$$(S(C|B) + S(C|D))_{\sigma}$$
 for BCD . (20)

Conveniently, this quantity is zero. To see why, note that

$$(S(C|B) + S(C|D))_{\sigma} = 0$$
 for BCD (21)

by our assumption [Eq. (8)]. A simple consequence of SSA is that

$$\left(S(C|B) + S(C|D)\right)_{\sigma} \ge \left(S(C|B) + S(C|DD')\right)_{\sigma}$$

$$\tag{22}$$

for any D'. Viewing the green faces in Eq. (20) that are colored in white in Eq. (21) as D', we can see that $(S(C|B) + S(C|D))_{\sigma} \leq 0$ for the choice B, C, and D shown in Eq. (20). At the same time, by the weak monotonicity, this quantity must be always zero. Thus the difference of $(S(C) + S(C|B))_{\sigma}$ appearing in Eq. (15) and Eq. (18) are exactly equal to each other, and in fact, both zero. This completes the proof.

We can generalize this procedure into an algorithm that extends our axioms. Without loss of generality, suppose our task is to prove $(S(C) + S(C|B))_{\sigma} = 0$, where C is a disk and B is an annulus surrounding B, so that BC together forms a disk. Pick any face $c \in C$ and let b = N(c). By assumption, $(S(c) + S(c|b))_{\sigma} = 0$. From SSA, it follows that $(S(c) + S(c|BC \setminus c))_{\sigma} = 0$. Now, the idea is to extend the disk c into C, adding

one face at a time. Specifically, consider a sequence of disks $\{C_i : i = 1, ..., n\}$ such that $c_1 = c$, $C_n = C$ and $C_i \subset C_{i+1}$.³ Letting $c_{i+1} := C_{i+1} \setminus C_i$ be a face, note the following identity:

$$(S(C_{i+1}) + S(C_{i+1}|BC \setminus C_{i+1}) - S(C_i) - S(C_i|BC \setminus C_i))_{\sigma} = (S(c_{i+1}|C_i) + S(c_{i+1}|BC \setminus C_{i+1}))_{\sigma}.$$
 (23)

Because $B \supset N(C)$ and $c_{i+1} \subset C$, $N(c_{i+1}) \subset BC$. Therefore, we can partition $N(c_{i+1})$ into $N(c_{i+1}) = (N(c_{i+1})_1 \cup N(c_{i+1})_2)$, where

$$N(c_{i+1})_1 := N(c_{i+1}) \cap C_i,$$

$$N(c_{i+1})_2 := N(c_{i+1}) \cap BC \setminus C_{i+1}.$$
(24)

Note that, by SSA, the following inequalities hold:

$$S(c_{i+1}|C_i) \le S(c_{i+1}|N(c_{i+1})_1),$$

$$S(c_{i+1}|BC \setminus C_{i+1}) \le S(c_{i+1}|N(c_{i+1})_2),$$
(25)

which together with weak monotoncity implies that Eq. (23) is zero. Iterating this argument from i = 1 to i = n - 1, we conclude that $(S(C) + S(C|B))_{\sigma} = 0$, proving our claim.

3.2 Extending A1

Though different in details, the argument for extending A1 is similar in spirit. Let us again work through a guiding example; below, we prove $(S(C|B) + S(C|D))_{\sigma} = 0$ for the choice of subsystems shown on right hand side of Eq. (26). This argument begins with $(S(C|B) + S(C|D))_{\sigma} = 0$ for the subsystems shown on the left hand side of Eq. (26) (which can be viewed as the assumption A1), followed by applications of A1 over other elementary disks.



Our argument is best explained in terms of the *deformations of the boundaries*. Specifically, for any subsystem $A \subset \Lambda$, let us define ∂A as the boundary of A, viewing A as a subset of \mathbb{R}^2 . These boundaries can be classified into the inner and the outer boundaries; the outer boundary is $\partial(BCD)$ and the inner boundaries are $\partial B \cap \partial C$, $\partial C \cap \partial D$, and $\partial B \cap \partial D$, drawn in green, blue, and red below:



We shall deform the shown boundaries, by deforming the subsystems (BC, CD, B, and D) by one face at a time whilst maintaining their topologies. The key point here is that the value of $(S(C|B) + S(C|D))_{\sigma}$ remains invariant under two types of moves that deform the boundaries. By combining these two moves together, one can conclude $(S(C|B) + S(C|D))_{\sigma}$ on the right hand side of Eq. (27) is zero.

The first move is to deform the outer boundaries, which entails enlarging/reducing either B or D. This move keeps the green and the blue boundary unchanged, but it may deform the red boundaries. Since the quantity at hand $(S(C|B) + S(C|D))_{\sigma}$ is symmetric under the exchange of B and D, an argument that works for one of them works for the other as well. We thus focus on enlarging/reducing D. Without loss of generality, let us consider the difference between $(S(C|B) + S(C|D))_{\sigma}$ and $(S(C|B) + S(C|Dd))_{\sigma}$, where $d \in \Lambda \setminus (BCD)$ is a face. The difference between the two quantities is $I(d : C|D)_{\sigma}$. By SSA,

$$I(d:C|D)_{\sigma} \le I(d:CD \setminus N(d)|D \cap N(d))_{\sigma},$$
(28)

³Such a sequence of disks can be constructed as follows. For each C_i , consider a face adjacent to C_i , which is still part of C, and then choose that to be c_i .

where we used the fact that C has a trivial intersection with N(d).⁴ Further,

$$I(d:CD \setminus N(d)|D \cap N(d))_{\sigma} \le (S(d|D \cap N(d)) + S(d|N(d) \setminus D))_{\sigma}.$$
(29)

By A1, the right hand side Eq. (29) is zero if $D \cap N(d)$ and $N(d) \setminus D$ are both disks.

The condition that these subsystems ought to be disks can be ensured for any face d as long as both Dand Dd are disks. Suppose otherwise. That is, there is a face d such that both D and Dd are disks but $N(d) \cap D$ is not a disk. Thus $N(d) \cap D$ must be a union of at least two disjoint disks. Moreover, there is a disk in $N(d) \setminus D$ — surrounded by the aforementioned two disks, d, and D — whose neighborhood is contained in Dd. Any loop that surrounds the aforementioned disk in $N(d) \setminus D$ is not contractible to a point, so Dd is not simply connected, which is a contradiction.

Therefore, provided that D and Dd are disks, the right hand side of Eq. (29) is at most 0. Since conditional mutual information is non-negative, $I(d:C|D)_{\sigma} = 0$. Thus

$$(S(C|B) + S(C|D))_{\sigma} = (S(C|B) + S(C|Dd))_{\sigma}.$$
(30)

The same argument can be employed to argue the invariance under deformation of B which retains its disk-like topology. Thus we conclude that $(S(C|B) + S(C|D))_{\sigma} = 0$ for the subsystems shown below:



Comparing this diagram to the right hand side of Eq. (27), it is clear now the main task is to deform the inner boundaries.

The argument for deforming the inner boundaries is exactly the same. Let us introduce a purifying system for BCD, denoted as E. Note the following identity:

$$(S(C|B) + S(C|D))_{\sigma} = (S(E|B) + S(E|D))_{\sigma}, \qquad (32)$$

which follows easily from the fact that the state over BCDE is a pure state. The role played by C in our previous argument can now be played by E. The new inner boundaries associated with B, E and D (as opposed to B, C, and D) is shown below:

$$(33)$$

E (Purifying system)

where the purifying system E is not specified explicitly⁵. Note that the outer boundaries with respect to B, E, and D are precisely the green and the blue boundaries with respect to B, C, and D. The argument for deforming these new outer boundaries is exactly the same as before, leading to

$$(S(E|B) + S(E|D))_{\sigma} = 0$$
 for E (Purifying system) (34)

⁴Recall that C is chosen in such a way that it is surrounded by B and D. Therefore, C cannot be in the neighborhood of d, which is in $\Lambda \setminus (BCD)$.

⁵Note that E is not necessarily $\Lambda \setminus (BCD)$, but rather an abstract system that purifies σ_{BCD} .

which is equivalent to $(S(C|B) + S(C|D))_{\sigma} = 0$ for the subsystems shown on the right hand side of Eq. (27).

The argument we employed in our example can be generalized to an algorithm. Without loss of generality, suppose our task is to show $(S(C|B) + S(C|D))_{\sigma}$ for B, C, and D such that BC, CD, B, and D are disks and BD is an annulus. First choose any $c \in C$. Now we choose $b, d \subset N(c)$ based on the red boundaries defined with respect to B, C, and D. If a red boundary has a nontrivial overlap with N(c), choose b and d such that the red boundaries defined with respect to b, c, and d overlap with the red boundaries defined with respect to B, C, and D. Then, we can grow the red boundaries defined with respect to b, c, and d to the extent that they include the red boundaries defined with respect to B, C, and D; this can be done by extending b and d by the blue and green faces adjacent to a unit interval of the red boundary. We can then trim the red boundary. This process ensures that we have a choice of B', C', D' such that their red boundaries match the red boundaries of B, C, and D. Now, the outer and the blue/green boundaries can be deformed using the following greedy algorithm. For D', at each time step, look at all the cells in N(D')and if $N(D') \cap D$ is non-empty and if there is an element whose addition does not change the topology, add that. The same applies to B'. Then remove the unnecessary faces.

While our results do not rely on showing this algorithm works unconditionally for any subsystems topologically equivalent to the ones appearing in A0 and A1, we expect that this is the case.⁶ Regardless, for the purpose of our paper, the description of the algorithm we provide shall be sufficient. This is because we simply need to extend A0 and A1 to a particular (finite) set of subsystems, for which one can verify that our algorithm works. These subsystems shall be introduced in Section 6 and their construction shall be modified from the discussion in Section 4, which we now turn our attention to.

4 LTQO and local uniqueness

In this section, we introduce the notion of local topological quantum order (LTQO), or having a locally unique ground state. Then we introduce a class of parent Hamiltonians for the reference state that satisfies this condition.

To produce a useful parent Hamiltonian, we want to establish some sense in which the ground state is unique. While models such as the toric code have degeneracies depending on the topology of the manifold, they still satisfy a certain "local uniqueness" property. In particular, for all topological disks A, every zero-energy state on A looks identical on the interior of A. In fact this is a reformulation of the condition introduced as "local topological quantum order" (LTQO) by [15]. They show that LTQO implies the gap is stable to local perturbations.

For simplicity, we define LTQO as a condition placed upon local, frustration-free Hamiltonians. Thus the condition will be a special case of ([15], Definition 4).

Definition 3 (Local topological quantum order). Consider a local frustration-free Hamiltonian, $H = \sum_X (1 - Q_X)$, summed over some O(1)-size disks X, with projections Q_X local to X. Let Q_A denote the projection onto $\{|\psi\rangle_A \in \mathcal{H}_A : (1 - Q_X)|\psi\rangle_A = 0 \ \forall X \subset A\}$, the space of states on A with zero energy. Let A be a ball of radius r and $A(\ell)$ be the set of sites that are distance at most ℓ away from A. Let

$$c_{\ell}(O_A) := \frac{\operatorname{Tr}(Q_{A(\ell)}O_A)}{\operatorname{Tr}(Q_{A(\ell)})}$$
(35)

for any operator O_A acting on \mathcal{H}_A . Then we say H satisfies the condition of *local topological quantum order* if there exists $\ell = O(1)$ independent of r, such that for all balls A and operators O_A ,

$$Q_{A(\ell)}O_A Q_{A(\ell)} = c_\ell(O_A) Q_{A(\ell)}.$$
(36)

⁶In fact, we suspect such an argument works even for tessellations of hyperbolic spaces.

The original definition of LTQO allows Eq. (36) to be satisfied approximately. We shall be content with this simplified version because we can show Eq. (36) exactly for H. In particular, if our version [Definition 3] holds, the original version of LTQO holds as well.

Perhaps a more physical formulation is the following.

Definition 4 (Locally unique ground state). Consider a Hamiltonian H as in Definition 3. Then we say H has a locally unique ground state if, for some $\ell = O(1)$ independent of r, for all balls A of any radius r, for all states $\rho_{A(\ell)}$ on enlarged region $A(\ell)$ that have zero energy on $A(\ell)$, the state

$$\tau_A := \operatorname{Tr}_{A(\ell) \setminus A} \rho_{A(\ell)} \tag{37}$$

is independent of the state $\rho_{A(\ell)}$. (We say that a state ρ_R on region R has zero energy if it is in the ground space of all the terms in H supported on R.)

These formulations are equivalent.

Proposition 1 (LTQO as local uniqueness). For local, frustration-free Hamiltonians, the LTQO condition stated in Definition 3 is equivalent to the local uniqueness condition of Definition 4.

The equivalence is known and straightforward. For completeness we include a proof of the less direct implication, that local uniqueness implies LTQO.

Proof. We assume H satisfies (37) and show (36). By (37), for any $|\Psi\rangle$, $|\Psi'\rangle \in \ker(I - Q_{A(\ell)})$ for a sufficiently large but constant ℓ ,

$$\langle \Psi | O_A | \Psi \rangle = \langle \Psi' | O_A | \Psi' \rangle. \tag{38}$$

Choosing $|\Psi\rangle = \frac{1}{\sqrt{2}}(|\Psi_1\rangle + |\Psi_2\rangle)$ and $|\Psi'\rangle = \frac{1}{\sqrt{2}}(|\Psi_1\rangle - |\Psi_2\rangle)$ for an orthogonal pair of states $|\Psi_1\rangle$ and $|\Psi_2\rangle$, we can conclude that

$$\langle \Psi_1 | O_A | \Psi_2 \rangle + \langle \Psi_2 | O_A | \Psi_1 \rangle = 0.$$
(39)

Choosing $|\Psi\rangle = \frac{1}{\sqrt{2}}(|\Psi_1\rangle + i|\Psi_2\rangle)$ and $|\Psi'\rangle = \frac{1}{\sqrt{2}}(|\Psi_1\rangle - i|\Psi_2\rangle)$, we can conclude that

$$\langle \Psi_1 | O_A | \Psi_2 \rangle - \langle \Psi_2 | O_A | \Psi_1 \rangle = 0, \tag{40}$$

implying $\langle \Psi_1 | O_A | \Psi_2 \rangle = \langle \Psi_2 | O_A | \Psi_1 \rangle = 0$. Therefore, the off-diagonal elements of O_A in ker $(I - Q_{A(\ell)})$ are identically zero. This immediately implies Eq. (36).

Now we turn to constructing a parent Hamiltonian for the reference state. We consider a Hamiltonian of the following form,

$$H = \sum_{X \in \mathcal{S}} (I - P_X), \tag{41}$$

where the elements of \mathcal{S} are disks of constant size and

$$P_X = \operatorname{proj}_{\operatorname{supp}(\rho_X)} \tag{42}$$

is the projector onto the support of σ_X . The main result of this section will be

Proposition 2. [Parent Hamiltonian satisfies LTQO] Assume the Hamiltonian defined in Eq. (41) uses a collection of regions S such that, for every elementary disk D, there exists a disk $X \in S$ such that $D \cup N(D) \subset X$. Then H satisfies LTQO [Definition 3].

One can think of the condition above as requiring that the regions X cover the plane with sufficient buffer. While this condition is sufficient for LTQO, later we will make further specifications of S. With these further properties, the resulting Hamiltonian will be frustration-free and commuting. Note "support" is a word used in two different ways: (i) the support of an operator is the subsystem X that the operator acts on, and (ii) the support of a density matrix σ_X is the orthogonal complement of the kernel, as a linear subspace of the Hilbert space on X. Since σ lives in the ground state subspace by definition and $H \ge 0$, the ground state energy is zero.

To prove Proposition 2, we will show the local uniqueness property of Definition 4, which is equivalent to LTQO [Definition 3] via Proposition 1. We begin with the following simple fact [6].

Lemma 1. Let ρ_{BC} be a density matrix such that $(S(C) + S(B|C))_{\rho} = 0$. For any density matrix τ_{BC} in the support of ρ_{BC} , $\tau_C = \rho_C$.

Proof. Note that $I(A:C)_{\psi_{\rho}} = 0$, where A is the purifying system. Here $\psi_{\rho} := |\psi_{\rho}\rangle_{ABC} \langle \psi_{\rho}|$ for some state vector $|\psi_{\rho}\rangle_{ABC}$. Thus $\rho_{AC} = \rho_A \otimes \rho_C$. By Uhlmann's theorem [19], this implies that there exists an isometry $V_{B_L B_R \to B} : \mathcal{H}_{B_L} \otimes \mathcal{H}_{B_R} \to \mathcal{H}_B$, where B_L and B_R are auxiliary systems purifying ρ_A and ρ_C ,

$$|\psi_{\rho}\rangle_{ABC} = V_{B_L B_R \to B} |\psi_1\rangle_{AB_L} \otimes |\psi_2\rangle_{B_R C}.$$
(43)

Thus the density matrices in the support of ρ_{BC} must be of the following form:

$$V_{B_L B_R \to B} \left(\lambda_{B_L} \otimes |\psi_2\rangle_{B_R C} \langle \psi_2 | \right) V_{B_L B_R \to B}^{\dagger}, \tag{44}$$

where λ_{B_L} is a density matrix acting on the support of $\operatorname{Tr}_A(|\psi_1\rangle_{AB_L}\langle\psi_1|)$. Upon tracing out *B* in Eq. (44), we obtain a density matrix $\operatorname{Tr}_{B_R}(|\psi_2\rangle_{B_RC}\langle\psi_2|)$, which is precisely ρ_C .

Using Lemma 1, we can conclude that any element in ker H, restricted to any elementary disk, ought to be exactly the marginal of σ defined on the same elementary disk. To see why, recall that we assumed that for any elementary disk $D, D \cup N(D)$ is included in some $X \subset S$. Therefore, any element in ker H, restricted to $D \cup N(D)$, must be supported on the support of $\sigma_{D \cup N(D)}$. Setting C = D and B = N(D), we recall that [Section 3.1]

$$(S(C) + S(B|C))_{\sigma} = 0. \tag{45}$$

By Lemma 1, $\rho_C = \sigma_C$, where C = D is an elementary disk. Thus for any elementary disk D, for any density matrix ρ supported in ker H,

$$\rho_D = \sigma_D. \tag{46}$$

This argument can be easily generalized to a Hamiltonian restricted to any finite disk. Consider a disk, denoted as $D \subset \Lambda$. Define

$$H_D := \sum_{\substack{X \in \mathcal{S}, \\ X \subset D}} (I - P_X).$$

$$\tag{47}$$

We claim that any element in ker H_D , restricted to any elementary disk whose distance from ∂D is larger than some constant of order unity, is equal to the reduced density matrix of σ on the same elementary disk. Recall that $X \subset S$ is a disk of constant size. Let r be the maximal radius of X. For any elementary disk u that has a distance (to ∂D) larger than 2r + 1, $u \cup N(u)$ cannot be included in $X \subset S$ that lies outside of D. On the other hand, by our construction, $u \cup N(u)$ ought to be a subset of some $X \subset S$. Therefore, for any elementary disk u whose distance from ∂D is larger than 2r + 1, there ought to be a $X \subset S$ such that $u \cup N(u) \subset X$. Again using Lemma 1, the reduced density matrix of any element of ker H_D over u is exactly σ_D .

Therefore, for any disk D, for any $D' \subset D$ separated from ∂D by at least some positive distance, the following property holds. Without loss of generality, consider an elementary disk associated with a face in D'. The reduced density matrix over this elementary disk (obtained from an element in ker H_D) is indistinguishable from the reduced density matrix of σ over the same elementary disk.

In fact, more can be said. Let $\rho_{D'}$ be a reduced density matrix of an element of ker H_D over D'. We claim

$$\rho_{D'} = \sigma_{D'}.\tag{48}$$

The proof is based on induction. Let $D_i \subset D'$ be a disk and suppose $\rho_{D_i} = \sigma_{D_i}$. Consider a face $f \in D'$, where $D_i f$ is a disk. From SSA, it follows that:

$$I(f: D_i \setminus N(f)|D_i \cap N(f))_{\sigma} \le (S(f|D_i \cap N(f)) + S(f|N(f) \setminus D_i))_{\sigma},$$

$$I(f: D_i \setminus N(f)|D_i \cap N(f))_{\rho} \le (S(f|D_i \cap N(f)) + S(f|N(f) \setminus D_i))_{\rho}.$$
(49)

Both ρ and σ are indistinguishable over D_i and $f \cup (N(f) \cap D_i)$ by assumption. Since two locally indistinguishable Markov chains are globally indistinguishable [Lemma 2], $\rho_{fD_i} = \sigma_{fD_i}$. This argument can be repeated until the enlarged disk becomes D'. This completes the proof of Proposition 2.

We conclude with the lemma used above,

Lemma 2. Let ρ_{ABC} and σ_{ABC} be quantum Markov chains. If $\rho_{AB} = \sigma_{AB}$ and $\rho_{BC} = \sigma_{BC}$, $\rho_{ABC} = \sigma_{ABC}$.

Proof. The following proof is based on Ref. [20]. Let $\tau_{ABC} = \frac{1}{2}(\rho_{ABC} + \sigma_{ABC})$.

$$S(\tau_{ABC}) - \frac{1}{2}(S(\rho_{ABC}) + S(\sigma_{ABC})) \le \frac{1}{2}(I(A:C|B)_{\rho} + I(A:C|B)_{\sigma}).$$
(50)

The right hand side is zero by our assumption. By the strict concavity of the von Neumann entropy, the left hand side must be zero. Moreover, this is possible if and only if $\rho_{ABC} = \sigma_{ABC}$.

5 Commuting projectors from quantum Markov chain

The Hamiltonian defined in Eq. (41) satisfies the LTQO condition [Proposition 2]. However, in order to use the stability of the gap proved in Ref. [15], one needs an additional ingredient: local gap condition. What this means is that H_X , for any disk X, has a spectral gap bounded from below between the ground state subspace of H_X and the rest of the spectrum. The purpose of this Section is to develop techniques that can establish such a result.

The main result of this Section is Proposition 3. This proposition states that, if ρ_{ABC} is a quantum Markov chain, the projectors onto the support of ρ_{AB} and ρ_{BC} commute.

Theorem 3. [21] A tripartite quantum state ρ_{ABC} , defined on a finite-dimensional Hilbert space $\mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$, satisfies $I(A:C|B)_{\rho} = 0$ if and only if there is a decomposition of B as

$$\mathcal{H}_B = \bigoplus_j \mathcal{H}_{b_j^L} \otimes \mathcal{H}_{b_j^R} \tag{51}$$

such that

$$\rho_{ABC} = \bigoplus_{j} q_j \rho_{Ab_j^L} \otimes \rho_{b_j^R C}$$
(52)

with states $\rho_{Ab_j^L}$ in $\mathcal{H}_{Ab_j^L}$ and $\rho_{b_j^R C}$ in $\mathcal{H}_{b_j^R C}$, and a probability distribution $\{q_j\}$.

Lemma 3. Consider a bipartite Hilbert space $\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B$ and let ρ_{AB} be a state that acts on it. Let P_{AB} and P_B be the projector onto the support of ρ_{AB} and ρ_B , respectively.

$$[P_B, P_{AB}] = 0. (53)$$

Proof. Consider the images of $I \otimes P_B$ and P_{AB} . The latter is a subspace of the former. Within the former subspace, $I \otimes P_B$ commutes with every element, and in particular, with P_{AB} . In the orthogonal complement, both operators annihilate every state, and as such, trivially commute.

Proposition 3. Let ρ_{ABC} be a state defined on a finite dimensional Hilbert space $\mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$ that satisfies $I(A:C|B)_{\rho} = 0$. Let P_{AB} and P_{BC} be the projector onto the support of ρ_{AB} and ρ_{BC} , respectively. Then

$$[P_{AB}, P_{BC}] = 0. (54)$$



Figure 5: Decomposition of the plane into 0-1-, and 2-cells, corresponding to the red, blue, and green balls.

Proof. Consider the decomposition in Eq. (52). Let $J = \{j : q_j \neq 0\}$. Define $P_{Ab_j^L}$ and $P_{b_j^R C}$ as projectors onto the support of $\rho_{Ab_j^L}$ and $\rho_{b_j^R C}$, respectively. Similarly, define $P_{b_j^L}$ and $P_{b_j^R}$ to be the projectors onto the support of $\rho_{b_j^L}$ and $\rho_{b_j^R}$, respectively. These projectors are related to P_{AB} and P_{BC} in the following way:

$$P_{AB} = \sum_{j \in J} P_{Ab_j^L} \otimes P_{b_j^R},$$

$$P_{BC} = \sum_{j \in J} P_{b_j^L} \otimes P_{b_j^R C}.$$
(55)

Since that $[P_{b_j^L}, P_{Ab_j^L}] = 0$ and $[P_{b_j^R}, P_{Ab_j^R}] = 0$, by Lemma 3, $[P_{AB}, P_{BC}] = 0$.

Using Proposition 3, with a judicious choice of subsystems used in defining the parent Hamiltonian [Eq. (41)], one can show that all the terms in Eq. (41) commute with each other.

However, it is a priori not obvious whether ensuring such commutation relation while ensuring LTQO is possible. We now turn to this issue in Section 6. An explicit construction that satisfies both shall be given, which establishes that there exists a local parent Hamiltonian for the reference state σ which moreover has a stable spectral gap.

6 Constructing the parent Hamiltonian

In Section 4, we showed that the Hamiltonian in Eq. (41) satisfies the LTQO condition. The key assumption was that the set of subsystems S in Eq. (41) is dense enough so that for every elementary disk D, there exists some $X \subset S$ such that $D \cup N(D) \subset X$. For any S that satisfies this condition, the LTQO condition follows. In Section 5, we established a sufficient condition under which two terms in the Hamiltonian [Eq. (41)] commute. This is possible if for any $X, Y \subset S$, there exists a set of subsystems A, B, and C such that X = AB, Y = BC, and $I(A : C|B)_{\sigma} = 0$ [Proposition 3]. What is not obvious is whether one can choose the set S such that the LTQO condition is satisfied while ensuring that the local terms commute with each other. The main purpose of this Section is to show that this is possible by constructing an explicit example. (Such a set S is not unique, and we provide another valid choice in Appendix A.) The Hamiltonian defined this way will have, by Ref. [15], a gap stable against any extensive but weak enough perturbations. Our construction is based on a decomposition of a two-dimensional plane into 0-, 1-, and 2-cells. We remark that, while we name these objects as cells of different dimensions, they are nevertheless all balls on the plane. In Fig. 5, the 0-, 1-, and 2-cells are denoted in red, blue, and green, respectively. A crucial property that we insist is that each *i*-cell is separated by other *i*-cells by some nonzero distance. We then define a cell c_i , associated with a 2-cell, in the following way. For a given 2-cell, collect all the 0- and 1-cells that neighbor this 2-cell, and take a union. We refer to this union as c_i , also colloquially referred to as a cell in this paper. Our Hamiltonian is defined as

$$H = \sum_{i:2-\text{cells}} (I - P_{c_i}).$$
(56)

There are two important properties of our decomposition. First, this decomposition ensures that the local terms in Eq. (56) commute. For any pair (c_i, c_j) , if c_i and c_j have trivial intersection, P_{c_i} and P_{c_j} trivially commute. When c_i and c_j do overlap, they overlap in such a way that the following condition is satisfied:

$$I(c_i \setminus c_j : c_j \setminus c_i | c_i \cap c_j)_{\sigma} = 0.$$
⁽⁵⁷⁾

(We shall prove this in Section 6.1.) Setting $AB = c_i$ and $BC = c_j$, we can invoke Proposition 3, concluding that the projectors P_{c_i} and P_{c_j} commute. Thus, Eq. (56) is a sum of local commuting projectors.

Second, for every elementary disk D, there exists at least one c_i such that $D \cup N(D) \subset c_i$. (This is a simple fact that can be verified from Fig. 5.) Importantly, this implies that the Hamiltonian Eq. (56) satisfies the LTQO condition [Proposition 2].

Thus, our decomposition serves two purposes: (i) ensuring that the local terms Eq. (56) commute with each other and (ii) the same Hamiltonian satisfies the LTQO condition. By the well-known sufficient condition for the gap stability [15], we can thus conclude that Eq. (56) has a *stable* spectral gap.

6.1 Conditional independence

In this Section, we seek to prove Eq. (57). Note that, when c_i and c_j have nontrivial intersection, there are three different ways in which this can happen. The examples are shown in Fig. 6, with the convention that the red, green, and the blue regions represent $c_i \setminus c_j$, $c_i \cap c_j$, and $c_j \setminus c_i$, respectively.



Figure 6: There are three different ways in which c_i and c_j can overlap. Here red, green, and blue regions represent $c_i \setminus c_j$, $c_i \cap c_j$, and $c_j \setminus c_i$, respectively.

In all three cases, we can prove the requisite conditional independence conditions [Eq. (57)] from an extension of the axiom A1 [Section 3.2]. For instance, suppose we seek to prove

$$I(A:C|B)_{\sigma} = 0 \quad \text{for} \quad A, B, C \text{ in Fig. 7(a)},$$
(58)

setting $A = c_i \setminus c_j$, $B = c_i \cap c_j$, and $C = c_j \setminus c_i$. (A note on convention: In Fig. 7, the subsystem A, B, C, and D are colored in red, green, blue, and yellow.) We can use

$$I(A:C|B)_{\sigma} \le S(C|B)_{\sigma} + S(C|D)_{\sigma} \quad \text{for } A, B, C, D \quad \text{in Fig. 7(a)}$$
(59)

The analogous bounds for the other cases uses the choice of A, B, C, and D shown in Fig. 7(b) and (c).



Figure 7: The choice of subsystems used for upper bounding conditional mutual information. Here red, green, blue, and yellow regions represent $A = c_i \setminus c_j$, $B = c_i \cap c_j$, $C = c_j \setminus c_i$, and D.

The key point is that, in all these cases, one can show $S(C|B)_{\sigma} + S(C|D)_{\sigma} = 0$; this immediately implies Eq. (57). So the natural question is how one can prove $S(C|B)_{\sigma} + S(C|D)_{\sigma} = 0$ for the subsystems shown in Fig. 7. This follows straightforwardly through a simple but repetitive applications of the argument presented in Section 3.2.

7 Gapped domain wall

Our discussion so far pertained to the construction of the parent Hamiltonian of the states satisfying the axioms A0 and A1 everywhere. However, there is a class of states for which A1 is violated along a line. In Ref. [9], some of us advocated viewing such a line as a (codimension-1) defect, also known as the *gapped domain wall*. Despite its name, it was previously unknown if the state in which a gapped domain wall is present is indeed stabilized by a gapped parent Hamiltonian. This is what we intend to prove in this Section.

In the language of Ref. [9], a gapped domain wall can be viewed as a one-dimensional line along which the axiom A1 may be violated. More precisely, let us consider a partition of a 2D plane into hexagonal cells, consisting of the ones that lie at the upper/lower half with respect to a line [Fig. 4]. While A0 continues to hold in the vicinity of every face, and A1 continues to hold in the vicinity of every face that does not touch the domain wall, A1 is modified for the faces that are in contact with the domain wall [Eq. (10), (11)]. To repeat what we said in Section 2, A1 may not hold if the boundaries between B and D are both away from the domain wall and lie on the same side with respect to the domain wall. Otherwise, the A1 continues to hold [9].

We remark that, just like the axioms can be extended in the absence of the gapped domain wall, the axioms for the gapped domain wall can also be extended to larger regions. The argument itself is identical to the one presented in Section 3, with details presented in Ref. [9]. As such, we omit the proof.

The main question we ask at this point is whether one can still show the existence of a gapped parent Hamiltonian with a stable spectral gap. We claim that this is indeed the case. The idea behind the proof is similar to the one presented in Section 6, accompanied by a modified argument for the subsystems that pass through the domain wall. More specifically, we propose to use Eq. (41) as the parent Hamiltonian. The location of the domain wall with respect to the 0-, 1-, and 2-cells are shown in Fig. 8.



Figure 8: Location of the gapped domain wall (red line) with respect to the 0-, 1-, and 2-cells, corresponding to the red, blue, and green balls.

Now the task at hand is to prove that the Hamiltonian Eq. (41), with respect to the decomposition in Fig. 8, has a stable spectral gap. As before, we can conclude so if we can prove the LTQO condition and the local gap condition.

For proving the LTQO condition, we follow the argument similar to the one in Section 4. Consider an arbitrary disk D which, if it intersects with the domain wall, the intersection is a single interval.⁷ Also, consider a disk $D' \subset D$ consisting of faces in D that are separated from ∂D by at least some positive distance, such that over any elementary disk associated with a face in D', any reduced density matrix over this elementary disk over ker H_D is indistinguishable from that of σ . By the discussion in Section 4, such a D' exists. The key question is whether $\rho_{D'} = \sigma_{D'}$ for any $\rho_{D'}$, which is a reduced density matrix of an element of ker H_D over D'. This follows from Lemma 1, by using A0. The proof of Proposition 2 applies verbatim at this point, establishing the LTQO condition.

For proving the local gap condition, it should be obvious that the terms that are away from the domain wall trivially commute with each other by the discussion in Section 5. What remains is to prove that even the terms that touch the domain wall commute with each other. To prove that, by Proposition 3, it suffices to prove $S(B|C)_{\sigma} + S(B|D)_{\sigma} = 0$ for the subsystems shown in Fig. 9. Luckily, these are all facts that follow from the extension of the axioms [9]. Therefore, the requisite conditional independence condition that ensures the commutativity of the local terms of the Hamiltonian [Proposition 3] follows.

To summarize, both the LTQO and the local gap condition continue to be satisfied for the Hamiltonian Eq. (41), even in the presence of gapped domain walls. Thus, we conclude that this Hamiltonian has a stable spectral gap, justifying the name "gapped domain wall."

8 Weight reduction

While the construction of the parent Hamiltonian in Section 6 and 7 are local in the sense that every term in the Hamiltonian acts nontrivially on a ball of O(1) size, that constant is rather large. One might wonder if it is possible to reduce each term's *weight*, defined as the number of faces contained in the support of the term.

To that end, we introduce a simple fact that lets us reduce the weight all the way down to three.

⁷Otherwise, the ensuing argument does not work. For instance, one can consider a disk D whose intersection with the domain wall consists of two intervals. For such a disk, the reduced density matrix of an element of ker H_D over D' may be different from $\sigma_{D'}$ [9].



Figure 9: The choice of subsystems used for upper bounding conditional mutual information in the presence of a gapped domain wall. Here, red, green, blue, and yellow regions represent $A = c_i \setminus c_j, B = c_i \cap c_j, C = c_j \setminus c_i$, and D.

Lemma 4. Let ρ_{ABC} be a state defined on a finite dimensional Hilbert space $\mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$ that satisfies $I(A: C|B)_{\rho} = 0$. Let P_{AB} , P_{BC} , and P_{ABC} be the projector onto the support of ρ_{AB} , ρ_{BC} , and ρ_{ABC} respectively. Then

$$P_{ABC} = P_{AB}P_{BC}.$$
 (60)

Proof. The proof follows immediately from the fact that $P_{b_j^L}P_{Ab_j^L} = P_{Ab_j^L}$, $P_{b_j^R}P_{b_j^RC} = P_{b_j^RC}$, and Eq. (55).

From Lemma 4, it follows that⁸

$$(I - P_{AB}) + (I - P_{BC}) \ge I - P_{ABC}.$$
 (61)

Using Eq. (61) recursively, we can obtain a Hamiltonian H' such that $H' \ge H$, where H is the Hamiltonian defined in Section 6 or in Section 7 and H' consists of terms of weight at most three.

Importantly, note that $\ker(H) = \ker(H')$. This is because the kernel of the left hand side and the right hand side of Eq. (61) are identical due to Lemma 4. Thus H' is also a parent Hamiltonian of the reference state that satisfies the local topological quantum order condition. Repeatedly applying Eq. (61), one can reduce the weight of the local terms all the way down to three. However, one disadvantage of this Hamiltonian is that the local terms may not commute with each other.

9 Discussion

In this paper, we constructed parent Hamiltonians for the states that satisfy the entanglement bootstrap axioms [6, 9]. This parent Hamiltonian is a local commuting projector Hamiltonian that satisfies the local topological quantum order condition. Therefore, this Hamiltonian has a stable spectral gap. Thus, our result establishes that the states that obey the entanglement bootstrap axiom define a stable phase of matter.

The main novel technical insight is Proposition 3, which relates the conditional independence of the ground state to the commutativity of the projectors onto the supports of different local reduced density matrices. Armed with this fact, constructing a parent Hamiltonian reduces to the problem of finding an appropriate decomposition of the lattice that satisfies a certain combinatorial property [Section 6].

⁸Explicitly, $Q \equiv 1 - P$ is a projector. Eq. (60) implies $1 - Q_{ABC} = (1 - Q_{AB})(1 - Q_{BC}) = (1 - Q_{BC})(1 - Q_{AB})$. Thus $Q_{AB} + Q_{BC} = Q_{ABC} + Q_{AB}Q_{BC} \ge Q_{ABC}$. The last inequality needs the fact that $Q_{AB}Q_{BC}$ is another projector, which follows from $Q_{AB}Q_{BC} = Q_{BC}Q_{AB}$.

We expect this insight to be useful for a number of applications, which we summarize below. First, it is likely that our construction continues to work in higher dimensions, such as in three spatial dimensions [10, 11]. One may also consider parent Hamiltonians for states that host topological excitations. It may be possible to also define a "refined" parent Hamiltonian whose excitations label the anyon excitations. We leave such studies for future work.

We also remark that there is an interesting tension between our work and the formula for the chiral central charge [12, 13]. This formula was derived under A1, modulo an extra physical assumption. However, what we showed was that, A1, together with A0, implies that there ought to be a parent Hamiltonian which is a local commuting projector. Such a system cannot have any edge energy current [22, 4], and as such, would be in tension with any nonzero value of the chiral central charge. More precisely, our result suggests that the entanglement bootstrap axioms cannot be satisfied *exactly* for ground states of systems that have nonzero chiral central charge, at least if each site is described by a finite-dimensional Hilbert space. Physically, this is because such a Hamiltonian necessarily has a vanishing energy current at the edge at all temperatures, which is possible only if the chiral central charge is zero.

There is, nonetheless, a lingering question concerning the recently proposed modular commutator formula [12] for chiral central charge c_{-} :

$$J(A, B, C) = \frac{\pi}{3}c_{-},$$

where A, B, C is a partition of a disk into three charts meeting at a point. Do we always have J(A, B, C) = 0, under exact **A0** and **A1**? We leave it as an open problem.

Our work raises several natural questions, which we leave for future work. First, we note that having a parent Hamiltonian with the local TQO condition is a property invariant under a constant-depth circuit, whereas the axiom A1 is not. Therefore, the existence of such a parent Hamiltonian is a strictly weaker assumption that holds for all states in the same phase as a state that satisfies A0 and A1.⁹ What can you prove from this weaker assumption? In particular, can one reproduce the results in Ref. [6]?

Second, it will be interesting to understand if our approach for constructing a commuting parent Hamiltonian works even if the local Hilbert space dimension is infinite. In fact, we think there are physical reasons to believe this is impossible, as we explain below. Note that our proof relied on the structure theorem for quantum Markov chains [21]. To the best of our knowledge, an infinite-dimensional analog of this result is not currently known. If such a result can be proven and if one accepts that the entanglement bootstrap axioms can be satisfied even if the chiral central charge is nonzero (by having infinite local Hilbert space dimension), we are led to the conclusion that there can be a commuting parent Hamiltonian, even for systems with nonzero chiral central charge! However, this seems unlikely because the energy current produced by such Hamiltonian must necessarily be zero. On this physical ground, one may expect that the structure theorem for quantum Markov chains breaks down in infinite dimensions. Making this argument more precise or coming up with a rigorous counterexample is left for future work.

Acknowledgement

IK acknowledges supports from NSF under award number PHY-2337931. DR acknowledges funding from NTT (Grant AGMT DTD 9/24/20). BS was supported by the Simons Collaboration on Ultra-Quantum Matter, a grant from the Simons Foundation (652264 JM). TCL is supported in part by funds provided by the U.S. Department of Energy (D.O.E.) under the cooperative research agreement DE-SC0009919 and by the Simons Collaboration on Ultra-Quantum Matter, which is a grant from the Simons Foundation (652264 JM).

⁹More precisely, while the local TQO condition is typically assumed only for square-shaped regions [18] or (geometric) balls [15], we may demand this condition for any region that is *topologically* a ball. Simply demanding local TQO for square-shaped regions or geometric balls would not reproduce Ref. [6]; such a condition is satisfied even for systems with gapped boundaries/domain walls, wherein the anyon contents are different on each side of the domain wall.

A Another explicit parent Hamiltonian

We constructed a commuting parent Hamiltonian Eq. (41) satisfying the local topological quantum order condition. Such a construction rests on the existence of a set S with a couple of special properties. In Section 6, we explicitly constructed a valid such set S. We emphasize that the specific form of the regions in the set S is not important. What is important is the following set of properties.

Definition 5 (Markov cover). We say the set of subsystems \mathcal{S} a Markov cover if it satisfies

- 1. (Markov condition) For $\forall X, Y \in \mathcal{S}$ such that $X \cap Y \neq \emptyset$, The conditional mutual information of the reference state $I(X \setminus Y, Y \setminus X | X \cap Y)_{\sigma} = 0$.
- 2. (cover condition) For every elementary disk D, there exists some $X \in S$ such that $D \cup N(D) \subset X$.

As one may expect, there are many alternative choices. In this Appendix, we provide another valid set S for illustration purposes. Consider a set of red hexagons on the hexagon lattice [Fig. 10]. Note that the



Figure 10: Partition used for a valid set \mathcal{S} .

distance between nearby red hexagons is equal to five.

Let us consider the following set:

 $\mathcal{S} = \{X | X = \text{a red face } f_{\text{red}} \cup \text{faces within distance 5 to } f_{\text{red}} \text{ that are not red}\}.$ (62)

It immediately follows that every $X \in S$ is identical in its shape, shown below:



As a matter of fact, the set S in Eq. (62) is a Markov cover. We need to check two properties, which explain below. First, the set S satisfies the Markov property; see the first item in Definition 5. The relative

positions between two regions $X, Y \in \mathcal{S}$, up to rotations, are the following



Thus, the Markov property can be checked explicitly, as we have done in Fig. 6 and 7. Second, we would like to check the cover condition (i.e., the second item of Definition 5). For any elementary disk D centered at a face f_D , we find its nearest red face f'_{red} .¹⁰ From the construction, we see that the distance between f_D and f'_{red} can only be 0, 1, 2, 3. We end up with a small set of cases to check. In all such cases, indeed $D \cup N(D)$ is contained in the cover X centered at the red face f'_{red} . Thus, the Hamiltonian Eq. (41) is a commuting Hamiltonian that satisfies LTQO if we choose the set S as Eq. (62).

A.1 Reference states with anyon excitations

Consider a known Hamiltonian with topological order and non-abelian anyons, such as a string-net model. Say we take our reference state to be not the ground state, but rather some state with non-abelian anyons present. Their presence can be diagnosed by violations of **A1**. Can we still construct a corresponding parent Hamiltonian? Or more generally, what happens if we consider a reference state with **A1** violations in some regions?

We comment on this briefly using the set S in Eq. (62). The nontrivial case is to have a non-Abelian anyon in the center of a disk, i.e. at the red hexagon in (63). Suppose the quantum dimension of the anyon is d_a ($d_a \ge \sqrt{2}$ for non-Abelian anyons), the constant sub-leading term in Eq. (1) is modified by $\log d_a$ if disk A contains an anyon [7]:

$$S(A) = \alpha |\partial A| - \gamma + \log d_a.$$
⁽⁶⁵⁾

Suppose there are multiple non-Abelian anyons on the plane, each is centered at some $X \in S$, then the parent Hamiltonian may have multiple ground states even on a topologically trivial manifold. Nonetheless, the steps leading to the stable energy gap (as in Section 6) still go through, provided that the anyons are well separated from each other. In particular, the resulting parent Hamiltonian Eq. (41) is commuting as long as those $X_i, X_j \in S$ containing anyons at the center are such that $X_i \cap X_j = \emptyset$. This is because, while **A1** may be violated, the conditional independence condition remains true. The additional contribution of log d_a to the entanglement entropy cancels out each other in the conditional mutual information.

References

- Xie Chen, Zheng-Cheng Gu, and Xiao-Gang Wen, Classification of gapped symmetric phases in onedimensional spin systems, Phys. Rev. B 83 (2011), 035107.
- [2] Norbert Schuch, David Pérez-García, and Ignacio Cirac, Classifying quantum phases using matrix product states and projected entangled pair states, Phys. Rev. B 84 (2011), 165139.
- [3] Lukasz Fidkowski and Alexei Kitaev, Topological phases of fermions in one dimension, Phys. Rev. B 83 (2011), 075103.

¹⁰If such a choice is not unique, pick one.

- [4] Alexei Kitaev, Anyons in an exactly solved model and beyond, Annals of Physics 321 (2006), no. 1, 2–111.
- [5] Jeongwan Haah, Local stabilizer codes in three dimensions without string logical operators, Phys. Rev. A 83 (2011), 042330.
- [6] Bowen Shi, Kohtaro Kato, and Isaac H Kim, Fusion rules from entanglement, Annals of Physics 418 (2020), 168164.
- [7] Alexei Kitaev and John Preskill, Topological Entanglement Entropy, Phys. Rev. Lett. 96 (2006), no. 11, 110404, hep-th/0510092.
- [8] Michael Levin and Xiao-Gang Wen, Detecting Topological Order in a Ground State Wave Function, Phys. Rev. Lett. 96 (2006), no. 11, 110405, cond-mat/0510613.
- Bowen Shi and Isaac H. Kim, Entanglement bootstrap approach for gapped domain walls, Phys. Rev. B 103 (2021), 115150.
- [10] Jin-Long Huang, John McGreevy, and Bowen Shi, Knots and entanglement, SciPost Phys. 14 (2023), 141.
- [11] Bowen Shi, Jin-Long Huang, and John McGreevy, Remote detectability from entanglement bootstrap i: Kirby's torus trick, 2023, 2301.07119.
- [12] Isaac H. Kim, Bowen Shi, Kohtaro Kato, and Victor V. Albert, Chiral central charge from a single bulk wave function, Phys. Rev. Lett. 128 (2022), 176402.
- [13] Isaac H. Kim, Bowen Shi, Kohtaro Kato, and Victor V. Albert, Modular commutator in gapped quantum many-body systems, Phys. Rev. B 106 (2022), 075147.
- [14] Ting-Chun Lin and John McGreevy, Conformal field theory ground states as critical points of an entropy function, 2023, 2303.05444.
- [15] Spyridon Michalakis and Justyna P. Zwolak, Stability of frustration-free hamiltonians, Communications in Mathematical Physics 322 (2013), no. 2, 277–302.
- [16] Dénes Petz, Sufficiency of channels over von Neumann algebras, The Quarterly Journal of Mathematics 39 (1988), no. 1, 97–108, https://academic.oup.com/qjmath/article-pdf/39/1/97/4559225/39-1-97.pdf.
- [17] Pieter Naaijkens, Quantum spin systems on infinite lattices, Springer, 2013.
- [18] Sergey Bravyi, Matthew B. Hastings, and Spyridon Michalakis, Topological quantum order: Stability under local perturbations, Journal of Mathematical Physics 51 (2010), no. 9, 093512.
- [19] A. Uhlmann, The "transition probability" in the state space of a *-algebra, Reports on Mathematical Physics 9 (1976), no. 2, 273–279.
- [20] Isaac H. Kim, On the informational completeness of local observables, 2014, 1405.0137.
- [21] Patrick Hayden, Richard Jozsa, Dénes Petz, and Andreas Winter, Structure of states which satisfy strong subadditivity of quantum entropy with equality, Communications in Mathematical Physics 246 (2004), no. 2, 359–374.
- [22] C. L. Kane and Matthew P. A. Fisher, Quantized thermal transport in the fractional quantum hall effect, Phys. Rev. B 55 (1997), 15832–15837.