THE FINE-GRAINED COMPLEXITY OF GRAPH HOMOMORPHISM PROBLEMS: TOWARDS THE OKRASA AND RZĄŜEWSKI CONJECTURE

AMBROISE BARIL, MIGUEL COUCEIRO, AND VICTOR LAGERKVIST

ABSTRACT. In this paper we are interested in the fine-grained complexity of deciding whether there is a homomorphism from an input graph G to a fixed graph H (the H-COLORING problem). The starting point is that these problems can be viewed as constraint satisfaction problems (CSPs), and that (partial) polymorphisms of binary relations are of paramount importance in the study of complexity classes of such CSPs.

Thus, we first investigate the expressivity of binary symmetric relations E_H and their corresponding (partial) polymorphisms $\operatorname{POl}(E_H)$. For irreflexive graphs we observe that there is no pair of graphs H and H' such that $\operatorname{POl}(E_H) \subseteq \operatorname{POl}(E_{H'})$, unless $E_{H'} = \emptyset$ or H = H'. More generally we show the existence of an *n*-ary relation R whose partial polymorphisms strictly subsume those of H and such that $\operatorname{CSP}(R)$ is NP-complete if and only if H contains an odd cycle of length at most n. Motivated by this we also describe the sets of total polymorphisms of nontrivial cliques, odd cycles, as well as certain cores, and we give an algebraic characterization of *projective cores*. As a by-product, we settle the *Okrasa and Rzążewski conjecture* for all graphs of at most 7 vertices.

1. INTRODUCTION

This paper aims to improve our understanding of fine-grained complexity of constraint satisfaction problems (CSPs) [11]. In a constraint satisfaction problem (CSP), given a set of variables X and a set of constraints of the form $R(\mathbf{x})$ for $\mathbf{x} \in X^k$ and some k-ary relation R, the objective is to assign values from X to a domain V such that every constraint in C is satisfied. This problem is usually denoted by $\text{CSP}(\Gamma)$, with the additional stipulation that every relation occurring in a constraint comes from the set of relations Γ , and it is typically phrased as the decision problem of verifying whether a solution exists.

In this article we take a particular interest in the restricted case when Γ is singleton and contains a binary, symmetric relation H, viewed as the edge relation of a graph¹, the H-COLORING problem [9]. This problem is arguably more naturally formulated as a homomorphism problem. Recall that a function $f: V_G \to V_H$ is said to be a homomorphism between the two graphs G and H if it "preserves" the edge relation, that is, if for every edge $(u, v) \in E_G$, we have $(f(u), f(v)) \in E_H$. In that case, we use the notation $f: G \to H$. For each graph H the H-COLORING problem can then succinctly be defined as follows.

H-COLORING PROBLEM. Given a graph G, decide whether there is a homomorphism $f: G \to H$.

Note that *H*-COLORING is the same problem as $CsP({E_H})$ (henceforth written $CsP(E_H)$). Clearly, the *H*-COLORING problem subsumes the well-known *k*-COLORING problem, $k \ge 1$, that asks for a coloring of the vertices of a graph using at most *k* colors such that each pair of adjacent vertices are assigned different colors. Indeed, it corresponds to the case where $H = K_k$, the complete graph (clique) of size *k*.

Hell and Nešetřil [9] showed that the *H*-COLORING problem is in P (the class of problems decidable in polynomial time) whenever *H* is bipartite, and it is NP-complete otherwise. Our goal is to bring some light into the (presumably) superpolynomial complexity of the *H*-COLORING problem when *H* is non-bipartite. On the one hand, there are already some strong upper-bounds results on the fine-grained complexity of *k*-COLORING for $k \ge 3$. Björklund *et al.* [2] proved that *k*-COLORING is solvable in time $O^*(2^n)$ (i.e., $O(2^n \times n^{O(1)})$) where *n* is the number of vertices of the input graph. Fomin *et al.* [7] also prove that C_{2k+1} -COLORING is solvable in time $O^*(\binom{n}{n/k}) = O^*((\alpha_k)^n)$, with $\alpha_k \xrightarrow[k\to\infty]{} 1$, and improved algorithms are also known when *H* has bounded *tree-width* or *clique-width* [7, 19]. On the other hand, lower bounds by Fomin *et al.* [6] rule out

Key words and phrases. Graph coloring, Graph homomorphism, Partial polymorphism, Clique, Projective graph, Core graph, Okrasa and Rzążewski conjecture .

¹Throughout, we assume that all graphs are finite, simple (no loops) and undirected. Every graph H is defined by its set V_H of vertices and its set E_H of edges.

the existence of a uniform $2^{O(n)}$ time algorithm under the *exponential-time hypothesis* (*i.e.*, that 3-SAT can not be solved in subexponential time).

We notice, however, that there is a lack of general tools for describing fine-grained properties of CSPs, and in particular we lack techniques for comparing NP-hard *H*-COLORING problems with each other, e.g., via size-preserving reductions. We explore these ideas through an algebraic approach, by investigating algebraic invariants of graphs. For this purpose, viewing *H*-COLORING as $CSP(E_H)$ is quite useful as it allows us to use the widely studied theory of the complexity of $CSP(\Gamma)$, since the former is just the special case when $\Gamma = \{R\}$ is a singleton containing a binary, symmetric relation. In particular, it was shown that the fine-grained complexity of CSP(R) only depends on the so-called *partial polymorphisms* of R [5, 11]. Briefly, a *polymorphism* is a higher-arity homomorphism from the relation to the relation itself. Additionally, a polymorphism that is not necessarily everywhere defined is known as a *partial polymorphism*, and we write Pol(R) (respectively, PPol(R)) for the set of all (partial) polymorphisms of a relation E_H . It is then known that partial polymorphisms correlate to fine-grained complexity in the sense that if $PPol(R) \subseteq PPol(R')$ and if CSP(R) is solvable in $O^*(c^n)$ time for some c > 1 then CSP(R') is also solvable in $O^*(c^n)$ time [11].

Thus, describing the inclusion structure between sets of the form $\operatorname{pPol}(H)$ would allow us to relate the fine-grained complexity of *H*-COLORING problems with each other, but, curiously, we manage to prove that *no* non-trivial inclusions of this form exist, suggesting that partial polymorphisms of graphs are not easy to relate via set inclusion. As a follow-up question we also study inclusions of the form $\operatorname{pPol}(H) \subseteq \operatorname{pPol}(R)$ when *R* is an arbitrary relation, and manage to give a non-trivial condition based on the length of the shortest odd cycle of *H*. Concretely, we prove that it is possible to find an *n*-ary relation *R* with $\operatorname{pPol}(E_H) \subsetneq \operatorname{pPol}(R)$ where $\operatorname{CSP}(R)$ is NP-complete, if and only if *H* contains an odd-cycle of length at most *n*. This result suggests that the size of the smallest odd-cycle is an interesting parameter when regarding the complexity of *H*-COLORING. As observed above, the smaller $\operatorname{pPol}(E_H)$ is, the harder $\operatorname{CSP}(E_H)$ (and thus *H*-COLORING) is. In other words, the greater the smallest odd-cycle of *H* is, the easier the *H*-COLORING problem is. This fact supports the already known algorithms presented in [7].

Despite this trivial inclusion structure, it could still be of great interest to provide a succinct description of pPol(H) for some noteworthy choices of non-bipartite, core H. As a first step in this project we concentrate on the total polymorphisms of H, and conclude that *projective graphs* [13] appear to be a reasonable class to target since the total polymorphisms of projective cores are *essentially at most unary*. Projective cores were studied by Okrasa and Rzążewski [15] in the context of fine-grained aspects of H-COLORING problems analyzed under tree-width. We have the following conjecture.

Okrasa and Rzążewski Conjecture ([15]). Let H be a connected non-trivial core on at least 3 vertices. Then H is projective if and only if it is indecomposable.

Thus, we should not hope to easily give a complete description of projective cores, but we do succeed in (1) proving that several well-known families of graphs, e.g., cliques, odd-cycles, and other core graphs, are projective, and (2) confirm the conjecture for *all* graphs with at most 7 vertices. Importantly, our proofs use the algebraic approach and are significantly simpler than existing proofs, and suggest that the algebraic approach might be a cornerstone in completely describing projective cores.

This paper is organized as follows. In Section 2, we recall the basic notions and preliminary results needed throughout the paper. We investigate the order structure of classes of graph (partial) polymorphisms in Section 3 where we show the aforementioned main results. In Section 4 we focus on projective and core graphs and present several general examples of projective cores, and settle the Okrasa and Rzążewski conjecture for graphs with at most 7 vertices. In Section 6 we discuss some consequences of our results and state a few noteworthy conjectures.

2. Preliminaries

Throughout the article we use the following notation.

For any $n \in \mathbb{N}$, [n] denotes the set $\{1, \ldots, n\}$. For every set $V, n \ge 1$ and $t = (t_1, \ldots, t_n) \in V^n$, t[i] denotes t_i , and given a relation $R \subseteq [k]^n$ for some $k \ge 1$, we write $\operatorname{ar}(R)$ for its arity n. For all $m \ge 1$ and $i \in [m]$, we write $\pi_i^m : V^m \to V$ for the *projection* on the *i*-th coordinate (the set V will always be implicit in the context).

For H a graph and $V \subseteq V_H$, we denote by H[V] the graph induced by V on H. We use the symbol \oplus to express the disjoint union of sets, and + to express the disjoint union of graphs.

For a unary function $f: V \to V$, and an *m*-ary function $g: V^m \to V$ we write $f \circ g$ for their composition that is defined by $(f \circ g)(x_1, \ldots, x_m) = f(g((x_1, \ldots, x_m)))$, for every $(x_1, \ldots, x_m) \in V^m$.

Also, for $k \ge 3$, K_k and C_k denote respectively a k-clique and a k-cycle.

2.1. Graph homomorphisms and cores. For two graphs G and H a function $f: V_G \to V_H$ is a homomorphism from G to H if $\forall (u, v) \in E_G$, $(f(u), f(v)) \in E_H$. In this case, f is also called an H-coloring of G, and we denote this fact by $f: G \to H$. The graph G is said to be H-colorable, which we denote by $G \to H$, if there exists $f: G \to H$. For a graph H, the H-COLORING problem thus asks whether a given graph G is H-colorable.

Theorem 1 ([9]). H-COLORING is in P whenever H is bipartite, and is NP-complete, otherwise.

A key notion in the proof of Theorem 1 is the notion of a graph core: let core(H) be the smallest induced subgraph H' of H such that $H \to H'$. The graph H is said to be a core if H = core(H). Note that the core of a graph H is unique up to isomorphism and that the problems H-COLORING and core(H)-COLORING are equivalent. Thus, for both classical and fine-grained complexity, it is sufficient to consider core(H)-COLORING. Moreover, it is not difficult to verify that cliques and odd-cycles are cores. Notice that a graph H is a core if and only if every H-coloring of H is bijective.

For two graphs G and H, we define their cross product $G \times H$ as the graph with $V_{G \times H} = V_G \times V_H$ and

$$E_{G \times H} = \{ ((u_1, v_1), (u_2, v_2)) \mid (u_1, u_2) \in E_G, (v_1, v_2) \in E_H \}.$$

Clearly, for graphs A, B and C, we have that $(V_A \times V_B) \times V_C$ and $V_A \times (V_B \times V_C)$ are in bijection and thus, up to isomorphism, the cross product is associative. Hence, for each $m \ge 1$, we can define $H^m = \underbrace{H \times \ldots \times H}_{m \text{ times}}$.

Last, we need the following graph parameter, defined with respect to the smallest odd-cycle in the graph.

Definition 2. The odd-girth of a non-bipartite graph H (denoted by og(H)) is the size of a smallest odd-cycle induced in H.

For a bipartite graph we define the odd-girth to be infinite.

2.2. Polymorphisms, pp/qfpp-definitions. Even though the previous definitions apply only to graphs, *i.e.*, binary symmetric and irreflexive relations, we will need to introduce the following notions for relations R of arbitrary arity.

Definition 3. Let V be a finite set, $n, m \ge 1$ be integers, and let $R \subseteq V^n$ be an n-ary relation on V. A partial function $f : dom(f) \to V$, with $dom(f) \subseteq V^m$, is said to be a partial polymorphism of R if for every $n \times m$ matrix $A = (A_{i,j}) \in V^{n \times m}$ such that for every $j \in [m]$, the j-th column $A_{*,j} \in R$ and for every $i \in [n]$, the i-th row $A_{i,*} \in dom(f)$, the column $(f(A_{1,*}), \ldots, f(A_{n,*}))^\top \in R$. In the case when $dom(f) = V^m$, f is a said to be a total polymorphism (or just a polymorphism) of R. We denote the sets of total and partial polymorphisms of R by Pol(R) and pPol(R), respectively.

Every (partial) function over a set V is a (partial) polymorphism of both the empty relation (denoted by \emptyset) and the equality relation $EQ_V = \{(x, x) \mid x \in V\}$ over V (or simply EQ when the domain is clear from the context).

For a graph H, we sometimes use POl(H) and Pol(H) instead of $\text{POl}(E_H)$ and $\text{Pol}(E_H)$, where E_H is viewed as a binary relation over the domain V_H . Note that Pol(H) is exactly the set of H-colorings of H^m for $m \ge 1$, and that POl(H) is exactly the set of H-colorings of the induced subgraphs of H^m for $m \in \mathbb{N}$.

Definition 4. Let R be a relation over a finite domain V. An n-ary relation R' over V is said to have a primitive positive-definition (pp-definition) w.r.t. R if there exists $m, m', n' \in \mathbb{N}$ such that

(1)
$$R'(x_1,\ldots,x_n) \equiv \exists x_{n+1},\ldots,x_{n+n'}$$
:

 $R(\mathbf{x}_1) \wedge \ldots \wedge R(\mathbf{x}_m) \wedge EQ(\mathbf{y}_1) \wedge \ldots \wedge EQ(\mathbf{y}_{m'})$

where each \mathbf{x}_i is an $\operatorname{ar}(R)$ -ary tuple and each \mathbf{y}_i is an binary tuple of variables from $x_1, \ldots, x_n, x_{n+1}, \ldots, x_{n+n'}$. Each term of the form $R(\mathbf{x}_i)$ or $EQ(\mathbf{y}_j)$ is called an atom or a constraint of the pp-definition (1).

In addition, if n' = 0, then (1) is called a *quantifier-free primitive positive-definition* (qfpp-definition) of R'. Let $\langle R \rangle_{\sharp}$ and $\langle R \rangle$ be the sets of qfpp-definable and of pp-definable, respectively, relations over R.

Theorem 5 ([17]). Let R and R' be two relations over the same finite domain. Then

- (1) $R' \in \langle R \rangle_{\frac{d}{2}}$ if and only if $pPol(R) \subseteq pPol(R')$ and (2) $R' \in \langle R \rangle$ if and only if $Pol(R) \subseteq Pol(R')$.

2.3. Csps and polymorphisms. We now recall the link between the complexity of CSPs and the algebraic tools described in the previous section (recall that the *H*-COLORING problem is the same problem as $\operatorname{Csp}(E_H)$).

Theorem 6 ([10]). Let R and R' be two relations over the same finite domain where $Pol(R) \subseteq Pol(R')$. Then $\operatorname{CSP}(R')$ is polynomial-time many-one reducible to $\operatorname{CSP}(R)$.

Let R be a relation over a finite domain V. Define:

 $\mathsf{T}(R) = \inf\{c > 1 : \operatorname{CSP}(R) \text{ is solvable in time } O^*(c^n)\}$

where n is the number of variables in a given CSP(R) instance, (with the notation $O^*(v_n) = O(v_n \times n^{O(1)})$ for all $(v_n)_{n \in \mathbb{N}} \in \mathbb{R}^{\mathbb{N}}$).

Theorem 7 ([11]). Let R and R' be relations over a finite domain V. If $pPol(R) \subseteq pPol(R')$, then $\mathsf{T}(R') \leq \mathsf{T}(R).$

These two theorems motivate our study of polymorphisms of graphs: since $Csp(E_H)$ is the same problem as H-COLORING, key information about the fine-grained complexity of H-COLORING is contained in the set $pPol(E_H).$

3. The inclusion structure of partial polymorphisms of graphs

In this section we study the inclusion structure of sets of the form pPol(H) when H is a graph with $V_H = V$ for a fixed, finite set V. In other words, we are interested in describing the set

$$\mathcal{H} = \{ \operatorname{pPol}(H) \mid H \text{ is a graph over } V \}$$

partially ordered by set inclusion. Here, one may observe that the requirement that $V_H = V_{H'} = V$ is not an actual restriction. Indeed, if $V_{H'} \subsetneq V$, then we can easily obtain a graph over V simply by adding isolated vertices, with no impact on the set of partial polymorphisms.

3.1. Trivial inclusion structure. Our starting point is to establish $pPol(H) \subseteq pPol(H')$ when H, H' are non-bipartite graphs, since it implies that (1) H-COLORING and H'-COLORING are both NP-complete, and (2) $\mathsf{T}(H') \leq \mathsf{T}(H)$, i.e., that H'-COLORING is not strictly harder than H-COLORING.

Inclusions of this kind e.g. raise the question whether there exist, for every fixed finite domain V, an NP-hard H-COLORING problem which is (1) maximally easy, or (2) maximally hard².

As we will soon prove, the set \mathcal{H} does not admit any non-trivial inclusions, in the sense that $pPol(\mathcal{H}) \subseteq$ pPol(H') implies that either H = H' or $E_{H'} = \emptyset$, for all pPol(H), $pPol(H') \in \mathcal{H}$.

Theorem 8. Let H and H' be two graphs with the same finite domain $V_H = V_{H'} = V$. Then $pPol(H) \subseteq$ pPol(H') if and only if H = H' or $E_{H'} = \emptyset$.

Proof. To prove sufficiency, assume that H = H' or that H' has no edges. Then $\operatorname{pPol}(H) = \operatorname{pPol}(H')$ or $pPol(H) \subseteq pPol(H')$ since in the latter case pPol(H') contains every partial function.

To prove necessity, assume that $pPol(H) \subseteq pPol(H')$. Then, by Theorem 5, E_H qfpp-defines $E_{H'}$. However, the only possible atoms using E_H and two variables x and y are: (1) $E_H(x,x)$ and $E_H(y,y)$, which cannot appear by irreflexivity, unless $E_{H'} = \emptyset$ and (2) $E_H(x, y)$ and $E_H(y, x)$, which are equivalent by symmetry. Also, if the qfpp-definition would contain an equality constraint EQ(x, y), then $E_{H'}$ would not be irreflexive, unless $E_{H'} = \emptyset$. Hence, any qfpp-definition of $E_{H'}$ either (1) contains $E_H(x,x)$, $E_H(y,y)$ or EQ(x,y), meaning that $E_{H'} = \emptyset$, or (2) only contains $E_H(x, y)$ or $E_H(y, x)$, meaning that H = H'.

²Here, "maximally" refers to the function T.

3.2. **Higher-arity inclusions.** As proven in Theorem 8, the expressivity of binary irreflexive symmetric relations is rather limited, in the sense that \mathcal{H} does not admit any non-trivial inclusions. It is thus natural to ask whether anything at all can be said concerning inclusions of the form $pPol(H) \subseteq pPol(R)$ when R is an arbitrary relation. In particular, under which conditions does there exist an n-ary R such that $pPol(H) \subsetneq pPol(R)$, given that H-COLORING and CSP(R) are both NP-complete? We give a remarkably sharp classification and, assuming that $\mathsf{P} \neq \mathsf{NP}$, we prove that an *n*-ary relation R with the stated properties exists if and only if H contains an odd-cycle of length $\leq n$. We first require the following auxiliary lemma (recall the definition of odd-girth from Section 2).

Lemma 9. Let H be a non-bipartite graph and let k := og(H). Let $(x_1, \ldots, x_k) \in (V_H)^k$. Then, $x_1x_2 \ldots x_kx_1$ forms an induced k-cycle in H if and only if $(x_1, x_2), (x_2, x_3), \dots, (x_{k-1}, x_k)$ and (x_k, x_1) are edges in H.

Proof. First, notice that if $x_1x_2\ldots x_kx_1$ forms an induced k-cycle in H, then $(x_1, x_2), (x_2, x_3), \ldots, (x_{k-1}, x_k)$ and (x_k, x_1) are edges in H, which proves sufficiency.

To prove necessity, assume that $(x_1, x_2), (x_2, x_3), \ldots, (x_{k-1}, x_k)$ and (x_k, x_1) are edges in H. Consider a smallest odd-cycle $C = x'_1 \dots x'_p x'_1$ (with p odd and $p \leq k$) that is a subgraph (not necessarily induced) of $H[\{x_1 \dots x_k\}]$. Such an odd-cycle exists because $x_1 \dots x_k x_1$ is an odd-cycle. We prove by contradiction that C is induced in H. If C is not induced in H, there exists an edge $(x'_i, x'_j) \in E_H$ with i < j and $j - i \notin \{1, p - 1\}$. Then either j - i is odd and $x'_1 \dots x'_{i-1} x'_i x'_j x'_{j+1} \dots x'_p x'_1$ is an odd-cycle (of size p - (j - i) + 1 < p) smaller than C that is a subgraph of $H[\{x_1 \dots x_k\}]$, which contradicts the definition of C, or j - i is even and $x'_i x'_{i+1} \dots x'_j x'_i$ is an odd-cycle (of size (j-i) + 1 < p) smaller than C that is a subgraph of $H[\{x_1 \dots x_k\}]$, which contradicts the definition of C.

By definition of k := og(H), the induced odd-cycle C has at least k vertices. Since C is an induced subgraph of $H[\{x_1 \ldots x_k\}]$ with k vertices that is an odd-cycle, $x_1 \ldots x_k x_1$ induces a k-cycle, which proves necessity.

The following definition and lemma are particularly usefull when establishing our classification.

Definition 10. Let $n, m \ge 1$ be integers, H be a graph, R be a relation of arity n over V_H , and let $M = (M_{i,j})$ be a $n \times m$ matrix of elements of V_H . We say that M is an R-wall for H if:

- (1) $\forall j \in [m], (M_{1,j}, ..., M_{n,j})^{\top} \in R, and$ (2) $\forall (i, i') \in [n]^2, \exists j \in [m], (M_{i,j}, M_{i',j})^{\top} \notin E_H.$

In the following lemma, we say, for a relation R, that CsP(R) is trivial if every instance of CsP(R) is satisfiable. Clearly, if CSP(R) is trivial, it is not NP-complete, even if P=NP.

Lemma 11. Let H be a graph and let R be an n-ary relation over V_H . Suppose that $pPol(H) \subseteq pPol(R)$ and that there exists an R-wall M for H. Then, Csp(R) is trivial.

Proof. Using property 2) of Definition 10, it is easy to check that any partial function f whose domain is the set of rows of M is in pPol(H). In particular, f can be chosen to be of constant value $a \in V_H$. Then, from $pPol(H) \subseteq pPol(R)$ it follows that $f \in pPol(R)$. Combining this with property 1) of Definition 10, we conclude that $(a, \ldots, a)^{\top} \in R$. Since the valuation sending all variables to a satisfies any instance of CsP(R), the proof is now complete.

We now propose a construction of an R-wall for a graph H with $n := \operatorname{ar}(R) < og(H)$, and such that $pPol(H) \subsetneq pPol(R).$

Lemma 12. Let H be a graph with og(H) > n, and let $R \neq \emptyset$ be an n-ary relation such that $pPol(H) \subseteq$ pPol(R). If $\forall (x_1, \ldots, x_n) \in (V_H)^n, R(x_1, \ldots, x_n) \implies E_H(x_1, x_2)$, then R approductions E_H .

Proof. Suppose that $R(x_1, \ldots, x_n) \implies E_H(x_1, x_2)$, and let $(a_1, \ldots, a_n) \in R \neq \emptyset$. Since H has no odd-cycle of size $\leq n$, $\{a_1, \ldots, a_n\}$ induces a bipartite graph in H: there exists a partition $A \oplus B$ of $\{a_1, \ldots, a_n\}$ such that $E_{H[\{a_1,\ldots,a_n\}]} \subseteq (A \times B) \oplus (B \times A). \text{ For } (x,y) \in E_H, \text{ define } f_{x,y} \colon \{a_1,\ldots,a_n\} \to V_H \text{ by } f_{x,y}(a_i) = x, \text{ if } a_i \in A, \text{ and } f_{x,y}(a_i) = y, \text{ if } a_i \in B. \text{ Since } (x,y) \in E_H, \text{ we have that } f_{x,y} \in \text{POl}(H), \text{ and since } \text{POl}(H) \subseteq \text{POl}(R), \text{ we also have that } f_{x,y} \in \text{POl}(R). \text{ As } (a_1,\ldots,a_n) \in R, \text{ it follows } (f_{x,y}(a_i))_{1 \leq i \leq n} \in R.$

This proves that $E_H(x,y) \implies R(\mathbf{x}_{A,B}(x,y))$, where $\mathbf{x}_{A,B}(x,y)[i] := f_{x,y}(a_i)$ equals x if $a_i \in A$ and $\mathbf{x}_{A,B}(x,y)[i] = y$ if $a_i \in B$.

Reversely, since $(a_1, \ldots, a_n) \in R$ and $R(x_1, \ldots, x_n) \implies E_H(x_1, x_2)$, we have $(a_1, a_2) \in E_H$. Recall that $E_H \subseteq (A \times B) \oplus (B \times A)$: it follows by definition of $\mathbf{x}_{A,B}(x, y)$ that for all vertices x and y of H, that $\{\mathbf{x}_{A,B}(x,y)[1], \mathbf{x}_{A,B}(x,y)[2]\} = \{x,y\}$. From the fact that E_H is symmetric and the hypothesis that $R(x_1, \ldots, x_n) \implies E_H(x_1, x_2)$, we deduce that $R(\mathbf{x}_{A,B}(x,y)) \implies E_H(x,y)$. Hence, $E_H(x,y) \equiv R(\mathbf{x}_{A,B}(x,y))$, and R qfpp-defines E_H .

Lemma 13. Let $n \ge 1$, H be a graph with og(H) > n, and let $R \ne \emptyset$ be an n-ary relation such that $pPol(H) \subsetneq pPol(R)$. Then, for all $(i,i') \in [n]^2$ with i < i', there is $(x_1^{(i,i')}, \ldots, x_n^{(i,i')})^\top \in R$ with $(x_i^{(i,i')}, x_{i'}^{(i,i')})^\top \notin E_H$.

Proof. We show only the existence for i = 1 and i' = 2; the other cases can be proven similarly. For the sake of a contradiction, suppose that $\forall (x_1, \ldots, x_n) \in (V_H)^n, (x_1, \ldots, x_n) \in R \implies (x_1, x_2) \in E_H$. By Lemma 12 we have $E_H \in \langle R \rangle_{\frac{1}{p}}$, and by Theorem 5, $\operatorname{pPol}(R) \subseteq \operatorname{pPol}(E_H)$. This contradicts our hypothesis that $\operatorname{pPol}(H) \subsetneq \operatorname{pPol}(R)$.

This leads to the following corollary whose proof provides a simple construction of an R-wall for graph H in the conditions of Lemma 13.

Corollary 14. Let $n \ge 1$, H be a graph with og(H) > n, and let $R \ne \emptyset$ be an n-ary relation such that $pPol(H) \subsetneq pPol(R)$. Then there is an R-wall for H.

Proof. Using the notation of Lemma 13, we can take the $n \times \frac{n(n-1)}{2}$ matrix M, whose $\frac{n(n-1)}{2}$ columns are the $(x_1^{(i,i')}, \ldots, x_n^{(i,i')})^{\top}$, for each $(i,i') \in [n]^2$ with i < i'.

We are now ready to prove the main result of this section.

Theorem 15. Let H be a graph, and let k := og(H). There exists an n-ary relation $R \neq \emptyset$ with $pPol(H) \subsetneq pPol(R)$ such that CsP(R) is NP-complete if and only if $k \le n$. Moreover, if k > n, any n-ary relation $R \neq \emptyset$ with $pPol(H) \subsetneq pPol(R)$ is such that CsP(R) is trivial.

Proof. We sketch the most important ideas.

Suppose first that k > n. In this case, H does not have an odd-cycle of length $\leq n$. Again for the sake of a contradiction, suppose that such a relation R exists. Note that $R \neq \emptyset$ since CsP(R) is NP-complete. Using Corollary 14, there exists an R-wall for H. Then, by Lemma 11, CsP(R) is trivial. This contradicts the fact that CsP(R) is NP-complete, and thus such a relation R does not exist.

Suppose now that $k \leq n$. Define $R(x_1, \ldots, x_n) \equiv E_H(x_1, x_2) \wedge E_H(x_2, x_3) \wedge \ldots \wedge E_H(x_{k-1}, x_k) \wedge E_H(x_k, x_1)$. Since k = og(H), it follows from Lemma 9 that $R = \{(x_1, \ldots, x_n) \mid (x_1, \ldots, x_k) \text{ forms a } k\text{-cycle}\}$ (the variables x_{k+1}, \ldots, x_n are inessential).

We then proceed as follows. Since E_H qfpp-defines R, we have that $pPol(H) \subseteq pPol(R)$, by Theorem 5. Also, the inclusion is strict since, for any edge (x, y) of H, the function $f : \{x, y\} \to V_H$ that maps both x and y to any $a \in V_H$, belongs to $pPol(R) \setminus pPol(H)$. Indeed, $f \in pPol(R)$ because $\{x, y\}^n \cap R = \emptyset$, since it is impossible to form an odd-cycle with only x and y.

To prove that $\operatorname{CSP}(R)$ is NP-complete, consider $C_k(H)$, the subgraph of H, with $V_{C_k(H)} = V_H$, and where each edge of H that does not belong to a cycle of length k has been removed. Note that as H contains a k-cycle, $C_k(H)$ also contains a k-cycle, and is therefore non-bipartite. Hence, $\operatorname{CSP}(E_{C_k(H)})$, which is the same problem as the $C_k(H)$ -COLORING problem, is NP-hard (by Theorem 1).

It is easy to see that R pp-defines $E_{C_k(H)}$, as

$$E_{C_k(H)}(x_1, x_2) \equiv \exists x_3, \dots, x_n, R(x_1, x_2, x_3, \dots, x_n).$$

From Theorem 5 and 6, $\operatorname{CSP}(E_{C_k(H)}) = C_k(H)$ -COLORING has a polynomial-time reduction to $\operatorname{CSP}(R)$, and $\operatorname{CSP}(R)$ is thus NP-hard. Clearly, it is also included in NP.

4. PROJECTIVE AND CORE GRAPHS

In this section we study the inclusion structure of sets of total polymorphisms. We are particularly interested in graphs H with small sets of polymorphisms since, intuitively, they correspond to the hardest H-COLORING problems. This motivates the following definitions.

 \square

An *m*-ary function f is said to be essentially at most unary if it is of the form $f = f' \circ \pi_i^m$ for some $i \in [m]$ and some unary function f'. Larose [12] says that a graph H is projective if every idempotent polymorphism (i.e., $f(x, \ldots, x) = x$ for every $x \in V_H$) is a projection. Okrasa and Rzążewski [15] showed that the polymorphisms of a core graph H are all essentially at most unary if and only if H is projective. Since it is sufficient to study cores in the context of H-COLORING, determining whether H is projective is particularly interesting.

In this section we use the algebraic approach for proving that a given graph is a projective core, that is, both projective and a core. As we will see, this enables simpler proofs than those of [12], and suggests the possibility of completely characterizing projective cores.

Using the following theorem, our proofs of projectivity can be seen as reductions from cliques.

Theorem 16 ([3, 14]). For $k \ge 3$, K_k is projective.

Let \mathfrak{S}_k be the set of permutations over [k]. It then follows that $\operatorname{Pol}(K_k) = \{\sigma \circ \pi_i^m \mid \sigma \in \mathfrak{S}_k, m \ge 1, i \in [m]\}$. The inclusion \supseteq is indeed clear. To justify \subseteq , note that if $f \in \operatorname{Pol}(K_k)$ with $\operatorname{ar}(f) = m$, the function $\sigma : x \mapsto f(x, \ldots, x)$ is a unary polymorphism of K_k , and is therefore bijective: σ is an automorphism of K_k ie. $\sigma \in \mathfrak{S}_k$. Then, since $\sigma^{-1} \circ f$ is a polymorphism of K_k (by composition of polymorphisms of K_k) that is idempotent, it is a projection π_i^m with $i \in [m]$ by Theorem 16, and then $f = \sigma \circ \pi_i^m$.

Corollary 17 below implies that the graphs we will consider in this subsection are projective cores.

Corollary 17. Let H be a graph on [k] with $k \ge 3$. Then E_H pp-defines the relation $NEQ_k = \{(x, x') \in V_H \mid x \ne x'\}$ if and only if H is a projective core.

Proof. First observe that $NEQ_k = E_{K_k}$. From Theorem 5 and using the definitions of cores and of projective graphs, we thus have that the following assertions are equivalent:

- (1) NEQ_k $\in \langle E_H \rangle$;
- (2) $\operatorname{Pol}(H) \subseteq \operatorname{Pol}(K_k);$

(3) all polymorphisms of H are essentially at most unary, and all unary polymorphisms of H are bijective;

(4) H is a projective core.

By following the steps in the proof of Corollary 17 we can obtain the following result.

Corollary 18. Let G and H be two graphs on the same set of vertices, with G projective (respectively, a core), and such that E_H that pp-defines E_G . Then H is also projective (respectively, a core).

Pp-definitions thus explains the property of being projective (respectively, a core). We hope that this viewpoint helps to discover new classes of projective graphs. For example, Corollary 17 enables a much simpler proof of the following theorem by Larose [12].

Theorem 19 ([12],[13]). Let $k \ge 3$ be an odd integer. The k-cycle C_k is a projective core.

Proof. We claim that

$$NEQ_{k}(x, x') \equiv \exists x_{2}, \dots, x_{k-2} \colon E_{C_{k}}(x, x_{2}) \land E_{C_{k}}(x_{2}, x_{3}) \land \dots \land E_{C_{k}}(x_{k-3}, x_{k-2}) \land E_{C_{k}}(x_{k-2}, x').$$

To see this, note that for any two vertices x and x' in C_k , $x \neq x'$ if and only if there exists an odd-path from x to x' of size < k (since k is odd). In other words, $x \neq x'$ if and only if there exists a (k-2)-path from x to x' (by going through the same edge as many times as necessary, k-2 being odd). By Corollary 17, it then follows that C_k is a projective core.

There are also other examples of cores that are projective, other than k-cliques for $k \ge 3$ and k-cycles. For instance, Okrasa and Rzążewski [15] proved that the so-called *Grötzsch graph* (see Figure 1) is a projective core.

Theorem 20. The Grötzsch and Petersen graph is a projective core³

 $^{^{3}}$ We acknowledge Mario Valencia-Pabon for pointing out that the proof for the Grötzsch graph also applies to the Petersen graph.



FIGURE 1. The Grötzsch graph (left) and the Petersen graph (right)

Proof. We provide an alternative proof using our algebraic framework. Let E_G be the set of edges of the Grötzsch graph. Note that the Grötzsch graph has 11 vertices. We can see that E_G pp-defines NEQ₁₁:

$$NEQ_{11}(x, x') \equiv \exists x_2, x_3 \colon E_G(x, x_2) \land E_G(x_2, x_3) \land E_G(x_3, x').$$

From Corollary 17 it follows that the Grötzsch graph is a projective core. The proof for the Petersen graph is analogous. \square

Complements $\overline{C_k}$ of odd-cycles of length $k \ge 5$ are also projective cores. Since $\overline{C_5} = C_5$ has already been studied, we take a look at $\overline{C_{2p+1}}$, for $p \ge 3$. The following result is an immediate corollary of Larose [12], but we give an algebraic proof using Corollary 17.

Theorem 21. $\overline{C_{2p+1}}$ is a projective core for $p \ge 3$.

Proof. It is not difficult to see that $NEQ_{2p+1}(x_1, x_4) \equiv \exists x_2, x_3, w_1, \ldots, w_{p-2} : R_1 \land R_2 \land R_3$, where

- (1) $R_1 = \bigwedge_{i \in [3]} E_{\overline{C_{2p+1}}}(x_i, x_{i+1}),$ (2) $R_2 = \bigwedge_{i \in [4], j \in [p-2]} E_{\overline{C_{2p+1}}}(x_i, w_j),$ and (3) $R_3 = \bigwedge_{(j,j') \in [p-2]^2, j < j'} E_{\overline{C_{2p+1}}}(w_j, w_{j'}).$

The result then follows from Corollary 17.

Moreover, we can prove by Corollary 17 that adding universal vertices to C_5 results in a projective core.

Theorem 22. Let $p \ge 0$. The graph $C_5 + p$, obtained from C_5 by adding p universal vertices⁴ is a projective core.

Proof. We can see that E_{C_5+p} pp-defines NEQ_{p+5} though the pp-definition: $NEQ_{p+5}(x_1, x_4) \equiv \exists x_2, x_3, w_1, \dots, w_p : R_1 \land R_2 \land R_3$, where

(1) $R_1 = \bigwedge_{i \in [3]} E_{C_5+p}(x_i, x_{i+1}),$ (2) $R_2 = \bigwedge_{i \in [4], j \in [p]} E_{C_5+p}(x_i, w_j),$ and (3) $R_3 = \bigwedge_{(j,j') \in [p]^2, j < j'} E_{C_5+p}(w_j, w_{j'}).$

which proves that $C_5 + p$ is a projective core by Corollary 17.

To see that the pp-definition is correct, notice that if $x_1 = x_4$, the pp-definition can not be satisfied, since it would imply the existence of a K_{p+3} (induced by $x_1, x_2, x_3, w_1, \ldots, w_p$) in $C_5 + p$, which is absurd. Note also that if $x_1 \neq x_4$ and x_1 and x_4 are adjacent, then there exists y_1, \ldots, y_p in $C_5 + 1$ such that $\{x_1, x_4, y_1, \ldots, y_p\}$ induces a K_{p+2} : taking $x_2 := x_4, x_3 := x_1$ and $w_j := y_j$ (for all $j \in [p]$) satisfies the pp-definition. Also, if $x_1 \neq x_4$ and x_1 and x_4 are not adjacent, we can assume by symetry that $x_1 = 1$ and $x_4 = 4$ (where the vertices of the C_5 induced in $C_5 + p$ are named 0, 1, 2, 3, 4 in order). Then, taking $x_2 := 2, x_3 := 3$ and w_1, \ldots, w_p the p universal vertex of $C_5 + p$ satisfies the pp-definition. The pp-definition is thus correct. \Box

⁴Formally $C_5 + p = (V_{C_5} \uplus V_{K_p}, E_{C_5} \uplus E_{K_p} \uplus (V_{C_5} \times V_{K_p}) \uplus (V_{K_p} \times V_{C_5})).$



FIGURE 2. Every core graph with at most 6 vertices

5. Verifying the conjecture on small graphs

Okrasa and Rzążewski [15] observed that a graph H that can be expressed as a disjoint union of two non-empty graphs H_1 and H_2 is not projective, since it admits the binary polymorphism f defined by $f|_{V_{H_1} \times V_H} = (\pi_1^2)|_{V_{H_1} \times V_H}$ and $f|_{V_{H_2} \times V_H} = (\pi_2^2)|_{V_{H_2} \times V_H}$. The same holds for the *cross-product* of non-trivial graphs $H = H_1 \times H_2$ (in which case the graph is said to be decomposable), with the binary polymorphism $f((x_1, x_2), (y_1, y_2)) \mapsto (x_1, y_2)$. Okrasa and Rzążewski also noticed the existence of disconnected cores, such as $G + K_3$ (indecomposable cores are much more difficult to study), where G is the Grötzsch graph from Figure 1. These observations resulted in the following conjecture.

Okrasa and Rzążewski Conjecture ([15]). Let H be a connected non-trivial core on at least 3 vertices. Then H is projective if and only if it is indecomposable.

The goal of this section is to apply the tools constructed in Section 4 to the verification of the conjecture below by Okrasa and Rzążewski [15]. First, in Section 5.1, we completely classify the cores on at most 6 vertices and verify that the Okrasa and Rzążewski Conjecture is true on each of these graphs. Then, in Section 5.2, after giving an exhaustive list of the cores on 7 vertices, we prove the projectivity of all these graphs.

This section aims at verifying the Okrasa and Rzążewski Conjecture on graphs with at most 7 vertices, and culminates with the proof of the following theorem:

Theorem 23. The Okrasa and Rzążewski Conjecture is true on graphs with at most 7 vertices

5.1. Core graphs with at most 6 vertices. In order to verify the conjecture on small graphs, we enumerate all the (indecomposable) small cores and check their projectivity. Recall from Theorem 16 that cliques are indecomposable, projective and core, and thus the Okrasa and Rzążewski Conjecture is true on cliques. We can therefore restrict to the non-clique core graphs. This motivates the definition of *proper cores*.

Definition 24. A proper core is a core graph that is not a clique.

Moreover, a proper core on $n \ge 0$ vertices is called a *proper n-core*.

Recall that a graph G is said to be *perfect* if for all induced subgraph G' of G, the size of the largest clique of G' equals the chromatic number of G'.

Remark 25. If a graph G is a proper core then it is not a perfect graph.

Proof. Assume by contradiction that G is a perfect graph, and let k be the chromatic number of G. Since G is a perfect graph, G has an induced K_k , and by definition of k, G is k-colorable. Since K_k is a core, it follows that $core(G) = K_k$, and since G is a core, $G = core(G) = K_k$. We have that G is a clique, which contradicts the hypothesis that G is a proper core.

Using the famous theorem of perfect graphs, we can drastically reduce the search space when trying to enumerate all proper 6-cores.



FIGURE 3. The "trivial" 7-cores. We prove in Theorem 31 that they are projective.

Theorem 26 (Theorem of perfect graphs). [4]

A graph G is perfect if and only if G does not contain any induced C_k or $\overline{C_k}$ for some k odd and $k \ge 5$.

We can immediately deduce the following corollary from Remark 25 and Theorem 26.

Corollary 27. Let G be a proper core on $n \ge 1$ vertices, then G contains an induced C_k or $\overline{C_k}$ for some k odd and $5 \le k \le n$.

Via Corollary 27 we can easily classify the cores on ≤ 5 vertices (remarking that $\overline{C_5} = C_5$).

Remark 28. The cliques K_1, \ldots, K_5 are cores. The other cores on ≤ 5 vertices are proper cores, so must contain an induced C_5 . It follows from Corollary 27 that the only proper n-core with $n \leq 5$ is C_5 .

We have completely classified the cores on ≤ 5 vertices. We now extend this classification to the cores on 6 vertices.

Theorem 29. The only core graphs on 6 vertices are K_6 and the graph $C_5 + 1$ presented in Figure 2. These two graphs are projective.

Proof. First, notice that K_6 and $C_5 + 1$ are projective cores by Theorem 16 and 22.

We now prove that K_6 and $C_5 + 1$ are the only cores on 6 vertices. Assume by contradiction that there exists a proper core G on 6 vertices different from $C_5 + 1$. Then by Corollary 27, five of the six vertices of G must induce a 5-cycle, call them a, b, c, d and e, and call u the sixth vertex.

Since G is not isomorphic to $C_5 + 1$, u must not be a neighbor to (at least) one the vertex in $\{a, b, c, d, e\}$. Assume by symmetry that u and a are not neighbors. Notice that G is 3-colorable by coloring a and u with the color 1; b and d with the color 2; and c and e with the color 3. We deduce that G has no triangle, otherwise we would have $core(G) = K_3$, contradicting the fact that G is a core. It follows that u has at most 2 non-adjacent neighbors, ie. the set of neighbors of u is contained in a set of the form $\{\alpha, \beta\}$ where α and β belong to $\{a, b, c, d, e\}$ and are non-adjacent. The vertices α and β have a common neighbor $\gamma \in \{a, b, c, d, e\}$. The function that maps u to γ and that leaves the rest of the graph unchanged is a homomorphism from G to the C_5 induced by $\{a, b, c, d, e\}$. This proves that $core(G) = C_5$, contradicting that G is a core.

We have proven by contradiction that the only cores on 6 vertices are K_6 and $C_5 + 1$, and that they are projective.

The completeness of the classification of cores on at most 6 vertices presented in Figure 2 follows from Remark 28 and Theorem 29. All of these graphs are projective, by Theorems 16 and 19 and 29 and thus are not counter-example to the Okrasa and Rzążewski Conjecture. Hence, we have proven that the Okrasa and Rzążewski Conjecture is true on graphs with at most 6 vertices, and we now continue with graphs on 7 vertices.

5.2. Cores on 7 vertices. In order to put the Okrasa and Rzążewski Conjecture to the test, we continue to enumerate the small cores. We provide the exhaustive list of cores on 7 vertices in Figure 3 and 4. The proof of the fact that this is indeed the exhaustive list of cores on 7 vertices is left to Appendix A.

Theorem 30. Up to isomorphism, there are exactly 10 cores graphs on 7 vertices. They are the graphs $K_7, C_7, \overline{C_7}, C_5 + 2$ presented in Figure 3, and the graphs G_1, \ldots, G_6 presented in Figure 4.

Proof. See Appendix A.

By Theorem 30, there are 10 cores on 7 vertices. We call the four 7-cores of Figure 3 "trivial" since it is very easy to prove that they are projective.



FIGURE 4. The "sporadic" 7-cores. We prove in Theorem 35 that they are projective.

Theorem 31. The graphs $K_7, C_7, \overline{C_7}$ and $C_5 + 2$ presented in Figure 3 are projective cores.

Proof. The cases of $K_7, C_7, \overline{C_7}$ and $C_5 + 2$ have been treated, respectively, in Theorems 16, 19, 21 and 22.

What remains is now to check the projectivity of the "sporadic" 7-cores presented in Figure 4. To simplify this we make use of Rosenberg's classification of minimal clones [18], here presented in a slightly condensed form specifically for projective graphs.

Theorem 32. [18] Let G be a non-projective graph. Then Pol(G) contains a function f of one of the following type:

- (1) $f: (x, y, z) \mapsto x + y + z$, where $(V_G, +)$ is the additive group of a \mathbb{F}_2 -vector space.
- (2) f is a ternary majority operation, i.e., $\forall (x,y) \in (V_G)^2$, f(x,x,y) = f(x,y,x) = f(y,x,x) = x.
- (3) f is a semiprojection of arity $m \ge 2$, i.e. f is not a projection, and there exists $i \in [m]$ such that $\forall (x_1, \ldots, x_m) \in (V_G)^m, |\{x_1, \ldots, x_m\}| < m \implies f(x_1, \ldots, x_m) = x_i.$

We know from the algebraic formulation of the CSP dichotomy theorem (see, e.g., the survey by Barto et al. [1]) that if G is a graph where Pol(G) contains a polymorphism of type 1 or 2, then G is bipartite. We therefore derive the following corollary.

Corollary 33. Let G be a core on at least 3 vertices such that Pol(G) does not contain any semiprojections. Then, G is a projective core.

In order to apply Corollary 33 to the graphs G_1, \ldots, G_6 , we carry out a reasoning in two steps:

- We verified by computer search [16] that $Pol(G_1), \ldots, Pol(G_6)$ do not contain any semiprojection of arity 2 and 3.
- We prove that $Pol(G_1), \ldots, Pol(G_6)$ do not contain any semiprojection of arity ≥ 4 .

The exclusion of non-trivial semiprojections of arity ≥ 4 is enabled by the following lemma.

Lemma 34. Let G be a core graph on at most $n \ge 3$ vertices, and denote by $\delta > 0$ the minimal degree of a vertex in G. Let $m := \lfloor \frac{n-1}{\delta} \rfloor + 1$ (m is an integer that satisfies $1 + m\delta > n$). Then, Pol(G) does not contain any semiprojection of arity $\ge m + 1$.

Proof. First, note that since G is a core with $G \neq K_1$, G has no isolated vertex. It follows that $\delta > 0$. Assume there is a semiprojection $f \in Pol(G)$ of arity $M \ge m + 1$. Thus, f is not a projection. We can assume, up to permute the coordinates, that f is a semiprojection on the first coordinate, i.e.,

$$\forall (x_1, \dots, x_M) \in (V_G)^M, |\{x_1, \dots, x_M\}| < M \implies f(x_1, \dots, x_M) = x_1.$$

Since f is not the projection on the first coordinate, there exists $(a_1, \ldots, a_M) \in (V_G)^M$ such that $f(a_1, \ldots, a_M) = a \neq a_1$. For each vertex u of G, let $N_G(u) := \{v \in V_G \mid (u, v) \in E_G\}$ be the open neighborhood of u in G. We claim that $N_G(a_1) \setminus N_G(a) \neq \emptyset$. Indeed if we assume by contradiction that $N_G(a_1) \subseteq N_G(a)$, then the function $h: V_G \mapsto V_G$ that maps a_1 to a and that leaves the rest of the graph unchanged would be a

non-bijective (since $a_1 \neq a$) *G*-coloring of *G*, contradicting the fact that *G* is a core. We can therefore take $x_1 \in N_G(a_1) \setminus N_G(a)$. Now, remark that, for cardinality reasons, the sets $\{x_1\}, N_G(a_2), \ldots, N_G(a_M)$ can not be pairwise disjoint: because they are all contained in V_G and because

$$|\{x_1\}| + \sum_{j=2}^{M} |N_G(a_j)| \ge 1 + (M-1)\delta \ge 1 + m\delta > n = |V_G|.$$

We can therefore consider $(x_2, \ldots, x_M) \in N_G(x_2) \times \cdots \times N_G(x_M)$, such that there exists $(j_0, j_1) \in [M]^2$ with $j_0 \neq j_1$ and $x_{j_0} = x_{j_1}$.

- On the one hand since $x_{j_0} = x_{j_1}$, we have $|\{x_1, \ldots, x_M\}| < M$. Since f is a semiprojection on the first coordinate, we deduce that $f(x_1, \ldots, x_M) = x_1$.
- On the other hand, since we have for all $j \in [M]$ that $(a_j, x_j)^\top \in E_G$ (because by definition, $x_j \in N_G(a_j)$). Since $f \in \text{Pol}(G)$, we deduce that $(f(a_1, \ldots, a_M), f(x_1, \ldots, x_M))^\top \in E_G$.

We obtain that $(a, x_1) \in E_G$ (recall that we defined $a := f(a_1, \ldots, a_M)$), contradicting the definition of x_1 (that $x_1 \notin N_G(a)$). Hence, a semiprojection of arity $\ge m + 1$ cannot exist.

By Lemma 34, observing that the minimal degree in the each of the core graphs G_1, \ldots, G_6 is 3, we deduce that there is no semiprojection in the $\text{Pol}(G_i)$ with $i \in [6]$ of arity $\ge (\lfloor \frac{7-1}{3} \rfloor + 1) + 1 = 4$.

Our main result in this theorem now follows by Theorem 30, Theorem 31, and Corollary 33.

Theorem 35. All (indecomposable) cores on 7 vertices are projectives.

Hence, we have verified the Okrasa and Rzążewski Conjecture for all graphs with at most 7 vertices, and thereby proved Theorem 23.

6. Conclusion and Future Research

In this paper, we have investigated the inclusion structure of the sets of partial polymorphisms of graphs, and proved that for all pairs of graphs H,H' on the same set of vertices, $\operatorname{pPol}(H) \subseteq \operatorname{pPol}(H')$ implies that H = H' or $E_{H'} = \emptyset$. Since this inclusion structure is trivial, it is natural to generalize the question and investigate inclusions of the form $\operatorname{pPol}(H) \subseteq \operatorname{pPol}(R)$, where H is a graph, but where R is an arbitrary relation. We deemed the case when $\operatorname{CSP}(R)$ was NP-complete to be of particular interest since the problem $\operatorname{CSP}(R)$ then bounds the complexity of H-COLORING from below, in a non-trivial way. We then identified a condition depending on the length of the shortest odd cycle in H (the odd-girth of H), and proved that there exists such an n-ary relation R if and only if the odd-girth of H is $\leq n$, otherwise, $\operatorname{CSP}(R)$ must be trivial. In an attempt to better understand the algebraic invariants of graphs, we then proceeded by studying total polymorphisms of graphs, with a particular focus on projective graphs, where we used the algebraic approach to obtain simplified and uniform proofs. Importantly, we used our algebraic tools to verify the Okrasa and Rzążewski Conjecture for all graphs of at most 7 vertices.

Concerning future research perhaps the most pressing question is whether we can use our algebraic results to prove (or disprove) the Okrasa and Rzążewski Conjecture for graphs with more than 7 vertices. By Corollary 17, the Okrasa and Rzążewski Conjecture is equivalent to the following statement.

Conjecture 36. Let *H* be a connected core on $k \ge 3$ vertices. Then, *H* is indecomposable if and only if $NEQ_k \in \langle E_H \rangle$.

To advance our understanding of the fine-grained complexity of H-COLORING, it would also be interesting to settle the following question.

Question 37. Let H be a projective core. Describe pPol(H).

For instance, is it possible to relate pPol(H) with the *treewidth* of H? More generally, are there structural properties of classes of (partial) polymorphisms that translate into bounded width classes of graphs [8]? These questions constitute topics that we are currently investigating.



FIGURE 5. Up to isomorphism, any "sporadic" 7-core (not $K_7, C_7, \overline{C_7}$ or $C_5 + 2$) ($\{a, b, c, d, e, u, v\}, E$) must satisfy this motif: $\{a, b, c, d, e\}$ induces a C_5 , $(b, v) \notin E$, all dashed edges are allowed (as long as $(u, v) \notin E$, or $(e, u) \notin E$), that $(c, u) \in E$ or $(c, v) \in E$, and that $(d, u) \in E$ or $(d, v) \in E$.

References

- Libor Barto, Andrei A. Krokhin, and Ross Willard. Polymorphisms, and how to use them. In Andrei A. Krokhin and Stanislav Zivný, editors, *The Constraint Satisfaction Problem: Complexity and Approximability*, volume 7 of *Dagstuhl Follow-Ups*, pages 1–44. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2017.
- [2] A. Björklund, T. Husfeldt, and M. Koivisto. Set partitioning via inclusion-exclusion. SIAM J. Computing, 39(2):546–563, 2009.
- [3] M. Bodirsky. Graph homomorphisms and universal algebra course notes. TU Dresden, 2015.
- Maria Chudnovsky, Neil Robertson, Paul D Seymour, and Robin Thomas. Progress on perfect graphs. Mathematical Programming, 97(1):405–422, 2003.
- [5] M. Couceiro, L. Haddad, and V. Lagerkvist. Fine-grained complexity of constraint satisfaction problems through partial polymorphisms: A survey. In 2019 IEEE 49th ISMVL 2019, pages 170–175. IEEE, 2019.
- [6] F. V. Fomin, A. Golovnev, A. S. Kulikov, and I. Mihajlin. Lower bounds for the graph homomorphism problem. In Proceedings of ICALP 2015, volume 9134 of Lecture Notes in Computer Science, pages 481–493. Springer, 2015.
- [7] F. V. Fomin, P. Heggernes, and D. Kratsch. Exact algorithms for graph homomorphisms. *Theory of Computing Systems*, 41(2):381–393, 2007.
- [8] S. Guillemot and D. Marx. Finding small patterns in permutations in linear time. In Proceedings of SODA 2014, pages 82–101. SIAM, 2014.
- [9] P. Hell and J. Nešetřil. On the complexity of h-coloring. Journal of Combinatorial Theory, Series B, 48(1):92–110, 1990.
- [10] P. Jeavons. On the algebraic structure of combinatorial problems. Theoretical Computer Science, 200:185–204, 1998.
- [11] P. Jonsson, V. Lagerkvist, G. Nordh, and B. Zanuttini. Strong partial clones and the time complexity of SAT problems. J. Computer and System Sciences, 84:52 – 78, 2017.
- [12] B. Larose. Families of strongly projective graphs. Discussiones Mathematicae Graph Theory, 22(2):271–292, 2002.
- [13] B. Larose and C. Tardif. Strongly rigid graphs and projectivity. J. Multiple-Valued Logic, 7:339-361, 2001.
- [14] T. Łuczak and J. Nešetřil. Note on projective graphs. J. Graph Theory, 47(2):81-86, 2004.
- [15] K. Okrasa and P. Rzążewski. Fine-grained complexity of the graph homomorphism problem for bounded-treewidth graphs. SIAM J. Comput., 50(2):487–508, 2021.
- [16] Jakub Oprsal. PCSPTools. https://github.com/jakub-oprsal/pcsptools, 2024. MIT License.
- [17] R Pöschel. Galois connections for operations. Galois Connections and Applications, 565:231, 2013.
- [18] Ivo G Rosenberg. Minimal clones i: the five types. In Lectures in universal algebra, pages 405-427. Elsevier, 1986.
- [19] M. Wahlström. New plain-exponential time classes for graph homomorphism. Theory of Computing Systems, 49(2):273-282, 2011.

APPENDIX A. CLASSIFICATION OF 7-CORES

The goal of this appendix is to prove Theorem 30.

Theorem 30. Up to isomorphism, there are exactly 10 cores graphs on 7 vertices. They are the graphs $K_7, C_7, \overline{C_7}, C_5 + 2$ presented in Figure 3, and the graphs G_1, \ldots, G_6 presented in Figure 4.

The high-level arguments are as follows.

- (1) Enumerate all the graphs compatible with the motif described in Figure 5, and keep only, among these graphs, the cores.
- (2) Keep exactly one representative for each class of isomorphism.

(3) Prove that all the sporadic 7-cores — core graphs on 7 vertices that are not $K_7, C_7, \overline{C_7}$ and $C_5 + 2$ have to be compatible with the motif described in Figure 5.

We begin with the first step. In Figure 6, 7 and 8, we do a case analysis for all compatible graphs, depending on the 3 possible cases for (u, v) and (u, e). We eliminate the non-cores by proving that their core is either K_3, K_4 or $C_5 + 1$, and we both show their core as an induced subgraph (represented by the thick edges), as well as giving a homomorphism to their core, represented by the colors on the vertices. Reciprocally, we ensure that the remaining graphs are cores due to Lemma 38.

Lemma 38. Let G be a graph on 7 vertices such that:

- G is not 3-colorable.
- G has no induced K_4 , and
- G has no vertex of degree ≥ 5 .

Then, G is a core.

Proof. Suppose, for the sake of contradiction, that G is not a core. This implies that core(G) must be a core graph with no more than 6 vertices. According to Remark 28 and Theorem 29 in Section 5.1, it follows that core(G) is one of the graphs in the set $\{K_n, n \in [6]\} \cup \{C_5, C_5 + 1\}$, as illustrated in Figure 2.

Given that G is not 3-colorable, we can conclude that $core(G) \notin \{K_1, K_2, K_3, C_5\}$. Furthermore, the absence of an induced K_4 in G means $core(G) \notin \{K_4, K_5, K_6\}$. Additionally, the fact that G does not have any vertex with degree 5 or higher eliminates the possibility of core(G) being $C_5 + 1$.

This leads to a contradiction: hence, G must indeed be a core.

We now continue with the second step, and study the isomorphisms between the obtained core graphs. Up to isomorphism, we find 6 "sporadic" 7-cores:

- 11 edges: only G_1 ,
- 12 edges: $G_2, G_4, G_5,$
- 13 edges: G_3, G_6 .

We discuss the question of possible isomorphisms here. Recall that isomorphic graphs always have the same number of edges.

- G_4 is not isomorphic to G_2 because if we keep only vertices of degree 4, G_4 becomes a triangle and $G_2 \ a \ P_3$.
- G_5 is not isomorphic to G_2 because if we keep only vertices of degree 4, G_5 becomes a triangle and $G_2 \ a \ P_3$.
- G_5 is not isomorphic to G_4 because if we keep only vertices of degree 3, G_4 becomes a P_4 and G_5 a star $K_{1,3}$.
- G_6 is not isomorphic to G_3 because in G_6 the 2 vertices of degree 3 are adjacent, and not in G_3 .
- G'_4 (see Figure 8) is isomorphic to G_4 , by the isomorphism from G'_4 to G_4 : $(a \ u \ b \ c \ d \ e) =$
- G'_4 (see Figure 6) is isomorphic to G'_4 , by the isomorphism from G'_4 to G'_4 . (a $u \ v \ c \ u \ c)$ G'_3 (see Figure 8) is isomorphic to G_3 through the isomorphism from G'_3 to G_3 : $(a \ d \ b \ c)(u \ v) = \begin{pmatrix} a \ b \ c \ d \ e \ u \ v \\ d \ e \ a \ b \ c \ v \ u \end{pmatrix}$.

The rest of this appendix focuses on the third step, i.e., proving that all the "sporadic" 7-cores (the 7-cores

that are not $K_7, C_7, \overline{C_7}$, or $C_5 + 2$) are necessarily compatible with the motif described in Figure 5. This is sufficient to prove that the list of 7-cores given by Figures 3 and 4 is exhaustive.

Lemma 39. Let C a core on at least 3 vertices, and let $v \in V_C$. Then $deg_C(v) \ge 2$.

Proof. We give a proof by contradiction by distinguishing between the following 3 cases.

• If $\deg_C(v) = 0$, the function that maps v to any other vertex and that leaves the rest of the graph unchanged is a non-bijective homomorphism. This contradicts that C is a core.



FIGURE 6. The 9 candidates with $(e, u) \notin E$ and $(u, v) \notin E$.



FIGURE 7. The 9 candidates with $(e, u) \notin E$ and $(u, v) \in E$.

- If v has a unique neighbor u, and if u has another neighbor w, the function that maps v to w and that leaves the rest of the graph unchanged is a non-bijective homomorphism. Again, this contradicts that C is a core.
- If v has a unique neighbor u, and if u has no other neighbor, then, since C has at least 3 vertices, there exists $w \in V_C \setminus (u, v)$. Since, by what precedes, $\deg_C(w) \neq 0$, w has a neighbor x. The function that maps v to w and u to x and that leaves the rest of the graph unchanged is a non-bijective homomorphism. Hence, this contradicts that C is a core.

We now continue by establishing the following necessary properties of sporadic 7-cores. In the following statements we implicitly assume that G is a sporadic 7-core with vertices named as in Figure 5.

Lemma 40. G has an induced C_5 .

Proof. Let G be a sporadic 7-core. Assume by contradiction that G has no induced C_5 . Since G has 7 vertices, G has no induced C_7 , nor has it an induced $\overline{C_7}$ (otherwise, we would have $G = C_7$ or $G = \overline{C_7}$). Thus, G has no induced C_{2k+1} nor has it an induced $\overline{C_{2k+1}}$ for any $k \ge 2$ (notice that $\overline{C_5} = C_5$). By the theorem of perfect graphs, G is a perfect graph, i.e., there exists $k \ge 1$ such that G has an induced K_k and G is k-colorable. In particular, $core(G) = K_k$ is a clique. Since G is a core, $G = core(G) = K_7$, leading to a contradiction.

By Lemma 40, we can assume without loss of generality that $V_G = \{a, b, c, d, e, u, v\}$ and that $\{a, b, c, d, e\}$ induce the C_5 : a - b - c - d - e - a.

Then, G depends only on the neighborhoods of u and v.

Lemma 41. *u* and *v* have a common neighbor.

Proof. Assume, with the aim of reaching a contradiction, that u and v do not have a common neighbor, and assume by symmetry that $\deg_G(u) \ge \deg_G(v)$. By Lemma 39, $\deg_G(v) \ge 2$. Consider the following case analysis.

- If G has no triangle, then the neighbors of u (respectively v) are non-adjacent, and u and v do not have a common neighbor. It follows that G is isomorphic to a subgraph of the graph presented in Figure 9. Thus there exists a homomorphism from G to C_5 . Since G has an induced C_5 by Lemma 40, we deduce that $core(G) = C_5$, contradicting that G is a core.
- If G has a triangle, then G is isomorphic to a subgraph of one of the three graphs presented in Figure 10, and is therefore 3-colorable (i.e., there is a homomorphism from G to K_3). Since G has an induced K_3 , it proves that $core(G) = K_3$, contradicting that G is a core.

In both cases, we have a contradiction. It follows that u and v have a common neighbor.

Lemma 42. There exists a vertex among $\{a, b, c, d, e\}$ that is not a common neighbor of u and v.

Proof. We prove it by contradiction. Note that if all vertices among $\{a, b, c, d, e\}$ are common neighbors of u and v, then G is one of the two graphs presented in Figure 11.

The first case is impossible, because the core of G would then be $C_5 + 1$, and hence G would not be a core. The second case is also impossible, because we assumed that G is a sporadic 7-core, so G is different from $C_5 + 2$. Clearly, there is a contradiction in both cases.

We remark that by Lemma 41, u and v have a common neighbor, and we can thus assume (without loss of generality, up to isomorphism) that the vertex "a" is a common neighbor of both u and v.

Lemma 43. G has no induced K_4 .



FIGURE 8. The 9 candidates with $(e, u) \in E$ and $(u, v) \notin E$.



FIGURE 9. Maximal case (up to isomorphism) where u and v do not have a common neighbor and G has no triangle. Even this maximal case is C_5 -colorable.



FIGURE 10. Maximal case (up to isomorphism) where u and v do not have a common neighbor and G has a triangle. Even this maximal case is 3-colorable.



FIGURE 11. The two possible graphs if all vertices in $\{a, b, c, d, e\}$ are joint neighbors of u and v.



FIGURE 12. Cases where G has no triangle. G is C_5 -colorable.

Proof. Note that G is a strict subgraph of $C_5 + 2$ obtained by removing at least an edge from one of the two universal vertices. We easily deduce that G is 4-colorable. It follows that G does not contain a K_4 , since otherwise $core(G) = K_4$.

Lemma 44. G has a triangle.

Proof. With the goal of reaching a contradiction: if G has no triangle, G is isomorphic to a subgraph of the graph presented in Figure 12. Indeed, recall that u and v have a common neighbor by Lemma 41. Thus, there is a homomorphism from G to C_5 , and then $core(G) = C_5$, which is a contradicts that G is a core. \Box

Corollary 45. G is not 3-colorable.

Proof. If G was 3-colorable, since, G has a triangle by Lemma 44, core(G) would be K_3 .

Lemma 46. For every vertex i in $\{a, b, c, d, e\}$, i is a neighbor of u or a neighbor of v.

Proof. Assume by contradiction that there exists a vertex i in $\{a, b, c, d, e\}$ that is not a neighbor of u and not a neighbor of v. The 5-cycle $\{a, b, c, d, e\}$ then becomes a $P_4 = \alpha - \beta - \gamma - \delta$ when i is removed, with α and δ being the two neighbors of i. First, note that if the two neighbors α and δ of i are also neighbors of u (respectively v), then the function that maps i to u and that leaves the rest of the graph G unchanged is a non-bijective homomorphism. This contradicts the fact that G is a core. We can now assume that the two neighbors α and δ of i are not also two neighbors of u, nor are they two neighbors of v.

With this in mind we prove that G is 3-colorable. It is sufficient to establish that G - i is 3-colorable, because since *i* has degree 2, we will be able to extend this 3-coloring to G by coloring *i* with (one of) the color(s) that is not taken by a neighbor of *i*. We have the following two cases.

• If u and v are both neighbors of α or δ : assume by symmetry that it is α . Then, by what precedes, δ is not a neighbor of u nor is it a neighbor of v. Since δ has only 1 neighbor among $\{\alpha, \beta, \gamma, u, v\}$ (it is



FIGURE 13. G - i is 3-colorable

 γ), it is sufficient to 3-color $(G-i) - \delta$ to prove that G-i is 3-colorable. Since G has no induced K_4 by Lemma 43, either $(u, v) \notin E_G$, $(\beta, u) \notin E_G$, or $(\beta, v) \notin E_G$, otherwise $\{\alpha, \beta, u, v\}$. In the first case, $(G-i) - \delta$ is 3-colorable by coloring α and γ with the color 1, u and v with the color 2, and β with the color 3. In the second case, color α and γ with the color 1, v and β with the color 2, and u with the color 3. In the third case, color α and γ with the color 1, v and β with the color 2, and u with the color 3.

The graph $(G - i) - \delta$ is 3-colorable, which implies that G is 3-colorable, contradicting Corollary 45.

- Since neither u nor v are common neighbors of α and δ , and since neither α nor δ are common neighbors of u and v, we can assume by symmetry that $(\alpha, v) \notin E_G$ and that $(\delta, u) \notin E_G$.
 - If $(u, v) \notin E_G$, color u and v with the same color, and 2-color the rest of G i (which is the P_4 $\alpha - \beta - \gamma - \delta$).
 - If $(u, v) \in E_G$, u and v can not have two adjacent common neighbors, otherwise G would have a K_4 which is forbidden by Lemma 43. Thus, either β or γ is not a common neighbor of u and v. We can assume by symmetry that $(\beta, v) \notin E_G$. G i is now 3-colorable by coloring α and γ , u and δ , and β and v with the same color.

For an illustration of the two previous cases, G - i is isomorphic to a subgraph of the two graphs presented in Figure 13.

Thus, G is 3-colorable, which contradicts Corollary 45.

Remark 47. By Lemma 41, u and v have a common neighbor among $\{a, b, c, d, e\}$, and by Lemma 42, not all vertices among $\{a, b, c, d, e\}$ are common neighbors of both u and v. It is thus possible to find two adjacent vertices, say a and b, such that a is a common neighbor of u and v, and such that b is not a common neighbor of u and v. By Lemma 46, b has at least one neighbor among $\{u, v\}$. By symmetry between u and v we can assume that b is a neighbor of u but not a neighbor of v.

Lemma 48. $deg_G(u) < 5$.

Proof. Assume, with the aim of reaching a contradiction, that $\deg(u) \ge 5$. If $(u, v) \notin E_G$, then u is a neighbor of a, b, c, d and e. The function that maps v to u and that leaves the rest of the graph unchanged is a non-bijective homomorphism. Contradiction with the fact that G is a core. Hence, assume now that $(u, v) \in E_G$.

- If $\deg_G(u) = 5$, then there exists $i \in \{a, b, c, d, e\}$ such that $(u, i) \notin E_G$. The function that maps i to u and that leaves the rest of the graph unchanged is a non-bijective homomorphism. This contradicts that G is a core.
- If $\deg_G(u) > 5$, then G is a neighbor of a, b, c, d and e. Since G does not contain a K_4 by Lemma 43, the neighbors of v in $\{a, b, c, d, e\}$ are non-adjacent. We deduce that the neighbors of v in $\{a, b, c, d, e\}$ are contained in a set of the form (α, β) , where α and β are non-adjacent. α and β have a common neighbor γ in $\{a, b, c, d, e\}$. The function that maps v to γ and that leaves the rest of the graph unchanged is a non-bijective homomorphism. Again, this contradicts that G is a core.

 \Box

Lemma 49. $(e, v) \in E_G$.

Proof. Assume by contradiction that $(e, v) \notin E_G$. Then by Lemma 46, $(e, u) \in E_G$. We have by definition of b and u that $(b, v) \notin E_G$ and $(b, u) \in E_G$. By definition of a, $(a, u) \in E_G$ and $(a, v) \in E_G$. By Lemma 48, u

has at least one non-neighbor among $\{a, b, c, d, e\}$. Since u is a neighbor of a, b and e, there are only two possible cases: either $(u, c) \notin E_G$ or $(u, d) \notin E_G$.

- Assume that $(u, c) \notin E_G$. Coloring b, v and e; a and d; and u and c with the same color results in a 3-coloring of G, which contradicts Corollary 45.
- Assume that $(u, d) \notin E_G$. Coloring b, v and e; a and c; and u and d with the same color results in a 3-coloring of G, which contradicts Corollary 45.

In either case there is a contradiction. Thus, $(e, v) \in E_G$.

For each vertex let us now summarizes the remaining possible cases.

- a: 0 choices: (u, a) and (v, a) are edges of G by Remark 47.
- b: 0 choices: (u, b) is an edge, and (v, b) is not an edge by Remark 47.
- c: 3 choices since c must be either a neighbor of u or of v by Lemma 46:
 - -(u,c) and (v,c) are edges of G.
 - (u, c) is an edge, and (v, c) is not an edge.
 - -(v,c) is an edge, and (u,c) is not an edge.
- d: 3 choices since d must be either a neighbor of u or of v by Lemma 46:
 - -(u,d) and (v,d) are edges of G.
 - (u, d) is an edge, and (v, d) is not an edge.
 - (v, d) is an edge, and (u, d) is not an edge.
- Concerning u, v and e, since (e, v) in an edge by Lemma 49, there are 3 possible cases:
 - Neither (u, v) nor (e, u) is an edge.
 - -(u, v) is an edge and (e, u) is not an edge.
 - -(e, u) is an edge and (u, v) is not an edge.

The case where both (u, v) and (e, u) is an edge is impossible, because otherwise $\{a, e, u, v\}$ would induce a K_4 , contradicting Lemma 43.

We conclude that, up to isomorphism, all the sporadic 7-cores belong to the list of the $3 \times 3 \times 3 = 27$ graphs presented in Figures 6, 7 and 8.

Corollary 50. All the sporadic 7-cores are contained in the graphs allowed in Figure 5.

(Ambroise Baril) UNIVERSITÉ DE LORRAINE, CNRS, LORIA, FRANCE Email address: ambroise.baril@loria.fr

(Miguel Couceiro) Université de Lorraine, CNRS, LORIA, France, INESC-ID, Instituto Superior Técnico, Universidade de Lisboa, Lisboa, Portugal

Email address: miguel.couceiro@loria.fr

(Victor Lagerkvist) DEPARTMENT OF COMPUTER AND INFORMATION SCIENCE, LINKÖPINGS UNIVERSITET, SWEDEN *Email address*: victor.lagerkvist@liu.se