

Convergence of Recursive Least Squares Based Input/Output System Identification with Model Order Mismatch

Brian Lai and Dennis S. Bernstein

Abstract—Discrete-time input/output models, also called infinite impulse response (IIR) models or autoregressive moving average (ARMA) models, are useful for online identification as they can be efficiently updated using recursive least squares (RLS) as new data is collected. Several works have studied the convergence of the input/output model coefficients identified using RLS under the assumption that the order of the identified model is the same as that of the true system. However, the case of model order mismatch is not as well addressed. This work begins by introducing the notion of *equivalence* of input/output models of different orders. Next, this work analyzes online identification of input/output models in the case where the order of the identified model is higher than that of the true system. It is shown that, given persistently exciting data, the higher-order identified model converges to the model equivalent to the true system that minimizes the regularization term of RLS.

I. INTRODUCTION

Least squares based methods are fundamental and widely used in identification, signal processing, and control [1], [2]. One useful application is online identification of a linear model for adaptive control [3, ch. 3]. A particular model structure that lends itself to online identification is a discrete-time input/output model [4], also called an infinite-impulse-response (IIR) model [5] or autoregressive moving-average (ARMA) model [3, p. 32]. An advantage of discrete-time input/output models is they can be efficiently updated in real time as new data is collected [3, ch. 3.2]. Note that in the single-input-single-output (SISO) case, the coefficients of a discrete-time input/output model directly give the discrete-time transfer function. However, the multi-input-multi-output (MIMO) case is considerably more complex [4], [6] and converting between input/output models and other linear model structures is nontrivial [7], [8]. As such, it may be beneficial to directly study the online identification of MIMO input/output models without converting to another model structure.

This paper focuses on the online identification of input/output models using recursive least squares (RLS), which has been used in adaptive model predictive control [9] and retrospective cost adaptive control [10] with various applications [11]–[13]. While related works have discussed convergence of model coefficients when identifying input/output models using RLS [3, ch. 3.4], [14], these results assume the order of the true model is known. A natural question is whether similar guarantees can be made if the order of the identified model and true system do not match. This work

will show that if the order of the identified model is higher than order of the true model, then the regressor of RLS is not persistently exciting, and hence standard convergence guarantees of RLS [3], [15], [16] do not apply. The main contribution of this work is developing new analysis of the case where the order of the identified input/output model is higher than that of the true input/output system.

This paper is organized as follows. Section II introduces discrete-time input/output models and a useful output transition equation. Next, Section III introduces the notion of *equivalence* of input/output models as models giving the same outputs under the same inputs and initial conditions. It is shown that a necessary and sufficient condition for equivalence can be written as a linear equation of the model coefficients. This section also introduces the notion of *reducibility* as the existence of a lower-order, equivalent input/output model. Finally, section IV discusses the online identification of input/output models using RLS. It is first shown that, in the case where the order of the identified model is the same as that of the true system, persistent excitation conditions guarantee global asymptotic stability of the coefficient estimation error. Next, in the case where the order of the identified model is higher than that of the true system, conditions are given under which the identified model converges to the higher-order model equivalent to the true system that minimizes the regularization term of RLS.

II. INPUT/OUTPUT MODELING

Let $k_0 \in \mathbb{N}$ be the initial time step and let $n \geq 0$ be the model order. Consider the input/output model where, for all $k \geq k_0$, $u_k \in \mathbb{R}^m$ is the input, $y_{k_0}, \dots, y_{k_0+n-1} \in \mathbb{R}^p$ are the initial conditions and, for all $k \geq k_0$, the output $y_{k+n} \in \mathbb{R}^p$ is given by

$$y_{k+n} = -\sum_{i=1}^n F_i y_{k+n-i} + \sum_{i=0}^n G_i u_{k+n-i}, \quad (1)$$

where $F_1, \dots, F_n \in \mathbb{R}^{p \times p}$, and $G_0, \dots, G_n \in \mathbb{R}^{p \times m}$ are the input/output model coefficients. It follows that, for all $k \geq k_0$,

$$y_{k+n} = -\mathcal{F}_n \mathcal{Y}_{k,n} + \mathcal{G}_n \mathcal{U}_{k,n}, \quad (2)$$

where $\mathcal{F}_n \in \mathbb{R}^{p \times p^n}$, $\mathcal{Y}_{k,n} \in \mathbb{R}^{p^n}$, $\mathcal{G}_n \in \mathbb{R}^{p \times m(n+1)}$, and $\mathcal{U}_{k,n} \in \mathbb{R}^{m(n+1)}$ are defined as

$$\mathcal{F}_n \triangleq [F_1 \quad \dots \quad F_n], \quad \mathcal{Y}_{k,n} \triangleq \begin{bmatrix} y_{k+n-1} \\ \vdots \\ y_k \end{bmatrix}, \quad (3)$$

$$\mathcal{G}_n \triangleq \begin{bmatrix} G_0 & \cdots & G_n \end{bmatrix}, \quad \mathcal{U}_{k,n} \triangleq \begin{bmatrix} u_{k+n} \\ \vdots \\ u_k \end{bmatrix}. \quad (4)$$

Proposition 1 shows that, for all $k \geq k_0$ and $j \geq 0$, y_{k+n+j} can be written as a linear combination of the inputs $u_k, \dots, u_{k+n+j-1}$ and the n outputs $y_k, \dots, y_{k+n-1}$. We call this the output transition equation.

Proposition 1. *Let $k_0 \in \mathbb{N}$. For all $k \geq k_0$, let $u_k \in \mathbb{R}^m$, let $y_{k_0}, \dots, y_{k_0+n-1} \in \mathbb{R}^p$, and, for all $k \geq k_0$, let $y_{k+n} \in \mathbb{R}^p$ be given by (1). Then, for all $k \geq k_0$ and $j \geq 0$,*

$$y_{k+n+j} = -\mathcal{F}_{n,j} \mathcal{Y}_{k,n} + \mathcal{G}_{n,j} \mathcal{U}_{k,n+j}, \quad (5)$$

where $\mathcal{F}_{n,0} \triangleq \mathcal{F}_n$, $\mathcal{G}_{n,0} \triangleq \mathcal{G}_n$, and, for all $j \geq 1$, $\mathcal{F}_{n,j} \in \mathbb{R}^{p \times pn}$ and $\mathcal{G}_{n,j} \in \mathbb{R}^{p \times m(n+1+j)}$ are defined

$$\mathcal{F}_{n,j} \triangleq \mathcal{S}(\mathcal{F}_{n,0}, j) - \sum_{i=1}^{\min\{j,n\}} F_i \mathcal{F}_{n,j-i}, \quad (6)$$

$$\mathcal{G}_{n,j} \triangleq \begin{bmatrix} \mathcal{G}_{n,0} & 0_{p \times jm} \end{bmatrix} - \sum_{i=1}^{\min\{j,n\}} F_i \begin{bmatrix} 0_{p \times im} & \mathcal{G}_{n,j-i} \end{bmatrix}, \quad (7)$$

and where $\mathcal{S}(\mathcal{F}_{n,0}, j) \in \mathbb{R}^{p \times pn}$ is the j step shift of $\mathcal{F}_{n,0}$, defined as

$$\mathcal{S}(\mathcal{F}_{n,0}, j) \triangleq \begin{cases} \begin{bmatrix} F_{j+1} & \cdots & F_n & 0_{p \times jp} \end{bmatrix} & j \leq n-1, \\ 0_{p \times np} & j \geq n. \end{cases} \quad (8)$$

Proof. Proof follows by strong induction on $j \geq 0$. Note that for all $k \geq k_0$ and $j = 0$, (5) simplifies to (2). Next, let $k \geq k_0$, let $j \geq 1$ and suppose that, for all $\hat{j} \leq j-1$, (5) holds. Note that by (1), y_{k+n+j} can be written as

$$y_{k+n+j} = -\sum_{i=1}^n F_i y_{k+n+j-i} + \sum_{i=0}^n G_i u_{k+n+j-i}.$$

Next, for all $1 \leq i \leq \min\{j, n\}$, it follows that $0 \leq j-i \leq j-1$ and by inductive hypothesis, $y_{k+n+j-i}$ can be written as

$$\begin{aligned} y_{k+n+j-i} &= -\mathcal{F}_{n,j-i} \mathcal{Y}_{k,n} + \mathcal{G}_{n,j-i} \mathcal{U}_{k,n+j-i} \\ &= -\mathcal{F}_{n,j-i} \mathcal{Y}_{k,n} + [0_{p \times im} \quad \mathcal{G}_{n,j-i}] \mathcal{U}_{k,n+j}. \end{aligned}$$

Furthermore, note that

$$\sum_{i=0}^n G_i u_{k+n+j-i} = \begin{bmatrix} \mathcal{G}_{n,0} & 0_{p \times jm} \end{bmatrix} \mathcal{U}_{k,n+j}.$$

Hence, y_{k+n+j} can be written as

$$\begin{aligned} y_{k+n-j} &= \sum_{i=1}^{\min\{j,n\}} F_i (\mathcal{F}_{n,j-i} \mathcal{Y}_{k,n} - [0_{p \times im} \quad \mathcal{G}_{n,j-i}] \mathcal{U}_{k,n+j}) \\ &\quad - \sum_{i=\min\{j,n\}+1}^n F_i y_{k+n+j-i} + [\mathcal{G}_{n,0} \quad 0_{p \times jm}] \mathcal{U}_{k,n+j}. \end{aligned}$$

Noting that $\sum_{i=\min\{j,n\}+1}^n F_i y_{k+n+j-i} = \mathcal{S}(\mathcal{F}_{n,0}, j) \mathcal{Y}_{k,n}$ and combining terms yields (5). $\square$

III. EQUIVALENCE AND REDUCIBILITY OF INPUT/OUTPUT MODELS

Consider an input/output model of order $\hat{n} \geq n$ where, for all $k \geq k_0$, $u_k \in \mathbb{R}^m$ is the input, $\hat{y}_{k_0}, \dots, \hat{y}_{k_0+\hat{n}-1} \in \mathbb{R}^p$ are the initial conditions and, for all $k \geq k_0$, the output $\hat{y}_{k+\hat{n}} \in \mathbb{R}^p$ is given by

$$\hat{y}_{k+\hat{n}} = -\sum_{i=1}^{\hat{n}} \hat{F}_i \hat{y}_{k+\hat{n}-i} + \sum_{i=0}^{\hat{n}} \hat{G}_i u_{k+\hat{n}-i}, \quad (9)$$

where $\hat{F}_1, \dots, \hat{F}_{\hat{n}} \in \mathbb{R}^{p \times p}$, and $\hat{G}_0, \dots, \hat{G}_{\hat{n}} \in \mathbb{R}^{p \times m}$ are the input/output model coefficients. Definition 1 defines models (1) and (9) as *equivalent* if they give the same outputs under the same inputs and initial conditions.

Definition 1. *Let $k_0 \in \mathbb{N}$. Consider input/output model (1) with order n and and input/output Model (9) with order $\hat{n} \geq n$. For all $k \geq k_0$, let $u_k \in \mathbb{R}^m$ be arbitrary. For all $k_0 \leq k \leq k_0 + n - 1$, let $y_k \in \mathbb{R}^p$ be arbitrary and, for all $k \geq k_0 + n$, let $y_k \in \mathbb{R}^p$ be given by (1). Next, for all $k_0 \leq k \leq k_0 + \hat{n} - 1$, let $\hat{y}_k = y_k$ and, for all $k \geq k_0 + \hat{n}$, let $\hat{y}_k \in \mathbb{R}^p$ be given by (9). Models (1) and (9) are **equivalent** if, for all $k \geq k_0 + \hat{n}$, $\hat{y}_k = y_k$.*

Proposition 2 shows that two input/output models of the same order are equivalent if and only if they have the same model coefficients.

Proposition 2. *Consider input/output models (1) and (9) with the same order $\hat{n} = n$. Models (1) and (9) are equivalent if and only if $\hat{G}_0 = G_0$ and, for all $1 \leq i \leq n$, $\hat{F}_i = F_i$ and $\hat{G}_i = G_i$.*

Proof. Proof of necessity follow immediately. To prove sufficiency, let $k_0 \in \mathbb{N}$. First consider that there exists $0 \leq i \leq n$ such that $\hat{G}_i \neq G_i$. Then, there exists $u \in \mathbb{R}^m$ such that $\hat{G}_i u \neq G_i u$. Next, consider $y_{k_0} = \dots = y_{k_0+n-1} = 0$, for all $k \geq k_0$ such that $k \neq k_0 + n - i$, $u_k = 0$, and $u_{k_0+n-i} = u$. Then, $\hat{y}_{k_0+n} = \hat{G}_i u$, $y_{k_0+n} = G_i u$, and $\hat{y}_{k_0+n} \neq y_{k_0+n}$.

Next, consider that there exists $1 \leq i \leq n$ such that $\hat{F}_i \neq F_i$. Similarly, there exists $y \in \mathbb{R}^p$ such that $\hat{F}_i y \neq F_i y$. Hence, if, for all $k_0 \leq k \leq k_0 + n - 1$, $k \neq k_0 + n - i$, $y_k = 0$, $y_{k_0+n-i} = y$, and, for all $k \geq k_0$, $u_k = 0$, then $\hat{y}_{k_0+n} = \hat{F}_i y$, $y_{k_0+n} = F_i y$, and $\hat{y}_{k_0+n} \neq y_{k_0+n}$. Thus, (1) and (9) are not equivalent and sufficiency is proven. $\square$

We next consider the equivalence of input/output models of different orders. Note that, for all $k \geq k_0$,

$$\hat{y}_{k+\hat{n}} = -\hat{\mathcal{F}}_{\hat{n}} \hat{\mathcal{Y}}_{k,\hat{n}} + \hat{\mathcal{G}}_{\hat{n}} \mathcal{U}_{k,\hat{n}}, \quad (10)$$

where $\mathcal{U}_{k,\hat{n}} \in \mathbb{R}^{m(\hat{n}+1)}$ is defined in (4) and $\hat{\mathcal{F}}_{\hat{n}} \in \mathbb{R}^{p \times p\hat{n}}$, $\hat{\mathcal{Y}}_{k,\hat{n}} \in \mathbb{R}^{p\hat{n}}$, and $\hat{\mathcal{G}}_{\hat{n}} \in \mathbb{R}^{p \times m(\hat{n}+1)}$ are defined as

$$\hat{\mathcal{F}}_{\hat{n}} \triangleq \begin{bmatrix} \hat{F}_1 & \cdots & \hat{F}_{\hat{n}} \end{bmatrix}, \quad \hat{\mathcal{Y}}_{k,\hat{n}} \triangleq \begin{bmatrix} \hat{y}_{k+\hat{n}-1} \\ \vdots \\ \hat{y}_k \end{bmatrix}, \quad (11)$$

$$\hat{\mathcal{G}}_{\hat{n}} \triangleq \begin{bmatrix} \hat{G}_0 & \cdots & \hat{G}_{\hat{n}} \end{bmatrix}. \quad (12)$$

Theorem 1 gives necessary and sufficient conditions for the equivalence of input/output models of different orders. We begin with two useful Lemmas.

Lemma 1. Let $k_0 \in \mathbb{N}$. Consider input/output model (1) with order n and input/output Model (9) with order $\hat{n} > n$. For all $k \geq k_0$, let $u_k \in \mathbb{R}^m$, for all $k_0 \leq k \leq k_0 + n - 1$, let $y_k \in \mathbb{R}^p$, and, for all $k \geq k_0 + n$, let $y_k \in \mathbb{R}^p$ be given by (1). Let $k^* \geq k_0$ and, for all $k_0 \leq k \leq k^* + \hat{n} - 1$, let $\hat{y}_k = y_k$. Finally, let $\hat{y}_{k^*+\hat{n}} \in \mathbb{R}^p$ be given by (9). Then, $\hat{y}_{k^*+\hat{n}}$ can be expressed as

$$\hat{y}_{k^*+\hat{n}} = -\hat{\mathcal{F}}_{n,\hat{n}-n}\mathcal{Y}_{k^*,n} + \hat{\mathcal{G}}_{n,\hat{n}-n}\mathcal{U}_{k^*,\hat{n}}, \quad (13)$$

where $\hat{\mathcal{F}}_{n,\hat{n}-n} \in \mathbb{R}^{p \times pn}$ and $\hat{\mathcal{G}}_{n,\hat{n}-n} \in \mathbb{R}^{p \times m(\hat{n}+1)}$ are defined

$$\hat{\mathcal{F}}_{n,\hat{n}-n} \triangleq [\hat{F}_{\hat{n}-n+1} \quad \cdots \quad \hat{F}_{\hat{n}}] - \sum_{i=1}^{\hat{n}-n} \hat{F}_i \mathcal{F}_{n,\hat{n}-n-i}, \quad (14)$$

$$\hat{\mathcal{G}}_{n,\hat{n}-n} \triangleq \hat{\mathcal{G}}_{\hat{n}} - \sum_{i=1}^{\hat{n}-n} \hat{F}_i [0_{p \times im} \quad \mathcal{G}_{n,\hat{n}-n-i}]. \quad (15)$$

Proof. Let $k_0 \in \mathbb{N}$ and let $k^* \geq k_0$. By (9), $\hat{y}_{k^*+\hat{n}}$ can be written as

$$\hat{y}_{k^*+\hat{n}} = - \sum_{i=1}^{\hat{n}-n} \hat{F}_i y_{k^*+\hat{n}-i} - \sum_{i=\hat{n}-n+1}^{\hat{n}} \hat{F}_i y_{k^*+\hat{n}-i} + \hat{\mathcal{G}}_{\hat{n}} \mathcal{U}_{k^*,\hat{n}}. \quad (16)$$

Note that the second term can be written as

$$\sum_{i=\hat{n}-n+1}^{\hat{n}} \hat{F}_i y_{k^*+\hat{n}-i} = [\hat{F}_{\hat{n}-n+1} \quad \cdots \quad \hat{F}_{\hat{n}}] \mathcal{Y}_{k^*,n}. \quad (17)$$

Next, for all $1 \leq i \leq \hat{n} - n$, note that $\hat{n} - n - i \geq 0$ and Proposition 1 (with $k = k^*$ and $j = \hat{n} - n - i$) implies that

$$y_{k^*+\hat{n}-i} = -\mathcal{F}_{n,\hat{n}-n-i}\mathcal{Y}_{k^*,n} + \mathcal{G}_{n,\hat{n}-n-i}\mathcal{U}_{k^*,\hat{n}-i}. \quad (18)$$

Substituting (17) and (18) into (16), it follows that

$$\begin{aligned} \hat{y}_{k^*+\hat{n}} &= - \sum_{i=1}^{\hat{n}-n} \hat{F}_i (-\mathcal{F}_{n,\hat{n}-n-i}\mathcal{Y}_{k^*,n} + \mathcal{G}_{n,\hat{n}-n-i}\mathcal{U}_{k^*,\hat{n}-i}) \\ &\quad - [\hat{F}_{\hat{n}-n+1} \quad \cdots \quad \hat{F}_{\hat{n}}] \mathcal{Y}_{k^*,n} + \hat{\mathcal{G}}_{\hat{n}} \mathcal{U}_{k^*,\hat{n}}. \end{aligned} \quad (19)$$

Finally, note that, for all $1 \leq i \leq \hat{n} - n$,

$$\mathcal{G}_{n,\hat{n}-n-i}\mathcal{U}_{k^*,\hat{n}-i} = [0_{p \times im} \quad \mathcal{G}_{n,\hat{n}-n-i}] \mathcal{U}_{k^*,\hat{n}}. \quad (20)$$

Substituting (19) into (20) and simplifying yields (13). $\square$

Lemma 2. For all $\hat{n} > n$,

$$[-\hat{\mathcal{F}}_{n,\hat{n}-n} \quad \hat{\mathcal{G}}_{n,\hat{n}-n}] = [-\hat{\mathcal{F}}_{\hat{n}} \quad \hat{\mathcal{G}}_{\hat{n}}] M_{n,\hat{n}}, \quad (21)$$

where $M_{n,\hat{n}} \in \mathbb{R}^{p\hat{n}+m(\hat{n}+1) \times pn+m(\hat{n}+1)}$ is defined

$$M_{n,\hat{n}} \triangleq \begin{bmatrix} M_{n,\hat{n}}^1 & M_{n,\hat{n}}^2 \\ I_{pn} & 0_{pn \times m(\hat{n}+1)} \\ 0_{m(\hat{n}+1) \times pn} & I_{m(\hat{n}+1)} \end{bmatrix} = \begin{bmatrix} [M_{n,\hat{n}}^1 \quad M_{n,\hat{n}}^2] \\ I_{pn+m(\hat{n}+1)} \end{bmatrix}, \quad (22)$$

where $M_{n,\hat{n}}^1 \in \mathbb{R}^{p(\hat{n}-n) \times pn}$ and $M_{n,\hat{n}}^2 \in \mathbb{R}^{p(\hat{n}-n) \times m(\hat{n}+1)}$ are defined

$$M_{n,\hat{n}}^1 \triangleq \begin{bmatrix} -\mathcal{F}_{n,\hat{n}-n-1} \\ -\mathcal{F}_{n,\hat{n}-n-2} \\ \vdots \\ -\mathcal{F}_{n,1} \\ -\mathcal{F}_{n,0} \end{bmatrix}, M_{n,\hat{n}}^2 \triangleq \begin{bmatrix} [0_{p \times m} \quad \mathcal{G}_{n,\hat{n}-n-1}] \\ [0_{p \times 2m} \quad \mathcal{G}_{n,\hat{n}-n-2}] \\ \vdots \\ [0_{p \times (\hat{n}-n-1)m} \quad \mathcal{G}_{n,1}] \\ [0_{p \times (\hat{n}-n)m} \quad \mathcal{G}_{n,0}] \end{bmatrix} \quad (23)$$

Proof. It follows from (14) that

$$\hat{\mathcal{F}}_{n,\hat{n}-n} = [\hat{F}_{\hat{n}-n+1} \quad \cdots \quad \hat{F}_{\hat{n}}] - [\hat{F}_1 \quad \cdots \quad \hat{F}_{\hat{n}-n}] M_{n,\hat{n}}^1.$$

Furthermore, it follows from (15) that

$$\hat{\mathcal{G}}_{n,\hat{n}-n} = \hat{\mathcal{G}}_{\hat{n}} - [\hat{F}_1 \quad \cdots \quad \hat{F}_{\hat{n}-n}] M_{n,\hat{n}}^2.$$

Combining these two expressions can yields (21). $\square$

Theorem 1. Consider input/output model (1) with order n and input/output model (9) with order $\hat{n} > n$. Models (1) and (9) are equivalent if and only if

$$\hat{\mathcal{F}}_{n,\hat{n}-n} = \mathcal{F}_{n,\hat{n}-n}, \quad \hat{\mathcal{G}}_{n,\hat{n}-n} = \mathcal{G}_{n,\hat{n}-n}, \quad (24)$$

which holds if and only if

$$[-\hat{\mathcal{F}}_{\hat{n}} \quad \hat{\mathcal{G}}_{\hat{n}}] M_{n,\hat{n}} = [-\mathcal{F}_{n,\hat{n}-n} \quad \mathcal{G}_{n,\hat{n}-n}]. \quad (25)$$

Proof. It follows from Lemma 2 that (24) and (25) are equivalent. Hence, it suffices to show that Models (1) and (9) are equivalent if and only if (24) holds. To prove sufficiency, assume models (1) and (9) are equivalent and let $k_0 \in \mathbb{N}$. Then, it follows from Definition 1 that, for all $\mathcal{Y}_{k_0,n} \in \mathbb{R}^{pn}$ and $\mathcal{U}_{k_0,\hat{n}} \in \mathbb{R}^{m(\hat{n}+1)}$, $y_{k_0+\hat{n}} = \hat{y}_{k_0+\hat{n}}$ holds, where, for all $k_0 + n \leq k \leq k_0 + \hat{n}$, $y_k \in \mathbb{R}^p$ is given by (1), for all $k_0 \leq k \leq k_0 + \hat{n} - 1$, $\hat{y}_k = y_k$, and where $\hat{y}_{k+\hat{n}} \in \mathbb{R}^p$ is given by (9). It follows from Proposition 1 and Lemma 1 that

$$\begin{aligned} y_{k+\hat{n}} &= -\mathcal{F}_{n,\hat{n}-n}\mathcal{Y}_{k,n} + \mathcal{G}_{n,\hat{n}-n}\mathcal{U}_{k,n}, \\ \hat{y}_{k+\hat{n}} &= -\hat{\mathcal{F}}_{n,\hat{n}-n}\mathcal{Y}_{k,n} + \hat{\mathcal{G}}_{n,\hat{n}-n}\mathcal{U}_{k,n}, \end{aligned}$$

Since $\mathcal{Y}_{k,n}$ and $\mathcal{U}_{k,\hat{n}}$ are chosen arbitrarily and $y_{k+\hat{n}} = \hat{y}_{k+\hat{n}}$, (24) follows.

To prove necessity, assume that (25) holds, let $k_0 \in \mathbb{N}$. For all $k \geq k_0$, let $u_k \in \mathbb{R}^m$ be arbitrary, for all $k_0 \leq k \leq k_0 + n - 1$, let $y_k \in \mathbb{R}^p$ be arbitrary and, for all $k \geq k_0 + n$, let $y_k \in \mathbb{R}^p$ be given by (1). Furthermore, for all $k_0 \leq k \leq k_0 + \hat{n} - 1$, let $\hat{y}_k = y_k$ and, for all $k \geq k_0 + \hat{n}$, let $\hat{y}_k \in \mathbb{R}^p$ be given by (9). We now show by strong induction that, for all $k^* \geq k_0$, $\hat{y}_{k^*+\hat{n}} = y_{k^*+\hat{n}}$. First, consider the base case $k^* = k_0$. It follows from (5) of Proposition 1 (with $k = k_0$ and $j = \hat{n} - n$) that $y_{k_0+\hat{n}} = -\mathcal{F}_{n,\hat{n}-n}\mathcal{Y}_{k_0,n} + \mathcal{G}_{n,\hat{n}-n}\mathcal{U}_{k_0,\hat{n}}$. Moreover, since by assumption, for all $k_0 \leq k \leq k_0 + \hat{n} - 1$, $\hat{y}_k = y_k$, it follows from (13) of Lemma 1 (with $k^* = k_0$) that $\hat{y}_{k_0+\hat{n}} = -\hat{\mathcal{F}}_{n,\hat{n}-n}\mathcal{Y}_{k_0,n} + \hat{\mathcal{G}}_{n,\hat{n}-n}\mathcal{U}_{k_0,\hat{n}}$. Finally, (24) implies that $y_{k_0+\hat{n}} = \hat{y}_{k_0+\hat{n}}$.

Next, let $k^* \geq k_0 + 1$ and assume for inductive hypothesis that, for all $k_0 \leq k \leq k^* - 1$, $\hat{y}_{k+\hat{n}} = y_{k+\hat{n}}$. It follows from (5) of Proposition 1 (with $k = k^*$ and $j = \hat{n} - n$)

that $y_{k^*+\hat{n}} = -\mathcal{F}_{n,\hat{n}-n}\mathcal{Y}_{k^*,n} + \mathcal{G}_{n,\hat{n}-n}\mathcal{U}_{k^*,\hat{n}}$. Moreover, it follows from inductive hypothesis and (13) of Lemma 1 that $\hat{y}_{k^*+\hat{n}} = -\hat{\mathcal{F}}_{n,\hat{n}-n}\mathcal{Y}_{k^*,n} + \hat{\mathcal{G}}_{n,\hat{n}-n}\mathcal{U}_{k^*,\hat{n}}$. Finally, (24) implies that $y_{k^*+\hat{n}} = \hat{y}_{k^*+\hat{n}}$. Thus, by strong induction, for all $k^* \geq k_0$, $\hat{y}_{k^*+\hat{n}} = y_{k^*+\hat{n}}$, and models (1) and (9) are equivalent. $\square$

Note that the matrix $M_{n,\hat{n}} \in \mathbb{R}^{p\hat{n}+m(\hat{n}+1) \times pn+m(\hat{n}+1)}$ will be of importance in the later discussion on online identification.

Example 1. To better illustrate the result of Theorem 1, consider the case $m = p = 1$, $n = 1$, and $\hat{n} = 2$. It follows from Theorem 1 that models (1) and (9) are equivalent if and only if

$$\begin{aligned} & [-\hat{F}_1 \quad -\hat{F}_2 \quad \hat{G}_0 \quad \hat{G}_1 \quad \hat{G}_2] \begin{bmatrix} -F_1 & 0 & G_0 & G_1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \\ &= [F_1^2 \quad G_0 \quad G_1 - F_1 G_0 \quad -F_1 G_1]. \end{aligned} \quad (26)$$

As a sanity check, note that the trivial case $\hat{F}_1 = F_1$, $\hat{G}_0 = G_0$, $\hat{G}_1 = G_1$ and $\hat{F}_2 = \hat{G}_2 = 0$ satisfies (26). $\diamond$

Definition 2 defines an input/output model as *reducible* if there exists an equivalent input/output model of lower-order, and *irreducible* otherwise. Theorem 2 provides necessary and sufficient conditions for reducibility of input/output models.

Definition 2. Consider input/output model (9) with order $\hat{n}$. The input/output model (9) is **reducible** if there exists an input/output model (1) with order $n \leq \hat{n} - 1$ such that (1) and (9) are equivalent. Otherwise, input/output model (9) is **irreducible**.

Theorem 2. The input/output model (9) is reducible if and only if there exists $F_1 \in \mathbb{R}^{p \times p}$ such that

$$(\hat{F}_1 - F_1)F_{\hat{n}-1} = \hat{F}_{\hat{n}}, \quad (27)$$

$$(\hat{F}_1 - F_1)G_{\hat{n}-1} = \hat{G}_{\hat{n}}, \quad (28)$$

where $G_0 \triangleq \hat{G}_0$ and, for all $2 \leq i \leq \hat{n}-1$ and $1 \leq j \leq \hat{n}-1$,

$$F_i \triangleq \hat{F}_i - (\hat{F}_1 - F_1)F_{i-1}, \quad (29)$$

$$G_j \triangleq \hat{G}_j - (\hat{F}_1 - F_1)G_{j-1}. \quad (30)$$

Moreover, if there exists $F_1 \in \mathbb{R}^{p \times p}$ satisfying (27) and (28), then input/output model (1) with coefficients $F_1, \dots, F_{\hat{n}-1}$ and $G_0, \dots, G_{\hat{n}-1}$ defined by (29) and (30) is equivalent to input/output model (9).

Proof. To begin, if input/output model (1) with order $n < \hat{n}-1$ is equivalent to input/output model (9), then there exists an input/output model with order $\hat{n}-1$ that is equivalent to (9), namely a model with the coefficients $F_1, \dots, F_n$, $G_0, \dots, G_n$, and $F_i = 0_{p \times p}$ and $G_i = 0_{p \times m}$ for all $n+1 \leq i \leq \hat{n}-1$. Hence, input/output model (9), with order $\hat{n}$ is reducible if and only if there exists an equivalent input/output model (1) with order $n = \hat{n}-1$.

Next, it follows from Theorem 1 that model (1) with order $n = \hat{n}-1$ and model (9) with order $\hat{n}$ are equivalent if and only if $\hat{\mathcal{F}}_{\hat{n}-1,1} = \mathcal{F}_{\hat{n}-1,1}$ and $\hat{\mathcal{G}}_{\hat{n}-1,1} = \mathcal{G}_{\hat{n}-1,1}$. Moreover, it follows from (6), (14), (7), and (15) that

$$\begin{aligned} \hat{\mathcal{F}}_{\hat{n}-1,1} &= [\hat{F}_2 \quad \hat{F}_3 \quad \dots \quad \hat{F}_{\hat{n}}] - \hat{F}_1 \mathcal{F}_{\hat{n}-1}, \\ \mathcal{F}_{\hat{n}-1,1} &= -F_1 \mathcal{F}_{\hat{n}-1} + [\hat{F}_2 \quad \hat{F}_3 \quad \dots \quad \hat{F}_{\hat{n}-1} \quad 0_{p \times p}], \\ \hat{\mathcal{G}}_{\hat{n}-1,1} &= \hat{\mathcal{G}}_{\hat{n}} - \hat{F}_1 [0_{p \times m} \quad \mathcal{G}_{\hat{n}-1}], \\ \mathcal{G}_{\hat{n}-1,1} &= -F_1 [0_{p \times m} \quad \mathcal{G}_{\hat{n}-1}] + [\mathcal{G}_{\hat{n}-1} \quad 0_{p \times m}]. \end{aligned}$$

Rearranging terms, it follows that $\hat{\mathcal{F}}_{\hat{n}-1,1} = \mathcal{F}_{\hat{n}-1,1}$ and $\hat{\mathcal{G}}_{\hat{n}-1,1} = \mathcal{G}_{\hat{n}-1,1}$ hold if and only if

$$\begin{aligned} (\hat{F}_1 - F_1)\mathcal{F}_{\hat{n}-1} &= [\hat{F}_2 - F_2 \quad \dots \quad \hat{F}_{\hat{n}-1} - F_{\hat{n}-1} \quad \hat{F}_{\hat{n}}], \\ (\hat{F}_1 - F_1)[0_{p \times m} \quad \mathcal{G}_{\hat{n}-1}] &= [\hat{G}_0 - G_0 \quad \dots \quad \hat{G}_{\hat{n}-1} - G_{\hat{n}-1} \quad \hat{G}_{\hat{n}}], \end{aligned}$$

which can be expanded as, for all $2 \leq i \leq \hat{n}-1$ and $1 \leq j \leq \hat{n}-1$,

$$(\hat{F}_1 - F_1)F_{i-1} = \hat{F}_i - F_i, \quad (31)$$

$$(\hat{F}_1 - F_1)F_{\hat{n}-1} = \hat{F}_{\hat{n}}, \quad (32)$$

$$0_{p \times m} = \hat{G}_0 - G_0. \quad (33)$$

$$(\hat{F}_1 - F_1)G_{j-1} = \hat{G}_j - G_j, \quad (34)$$

$$(\hat{F}_1 - F_1)G_{\hat{n}-1} = \hat{G}_{\hat{n}}. \quad (35)$$

Finally, note that, for all $2 \leq i \leq \hat{n}-1$, (31) holds if and only if (29), for all $1 \leq j \leq \hat{n}-1$, (34) holds if and only if (30), and (33) holds if and only if $G_0 = \hat{G}_0$. Hence, model (1) with order $n = \hat{n}-1$ and model (9) with order $\hat{n}$ are equivalent if and only if F_1 satisfies (32) and (35). $\square$

IV. ONLINE IDENTIFICATION OF INPUT/OUTPUT MODELS USING RECURSIVE LEAST SQUARES

Next, we discuss the online identification of input/output models using recursive least squares. We consider input/output model (1) of order n to be the true system. We define $\theta_{n,\text{true}} \in \mathbb{R}^{p \times pn+m(n+1)}$ as

$$\theta_{n,\text{true}} \triangleq [F_1 \quad \dots \quad F_n \quad G_0 \quad \dots \quad G_n] = [\mathcal{F}_n \quad \mathcal{G}_n]. \quad (36)$$

It follows from (2) that, for all $k \geq 0$, $y_k \in \mathbb{R}^p$ is given by

$$y_k = \theta_{n,\text{true}} \phi_{n,k}, \quad (37)$$

where, for all $k \geq 0$ and $n \geq 0$, $\phi_{n,k} \in \mathbb{R}^{pn+m(n+1)}$ is defined as

$$\phi_{n,k} \triangleq \begin{bmatrix} -y_{k-1} \\ \vdots \\ -y_{k-n} \\ u_k \\ \vdots \\ u_{k-n} \end{bmatrix} = \begin{bmatrix} \mathcal{Y}_{k-n,n} \\ \mathcal{U}_{k-n,n} \end{bmatrix}. \quad (38)$$

The objective of online identification is to identify the coefficients an input/output model (9) of order $\hat{n}$ using measurements of the inputs u_k and outputs y_k generated

from (1). This can be accomplished by minimizing the cost function $J_{\hat{n},k}: \mathbb{R}^{p \times p\hat{n}+m(\hat{n}+1)} \rightarrow \mathbb{R}$, defined as

$$J_{\hat{n},k}(\hat{\theta}_{\hat{n}}) = \sum_{i=0}^k z_{\hat{n},i}^T(\hat{\theta}_{\hat{n}}) z_{\hat{n},i}(\hat{\theta}_{\hat{n}}) + \text{tr} \left[(\hat{\theta}_{\hat{n}} - \theta_{\hat{n},0}) P_{\hat{n},0}^{-1} (\hat{\theta}_{\hat{n}} - \theta_{\hat{n},0})^T \right], \quad (39)$$

where $\hat{\theta}_{\hat{n}} \in \mathbb{R}^{p \times p\hat{n}+m(\hat{n}+1)}$ are the coefficients to be identified, defined as

$$\hat{\theta}_{\hat{n}} \triangleq [\hat{F}_1 \quad \cdots \quad \hat{F}_{\hat{n}} \quad \hat{G}_0 \quad \cdots \quad \hat{G}_{\hat{n}}] = [\hat{\mathcal{F}}_{\hat{n}} \quad \hat{\mathcal{G}}_{\hat{n}}], \quad (40)$$

where the residual error function $z_{\hat{n},k}: \mathbb{R}^{p \times p\hat{n}+m(\hat{n}+1)} \rightarrow \mathbb{R}^p$ is defined as

$$z_{\hat{n},k}(\hat{\theta}_{\hat{n}}) \triangleq y_k - \hat{\theta}_{\hat{n}} \phi_{\hat{n},k}, \quad (41)$$

where $\phi_{\hat{n},k} \in \mathbb{R}^{p\hat{n}+m(\hat{n}+1)}$ is defined in (38), and where $\theta_{\hat{n},0} \in \mathbb{R}^{p \times p\hat{n}+m(\hat{n}+1)}$ is an initial guess of the coefficients and $\bar{P}_{\hat{n},0}^{-1} \in \mathbb{R}^{[p\hat{n}+pm(\hat{n}+1)] \times [p\hat{n}+pm(\hat{n}+1)]}$ is the positive-definite regularization matrix. The following algorithm from [9] uses recursive least squares to minimize $J_{\hat{n},k}$.

Proposition 3. *For all $k \geq -\hat{n}$, let $u_k \in \mathbb{R}^m$, $y_k \in \mathbb{R}^p$. Furthermore, let $\theta_{\hat{n},0} \in \mathbb{R}^{p \times p\hat{n}+m(\hat{n}+1)}$ and let $P_{\hat{n},0} \in \mathbb{R}^{[p\hat{n}+m(\hat{n}+1)] \times [p\hat{n}+m(\hat{n}+1)]}$ be positive definite. Then, for all $k \geq 0$, $J_{\hat{n},k}$, defined in (39), has a unique global minimizer, denoted*

$$\theta_{\hat{n},k+1} \triangleq \arg \min_{\hat{\theta}_{\hat{n}} \in \mathbb{R}^{p \times p\hat{n}+m(\hat{n}+1)}} J_k(\hat{\theta}_{\hat{n}}). \quad (42)$$

which, for all $k \geq 1$, is given by

$$\theta_{\hat{n},k} = (\mathcal{Y}_{0,k} \Phi_{\hat{n},k}^T + \theta_{\hat{n},0} P_{\hat{n},0}^{-1}) (\Phi_{\hat{n},k} \Phi_{\hat{n},k}^T + P_{\hat{n},0}^{-1})^{-1}, \quad (43)$$

and where $\Phi_{\hat{n},k} \in \mathbb{R}^{p\hat{n}+m(\hat{n}+1) \times k}$ is defined

$$\Phi_{\hat{n},k} \triangleq [\phi_{\hat{n},k-1} \quad \cdots \quad \phi_{\hat{n},0}]. \quad (44)$$

Moreover, for all $k \geq 0$, $\bar{\theta}_{\hat{n},k+1}$ is given recursively as

$$P_{\hat{n},k+1} = P_{\hat{n},k} - \frac{P_{\hat{n},k} \phi_{\hat{n},k} \phi_{\hat{n},k}^T P_{\hat{n},k}}{1 + \phi_{\hat{n},k}^T P_{\hat{n},k} \phi_{\hat{n},k}}, \quad (45)$$

$$\theta_{\hat{n},k+1} = \theta_{\hat{n},k} + (y_k - \theta_{\hat{n},k} \phi_{\hat{n},k}) \phi_{\hat{n},k}^T P_{\hat{n},k+1}. \quad (46)$$

Proof. See [9] and [17]. $\square$

In practice, the recursive formulation (46) is used to update the coefficient identification in real time as new measurements are obtained. We've included the batch formulation (43) to aid in subsequent analysis. It will also be beneficial for analysis to note that using Lemma A.1, for all $k \geq 0$, (45) can be rewritten as

$$P_{\hat{n},k+1}^{-1} = P_{\hat{n},k}^{-1} + \phi_{\hat{n},k} \phi_{\hat{n},k}^T. \quad (47)$$

A. Convergence with Correct Model Order

We begin by considering the case $\hat{n} = n$, where the correct model order is known. In this case, the identified coefficients $\theta_{n,k}$ and true model coefficients $\theta_{n,\text{true}}$ are the same dimension, and it is natural to define the estimation error $\tilde{\theta}_{n,k} \in \mathbb{R}^{p \times pn+m(n+1)}$ as

$$\tilde{\theta}_{n,k} \triangleq \theta_{n,k} - \theta_{n,\text{true}}. \quad (48)$$

It then follows from (37), (46), and (47) that, for all $k \geq 0$, the estimation error dynamics can be written as

$$\tilde{\theta}_{n,k+1} = \tilde{\theta}_{n,k} P_{n,k}^{-1} P_{n,k+1}. \quad (49)$$

It then follows from (47) that, for all $k \geq 0$,

$$\tilde{\theta}_{n,k} = \tilde{\theta}_{n,0} P_{n,0}^{-1} P_{n,k} = \tilde{\theta}_{n,0} P_{n,0}^{-1} (\Phi_{n,k-1} \Phi_{n,k-1}^T + P_{n,0}^{-1})^{-1}. \quad (50)$$

We now define the notions of weak persistent excitation and persistent excitation. Note that persistent excitation implies weak persistent excitation.

Definition 3. $(\phi_k)_{k=0}^\infty \subset \mathbb{R}^{p \times n}$ is **weakly persistently exciting** if

$$\lim_{k \rightarrow \infty} \lambda_{\min} \left[\sum_{i=0}^k \phi_i^T \phi_i \right] = \infty. \quad (51)$$

$(\phi_k)_{k=0}^\infty \subset \mathbb{R}^{p \times n}$ is **persistently exciting** if

$$C \triangleq \lim_{k \rightarrow \infty} \frac{1}{k} \sum_{i=0}^{k-1} \phi_i^T \phi_i \quad (52)$$

exists and is positive definite.

Theorem 3 shows that weak persistent excitation is necessary and sufficient conditions for the global asymptotic stability (GAS) of the error dynamics (49). For definition of GAS and further discussion on weak persistent excitation, see [15]. Moreover, Theorem 3 shows that persistent excitation implies that the convergence of $\tilde{\theta}_{n,k}$ to zero is asymptotically proportional to $1/k$.

Theorem 3. *Consider the assumptions and notation of Proposition 3. (49) is GAS if and only if $(\phi_{n,k}^T)_{k=0}^\infty$ is weakly persistently exciting. Moreover, if $(\phi_{n,k}^T)_{k=0}^\infty$ is persistently exciting, then*

$$\lim_{k \rightarrow \infty} k \tilde{\theta}_{n,k} = \tilde{\theta}_{n,0} P_{n,0}^{-1} C_n^{-1}, \quad (53)$$

where $C_n \triangleq \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=0}^{N-1} \phi_{n,i} \phi_{n,i}^T$.

Proof. Note that, for all $1 \leq i \leq p$,

$$(\tilde{\theta}_{n,k+1}^i)^T = P_{n,k+1}^{-1} (\tilde{\theta}_{n,k}^i)^T, \quad (54)$$

where $\tilde{\theta}_{n,k}^i \in \mathbb{R}^{1 \times pn+m(n+1)}$ is the i^{th} row of $\tilde{\theta}_{n,k}$. It follows from Theorem 3 of [15] that (54) is GAS if and only if $(\phi_{n,k}^T)_{k=0}^\infty$ is weakly persistently exciting. Hence, (49) is GAS if and only if $(\phi_{n,k}^T)_{k=0}^\infty$ is weakly persistently exciting.

Next, if $(\phi_{n,k}^T)_{k=0}^\infty$ is persistently exciting, it follows from (50) that

$$\lim_{k \rightarrow \infty} k\tilde{\theta}_{n,k} = \lim_{k \rightarrow \infty} \tilde{\theta}_{n,0} P_{n,0}^{-1} \left(\frac{1}{k} \Phi_{n,k-1} \Phi_{n,k-1}^T + \frac{1}{k} P_{n,0}^{-1} \right)^{-1} = \tilde{\theta}_{n,0} P_{n,0}^{-1} C_n^{-1}. \quad \square$$

Proposition 4 show that for $(\phi_{n,k}^T)_{k=0}^\infty$ to be weakly persistently exciting, it is necessary that (1) be irreducible.

Proposition 4. *If (1) is reducible, then $(\phi_{n,k}^T)_{k=0}^\infty$ is not weakly persistently exciting.*

Proof. If (1) is reducible, then there exists an equivalent input/output model of order $n_r < n$. Lemma 3 implies that

$$\phi_{n,k} = M_{n_r,n} \begin{bmatrix} \mathcal{Y}_{k-n,n_r} \\ \mathcal{U}_{k-n,n} \end{bmatrix}.$$

Hence, for all $N \geq 0$,

$$\text{rank} \left(\sum_{k=0}^N \phi_{n,k} \phi_{n,k}^T \right) \leq p n_r + m(n+1).$$

but $\sum_{k=0}^N \phi_{n,k} \phi_{n,k}^T \in \mathbb{R}^{pn+m(n+1) \times pn+m(n+1)}$. Therefore, for all $N \geq 0$, $\lambda_{\min} \left[\sum_{k=0}^N \phi_{n,k} \phi_{n,k}^T \right] = 0$. $\square$

Note that if (1) is reducible, then there exists an equivalent input/output model of a lower-order which is irreducible. In other words, (1) being reducible can be viewed as the order of the identified model being higher than the order of the true model. This case is addressed in the following subsection.

B. Convergence with Higher Model Order

Next, we address the case $\hat{n} > n$, where the identified model order is higher than the model order of the true system. To begin, note that a model of order $\hat{n}$ which is equivalent to (1) is given by the coefficients¹

$$\theta_{\hat{n},\text{true}} = \begin{bmatrix} \mathcal{F}_n & 0_{p \times p(\hat{n}-n)} & \mathcal{G}_n & 0_{p \times m(\hat{n}-n)} \end{bmatrix}. \quad (55)$$

In particular, it holds that, for all $k \geq 0$,

$$y_k = \theta_{\hat{n},\text{true}} \phi_{\hat{n},k}. \quad (56)$$

Since $\hat{n} > n$, it follows from Proposition 4 that $(\phi_{\hat{n},k}^T)_{k=0}^\infty$ is not weakly persistently exciting. However, it is possible that $(\phi_{n,\hat{n},k}^T)_{k=0}^\infty$ is weakly persistently exciting, where, for all $k \geq 0$, $\phi_{n,\hat{n},k} \in \mathbb{R}^{pn+m(\hat{n}+1)}$ is defined

$$\phi_{n,\hat{n},k} \triangleq \begin{bmatrix} -y_{k-\hat{n}+n-1} \\ \vdots \\ -y_{k-\hat{n}} \\ u_k \\ \vdots \\ u_{k-\hat{n}} \end{bmatrix} = \begin{bmatrix} \mathcal{Y}_{k-\hat{n},n} \\ \mathcal{U}_{k-\hat{n},\hat{n}} \end{bmatrix}. \quad (57)$$

By similar reasoning to Proposition 4, for $(\phi_{n,\hat{n},k}^T)_{k=0}^\infty$ to be weakly persistently exciting, it is necessary that (1) be

¹The trivial equivalent model (55) is chosen because it can be written in terms of only the true model coefficients.

irreducible. Lemma 3 shows that $\phi_{\hat{n},k} = M_{n,\hat{n}} \phi_{n,\hat{n},k}$ where $M_{n,\hat{n}}$ is defined in (22).

Lemma 3. *For all $k \geq 0$ and $\hat{n} > n$,*

$$\phi_{\hat{n},k} = M_{n,\hat{n}} \phi_{n,\hat{n},k}. \quad (58)$$

Proof. Note that $\phi_{\hat{n},k}$ can be written as

$$\phi_{\hat{n},k} = \begin{bmatrix} \mathcal{Y}_{k-\hat{n},\hat{n}} \\ \mathcal{U}_{k-\hat{n},\hat{n}} \end{bmatrix} = \begin{bmatrix} \mathcal{Y}_{k-\hat{n}+n,\hat{n}-n} \\ \mathcal{Y}_{k-\hat{n},n} \\ \mathcal{U}_{k-\hat{n},\hat{n}} \end{bmatrix}$$

For all $1 \leq i \leq \hat{n}-n$, note that $k-i = (k-\hat{n})+n+(\hat{n}-n-i)$ and it follows from Proposition 1 that

$$\begin{aligned} y_{k-i} &= -\mathcal{F}_{n,\hat{n}-n-i} \mathcal{Y}_{k-\hat{n},n} + \mathcal{G}_{n,\hat{n}-n-i} \mathcal{U}_{k-\hat{n},\hat{n}-i} \\ &= -\mathcal{F}_{n,\hat{n}-n-i} \mathcal{Y}_{k-\hat{n},n} + [0_{p \times im} \quad \mathcal{G}_{n,\hat{n}-n-i}] \mathcal{U}_{k-\hat{n},\hat{n}-i}. \end{aligned}$$

Hence,

$$\mathcal{Y}_{k-\hat{n}+n,\hat{n}-n} = \begin{bmatrix} M_{n,\hat{n}}^1 & M_{n,\hat{n}}^2 \end{bmatrix} \begin{bmatrix} \mathcal{Y}_{k-\hat{n},n} \\ \mathcal{U}_{k-\hat{n},\hat{n}} \end{bmatrix},$$

and (58) follows from (22). $\square$

Proposition 5 shows that input/output model (9) with coefficients $\hat{\theta}_{\hat{n}}$ is equivalent to (1) if and only if (59) holds. Next, Proposition 6 give an explicit formulation for the equivalent model of order $\hat{n}$ which minimizes the regularization term $\text{tr} \left[(\hat{\theta}_{\hat{n}} - \theta_{\hat{n},0}) P_{\hat{n},0}^{-1} (\hat{\theta}_{\hat{n}} - \theta_{\hat{n},0})^T \right]$ of the of cost function $J_{\hat{n},k}$, defined in (39).

Proposition 5. *Input/Output models (1) and (9) are equivalent if and only if*

$$(\hat{\theta}_{\hat{n}} - \theta_{\hat{n},\text{true}}) M_{n,\hat{n}} = 0_{p \times pn+m(\hat{n}+1)}. \quad (59)$$

Proof. Note that an input/output model of order $\hat{n}$ with coefficients $\theta_{\hat{n},\text{true}}$ is trivially equivalent to (1). Hence, by Theorem 1, $\theta_{\hat{n},\text{true}} M_{n,\hat{n}} = [-\mathcal{F}_{n,\hat{n}-n} \quad \mathcal{G}_{n,\hat{n}-n}]$. Then, it also follows from Theorem 1 that (9) is equivalent to (1) if and only of $\hat{\theta}_{\hat{n}} M_{n,\hat{n}} = [-\mathcal{F}_{n,\hat{n}-n} \quad \mathcal{G}_{n,\hat{n}-n}] = \theta_{\hat{n},\text{true}} M_{n,\hat{n}}$. $\square$

Proposition 6. *The constrained optimization problem*

$$\begin{aligned} \min_{\hat{\theta}_{\hat{n}} \in \mathbb{R}^{p \times \hat{n}(m+p)+m}} \quad & \text{tr} \left[(\hat{\theta}_{\hat{n}} - \theta_{\hat{n},0}) P_{\hat{n},0}^{-1} (\hat{\theta}_{\hat{n}} - \theta_{\hat{n},0})^T \right], \quad (60) \\ \text{such that} \quad & (\hat{\theta}_{\hat{n}} - \theta_{\hat{n},\text{true}}) M_{n,\hat{n}} = 0_{p \times pn+m(\hat{n}+1)}, \end{aligned}$$

has the unique solution

$$\theta_{\hat{n}}^* \triangleq \theta_{\hat{n},0} + (\theta_{\hat{n},\text{true}} - \theta_{\hat{n},0}) H_{\hat{n}}, \quad (61)$$

where hat matrix $H_{\hat{n}} \in \mathbb{R}^{p\hat{n}+m(\hat{n}+1) \times p\hat{n}+m(\hat{n}+1)}$ is defined

$$H_{\hat{n}} \triangleq M_{n,\hat{n}} (M_{n,\hat{n}}^T P_{n,0} M_{n,\hat{n}})^{-1} M_{n,\hat{n}}^T P_{n,0}. \quad (62)$$

Proof. Note that

$$\begin{aligned} \text{tr} \left[(\hat{\theta}_{\hat{n}} - \theta_{\hat{n},0}) P_{\hat{n},0}^{-1} (\hat{\theta}_{\hat{n}} - \theta_{\hat{n},0})^T \right] \\ = \sum_{i=1}^p (\hat{\theta}_{\hat{n},i} - \theta_{\hat{n},0,i}) P_{\hat{n},0}^{-1} (\hat{\theta}_{\hat{n},i} - \theta_{\hat{n},0,i})^T, \end{aligned}$$

where, for all $1 \leq i \leq p$, $\hat{\theta}_{\hat{n},i} \in \mathbb{R}^{1 \times \hat{n}(m+p)+m}$ and $\theta_{\hat{n},0,i} \in \mathbb{R}^{1 \times \hat{n}(m+p)+m}$ are the i^{th} row of $\hat{\theta}_{\hat{n}}$ and $\theta_{\hat{n},0}$, respectively. It then follows that (60) can be written as p separate optimization problems given by, for all $1 \leq i \leq p$,

$$\min_{\hat{\theta}_{\hat{n},i} \in \mathbb{R}^{1 \times \hat{n}(m+p)+m}} (\hat{\theta}_{\hat{n},i} - \theta_{\hat{n},0,i}) P_{\hat{n},0}^{-1} (\hat{\theta}_{\hat{n},i} - \theta_{\hat{n},0,i})^T, \quad (63)$$

such that $M_{n,\hat{n}}^T \hat{\theta}_{\hat{n},i}^T = M_{n,\hat{n}}^T \theta_{\hat{n},\text{true},i}^T$,

where, for all $1 \leq i \leq p$, $\theta_{\hat{n},\text{true},i} \in \mathbb{R}^{1 \times \hat{n}(m+p)+m}$ is the i^{th} row of $\theta_{\hat{n},\text{true}}$. It then follows from Lemma A.2 that, for all $1 \leq i \leq p$, (63) has a unique solution $\theta_{\hat{n},i}^*$, given by

$$M_{n,\hat{n}}^T \theta_{\hat{n},i}^{*T} = M_{n,\hat{n}}^T \theta_{\hat{n},\text{true},i}^T, \quad (64)$$

$$P_{\hat{n},0}^{-1} \theta_{\hat{n},i}^{*T} + M_{n,\hat{n}} \nu_i^{*T} = P_{\hat{n},0}^{-1} \theta_{\hat{n},0,i}^T. \quad (65)$$

It follows from (65) that

$$\theta_{\hat{n},i}^{*T} = \theta_{\hat{n},0,i}^T - P_{\hat{n},0} M_{n,\hat{n}} \nu_i^{*T}. \quad (66)$$

Substituting (66) into (64), it follows that

$$M_{n,\hat{n}}^T (\theta_{\hat{n},0,i}^T - P_{\hat{n},0} M_{n,\hat{n}} \nu_i^{*T}) = M_{n,\hat{n}}^T \theta_{\hat{n},\text{true},i}^T. \quad (67)$$

Hence, ν_i^{*T} is given as

$$\nu_i^{*T} = (M_{n,\hat{n}}^T P_{\hat{n},0} M_{n,\hat{n}})^{-1} M_{n,\hat{n}}^T (\theta_{\hat{n},0,i}^T - \theta_{\hat{n},\text{true},i}^T). \quad (68)$$

Substituting (68) into (66), it follows that $\theta_{\hat{n},i}^{*T} = \theta_{\hat{n},0,i}^T - P_{\hat{n},0} M_{n,\hat{n}} (M_{n,\hat{n}}^T P_{\hat{n},0} M_{n,\hat{n}})^{-1} M_{n,\hat{n}}^T (\theta_{\hat{n},0,i}^T - \theta_{\hat{n},\text{true},i}^T)$. Finally, taking the transpose and noting that $\theta_{\hat{n},i}^*$ is the i^{th} row of $\theta_{\hat{n}}^*$ yields (61). $\square$

As mentioned previously, the input/output model (9) with coefficients given by $\theta_{\hat{n}}^*$ minimizes the regularization term of $J_{\hat{n},k}$ over the set of input/output models of order $\hat{n}$ equivalent to (1). Theorem 4 show that if $(\phi_{n,\hat{n},k}^T)_{k=0}^\infty$ is weakly persistently exciting, then $\theta_{\hat{n},k}$ converges to $\theta_{\hat{n}}^*$. Moreover, if $(\phi_{n,\hat{n},k}^T)_{k=0}^\infty$ is persistently exciting, then the convergence of $\theta_{\hat{n},k} - \theta_{\hat{n}}^*$ to zero is asymptotically proportional to $1/k$.

Theorem 4. Consider the assumptions and notation of Proposition 3. If $(\phi_{n,\hat{n},k}^T)_{k=0}^\infty$ is weakly persistently exciting, then

$$\lim_{k \rightarrow \infty} \theta_{\hat{n},k} = \theta_{\hat{n}}^*, \quad (69)$$

where $\theta_{\hat{n}}^*$ is defined in (61). If, additionally, $(\phi_{n,\hat{n},k}^T)_{k=0}^\infty$ is persistently exciting, then

$$\begin{aligned} & \lim_{k \rightarrow \infty} k(\theta_{\hat{n},k} - \theta_{\hat{n}}^*) \\ &= (\theta_{\hat{n},0} - \theta_{\hat{n},\text{true}}) M_{n,\hat{n}} W_{n,\hat{n}}^{-1} C_{n,\hat{n}}^{-1} W_{n,\hat{n}}^{-1} M_{n,\hat{n}}^T P_{\hat{n},0}, \end{aligned} \quad (70)$$

where $C_{n,\hat{n}} \triangleq \lim_{k \rightarrow \infty} \frac{1}{k} \sum_{k=0}^{k-1} \phi_{n,\hat{n},i} \phi_{n,\hat{n},i}^T$ and $W_{n,\hat{n}} \triangleq M_{n,\hat{n}}^T P_{\hat{n},0} M_{n,\hat{n}}$.

Proof. For brevity, denote $P_0 \triangleq P_{\hat{n},0}$, $\theta^* \triangleq \theta_{\hat{n}}^*$, $\theta_0 \triangleq \theta_{\hat{n},0}$, $\theta \triangleq \theta_{\hat{n},\text{true}}$, $M \triangleq M_{n,\hat{n}}$, $H \triangleq H_{\hat{n}}$, $W \triangleq W_{n,\hat{n}}$, and, for all $k \geq 0$, $\theta_k \triangleq \theta_{\hat{n},k}$, $\Phi_k \triangleq \Phi_{\hat{n},k}$ and $\bar{\Phi}_k \triangleq [\phi_{n,\hat{n},k-1} \cdots \phi_{n,\hat{n},0}]$. It follows from (43), (56), and Lemma 3 that, for all $k \geq 0$,

$$\theta_{k+1} = (\theta \Phi_k \Phi_k^T + \theta_0 P_0^{-1}) (\Phi_k \Phi_k^T + P_0^{-1})^{-1}$$

$$\begin{aligned} &= \theta + (\theta_0 - \theta) P_0^{-1} (\Phi_k \Phi_k^T + P_0^{-1})^{-1} \\ &= \theta + (\theta_0 - \theta) P_0^{-1} (M \bar{\Phi}_k \bar{\Phi}_k^T M^T + P_0^{-1})^{-1}. \end{aligned}$$

Subtracting both sides by θ^* and substituting (61) yields

$$\begin{aligned} \theta_{k+1} - \theta^* &= (\theta_0 - \theta) [-I + P_0^{-1} (M \bar{\Phi}_k \bar{\Phi}_k^T M^T + P_0^{-1})^{-1} + H]. \end{aligned} \quad (71)$$

Since $(\phi_{n,\hat{n},k}^T)_{k=0}^\infty$ is weakly persistently exciting, there exists N such that, for all $k \geq N$, $\bar{\Phi}_k \bar{\Phi}_k^T$ is nonsingular. Then, it follows from Lemma A.1 that, for all $k \geq N$,

$$\begin{aligned} & (M \bar{\Phi}_k \bar{\Phi}_k^T M^T + P_0^{-1})^{-1} \\ &= P_0 - P_0 M [(\bar{\Phi}_k \bar{\Phi}_k^T)^{-1} + W]^{-1} M^T P_0, \end{aligned}$$

and hence

$$\begin{aligned} & -I + P_0^{-1} (M \bar{\Phi}_k \bar{\Phi}_k^T M^T + P_0^{-1})^{-1} \\ &= -M [(\bar{\Phi}_k \bar{\Phi}_k^T)^{-1} + W]^{-1} M^T P_0. \end{aligned} \quad (72)$$

Substituting (72) and $H = M(M^T P_0 M)^{-1} M^T P_0$ into (71) implies that, for all $k \geq N$,

$$\begin{aligned} \theta_{k+1} - \theta^* &= (\theta_0 - \theta) M [-[(\bar{\Phi}_k \bar{\Phi}_k^T)^{-1} + W]^{-1} + W^{-1}] M^T P_0. \end{aligned}$$

Again applying Lemma A.1, it follows that

$$-[(\bar{\Phi}_k \bar{\Phi}_k^T)^{-1} + W]^{-1} + W^{-1} = W^{-1} [\bar{\Phi}_k \bar{\Phi}_k^T - W^{-1}]^{-1} W^{-1}$$

which yields

$$\theta_{k+1} - \theta^* = (\theta_0 - \theta) M W^{-1} [\bar{\Phi}_k \bar{\Phi}_k^T - W^{-1}]^{-1} W^{-1} M^T P_0.$$

Since $(\phi_{n,\hat{n},k}^T)_{k=0}^\infty$ is weakly persistently exciting, it follows that $\lim_{k \rightarrow \infty} \lambda_{\min}(\bar{\Phi}_k \bar{\Phi}_k^T) = \infty$ and $\lim_{k \rightarrow \infty} \theta_{k+1} - \theta^* = 0$ and (69) follows. If additionally, $(\phi_{n,\hat{n},k}^T)_{k=0}^\infty$ is persistently exciting, then

$$\begin{aligned} & \lim_{k \rightarrow \infty} k(\theta_{k+1} - \theta^*) \\ &= (\theta_0 - \theta) M W^{-1} \left[\lim_{k \rightarrow \infty} \frac{1}{k} \bar{\Phi}_k \bar{\Phi}_k^T - \frac{1}{k} W^{-1} \right]^{-1} W^{-1} M^T P_0. \end{aligned}$$

Since $\lim_{k \rightarrow \infty} \frac{1}{k} \bar{\Phi}_k \bar{\Phi}_k^T - \frac{1}{k} W^{-1} = C_{n,\hat{n}}$, (70) follows. $\square$

Finally, since the input/output model (9) with coefficients given by $\theta_{\hat{n}}^*$ is equivalent to (1) and $\lim_{k \rightarrow \infty} \theta_{\hat{n},k} = \theta_{\hat{n}}^*$, an immediate corollary is that the residual error $z_{\hat{n},k}(\theta_{\hat{n},k})$ converges to 0, where $z_{\hat{n},k}$ is defined in (41).

Corollary 1. Consider the assumptions and notation of Proposition 3. If $(\phi_{n,\hat{n},k}^T)_{k=0}^\infty$ is weakly persistently exciting, then

$$\lim_{k \rightarrow \infty} z_{\hat{n},k}(\theta_{\hat{n},k}) = 0_{p \times 1}. \quad (73)$$

V. CONCLUSIONS

This work shows that when using RLS to identify an input/output model of a higher-order than that of the true input/output system, the identified coefficients still converge to predictable values, given weakly persistently exciting data. In particular, we obtain the very natural result that the higher-order identified model converges to the model equivalent to the true system that minimizes the regularization term of RLS.

One outstanding question in our analysis is under what conditions the persistent excitation condition of Theorem 4 is met. This itself is a deep question with sufficient conditions developed in [18] and a relaxation of those conditions presented in [19]. These conditions require that the input be persistently exciting of a sufficiently high order, see [18] for further details. One method to guarantee the input is persistently exciting of arbitrarily high order is to introduce random inputs in, for example, an ε -greedy control strategy [20]. Designing a suitable input that is persistently exciting of sufficiently high order relates to the classical dilemma of exploration and exploitation [21] and is beyond the scope of this work.

Another issue not addressed explicitly is the case where the order of the identified input/output model is lower than that of the true input/output system. One immediate result however, is that if the true system is irreducible, then the identified model cannot converge to an equivalent model, as no lower-order equivalent model exists. As a result, the residual error in the RLS cost will not converge to zero, unlike in the higher-order case (see Corollary 1).

An important application of this work is the selection of a model order for online identification. Many criterion rules for model-order selection balance the trade-off of maximizing goodness of fit of the model (minimizing residual error in this case) and the simplicity of the model (minimizing model order in this case) [22]. This work shows how model identification of different orders perform under idealized conditions, which may help inform model order selection in more realistic conditions.

REFERENCES

- [1] K. J. Astrom and B. Wittenmark, *Adaptive Control*, 2nd ed. USA: Addison-Wesley Longman Publishing Co., Inc., 1994.
- [2] L. Ljung and T. Söderström, *Theory and practice of recursive identification*. MIT press, 1983.
- [3] G. C. Goodwin and K. S. Sin, *Adaptive filtering prediction and control*. Courier Corporation, 2014.
- [4] J. C. Willems, "From time series to linear system—part i. finite dimensional linear time invariant systems," *Automatica*, vol. 22, no. 5, pp. 561–580, 1986.
- [5] E. Cuevas, J. Gálvez, S. Hinojosa, O. Avalos, D. Zaldívar, M. Pérez-Cisneros *et al.*, "A comparison of evolutionary computation techniques for iir model identification," *Journal of Applied Mathematics*, vol. 2014, 2014.
- [6] J. C. Willems and J. W. Polderman, *Introduction to mathematical systems theory: a behavioral approach*. Springer Science & Business Media, 1997, vol. 26.
- [7] L. Shieh and Y. Tsay, "Transformations of a class of multivariable control systems to block companion forms," *IEEE Transactions on Automatic Control*, vol. 27, no. 1, pp. 199–203, 1982.

- [8] N. Al-Muthairi, S. Bingulac, and M. Zribi, "Identification of discrete-time mimo systems using a class of observable canonical-form," *IEEE Proceedings-Control Theory and Applications*, vol. 149, no. 2, pp. 125–130, 2002.
- [9] T. W. Nguyen, S. A. U. Islam, D. S. Bernstein, and I. V. Kolmanovsky, "Predictive cost adaptive control: A numerical investigation of persistency, consistency, and exigency," *IEEE Control Systems Magazine*, vol. 41, no. 6, pp. 64–96, 2021.
- [10] S. A. U. Islam, T. W. Nguyen, I. V. Kolmanovsky, and D. S. Bernstein, "Data-driven retrospective cost adaptive control for flight control applications," *Journal of Guidance, Control, and Dynamics*, vol. 44, no. 10, pp. 1732–1758, 2021.
- [11] B. Y. Lai, S. A. U. Islam, S. Nivison, and D. S. Bernstein, "Data-driven retrospective cost adaptive control of a quadrotor uav," in *AIAA SCITECH 2023 Forum*, 2023, p. 0694.
- [12] N. Mohseni, D. S. Bernstein, and M. B. Quadrelli, "Adaptive force control for small celestial body sampling," in *2022 American Control Conference (ACC)*. IEEE, 2022, pp. 2515–2520.
- [13] A. Farahmandi and B. Reitz, "Predictive cost adaptive control of a planar missile with unmodeled aerodynamics," in *AIAA SCITECH 2024 Forum*, 2024, p. 2218.
- [14] F. Ding, P. X. Liu, and G. Liu, "Multiinnovation least-squares identification for system modeling," *IEEE Transactions on Systems, Man, and Cybernetics, Part B (Cybernetics)*, vol. 40, no. 3, pp. 767–778, 2009.
- [15] A. L. Bruce, A. Goel, and D. S. Bernstein, "Necessary and sufficient regressor conditions for the global asymptotic stability of recursive least squares," *Systems & Control Letters*, vol. 157, p. 105005, 2021.
- [16] B. Lai, S. A. U. Islam, and D. S. Bernstein, "Regularization-induced bias and consistency in recursive least squares," in *2021 American Control Conference (ACC)*. IEEE, 2021, pp. 3987–3992.
- [17] S. A. U. Islam and D. S. Bernstein, "Recursive least squares for real-time implementation [lecture notes]," *IEEE Control Systems Magazine*, vol. 39, no. 3, pp. 82–85, 2019.
- [18] J. C. Willems, P. Rapisarda, I. Markovsky, and B. L. De Moor, "A note on persistency of excitation," *Systems & Control Letters*, vol. 54, no. 4, pp. 325–329, 2005.
- [19] H. J. Van Waarde, C. De Persis, M. K. Camlibel, and P. Tesi, "Willems' fundamental lemma for state-space systems and its extension to multiple datasets," *IEEE Control Systems Letters*, vol. 4, no. 3, pp. 602–607, 2020.
- [20] C. J. C. H. Watkins, "Learning from delayed rewards," 1989.
- [21] R. S. Sutton and A. G. Barto, *Reinforcement learning: An introduction*. MIT press, 2018.
- [22] P. Stoica and Y. Selen, "Model-order selection: a review of information criterion rules," *IEEE Signal Processing Magazine*, vol. 21, no. 4, pp. 36–47, 2004.
- [23] S. P. Boyd and L. Vandenberghe, *Convex optimization*. Cambridge university press, 2004.

APPENDIX

Lemma A.1. *Let $A \in \mathbb{R}^{n \times n}$, $U \in \mathbb{R}^{n \times p}$, $C \in \mathbb{R}^{p \times p}$, $V \in \mathbb{R}^{p \times n}$. Assume A , C , and $A + UCV$ are nonsingular. Then, $(A + UCV)^{-1} = A^{-1} - A^{-1}U(C^{-1} + VA^{-1}U)^{-1}VA^{-1}$.*

Lemma A.2. *Consider the quadratic minimization problem with affine constraints*

$$\min_{x \in \mathbb{R}^n} \frac{1}{2} x^T P x + q^T x + r, \quad (74)$$

such that $Ax = b$,

where $P \in \mathbb{R}^{n \times n}$ is positive definite and $A \in \mathbb{R}^{p \times n}$. Then, (74) has a unique solution (x^*, ν^*) given by the system

$$Ax^* = b, \quad Px^* + A^T \nu^* = -q. \quad (75)$$

Proof. See section 10.1.1 of [23]. $\square$