

Turbulence revealed by wavelet transform: power spectrum and intermittency for the velocity field of the cosmic baryonic fluid

YUN WANG (王云)¹ AND PING HE (何平)^{1,2}

¹College of Physics, Jilin University, Changchun 130012, China.

²Center for High Energy Physics, Peking University, Beijing 100871, China.

ABSTRACT

We use continuous wavelet transform techniques to construct the global and environment-dependent wavelet statistics, such as energy spectrum and kurtosis, to study the fluctuation and intermittency of the turbulent motion in the cosmic fluid velocity field with the IllustrisTNG simulation data. We find that the peak scales of the energy spectrum and the spectral ratio define two characteristic scales, which can be regarded as the integral scale and the dissipation scale of turbulence, respectively, so that the energy spectrum can be divided into the energy-containing range, the inertial range and the dissipation range of turbulence. The wavelet kurtosis is an increasing function of the wavenumber k , first grows rapidly then slowly with k , indicating that the cosmic fluid becomes increasingly intermittent with k . In the energy-containing range, the energy spectrum increases significantly from $z = 2$ to 1, but remains almost unchanged from $z = 1$ to 0. We find that both the environment-dependent spectrum and kurtosis are similar to the global ones, and the magnitude of the spectrum is smallest in the lowest-density and largest in the highest-density environment, suggesting that the cosmic fluid is more turbulent in a high-density than in a low-density environment. In the inertial range, the exponent of the energy spectrum is steeper than not only the Kolmogorov but also the Burgers exponent, suggesting that there may be more complex mechanisms for energy transfer than Kolmogorov and Burgers turbulence.

Unified Astronomy Thesaurus concepts: [Intracluster medium \(858\)](#); [Large-scale structure of the universe \(902\)](#); [Wavelet analysis \(1918\)](#); [Astrophysical fluid dynamics \(101\)](#); [Hydrodynamical simulations \(767\)](#)

1. INTRODUCTION

The turbulent motion of the cosmic baryonic fluid in large-scale structures of the universe has attracted increasing attention in cosmological studies over the last several decades. The physical origin of turbulence in cosmic baryonic fluids is complex and diverse, mainly including the following. Accretion through structure formation or hierarchical mergers should be able to generate and sustain turbulence in the intergalactic medium (IGM) and in galaxy clusters (Subramanian et al. 2006; Bauer & Springel 2012; Iapichino et al. 2017), and the associated physical processes, such as the injection and amplification of vorticity by shock waves (e.g. Ryu et al. 2008; Porter et al. 2015; Vazza et al. 2017) or ram pressure stripping (e.g. Cassano & Brunetti 2005; Subramanian et al. 2006; Roediger & Brüggen 2007), can also generate turbulence in clusters of galaxies. In addition, outflows or feed-

backs from active galactic nucleus (AGN) jets, which inflate buoyant bubbles and eventually stir the intracluster medium (ICM), are expected to drive turbulence and heat cluster cores (Brüggen et al. 2005; Iapichino & Niemeyer 2008a; Gaspari et al. 2011; Banerjee & Sharma 2014; Angelinelli et al. 2020), and the supernova (SN)-driven galactic winds are also expected to generate turbulence in and around galaxies (e.g. Evoli & Ferrara 2011; Iapichino et al. 2013).

Observational evidence for turbulent motions in the ICM has been provided by direct detection of non-thermal broadening of the X-ray emission lines by the Hitomi satellite (Hitomi Collaboration et al. 2016), and by indirect observations such as fluctuations in the magnetic field in the diffuse cluster radio sources (Vogt & Enßlin 2003; Murgia et al. 2004; Vogt & Enßlin 2005; Enßlin & Vogt 2006; Bonafede et al. 2010; Vacca et al. 2010, 2012), fluctuations in the X-ray surface brightness or in pressure inferred from X-ray and Sunyaev-Zel'dovich effect maps (Schuecker et al. 2004; Churazov et al. 2012; Gaspari et al. 2014; Zhuravleva et al. 2014; Walker et al. 2015; Khatri & Gaspari 2016; Zhuravleva et al. 2018), and the suppression of resonant line scatter-

ing in the X-ray spectra (Churazov et al. 2004; Zhuravleva et al. 2013; Hitomi Collaboration et al. 2018; Shi & Zhang 2019). These effects can be used to detect and measure turbulence in the cosmic baryonic fluids. In addition to observations, the origin and evolution of turbulence in cosmic baryonic matter has been extensively studied using various cosmological hydrodynamical simulations (e.g. Norman & Bryan 1999; Dolag et al. 2005; Iapichino & Niemeyer 2008b; Lau et al. 2009; Zhu et al. 2010; Vazza et al. 2011; Gaspari et al. 2014; Miniati 2014; Brüggén & Vazza 2015; Angelinelli et al. 2020).

There are a large number of problems relevant to the study of the formation and evolution of galaxies and galaxy clusters. Among these, the study of the heating mechanisms in the ICM and IGM is particularly important. If there is no sufficient heating mechanism, then there will be a phenomenon called cooling flow within the galaxy clusters, and there will also be an excess of star birth in the galaxy, which is known as the overcooling problem (Voit 2005). In addition, a heating mechanism may also be responsible for the missing baryon problem (Bregman 2007).

The main proposals to overcome this overcooling problem are the heating mechanisms such as SN or AGN feedback, and turbulence in the IGM or ICM. In the popular semi-analytical models of galaxy formation, the heating mechanism is mainly based on SN and AGN feedback, and turbulent heating is not considered (Guo et al. 2011; Henriques et al. 2015). However, some studies suggest that turbulent heating has a significant effect and should not be ignored. For example, Zhu et al. (2010) suggests that the turbulent pressure can be compared to the thermodynamic pressure of the baryonic gas. Furthermore, Zhuravleva et al. (2014) suggests that turbulent heating is sufficient to offset radial cooling, and indeed appears to be locally balanced at each radius, which may be a key element in solving the gas cooling problem in cluster cores.

In a series of papers, we have carried out a number of studies on the turbulence of the cosmic baryon fluid, briefly summarised as follows. Kim et al. (2005) explores the dynamics and evolution of the baryonic gas in the universe, focusing on its velocity field and its interaction with the dark matter gravity, and discusses the implications of a Burgers fluid model for describing the dynamics of the IGM, highlighting the importance of considering heating and cooling processes in understanding the evolution of the IGM. Using the WIGEON hydrodynamical simulation (Feng et al. 2004), He et al. (2006) shows that the intermittency of the velocity field of the cosmic baryonic fluid at redshift $z = 0$ in the scale range from the Jeans length to about $16 h^{-1} \text{Mpc}$ can be described extremely well by She-Leveque's universal scaling formula, and these results imply that the motion of the highly evolved cosmic baryonic fluid is similar to a fully de-

veloped turbulence. Yang et al. (2020) use updated WIGEON data (Zhu et al. 2013) and study the turbulence-induced deviation of the spatial distributions between baryons and dark matter. They find that at $z = 0$, at the 1% deviation level, the deviation scale is about $3.7 h^{-1} \text{Mpc}$ for the density field, while it is as large as $23 h^{-1} \text{Mpc}$ for the velocity field, a scale that falls into the weakly nonlinear regime for the structure formation paradigm. Their results also suggest that the effect of turbulence heating is comparable to that of these processes such as SN and AGN feedback. Yang et al. (2022) compare the results derived from IllustrisTNG and WIGEON simulations, and find that for the ratio of the density power spectrum between dark matter and baryonic matter, as scales become smaller and smaller, the power spectra for baryons are increasingly suppressed for WIGEON simulations, while for TNG simulations, the suppression stops at $k = 15 - 20 h \text{Mpc}^{-1}$, and the power spectrum ratios increase when $k > 20 h \text{Mpc}^{-1}$. These results indicate that turbulent effects can also have the consequence to suppress the power ratio between baryons and dark matter. These are of great importance for understanding the distribution and evolution of baryonic matter in the universe, as well as issues related to galaxy formation, and therefore turbulence in the cosmic baryonic fluid is worthy of in-depth study.

The wavelet analysis technique is a powerful tool, and the application of the wavelet transform technique to turbulence can be traced back more than three decades. In a review article, Farge (1992) extensively discusses wavelet transforms as a powerful tool for achieving scale localization, allowing for the analysis of different scales of motion in turbulent flows. The article delves into the computation of local energy spectra, wavelet coefficients, and statistical properties, showing how wavelet analysis facilitates the investigation of coherent structures, intermittency, and other key aspects of turbulence. The paper details the application of wavelet transforms to turbulence analysis, emphasizing their ability to provide a localized and multi-scale view of the flow, leading to insights into coherent structures, intermittency, and statistical properties. Farge (1992) emphasizes the advantages of wavelet analysis over traditional methods and provides concrete examples of its application to turbulent phenomena, providing a valuable resource for fluid mechanics researchers seeking a multi-scale perspective on turbulent flows.

The wavelet analysis method can also find its application in the study of cosmological turbulence. To name just a few that are relevant to our current interests. Schuecker et al. (2004) construct two-dimensional pressure maps from XMM-Newton observations of the Coma galaxy cluster using wavelet methods, which effectively suppress noise and reveal small-scale turbulence structures, helping to isolate and analyze pressure fluctuations, enabling the computation of their power spectra and providing evidence for turbulence

in the ICM. Kowal & Lazarian (2010) use wavelet analysis to decompose the velocity field of compressible magnetohydrodynamic turbulence into Alfvén, slow, and fast modes, and investigate in detail the turbulence properties such as spectra, anisotropy, scaling exponents and intermittency, showing that these properties are influenced by both the sonic and Alfvén Mach numbers. Shi et al. (2018) use a novel wavelet analysis method to study the radial dependence of the ICM turbulence spectrum, and find that faster turbulence dissipation in the inner high-density regions causes the turbulence amplitude to increase with radius. They also find that the ICM turbulence at all radii decays in two phases after a major merger, i.e. an early fast-decay phase and a slow secular-decay phase.

In a series of works, we have used the continuous wavelet transform (CWT) to construct the wavelet statistics such as the wavelet power spectrum (WPS), wavelet cross-correlation and wavelet bicoherence, as well as the environment-dependent WPS (env-WPS), and analyzed the clustering and non-Gaussian properties of large-scale structures in the universe using the cosmological simulation data (Wang & He 2021; Wang et al. 2022; Wang & He 2022). In a recent paper (Wang & He 2024), we mainly use the env-WPS to study how baryonic effects vary with scale and local density environment. In addition, we also complete a comparative study of several fast algorithms for one-dimensional (1D) CWT (Wang & He 2023).

In this work, we apply the CWT techniques to the IllustrisTNG simulation data to study the turbulent motion of the cosmic baryonic fluid in the large-scale structures of the universe. The paper is organized as follows. In Section 2, we briefly introduce our methods and the simulation data used in this work. In Section 3, we present the results of our work. In Section 4, we present the summary and conclusions of the paper.

2. METHODS AND DATA

2.1. Continuous Wavelet Transform and Power Spectrum

For a random field with the zero mean value, say the 3-dimensional (3D) velocity field of the cosmic baryonic fluid \mathbf{u} , its isotropic CWT $\tilde{\mathbf{u}}(w, \mathbf{x})$ is obtained by convolution with the wavelet function Ψ as

$$\tilde{\mathbf{u}}(w, \mathbf{x}) = \int \mathbf{u}(\boldsymbol{\tau}) \Psi(w, \mathbf{x} - \boldsymbol{\tau}) d^3\boldsymbol{\tau}. \quad (1)$$

Throughout this work, we use the so-called 3D isotropic cosine-weighted Gaussian-derived wavelet (CW-GDW), which can achieve good localization in both spatial and frequency space simultaneously (Wang & He 2022; Wang & He

2024). It is defined in real space as

$$\begin{aligned} \Psi(w, \mathbf{x}) &= w^{\frac{3}{2}} \Psi(wr) \\ &= C_N w^{\frac{3}{2}} \left[(4 - w^2 r^2) \cos wr \right. \\ &\quad \left. + 2 \left(\frac{1}{wr} - wr \right) \sin wr \right] e^{-\frac{1}{2} w^2 r^2}, \end{aligned} \quad (2)$$

with its Fourier transform

$$\begin{aligned} \hat{\Psi}(w, \mathbf{k}) &= w^{-\frac{3}{2}} \hat{\Psi}(k/w) \\ &= (2\pi)^{\frac{3}{2}} C_N w^{-\frac{3}{2}} \left(\frac{k}{w} \right) \left[\left(\frac{k}{w} \right) \cosh \frac{k}{w} \right. \\ &\quad \left. - \sinh \frac{k}{w} \right] e^{-\frac{1}{2} (1+k^2/w^2)}, \end{aligned} \quad (3)$$

where $r = |\mathbf{x}|$, $k = |\mathbf{k}|$, and $C_N = \frac{2}{\pi^{3/4}} \sqrt{2e/(9+55e)}$ is the normalization constant such that $\int |\Psi(\mathbf{x})|^2 d^3\mathbf{x} = 1$.

In many cases, we compare the results of the CWT with those of the Fourier transform, such as comparing the global WPS with the Fourier power spectrum (FPS). However, the wavelet scale w and the Fourier wavenumber k are not a simple one-to-one correspondence. Following the suggestions by Meyers et al. (1993) and Torrence & Compo (1998), this correspondence $w = c_w k$ can be determined by calculating the maximum of the modulus square for the CWT of the cosine wave $\cos(kx)$. For the 3D isotropic CW-GDW, $c_w \simeq 0.3883$ (Wang & He 2024).

With the CWT $\tilde{\mathbf{u}}(w, \mathbf{x})$, we can define the env-WPS of the velocity field as

$$\tilde{P}_{\mathbf{u}}(w, \delta) \equiv \left\langle |\tilde{\mathbf{u}}(w, \mathbf{x})|^2 \right\rangle_{\delta(\mathbf{x})=\delta}, \quad (4)$$

where the environment is specified with the density contrast δ and the average $\langle \dots \rangle$ is performed over all the spatial points where the condition $\delta(\mathbf{x}) = \delta$ is satisfied. The global WPS is obtained by averaging over all spatial positions,

$$\tilde{P}_{\mathbf{u}}(w) \equiv \left\langle |\tilde{\mathbf{u}}(w, \mathbf{x})|^2 \right\rangle_{\text{all } \mathbf{x}}. \quad (5)$$

It can be shown that the global WPS $\tilde{P}_{\mathbf{u}}(w)$ is related to the FPS $P_{\mathbf{u}}(k)$ as (Wang & He 2021)

$$\tilde{P}_{\mathbf{u}}(w) = \frac{1}{2\pi^2} \int_0^{+\infty} P_{\mathbf{u}}(k) \hat{\Psi}(w, k)^2 k^2 dk. \quad (6)$$

One can see that the global WPS is the wavelet-weighted average of the FPS in k space. The former therefore has a similar but smoother shape to the latter. Obviously, the wider the bandwidth of the wavelet function, the smoother the global WPS. If the bandwidth is narrower, the global WPS is closer to the FPS.

Integrating both sides of Equation (6) with respect to w , we can obtain an approximate relationship with the correspondence $w = c_w k$ as

$$\tilde{P}_{\mathbf{u}}(k) \approx \frac{I_{\Psi}}{2\pi^2 c_w} P_{\mathbf{u}}(k), \quad (7)$$




Figure 1. The two-dimensional slice of the IllustrisTNG100-1 dark matter density field at $z = 0$ (top left), and spatial regions corresponding to the density ranges labelled in these panels. The slice covers a $75 \times 75 h^{-2} \text{Mpc}^2$ area with a thickness of $0.3 h^{-1} \text{Mpc}$. The grey background indicates the region outside the specified density range. Fourier analysis can only be applied to the regular region, such as a square or a cube (top left panel), whereas wavelet analysis can be applied to regions of any shape (all four panels).

where $I_\Psi = \int_0^{+\infty} k |\hat{\Psi}(k)|^2 dk$ is a constant depending on the wavelet function. Therefore, the global WPS and the FPS differ by only a scaling factor that depends on the specific wavelet function selected. Obviously, for the power-law power spectrum $P_u(k) \propto k^n$, Equation (7) is strictly valid and is no longer an approximation. For the 3D isotropic CWGDW used in this work, $I_\Psi / (2\pi^2 c_w) \approx 1.0754$, at which point the shape and size of the global WPS and FPS are very close. However, due to the simultaneous spatial and frequency localization of the CWT, we can specify any shape of the region for integration, such as a density-bounded region. Therefore, we can define the env-WPS. The techniques of wavelet analysis are therefore superior to those of Fourier analysis, as can be seen in Figure 1.

For details of the CWT techniques that we have developed, please refer to Wang & He (2021), Wang et al. (2022), Wang & He (2022, 2023) and (Wang & He 2024).

2.2. Turbulence in the Cosmic Baryonic Fluid

As mentioned in Section 1, the cosmic baryonic fluid is characterized by turbulent motion. In this work, we investigate in which regions, to what extent, and on which scales the cosmic baryonic fluid is turbulent.

2.2.1. Filtering Out the Bulk Flow

Although it is difficult to define turbulence (Davidson 2015), we can still provide a description of turbulent flow in fluids and distinguish between turbulent and bulk flow.




Figure 2. Both Fourier and global wavelet energy spectra at $z = 0$ for the velocity field \mathbf{u} , its ‘ \parallel ’ and ‘ \perp ’ component of the cosmic fluid (left column), with the corresponding spectral ratios (right column). From top to bottom are the TNG300, TNG100, and TNG50 data, respectively. To aid the eye, the relevant power-laws are indicated in the figure. The grey area shows the $k > 0.4k_{\text{Nyquist}}$ region, where the numerical effects, such as smearing, aliasing and shot noise, are significant (Wang & He 2024).

In general, a velocity field of the fluid \mathbf{u} is a superposition of the turbulent velocity and the bulk velocity, i.e. $\mathbf{u} = \mathbf{u}_{\text{turb}} + \mathbf{u}_{\text{bulk}}$. Bulk flow is a flow state in which the fluid particles move in a relatively orderly manner, and the flow velocity and direction are relatively uniform in a given cross-section. In contrast, turbulent flow refers to a chaotic, disordered flow state in which the fluid particles move irregularly and randomly, and usually occurs when the fluid velocity is high, the pressure gradient is large, or the Reynolds number exceeds a certain critical value. Turbulent flow has the characteristics of rapid mixing, strong diffusion, and good heat transfer. Therefore, it is necessary to develop an approach to separate the turbulent flow from the bulk motion (e.g. Dolag et al. 2005; Vazza et al. 2012; ZuHone et al. 2013; Shi et al. 2018; Angelinelli et al. 2020; Vallés-Pérez et al. 2021).

In this work, we use the iterative multi-scale filtering approach developed by Vazza et al. (2012) to extract turbulent motions from the velocity field of the cosmic fluid.

2.2.2. Intermittency of Turbulence

In general, there are two types of turbulence in fluids, Kolmogorov turbulence and Burgers turbulence. Kolmogorov turbulence is essentially excited by subsonic motion of the flow, and characterized by eddies of different scales (Bauer & Springel 2012). In the fully developed state, the fluid satisfies $\nabla \cdot \mathbf{u} = 0$, and its kinetic energy spectrum follows the Kolmogorov's $k^{-5/3}$ scaling-law. Burgers turbulence is essentially excited by supersonic motion of the flow (Boldyrev et al. 2004; Federrath et al. 2010; Bauer & Springel 2012) and characterized by shock waves (Konstandin et al. 2015), which gives k^{-2} for the kinetic energy spectrum at steady state when Re , the Reynolds number, satisfies $\text{Re} \gg kL_c \gg 1$, where L_c is some correlation length (Balkovsky et al. 1997). The turbulence in the cosmic baryonic fluid is composed of both Kolmogorov turbulence and Burgers turbulence, but it is not simply a mixture of the two types of turbulence, as will be discussed later.

In turbulence, the presence of shocks leads to a strong intermittency, and therefore the turbulent energy dissipation in space is also intermittent. Intermittency in turbulence refers to the phenomenon where fluctuations in velocity, pressure or other flow properties occur in a sporadic or irregular manner. Intermittency arises because these fluctuations are not uniformly distributed in space and time. Usually, the deviation of the tails of the probability density function (PDF) from Gaussian is seen as a manifestation of intermittency, with a highly unusual scaling for the moments of the velocity differences (Balkovsky et al. 1997). Therefore, to characterize the intermittency of turbulence in the cosmic fluid, as in Meneveau (1991); Angulo & Madrid (2023), we define the wavelet

global kurtosis (or flatness) of the cosmic velocity field as

$$\tilde{K}_{\mathbf{u}}(k) \equiv \frac{\langle \Delta \tilde{u}(k, \mathbf{x})^4 \rangle_{\text{all } \mathbf{x}}}{\langle \Delta \tilde{u}(k, \mathbf{x})^2 \rangle_{\text{all } \mathbf{x}}^2}, \quad (8)$$

and the wavelet environment-dependent (env-) kurtosis as

$$\tilde{K}_{\mathbf{u}}(k, \delta) \equiv \frac{\langle \Delta \tilde{u}(k, \mathbf{x})^4 \rangle_{\delta(\mathbf{x})=\delta}}{\langle \Delta \tilde{u}(k, \mathbf{x})^2 \rangle_{\delta(\mathbf{x})=\delta}^2}, \quad (9)$$

in which $\tilde{u}(k, \mathbf{x})$ is the CWT of the velocity component along a given axis, and $\Delta \tilde{u}(k, \mathbf{x}) = \tilde{u}(k, \mathbf{x}) - \langle \tilde{u}(k, \mathbf{x}) \rangle$. For a scalar Gaussian field, the kurtosis = 3, whereas for the turbulent velocity field of fluids, its PDF is generally a long-tailed distribution, with kurtosis > 3 .

Kurtosis can be thought of as a measure of the intermittency of the turbulence. The greater the kurtosis, the more intermittent the cosmic flow should be.

Note that Farge (1992) use the local wavelet energy spectrum of the velocity field to define the local intermittency of turbulent flow, see their Equation-(59). We do not use this definition as we have already used env-WPS to perform the spectral analyses throughout this work.

2.2.3. Ratio of Power Spectra of the Two Modes

The velocity field of the cosmic baryonic fluid \mathbf{u} , can be separated by the Helmholtz-Hodge decomposition (Arfken & Weber 2005) into a compressive (or longitudinal) part ' \parallel ' and a solenoidal (or transverse) part ' \perp ', as $\mathbf{u} = \mathbf{u}_{\parallel} + \mathbf{u}_{\perp}$, where the compressive part \mathbf{u}_{\parallel} satisfies $\nabla \times \mathbf{u}_{\parallel} = 0$, and the solenoidal part \mathbf{u}_{\perp} satisfies $\nabla \cdot \mathbf{u}_{\perp} = 0$, respectively.

For $d \equiv \nabla \cdot \mathbf{u}$ and $\omega \equiv \nabla \times \mathbf{u}$, it is easy to show that

$$P_d(k) = k^2 P_{\parallel}(k), \quad P_{\omega}(k) = k^2 P_{\perp}(k), \quad (10)$$

in which $P_d(k)$, $P_{\omega}(k)$, $P_{\parallel}(k)$ and $P_{\perp}(k)$ are the FPS of d , ω , the velocity component \mathbf{u}_{\parallel} and \mathbf{u}_{\perp} , respectively.

The small-scale compression ratio can be defined as follows (Kida & Orszag 1990; Schmidt et al. 2009)

$$r_{\text{CS}} \equiv \frac{\langle d^2 \rangle}{\langle d^2 \rangle + \langle \omega^2 \rangle}. \quad (11)$$

This ratio quantifies the relative importance of the compressive and solenoidal modes in a flow (Iapichino et al. 2011). Inspired by this compression ratio, we define the FPS ratio of the solenoidal mode to the total turbulence as

$$Q(k) \equiv \frac{P_{\omega}(k)}{P_d(k) + P_{\omega}(k)} = \frac{P_{\perp}(k)}{P_{\parallel}(k) + P_{\perp}(k)}. \quad (12)$$

It can be seen that $Q(k)$ is superior to r_{CS} as it is scale dependent. Similar to the FPS ratio in Equation (12), we can also define the global WPS ratio as

$$\tilde{Q}(k) \equiv \frac{\tilde{P}_{\omega}(k)}{\tilde{P}_d(k) + \tilde{P}_{\omega}(k)} = \frac{\tilde{P}_{\perp}(k)}{\tilde{P}_{\parallel}(k) + \tilde{P}_{\perp}(k)}, \quad (13)$$




Figure 3. The z -evolution of the energy spectrum (left panel) and the spectral ratio (right panel) for the TNG50 simulation. Similar to Figure 2, the relevant power-laws are indicated in the figure. The grey area shows the $k > 0.4k_{\text{Nyquist}}$ region.




Figure 4. The global kurtosis of the velocity field \mathbf{u} , \mathbf{u}_{\parallel} and \mathbf{u}_{\perp} of the TNG50 data at $z = 0$, respectively. The curves show the x -, y -, z -direction of \mathbf{u} and the z -direction of \mathbf{u}_{\parallel} and \mathbf{u}_{\perp} respectively. The horizontal dotted line shows the kurtosis = 3, and the grey area shows the $k > 0.4k_{\text{Nyquist}}$ region.

in which $\tilde{P}_d(k)$, $\tilde{P}_\omega(k)$, $\tilde{P}_{\parallel}(k)$, and $\tilde{P}_{\perp}(k)$ are global WPS of d , ω , the velocity component \mathbf{u}_{\parallel} and \mathbf{u}_{\perp} , respectively, computed in the same way as Equation (5), and the correspondence $w = c_w k$ is used to convert w to k .

As mentioned above, the turbulence in the cosmic fluid consists not only of Kolmogorov turbulence, but also of Burgers turbulence. Therefore, it is not proper to treat the turbulent energy spectrum as consisting only of the solenoidal component (cf. Ryu et al. 2008), and it is better to take $P_d(k) + P_\omega(k)$, or its wavelet counterpart $\tilde{P}_d(k) + \tilde{P}_\omega(k)$, as the total energy spectrum. Thus $Q(k)$ and $\tilde{Q}(k)$ reflect the solenoidal fraction of the total turbulence energy as a function of k .

Figure 5. z -evolution of the global kurtosis for the TNG50 data. The horizontal dotted line shows the kurtosis = 3, and the grey area shows the $k > 0.4k_{\text{Nyquist}}$ region.

With the env-WPS of the velocity field in Equation (4), we can also define the env-WPS ratio as

$$\tilde{Q}(k, \delta) \equiv \frac{\tilde{P}_\omega(k, \delta)}{\tilde{P}_d(k, \delta) + \tilde{P}_\omega(k, \delta)} = \frac{\tilde{P}_{\perp}(k, \delta)}{\tilde{P}_{\parallel}(k, \delta) + \tilde{P}_{\perp}(k, \delta)}, \quad (14)$$

in which $\tilde{P}_d(k, \delta)$, $\tilde{P}_\omega(k, \delta)$, $\tilde{P}_{\parallel}(k, \delta)$ and $\tilde{P}_{\perp}(k, \delta)$ are the env-WPS of d , ω , the velocity component \mathbf{u}_{\parallel} and \mathbf{u}_{\perp} , respectively. $\tilde{Q}(k, \delta)$ is the ratio of the solenoidal component to the total energy as a function of k and the environment δ .

2.3. Simulation Data

We utilize data from the IllustrisTNG simulations (TNG for short) (Pillepich et al. 2018a; Springel et al. 2018; Marinacci et al. 2018; Nelson et al. 2018; Naiman et al.




Figure 6. The environment-dependent wavelet energy spectra of the TNG50 simulation at $z = 0$ for the velocity field \mathbf{u} of the cosmic fluid (left column), with the corresponding spectral ratios (right column). The five environments are indicated on the color scale on the right side. The grey area shows the $k > 0.4k_{\text{Nyquist}}$ region.

2018; Nelson et al. 2019), focusing on the samples TNG300-1, TNG100-1, and TNG50-1. These samples correspond to simulated universes of sizes $205 h^{-1}\text{Mpc}$, $75 h^{-1}\text{Mpc}$ and $35 h^{-1}\text{Mpc}$, respectively. TNG simulations are comprehensive, encompassing large-scale gravo-magnetohydrodynamical processes within the cosmos, and are executed using the moving-mesh code AREPO (Springel 2010). This code employs a second-order accurate Godunov-type scheme for the hydrodynamic equations on a dynamically unstructured mesh, improving the fidelity of both supersonic and subsonic fluid turbulence simulations. Bauer & Springel (2012) demonstrated AREPO’s superiority in accurately replicating turbulence, achieving Kolmogorov-like scaling laws for density, velocity, and vorticity power spectra, aligning with theoretical expectations from the fully developed isotropic turbulence.

In addition to gravitational and hydrodynamic calculations, the TNG simulations incorporate a comprehensive set of physical processes (Pillepich et al. 2018b; Nelson et al. 2019), which include: (1) the formation and evolution of stars, (2) the associated metal enrichment and loss of mass, (3) cooling processes for primordial and metal-enriched gases, (4) the impact of SN on the pressurization of the interstellar medium, without resolving individual SN events, (5) feedback from stars, manifesting as galactic winds powered either by SN or stars in the asymptotic giant branch (AGB), (6) the formation and growth of supermassive black holes, accompanied by energy feedback from AGN in both high-accretion quasar and low-accretion kinetic wind phases, and (7) the role of magnetic fields in cosmic structures.

Given the abundance of the physical processes included in the TNG simulations and AREPO’s excellent performance in hydrodynamical computations, the TNG data are well suited for studying cosmic fluid turbulence.

3. RESULTS

We use the piecewise cubic spline (PCS) scheme (Sefusatti et al. 2016) to assign all the simulation particles into a 1536^3 grid, and with this grid, we can use FFT to compute the FPS and WPS (see Wang & He 2024, for the numerical pipeline).

3.1. The Global Results

As mentioned in Section 2.2.1, in all the following computations, the bulk flow has been removed from the velocity field using the approach of Vazza et al. (2012).

In Figure 2, we show both the FPS and the global WPS at $z = 0$ for the velocity field \mathbf{u} and its ‘ \parallel ’ and ‘ \perp ’ components of the cosmic fluid, with the corresponding spectral ratios $Q(k)$ and $\tilde{Q}(k)$. From top to bottom are the results for the TNG300, TNG100, and TNG50 simulations, respectively. As analyzed in Section 2.1, the global WPS is almost the same as the FPS, just with a slightly higher factor. The ratios $Q(k)$ and $\tilde{Q}(k)$ also give almost the same results. Indeed, we observe that the Fourier analysis and the global wavelet analysis give consistent results, suggesting that the wavelet analysis is valid.

It can be seen that there are peaks in all energy spectra of the velocity \mathbf{u} of the cosmic fluid, and the peak scales are $k_{\text{S-peak}} = 6.0, 8.2$ and $8.8 h\text{Mpc}^{-1}$ for TNG300, TNG100 and TNG50 respectively. At large scales, i.e. $k < k_{\text{S-peak}}$, the energy spectra are increasing functions of k . The velocity field undergoes structure collapse and is mostly dominated by its compressive component. Within this scale range, the gravitational collapse converts the potential energy of the structure into kinetic energy of the bulk flow, and the bulk flow further decays into turbulent flows - first in the compressive mode and, with increasing k , mostly in the solenoidal mode, as can also be seen from the spectral ratios $Q(k)$ and $\tilde{Q}(k)$. The transition scales of the two modes can be determined




Figure 7. z -evolution of the energy spectrum (left column) and the spectral ratio (right column) for the TNG50 simulation of the three environments Δ_0 , Δ_2 and Δ_4 , respectively. The four redshifts are shown in the figure. As in Figure 2, the relevant power-laws are indicated in the figure, and the grey area shows the $k > 0.4k_{\text{Nyquist}}$ region.




Figure 8. Environment-dependent kurtosis for the TNG50 data at $z = 0$. The five environments are indicated on the color scale on the right side. The horizontal dotted line shows the kurtosis = 3, and the grey area shows the $k > 0.4k_{\text{Nyquist}}$ region.

by setting $\tilde{Q}(k_{\text{trans}})$ or $Q(k_{\text{trans}}) = 1/2$, and thus we get $k_{\text{trans}} = 1.9, 2.9$ and 2.7 hMpc^{-1} for TNG300, TNG100 and TNG50 respectively.

One can see that the energy spectra drop rapidly at $k > k_{\text{S-peak}}$. As shown in the figure, the exponent of the energy spectrum within this scale range is $-2.59, -2.56$ and -2.41 for TNG300, TNG100 and TNG50 respectively, which are all steeper than not only the Kolmogorov exponent $-5/3$ but also the Burgers turbulence exponent -2 . As discussed in Section 2.2.2, Kolmogorov turbulence is excited by subsonic motion of the flow, characterized by eddies of different scales, and hence the \mathbf{u}_\perp component is mostly responsible for Kolmogorov turbulence, while Burgers turbulence is excited by supersonic motion, characterized by shock waves, and hence the \mathbf{u}_\parallel component is mostly responsible for Burgers turbulence. The mechanism of energy dissipation is different for Kolmogorov and Burgers turbulence. In Kolmogorov turbulence, kinetic energy is transferred from larger vortices to smaller and smaller vortices and then dissipated into thermal energy below the dissipation scale, whereas in Burgers turbulence, kinetic energy can be directly dissipated into thermal energy by shock waves at any scale. As mentioned earlier, the turbulence in the cosmic baryonic fluid is not simply a mixture of Kolmogorov turbulence and Burgers turbulence, and hence the steeper exponent of the energy spectrum suggests that there may be more complex mechanisms for energy transfer than Kolmogorov and Burgers turbulence.

We notice that there are also peaks in the spectral ratios $Q(k)$ and $\tilde{Q}(k)$. The peak scales $k_{\text{Q-peak}} = 14.9, 25.7$ and 34.9 hMpc^{-1} for TNG300, TNG100 and TNG50 respectively. The power of the \mathbf{u}_\perp component drops rapidly beyond $k_{\text{Q-peak}}$.

According to the general theory of turbulence, the turbulence energy spectrum can be divided into the energy-containing range, the inertial range and the dissipation range, using two characteristic scales - the integral scale and the dissipation scale (e.g. Davidson 2015). In our current work, the peak scale $k_{\text{S-peak}}$ can be treated as the integral scale, and the scale range $k < k_{\text{S-peak}}$ can be considered as the energy-containing range. The peak scale $k_{\text{Q-peak}}$ can be treated as the dissipation scale, determined not by physical but by numerical viscosity. Therefore, the scale range $k_{\text{S-peak}} < k < k_{\text{Q-peak}}$ can be considered as the inertial range, and the scale range $k > k_{\text{Q-peak}}$ can be considered as the dissipation range of turbulence.

Note that, taking into account the numerical viscosity, the effective Reynolds number of the numerically computed flow scales as $\sim (L_c/\Delta_g)^{4/3}$, where L_c is some correlation length, say the integral scale, and Δ_g is the grid scale (Schmidt 2015). Due to the smaller Δ_g and therefore larger effective Reynolds number for TNG50, this is the reason why the dissipation scale for TNG50 is the smallest. Therefore, the TNG50 simulation is less affected by numerical viscosity, and is more suitable for turbulence studies. In the following, we will use only TNG50 data for the subsequent work.

However, it should be noted that when $k > 0.4k_{\text{Nyquist}}$ (indicated by the grey area in the figure), where k_{Nyquist} is the Nyquist frequency, the numerical effects such as smearing, aliasing and shot noise, are significant, the conclusions are not very reliable (Wang & He 2024), and therefore we restrict ourselves to consider only the scale range of $k < 0.4k_{\text{Nyquist}}$.

Figure 3 shows the z -evolution of both the global energy spectrum and the spectral ratio \tilde{Q} for the TNG50 simulation. We observe that the spectrum grows slightly from $z = 4$ to 2, but increases significantly from $z = 2$ to 1 in the $k < k_{\text{S-peak}}$ scale range, indicating that turbulence induced by structure formation is increasingly injected into the cosmic fluid with time. However, the spectrum remains almost unchanged from $z = 1$ to 0, indicating that the turbulence is close to saturation, or that the injection of turbulence by structure formation is balanced by the transfer from the larger to the smaller scales and the dissipation of turbulence into heat. In the $k > k_{\text{S-peak}}$ range, the energy spectrum is also enhanced from $z = 4$ to 2, but the magnitude of the enhancement is less than for $k < k_{\text{S-peak}}$.

From $z = 4$ to $z = 0$, the peak scale of the spectrum increases from $k_{\text{S-peak}} = 27.7 \text{ hMpc}^{-1}$ to $k_{\text{S-peak}} = 8.8 \text{ hMpc}^{-1}$, and the transition scale from the compressive mode to the solenoidal mode also increases from $k_{\text{trans}} = 14.2 \text{ hMpc}^{-1}$ to $k_{\text{trans}} = 2.7 \text{ hMpc}^{-1}$, while the peak scale of \tilde{Q} decreases from $k_{\text{Q-peak}} = 27.7 \text{ hMpc}^{-1}$ to $k_{\text{Q-peak}} = 34.9 \text{ hMpc}^{-1}$. The z -evolution of these characteristic scales indicates that the inertial range for the turbulence of the cosmic fluid gradually increases with decreasing z .




Figure 9. z -evolution of the env-kurtosis for the TNG50 simulation of the three environments Δ_0 , Δ_2 and Δ_4 , respectively. The horizontal dotted line shows the kurtosis = 3, and the grey area shows the $k > 0.4k_{\text{Nyquist}}$ region.

In Figure 4, we show the global kurtosis as a function of k for the $z = 0$ TNG50 data. We can see that the three directions give almost the same results, so we will only use the z -direction data in the following calculation. For the \mathbf{u} field, it can be seen that the kurtosis = 3 at about $k = 0.2 h\text{Mpc}^{-1}$, then grows rapidly with k to about $k_{\text{trans}} = 2.7 h\text{Mpc}^{-1}$, and then grows slowly to the large- k end, indicating that the cosmic fluid becomes increasingly intermittent as k increases. As can be seen from the results for \mathbf{u}_\perp and \mathbf{u}_\parallel , the kurtosis for \mathbf{u}_\perp is generally greater than that for \mathbf{u}_\parallel . Therefore, combining the energy spectra of \mathbf{u}_\perp and \mathbf{u}_\parallel , we find that the global kurtosis is biased towards that for \mathbf{u}_\parallel if $k < k_{\text{trans}}$, whereas the global kurtosis is dominated by that for \mathbf{u}_\perp if $k > k_{\text{trans}}$, and thus the transition scale k_{trans} can be determined by $\tilde{Q}(k_{\text{trans}}) = 1/2$. We see that k_{trans} just happens to be the scale at which the fast growing mode transitions to the slow growing mode for the global kurtosis.

In Figure 5, we show the z -evolution of the global kurtosis as a function of k for the TNG50 data. One can see that for all four redshifts the kurtosis increases with k , just as for $z = 0$, first in a fast mode, then in a slow growth mode. As previously analyzed, the transition scales $k_{\text{trans}} = 14.2, 7.9, 5.0$ and $2.7 h\text{Mpc}^{-1}$ for $z = 4, 2, 1$ and 0 respectively, corresponding to scales where the fast growing mode switches to the slow mode.

At large scales of $k < 3 h\text{Mpc}^{-1}$ the kurtosis increases monotonically with decreasing z , whereas at small scales of $k > 3 h\text{Mpc}^{-1}$ the kurtosis does not change monotonically with decreasing redshift - from $z = 4$ to 2 the global kurtosis first increases and then decreases at $z < 2$. These results show how the intermittency of turbulence for cosmic fluid evolves with redshift.

3.2. The Environment-dependent Results

We divide the simulation space of TNG50 into five different environments according to the dark matter density, denoted as Δ_i with $i = 0, 1, \dots, 4$ and listed in Table 1. Among

Table 1. The local density environments, specified with $\Delta_{\text{dm}} = \rho_{\text{dm}}/\bar{\rho}_{\text{dm}}$ of dark matter.

	Δ_0	Δ_1	Δ_2	Δ_3	Δ_4
$\Delta_{\text{dm}} \in$	$[0, 1/8)$	$[1/8, 1/2)$	$[1/2, 2)$	$[2, 8)$	$[8, +\infty)$

them, Δ_0 and Δ_1 can be regarded as voids or low-density regions, and Δ_4 as various high-density structures, such as clusters, filaments, sheets and their outskirts (see the bottom-right panel of Figure 1).

In Figure 6, we show the $z = 0$ energy spectra of the velocity field \mathbf{u} for the five environments of TNG50. We see that, similar to the global energy spectra in Figure 2, there also exist peaks in these env-dependent spectra. As the density increases, there is a tendency for the peak position $k_{\text{S-peak}}$ to shift towards the larger k -end. These five env-dependent energy spectra, starting from the same k , increase with the increasing k , reach their peaks, and then decrease rapidly, with the exponent also steeper than the Burgers exponent. These spectra are regularly distributed - the spectrum with the lowest density at the bottom and the spectrum with the highest density at the top. These results suggest that the cosmic fluid is more turbulent in a high-density environment than in a low-density environment.

From the env-dependent $\tilde{Q}(k, \Delta_i)$, we see that the transition scales k_{trans} for the five environments are between $2 - 3 h\text{Mpc}^{-1}$. When $k > 20 h\text{Mpc}^{-1}$, \tilde{Q} increases from low- to high-density environments, implying that the solenoidal mode of the velocity field is more significant in high-density environments than in low-density environments.

In Figure 7, we show the z -evolution of the env-dependent energy spectra and the spectral ratios $\tilde{Q}(k, \Delta_i)$ for TNG50 data from the three environments Δ_0 , Δ_2 and Δ_4 respectively. In $k < k_{\text{S-peak}}$, all env-dependent spectra grow slightly from $z = 4$ to 2 , but are significantly enhanced from $z = 2$ to 1 , and the spectra remain almost unchanged from

$z = 1$ to 0 . In $k > k_{S-\text{peak}}$, these energy spectra are also enhanced from $z = 2$ to 1 , but the enhancement is less than for $k < k_{S-\text{peak}}$. From $z = 4$ to $z = 0$, both the peak scales of the spectra and the transition scales increase, while the peak scales of $\tilde{Q}(k, \Delta_i)$ decrease, and thus the inertial ranges increase with decreasing redshift. In general, the z -evolution of the env-dependent energy spectra and spectral ratios is similar to the z -evolution of the global energy spectra and spectral ratios in Figure 3.

In Figure 8, we show the env-kurtosis for $z = 0$ TNG50 data. It can be seen that these env-dependent results are all similar to the global kurtosis - a fast growing mode followed by a slow growing mode at the transition scale k_{trans} as k increases. However, the kurtosis is lowest for the densest environment Δ_4 and increases towards the lowest density environment Δ_0 . Such results indicate that the intermittency of the velocity field in the region $k > k_{\text{trans}}$ becomes increasingly strong as the environment density decreases.

In Figure 9, we show the z -evolution of the env-kurtosis for the three environments Δ_0 , Δ_2 and Δ_4 . In general, these results are similar to the z -evolution of the global kurtosis shown in Figure 5, except for the cases of Δ_0 and Δ_4 , where the kurtosis does not change monotonically from $z = 4$ to 2 at small scales of $k > 20 h\text{Mpc}^{-1}$. The puzzling nature of the z -evolution for the env-kurtosis as such suggests the existence of complex physical mechanisms behind it.

4. SUMMARY AND CONCLUSIONS

The turbulent motion of the cosmic baryonic fluid, an important topic in the study of the large-scale structures of the universe, has attracted increasing attention in cosmological studies over the last few decades. In this study, we first use CWT techniques to construct the global and environment-dependent wavelet energy spectra of the cosmic baryonic fluid, which are used to characterize the intensity of the turbulence as a function of the wavenumber k . The presence of shocks leads to a strong intermittency in the baryonic fluid, so that the turbulent energy dissipation in space is also intermittent. The deviation of the tails of the PDF from Gaussian is usually considered as a manifestation of intermittency. We then define the wavelet global and environment-dependent kurtosis (or flatness) of the velocity field to characterize the intermittency of turbulence in the cosmic fluid.

We use the velocity field data from the IllustrisTNG simulation. The velocity of the baryonic fluid is decomposed into turbulent and bulk flow using the iterative multiscale filtering approach of Vazza et al. (2012), and the turbulent flow is further separated into compressive and solenoidal modes using the Helmholtz-Hodge decomposition.

We define five environments according to the dark matter density. We then compute both the global and environment-dependent energy spectrum and kurtosis of the turbulent ve-

locity for $z = 0$ TNG data, and present the z -evolution of the turbulent energy spectrum and kurtosis. To study the contribution of the solenoidal (or compressive) mode to the total turbulence energy, we also define the spectral ratio, either the Fourier Q or the wavelet \tilde{Q} , of the two modes of the velocity field. Our main findings and conclusions are as follows:

1. There are peaks in all the energy spectra at $k_{S-\text{peak}}$ and in the spectral ratios $Q(k)$ and $\tilde{Q}(k)$ at $k_{Q-\text{peak}}$ for the velocity field of the cosmic baryonic fluid. For the two characteristic scales, $k_{S-\text{peak}}$ and $k_{Q-\text{peak}}$ can be treated as the integral scale and the dissipation scale, and the scale range $k < k_{S-\text{peak}}$, $k_{S-\text{peak}} < k < k_{Q-\text{peak}}$ and $k > k_{Q-\text{peak}}$ can be considered as the energy-containing range, the inertial range, and the dissipation range of the turbulence, respectively. In the energy-containing range, the velocity field undergoes structure collapse and is mostly dominated by its compressive component. The gravitational collapse converts the potential energy of the structure into kinetic energy of the bulk flow, and further decays into turbulent flows, initially in the compressive mode and, as k increases, mostly in the solenoidal mode. In the inertial range, the energy passes from large-scale to small-scale eddies by the solenoidal mode or is dissipated directly into thermal energy by the compressive mode. The exponent of the energy spectrum is steeper than not only the Kolmogorov exponent but also the Burgers exponent, suggesting that there may be more complex mechanisms for energy transfer than Kolmogorov and Burgers turbulence.
2. The global energy spectrum increases significantly from $z = 2$ to 1 in the energy-containing range, indicating that structure formation-induced turbulence is increasingly injected into the cosmic fluid with time. However, the spectrum remains almost unchanged from $z = 1$ to 0 , indicating that the injection of turbulence by structure formation is balanced by the transfer from the larger to the smaller scales and the dissipation of turbulence into heat. In the inertial range, the energy spectrum is also enhanced from $z = 2$ to 1 , but the magnitude of the enhancement is less than for the energy-containing range. We also find that the inertial range for the turbulence of the cosmic fluid gradually increases with decreasing z .
3. Similar to the global energy spectra, the environment-dependent spectra can also be classified into the energy-containing, the inertial and the dissipation range based on the two characteristic scale, $k_{S-\text{peak}}$ and $k_{Q-\text{peak}}$. The spectral magnitude is lowest in the lowest-density environment and highest in the highest-density environment, suggesting that the cosmic fluid

is more turbulent in a high-density environment than in a low-density environment. In $k > 20 h\text{Mpc}^{-1}$, \tilde{Q} increases from low- to high-density environments, implying that the solenoidal mode of the velocity field is more significant in high-density environments than in low-density environments. The z -evolution of the environment-dependent energy spectra and spectral ratios is generally similar to that of the global energy spectra and spectral ratios.

4. The kurtosis (or flatness) of the cosmic velocity field is an appropriate statistic to characterize the intermittency of turbulence in the cosmic fluid. A characteristic scale k_{trans} can be determined by $\tilde{Q}(k_{\text{trans}})$ or $Q(k_{\text{trans}}) = 1/2$. For the velocity field, the wavelet global kurtosis grows rapidly with k in the $k < k_{\text{trans}}$ range, and then grows slowly in the $k > k_{\text{trans}}$ range, indicating that the cosmic fluid becomes increasingly intermittent as k increases. The kurtosis for the solenoidal component is generally greater than that of the compressive component. As z decreases, the kurtosis increases in the $k < 3 h\text{Mpc}^{-1}$ range, whereas in $k > 3 h\text{Mpc}^{-1}$ the global kurtosis first increases from $z = 4$ to 2 and then decreases at $z < 2$.
5. The environment-dependent kurtosis is almost completely similar to the global one - a fast growing mode followed by a slow growing mode at the transition scale k_{trans} as k increases. The kurtosis is lowest for the densest environment and increases towards the lowest-density environment, indicating that the intermittency of the velocity field in the $k > k_{\text{trans}}$ range becomes increasingly strong as the density of the environments decreases. The z -evolution of the env-kurtosis is similar to that of the global kurtosis, except for the cases of the lowest- and highest-density environment, where the kurtosis does not change monotonically from $z = 4$ to 2 in the $k > 20 h\text{Mpc}^{-1}$ range. The puzzling nature of the z -evolution for the env-kurtosis as such suggests the existence of complex physical mechanisms behind it.

Comparisons of the global WPS and the spectral ratios $\tilde{Q}(k)$ with the corresponding results of the Fourier analysis show that the Fourier analysis and the global wavelet analysis give almost the same results, demonstrating that the wavelet analysis techniques are reliable and trustworthy. However, the environment-dependent wavelet analysis proves to be more powerful than the Fourier analysis in that the wavelet statistics such as the env-WPS or the env-spectral ratios $\tilde{Q}(k, \Delta_i)$ cannot be implemented by the Fourier analysis. In this work, the environments are simply defined by the dark matter density. In fact, one can design various complex

environments by specifying, for example, a particular spatial region or a specific cosmic structure.

The simple theoretical framework of turbulence is based on Kolmogorov turbulence, which is homogeneous and isotropic in space and is characterized by eddies of different scales. However, this picture is problematic for the turbulence of the cosmic baryonic fluid, which is subject to structure formation driven by gravity in the context of cosmic expansion. Structure formation leads, for example, to the density stratification of the cosmic fluid, where buoyancy forces resist radial motions, making the turbulence anisotropic (e.g. Shi et al. 2018; Shi & Zhang 2019; Mohapatra et al. 2020; Simonte et al. 2022; Wang et al. 2023). In this work, we have not yet considered the anisotropic turbulence caused by the density stratification, which will be left for future studies.

Following the general theory of turbulence, we also divide the turbulent energy spectrum of the cosmic baryonic fluid into three ranges, the energy-containing range, the inertial range and the dissipation range. Note that these authors (Schmidt et al. 2009, 2010, 2016; Shi et al. 2018; Shi & Zhang 2019; Shi et al. 2020) also refer to these terms, but they should adopt different definitions from ours. Indeed, as Shi et al. (2018) points out, these concepts are not easy to identify and define. In fact, the real situation is much more complex than this general picture. Since the compressive mode exists at all scales, the kinetic energy of the turbulent flow, even of the bulk flow, can be directly dissipated into thermal energy at all scales.

In addition, in the inertial and dissipation range, both kinetic and thermal energy may also be injected by feedback from SN-driven galactic winds or AGN outflows, and these physical processes deserve to be studied in detail. Thus, the fact that the spectral exponent in the inertial range is steeper than that of Kolmogorov or Burgers turbulence, indicates that the simple theory of Kolmogorov or Burgers turbulence does not hold for the cosmic baryonic fluid. One can speculate that a steeper energy spectrum may be of great significance for structure formation, which, for example, can lead to more efficient transfer of energy to smaller scales, potentially affecting the distribution of galaxies and galaxy clusters.

ACKNOWLEDGMENTS

We acknowledge the use of the data from IllustrisTNG simulation for this work. We also acknowledge the support by the National Science Foundation of China (No. 12147217, 12347163), and by the Natural Science Foundation of Jilin Province, China (No. 20180101228JC).

Software: NumPy (<https://numpy.org/>), nbbodykit (<https://nbbodykit.readthedocs.io/en/latest/>), SciPy (<https://scipy.org/>), Matplotlib (<https://matplotlib.org/>), Jupyter Notebook (<https://jupyter.org/>), WPSmesh (<https://github.com/WangYun1995/WPSmesh>)

REFERENCES

- Angelinelli, M., Vazza, F., Giocoli, C., et al. 2020, MNRAS, 495, 864, doi: [10.1093/mnras/staa975](https://doi.org/10.1093/mnras/staa975)
- Angulo, J. M., & Madrid, A. E. 2023, Entropy, 25, 1080, doi: [10.3390/e25071080](https://doi.org/10.3390/e25071080)
- Arfken, G. B., & Weber, H. J. 2005, Mathematical methods for physicists 6th ed. (Elsevier)
- Balkovsky, E., Falkovich, G., Kolokolov, I., & Lebedev, V. 1997, PhRvL, 78, 1452, doi: [10.1103/PhysRevLett.78.1452](https://doi.org/10.1103/PhysRevLett.78.1452)
- Banerjee, N., & Sharma, P. 2014, MNRAS, 443, 687, doi: [10.1093/mnras/stu1179](https://doi.org/10.1093/mnras/stu1179)
- Bauer, A., & Springel, V. 2012, MNRAS, 423, 2558, doi: [10.1111/j.1365-2966.2012.21058.x](https://doi.org/10.1111/j.1365-2966.2012.21058.x)
- Boldyrev, S., Linde, T., & Polyakov, A. 2004, PhRvL, 93, 184503, doi: [10.1103/PhysRevLett.93.184503](https://doi.org/10.1103/PhysRevLett.93.184503)
- Bonafede, A., Feretti, L., Murgia, M., et al. 2010, A&A, 513, A30, doi: [10.1051/0004-6361/200913696](https://doi.org/10.1051/0004-6361/200913696)
- Bregman, J. N. 2007, ARA&A, 45, 221, doi: [10.1146/annurev.astro.45.051806.110619](https://doi.org/10.1146/annurev.astro.45.051806.110619)
- Brüggen, M., Hoeft, M., & Ruszkowski, M. 2005, ApJ, 628, 153, doi: [10.1086/430732](https://doi.org/10.1086/430732)
- Brüggen, M., & Vazza, F. 2015, in Astrophysics and Space Science Library, Vol. 407, Magnetic Fields in Diffuse Media, ed. A. Lazarian, E. M. de Gouveia Dal Pino, & C. Melioli, 599
- Cassano, R., & Brunetti, G. 2005, MNRAS, 357, 1313, doi: [10.1111/j.1365-2966.2005.08747.x](https://doi.org/10.1111/j.1365-2966.2005.08747.x)
- Churazov, E., Forman, W., Jones, C., Sunyaev, R., & Böhringer, H. 2004, MNRAS, 347, 29, doi: [10.1111/j.1365-2966.2004.07201.x](https://doi.org/10.1111/j.1365-2966.2004.07201.x)
- Churazov, E., Vikhlinin, A., Zhuravleva, I., et al. 2012, MNRAS, 421, 1123, doi: [10.1111/j.1365-2966.2011.20372.x](https://doi.org/10.1111/j.1365-2966.2011.20372.x)
- Davidson, P. A. 2015, TURBULENCE - An Introduction for Scientists and Engineers, 2nd ed. (Oxford University Press)
- Dolag, K., Vazza, F., Brunetti, G., & Tormen, G. 2005, MNRAS, 364, 753, doi: [10.1111/j.1365-2966.2005.09630.x](https://doi.org/10.1111/j.1365-2966.2005.09630.x)
- Enßlin, T. A., & Vogt, C. 2006, A&A, 453, 447, doi: [10.1051/0004-6361:20053518](https://doi.org/10.1051/0004-6361:20053518)
- Evoli, C., & Ferrara, A. 2011, MNRAS, 413, 2721, doi: [10.1111/j.1365-2966.2011.18343.x](https://doi.org/10.1111/j.1365-2966.2011.18343.x)
- Farge, M. 1992, Annual Review of Fluid Mechanics, 24, 395, doi: [10.1146/annurev.fl.24.010192.002143](https://doi.org/10.1146/annurev.fl.24.010192.002143)
- Federrath, C., Roman-Duval, J., Klessen, R. S., Schmidt, W., & Mac Low, M. M. 2010, A&A, 512, A81, doi: [10.1051/0004-6361/200912437](https://doi.org/10.1051/0004-6361/200912437)
- Feng, L.-L., Shu, C.-W., & Zhang, M. 2004, ApJ, 612, 1, doi: [10.1086/422513](https://doi.org/10.1086/422513)
- Gaspari, M., Churazov, E., Nagai, D., Lau, E. T., & Zhuravleva, I. 2014, A&A, 569, A67, doi: [10.1051/0004-6361/201424043](https://doi.org/10.1051/0004-6361/201424043)
- Gaspari, M., Melioli, C., Brighenti, F., & D’Ercole, A. 2011, MNRAS, 411, 349, doi: [10.1111/j.1365-2966.2010.17688.x](https://doi.org/10.1111/j.1365-2966.2010.17688.x)
- Guo, Q., White, S., Boylan-Kolchin, M., et al. 2011, MNRAS, 413, 101, doi: [10.1111/j.1365-2966.2010.18114.x](https://doi.org/10.1111/j.1365-2966.2010.18114.x)
- He, P., Liu, J., Feng, L.-L., Shu, C.-W., & Fang, L.-Z. 2006, PhRvL, 96, 051302, doi: [10.1103/PhysRevLett.96.051302](https://doi.org/10.1103/PhysRevLett.96.051302)
- Henriques, B. M. B., White, S. D. M., Thomas, P. A., et al. 2015, MNRAS, 451, 2663, doi: [10.1093/mnras/stv705](https://doi.org/10.1093/mnras/stv705)
- Hitomi Collaboration, Aharonian, F., Akamatsu, H., et al. 2016, Nature, 535, 117, doi: [10.1038/nature18627](https://doi.org/10.1038/nature18627)
- . 2018, PASJ, 70, 10, doi: [10.1093/pasj/psx127](https://doi.org/10.1093/pasj/psx127)
- Iapichino, L., Federrath, C., & Klessen, R. S. 2017, MNRAS, 469, 3641, doi: [10.1093/mnras/stx882](https://doi.org/10.1093/mnras/stx882)
- Iapichino, L., & Niemeyer, J. C. 2008a, MNRAS, 388, 1089, doi: [10.1111/j.1365-2966.2008.13518.x](https://doi.org/10.1111/j.1365-2966.2008.13518.x)
- . 2008b, MNRAS, 388, 1089, doi: [10.1111/j.1365-2966.2008.13518.x](https://doi.org/10.1111/j.1365-2966.2008.13518.x)
- Iapichino, L., Schmidt, W., Niemeyer, J. C., & Merklein, J. 2011, MNRAS, 414, 2297, doi: [10.1111/j.1365-2966.2011.18550.x](https://doi.org/10.1111/j.1365-2966.2011.18550.x)
- Iapichino, L., Viel, M., & Borgani, S. 2013, MNRAS, 432, 2529, doi: [10.1093/mnras/stt611](https://doi.org/10.1093/mnras/stt611)
- Khatri, R., & Gaspari, M. 2016, MNRAS, 463, 655, doi: [10.1093/mnras/stw2027](https://doi.org/10.1093/mnras/stw2027)
- Kida, S., & Orszag, S. A. 1990, Journal of Scientific Computing, 5, 85
- Kim, B., He, P., Pando, J., Feng, L.-L., & Fang, L.-Z. 2005, ApJ, 625, 599, doi: [10.1086/429556](https://doi.org/10.1086/429556)
- Konstandin, L., Shetty, R., Girichidis, P., & Klessen, R. S. 2015, MNRAS, 446, 1775, doi: [10.1093/mnras/stu2154](https://doi.org/10.1093/mnras/stu2154)
- Kowal, G., & Lazarian, A. 2010, ApJ, 720, 742, doi: [10.1088/0004-637X/720/1/742](https://doi.org/10.1088/0004-637X/720/1/742)
- Lau, E. T., Kravtsov, A. V., & Nagai, D. 2009, ApJ, 705, 1129, doi: [10.1088/0004-637X/705/2/1129](https://doi.org/10.1088/0004-637X/705/2/1129)
- Marinacci, F., Vogelsberger, M., Pakmor, R., et al. 2018, MNRAS, 480, 5113, doi: [10.1093/mnras/sty2206](https://doi.org/10.1093/mnras/sty2206)
- Meneveau, C. 1991, Journal of Fluid Mechanics, 232, 469, doi: [10.1017/S0022112091003786](https://doi.org/10.1017/S0022112091003786)
- Meyers, S. D., Kelly, B. G., & O’Brien, J. J. 1993, Monthly Weather Review, 121, 2858, doi: [10.1175/1520-0493\(1993\)121<2858:AITWAI>2.0.CO;2](https://doi.org/10.1175/1520-0493(1993)121<2858:AITWAI>2.0.CO;2)
- Miniati, F. 2014, ApJ, 782, 21, doi: [10.1088/0004-637X/782/1/21](https://doi.org/10.1088/0004-637X/782/1/21)
- Mohapatra, R., Federrath, C., & Sharma, P. 2020, MNRAS, 493, 5838, doi: [10.1093/mnras/staa711](https://doi.org/10.1093/mnras/staa711)
- Murgia, M., Govoni, F., Feretti, L., et al. 2004, A&A, 424, 429, doi: [10.1051/0004-6361:20040191](https://doi.org/10.1051/0004-6361:20040191)
- Naiman, J. P., Pillepich, A., Springel, V., et al. 2018, MNRAS, 477, 1206, doi: [10.1093/mnras/sty618](https://doi.org/10.1093/mnras/sty618)
- Nelson, D., Pillepich, A., Springel, V., et al. 2018, MNRAS, 475, 624, doi: [10.1093/mnras/stx3040](https://doi.org/10.1093/mnras/stx3040)

- Nelson, D., Springel, V., Pillepich, A., et al. 2019, *Computational Astrophysics and Cosmology*, 6, 2, doi: [10.1186/s40668-019-0028-x](https://doi.org/10.1186/s40668-019-0028-x)
- Norman, M. L., & Bryan, G. L. 1999, in *The Radio Galaxy Messier 87*, ed. H.-J. Röser & K. Meisenheimer, Vol. 530 (Springer-Verlag Berlin Heidelberg), 106
- Pillepich, A., Springel, V., Nelson, D., et al. 2018a, *MNRAS*, 473, 4077, doi: [10.1093/mnras/stx2656](https://doi.org/10.1093/mnras/stx2656)
- Pillepich, A., Nelson, D., Hernquist, L., et al. 2018b, *MNRAS*, 475, 648, doi: [10.1093/mnras/stx3112](https://doi.org/10.1093/mnras/stx3112)
- Porter, D. H., Jones, T. W., & Ryu, D. 2015, *ApJ*, 810, 93, doi: [10.1088/0004-637X/810/2/93](https://doi.org/10.1088/0004-637X/810/2/93)
- Roediger, E., & Brüggen, M. 2007, *MNRAS*, 380, 1399, doi: [10.1111/j.1365-2966.2007.12241.x](https://doi.org/10.1111/j.1365-2966.2007.12241.x)
- Ryu, D., Kang, H., Cho, J., & Das, S. 2008, *Science*, 320, 909, doi: [10.1126/science.1154923](https://doi.org/10.1126/science.1154923)
- Schmidt, W. 2015, *Living Reviews in Computational Astrophysics*, 1, 2, doi: [10.1007/lrca-2015-2](https://doi.org/10.1007/lrca-2015-2)
- Schmidt, W., Engels, J. F., Niemeyer, J. C., & Almgren, A. S. 2016, *MNRAS*, 459, 701, doi: [10.1093/mnras/stw632](https://doi.org/10.1093/mnras/stw632)
- Schmidt, W., Federrath, C., Hupp, M., Kern, S., & Niemeyer, J. C. 2009, *A&A*, 494, 127, doi: [10.1051/0004-6361:200809967](https://doi.org/10.1051/0004-6361/200809967)
- Schmidt, W., Kern, S. A. W., Federrath, C., & Klessen, R. S. 2010, *A&A*, 516, A25, doi: [10.1051/0004-6361/200913904](https://doi.org/10.1051/0004-6361/200913904)
- Schuecker, P., Finoguenov, A., Miniati, F., Böhringer, H., & Briel, U. G. 2004, *A&A*, 426, 387, doi: [10.1051/0004-6361:20041039](https://doi.org/10.1051/0004-6361:20041039)
- Sefusatti, E., Crocce, M., Scoccimarro, R., & Couchman, H. M. P. 2016, *MNRAS*, 460, 3624, doi: [10.1093/mnras/stw1229](https://doi.org/10.1093/mnras/stw1229)
- Shi, X., Nagai, D., Aung, H., & Wetzel, A. 2020, *MNRAS*, 495, 784, doi: [10.1093/mnras/staa1221](https://doi.org/10.1093/mnras/staa1221)
- Shi, X., Nagai, D., & Lau, E. T. 2018, *MNRAS*, 481, 1075, doi: [10.1093/mnras/sty2340](https://doi.org/10.1093/mnras/sty2340)
- Shi, X., & Zhang, C. 2019, *MNRAS*, 487, 1072, doi: [10.1093/mnras/stz1392](https://doi.org/10.1093/mnras/stz1392)
- Simonte, M., Vazza, F., Brighenti, F., et al. 2022, *A&A*, 658, A149, doi: [10.1051/0004-6361/202141703](https://doi.org/10.1051/0004-6361/202141703)
- Springel, V. 2010, *MNRAS*, 401, 791, doi: [10.1111/j.1365-2966.2009.15715.x](https://doi.org/10.1111/j.1365-2966.2009.15715.x)
- Springel, V., Pakmor, R., Pillepich, A., et al. 2018, *MNRAS*, 475, 676, doi: [10.1093/mnras/stx3304](https://doi.org/10.1093/mnras/stx3304)
- Subramanian, K., Shukurov, A., & Haugen, N. E. L. 2006, *MNRAS*, 366, 1437, doi: [10.1111/j.1365-2966.2006.09918.x](https://doi.org/10.1111/j.1365-2966.2006.09918.x)
- Torrence, C., & Compo, G. P. 1998, *Bulletin of the American Meteorological Society*, 79, 61, doi: [10.1175/1520-0477\(1998\)079<0061:APGTWA>2.0.CO;2](https://doi.org/10.1175/1520-0477(1998)079<0061:APGTWA>2.0.CO;2)
- Vacca, V., Murgia, M., Govoni, F., et al. 2010, *A&A*, 514, A71, doi: [10.1051/0004-6361/200913060](https://doi.org/10.1051/0004-6361/200913060)
- . 2012, *A&A*, 540, A38, doi: [10.1051/0004-6361/201116622](https://doi.org/10.1051/0004-6361/201116622)
- Vallés-Pérez, D., Planelles, S., & Quilis, V. 2021, *MNRAS*, 504, 510, doi: [10.1093/mnras/stab880](https://doi.org/10.1093/mnras/stab880)
- Vazza, F., Brunetti, G., Gheller, C., Brunino, R., & Brüggen, M. 2011, *A&A*, 529, A17, doi: [10.1051/0004-6361/201016015](https://doi.org/10.1051/0004-6361/201016015)
- Vazza, F., Jones, T. W., Brüggen, M., et al. 2017, *MNRAS*, 464, 210, doi: [10.1093/mnras/stw2351](https://doi.org/10.1093/mnras/stw2351)
- Vazza, F., Roediger, E., & Brüggen, M. 2012, *A&A*, 544, A103, doi: [10.1051/0004-6361/201118688](https://doi.org/10.1051/0004-6361/201118688)
- Vogt, C., & Enßlin, T. A. 2003, *A&A*, 412, 373, doi: [10.1051/0004-6361:20031434](https://doi.org/10.1051/0004-6361:20031434)
- . 2005, *A&A*, 434, 67, doi: [10.1051/0004-6361:20041839](https://doi.org/10.1051/0004-6361:20041839)
- Voit, G. M. 2005, *Reviews of Modern Physics*, 77, 207, doi: [10.1103/RevModPhys.77.207](https://doi.org/10.1103/RevModPhys.77.207)
- Walker, S. A., Sanders, J. S., & Fabian, A. C. 2015, *MNRAS*, 453, 3699, doi: [10.1093/mnras/stv1929](https://doi.org/10.1093/mnras/stv1929)
- Wang, C., Oh, S. P., & Ruszkowski, M. 2023, *MNRAS*, 519, 4408, doi: [10.1093/mnras/stad003](https://doi.org/10.1093/mnras/stad003)
- Wang, Y., & He, P. 2021, *Communications in Theoretical Physics*, 73, 095402, doi: [10.1088/1572-9494/ac10be](https://doi.org/10.1088/1572-9494/ac10be)
- Wang, Y., & He, P. 2022, *ApJ*, 934, 112, doi: [10.3847/1538-4357/ac7a3d](https://doi.org/10.3847/1538-4357/ac7a3d)
- . 2023, *RAS Techniques and Instruments*, 2, 307, doi: [10.1093/rasti/rzad020](https://doi.org/10.1093/rasti/rzad020)
- Wang, Y., & He, P. 2024, *MNRAS*, 528, 3797, doi: [10.1093/mnras/stae229](https://doi.org/10.1093/mnras/stae229)
- Wang, Y., Yang, H.-Y., & He, P. 2022, *ApJ*, 934, 77, doi: [10.3847/1538-4357/ac752c](https://doi.org/10.3847/1538-4357/ac752c)
- Yang, H.-Y., He, P., Zhu, W., & Feng, L.-L. 2020, *MNRAS*, 498, 4411, doi: [10.1093/mnras/staa2666](https://doi.org/10.1093/mnras/staa2666)
- Yang, H.-Y., Wang, Y., He, P., Zhu, W., & Feng, L.-L. 2022, *MNRAS*, 509, 1036, doi: [10.1093/mnras/stab3062](https://doi.org/10.1093/mnras/stab3062)
- Zhu, W., Feng, L.-L., & Fang, L.-Z. 2010, *ApJ*, 712, 1, doi: [10.1088/0004-637X/712/1/1](https://doi.org/10.1088/0004-637X/712/1/1)
- Zhu, W., Feng, L.-L., Xia, Y., et al. 2013, *ApJ*, 777, 48, doi: [10.1088/0004-637X/777/1/48](https://doi.org/10.1088/0004-637X/777/1/48)
- Zhuravleva, I., Allen, S. W., Mantz, A., & Werner, N. 2018, *ApJ*, 865, 53, doi: [10.3847/1538-4357/aadae3](https://doi.org/10.3847/1538-4357/aadae3)
- Zhuravleva, I., Churazov, E., Sunyaev, R., et al. 2013, *MNRAS*, 435, 3111, doi: [10.1093/mnras/stt1506](https://doi.org/10.1093/mnras/stt1506)
- Zhuravleva, I., Churazov, E., Schekochihin, A. A., et al. 2014, *Nature*, 515, 85, doi: [10.1038/nature13830](https://doi.org/10.1038/nature13830)
- ZuHone, J. A., Markevitch, M., Brunetti, G., & Giacintucci, S. 2013, *ApJ*, 762, 78, doi: [10.1088/0004-637X/762/2/78](https://doi.org/10.1088/0004-637X/762/2/78)