Momentum dependent nucleon-nucleon contact interactions and their effect on p-d scattering observables

- E. Filandri $^{1,2*},\;\;$ L. Girlanda $^{3,4},\;\;$ A. Kievsky $^{1,2},\;\;$ L.E. Marcucci $^{1,2},\;\;$ M. Viviani 1,2
- ^{1*}Department of Physics, University of Pisa, Largo Bruno Pontecorvo, Pisa, 56127, Italy.
 - ² INFN Sezione Pisa, Largo Bruno Pontecorvo, 3/Edificio C, Pisa, 56127, Italy.
- ³Department of Mathematics and Physics, University of Salento, Via per Arnesano, Lecce, I-73100, Italy.

 ²INFN, Sez. di Lecce, Lecce, I-73100, Italy.
 - *Corresponding author(s). E-mail(s): elena.filandri@df.unipi.it;

Abstract

Starting from a complete set of relativistic nucleon-nucleon contact operators up to order $O(p^4)$ of the expansion in the soft (relative or nucleon) momentum p, we show that non-relativistic expansions of relativistic operators involve twenty-six independent combinations, two starting at $O(p^0)$, seven at order $O(p^2)$ and seventeen at order $O(p^4)$. This demonstrates the existence of two low-energy free constants that parameterize interactions dependent on the total momentum of the pair of nucleons P. The latter, through the use of a unitary transformation, can be removed in the two-nucleon fourth-order contact interaction of the Chiral Effective Field Theory, generating a three-nucleon interaction at the same order. Within a hybrid approach in which this interaction is considered together with the phenomenological potential AV18, we show that the LECs involved can be used to fit very accurate data on the polarization observables of the low-energy p-d scattering, in particular the A_p asymmetry.

Keywords: Unitarity Transformation, Effective field theory, Three-body contact interactions

1 Introduction

Effective Field Theories (EFTs)[1–7] have established themselves as the preferred systematic approach for tackling the complex problem of nuclear interactions. This approach rests on several fundamental principles. It starts with the identification of the most general effective Lagrangian, respecting all pertinent symmetries, including the approximate chiral symmetry of Quantum Chromodynamics (QCD). The ordering of interactions is accomplished through a power-counting scheme. Consequently, this framework yields a predictive context in which physical observables, at each stage of the low-energy expansion, are expressed in terms of a finite set of low-energy constants (LECs). These LECs serve as fitting parameters and are determined through experimental data.

One essential task is the precise determination of the required number of parameters, both necessary and sufficient, at each stage of the expansion. This task not only rigorously scrutinizes the theory but also aids in estimating the theoretical uncertainty arising from unaccounted higher-order interactions [8–11]. In the realm of nuclear forces, these fitting parameters pertain to LECs associated with contact interactions between nucleons, which are not constrained by chiral symmetry. However, they are subject to constraints imposed by Poincaré symmetry [12, 13]. Despite the common non-relativistic quantum-mechanical framework used in nuclear physics, Poincaré symmetry must ultimately be respected. This requires the reconciliation of various relativistic effects arising from different sources, such as recoil corrections in energy denominators and vertex corrections from the heavy baryon expansion. Given that relativistic effects scale with the soft nucleon momenta, they can be systematically examined in the low-energy expansion, and the constraints on interactions can be imposed algebraically.

In the present work, starting from a manifestly Lorentz-invariant two-nucleon (2N) contact Lagrangian density and performing non-relativistic reductions up to the order $1/m^4$, where m represents the mass of the nucleon we retrace the results already obtained in Refs. [14, 15].

Furthermore, we underline the importance of two additional 2N contact LECs at N3LO, which characterize momentum-dependent interactions allowed by Poincaré symmetry. These LECs can be transformed into a three-nucleon (3N) interaction through a unitary transformation. This finding may explain the challenges faced when attempting to enhance accuracy in 3N systems, particularly in the context of scattering observables, during the transition from N2LO to N3LO [16].

The inclusion of the N4LO 3N contact interaction has proven to be of significant importance in reducing existing discrepancies between theoretical predictions and experimental data [17].

This paper offers quantitative evidence that the additional two LECs D_{16} and D_{17} at N3LO introduce the necessary flexibility to substantially enhance the description of low-energy p-d scattering polarization observables, with a particular focus on the A_y asymmetry. This aspect has long posed challenges for most nuclear interaction models.

Our approach is hybrid in nature, involving the consideration of the 3N force induced by D_{16} and D_{17} potential terms in conjunction with the phenomenological

Table 1 Transformation proprieties of the different elements of the Clifford algebra, metric tensor, Levi-Civita tensor and derivative operators under parity (\mathcal{P}) , charge conjugation (\mathcal{C}) and Hermitian conjugation (h.c.)

		1	γ_5	γ_{μ}	$\gamma_{\mu}\gamma_{5}$	$\sigma_{\mu\nu}$	$g_{\mu\nu}$	$\epsilon_{\mu\nu\rho}$	$\sigma \overleftrightarrow{\partial}_{\mu}$	∂_{μ}	$ au^a$
Ī	\mathcal{P}				_			_		+	+
	\mathcal{C}	+	+	_	+	_	+	+	_	+	$(-1)^{a+1}$
	h.c	. +	_	+	+	+	+	+	_	+	+

AV18 2N potential. A more comprehensive calculation in Chiral Effective Field Theory (ChEFT) is deferred to future research.

The structure of the paper is as follows. In Sec. 2 we show the basic steps of non-relativistic reduction to order p^4 of a covariant 2N Lagrangian, emphasizing the existence of two interactions dependent by the total momentum P accompanied by two free LECs D_{16} and D_{17} . In Sec. 3 we explain how these two off-shell interactions are related by unitary transformation to a three-body force and how they can be used for a fit of p-d polarization observables. In Sec. 4 we show the fit results and in Sec. 5 final conclusions are drawn.

2 Two extra interactions from the non relativistic reduction of 2N Contact Lagrangian up to N3LO

The general expression of the relativistic 2N contact Lagrangian is derived following the approach of Ref. [14, 15, 18, 19]. It consists of products of fermion bilinears, such as

$$(\bar{\psi} \overleftrightarrow{\partial}_{\mu_1} \cdots \overleftrightarrow{\partial}_{\mu_i} \Gamma_A \psi) \partial_{\lambda_1} \cdots \partial_{\lambda_k} (\bar{\psi} \overleftrightarrow{\partial}_{\nu_1} \cdots \overleftrightarrow{\partial}_{\nu_j} \Gamma_B \psi), \tag{1}$$

where ψ indicates the relativistic nucleon field, a doublet in isospin space, and $\Gamma_{A,B}$ are generic elements of the Clifford algebra.

To construct the covariant Lagrangian, various symmetries must be satisfied, including Lorentz invariance, isospin, parity, charge conjugation, and time reversal symmetry. According to the CPT theorem, time reversal symmetry is automatically satisfied if charge conjugation and parity symmetries are fulfilled.

Table 2 lists the transformation properties of different elements of the Clifford algebra, metric tensor, Levi-Civita tensor, and derivative operators under parity (\mathcal{P}) , charge conjugation (\mathcal{C}) , and Hermitian conjugation (h.c.). These properties play a crucial role in the construction of the Lagrangian.

Regarding the isospin degrees of freedom, both flavor structures $1 \otimes 1$ and $\tau^a \otimes \tau^a$ are allowed, but the latter can be disregarded thanks to Fierz identities. To specify the chiral order of each building block, it is necessary to identify the powers of the soft relative momentum \mathbf{p} . The derivatives ∂ acting on the entire bilinear are of order p, while the derivative $\overrightarrow{\partial}$ acting inside a bilinear is of order p^0 due to the presence of the heavy fermion mass scale. The fields equations of motion can be used to reduce the number of cosidered terms. Further criteria in specifying the power counting of the operators regard the Dirac matrix γ_5 , which can be thought of as O(p) since it mixes the large and small components of the Dirac spinor, and the Levi-Civita tensor

 $\epsilon_{\mu\nu\rho\sigma}$, which raises the chiral order by n-1, when contracted with n derivatives acting inside a bilinear.

These guidelines lead to the complete set of relativistic contact operators displayed in Table 2 of Ref.[15]. The last column contains, for each one of the Dirac structures, the additional combination of four-gradients arising up to $O(p^4)$. This construction differs from the one conducted in Ref. [14], due to a different choice of operators reduced by the equations of motion.

The next step is the non-relativistic reduction of these operators in terms of a minimal basis of non-relativistic 2N contact operators, involving up to 4 powers of three-gradients in terms of non-relativistic nucleon fields. It is important to note that the 2N contact Hamiltonian density takes the form of $\mathcal{H}_{2N} = \mathcal{H}^{(0)} + \mathcal{H}^{(2)} + \mathcal{H}^{(4)}$, with $\mathcal{H}^{(0)}$, $\mathcal{H}^{(2)}$, and $\mathcal{H}^{(4)}$ defined with the corresponding LECs.

Table 3 of Ref. [15] provides a complete basis of non-relativistic operators computed between states of two nucleons with initial and final momenta. It includes both LECs and operators for $O(p^4)$. Notably, this basis accounts for the general reference frame, and indeed some of the operators are P-dependent, where P denotes the overall momentum of the nucleon pairs.

The operators related to the constants D_{16} and D_{17} are introduced in this basis, representing new LECs that parametrize the P-dependent 2N interaction in the general reference frame. These LECs do not contribute in the center-of-mass frame.

3 Influence of two-body off-shell forces on the A_y puzzle

The N3LO 2N contact potential was originally considered in Refs. [20, 21] as consisting of 15 LECs. After careful scrutiny of the constraints imposed by relativity, two further LECs emerge, leading to the following expression in the general reference frame,

$$V_{2N}^{(4)} = D_{1}k^{4} + D_{2}Q^{4} + D_{3}k^{2}Q^{2} + D_{4}(\mathbf{k} \times \mathbf{Q})^{2} + [D_{5}k^{4} + D_{6}Q^{4} + D_{7}k^{2}Q^{2} + D_{8}(\mathbf{k} \times \mathbf{Q})^{2}] (\boldsymbol{\sigma}_{1} \cdot \boldsymbol{\sigma}_{2})$$

$$+ \frac{i}{2} (D_{9}k^{2} + D_{10}Q^{2}) (\boldsymbol{\sigma}_{1} + \boldsymbol{\sigma}_{2}) \cdot (\mathbf{Q} \times \mathbf{k})$$

$$+ (D_{11}k^{2} + D_{12}Q^{2}) (\boldsymbol{\sigma}_{1} \cdot \mathbf{k}) (\boldsymbol{\sigma}_{2} \cdot \mathbf{k})$$

$$+ (D_{13}k^{2} + D_{14}Q^{2}) (\boldsymbol{\sigma}_{1} \cdot \mathbf{Q}) (\boldsymbol{\sigma}_{2} \cdot \mathbf{Q})$$

$$+ D_{15} \boldsymbol{\sigma}_{1} \cdot (\mathbf{k} \times \mathbf{Q}) \boldsymbol{\sigma}_{2} \cdot (\mathbf{k} \times \mathbf{Q})$$

$$+ iD_{16} \boldsymbol{k} \cdot \mathbf{Q} \boldsymbol{Q} \times \boldsymbol{P} \cdot (\boldsymbol{\sigma}_{1} - \boldsymbol{\sigma}_{2})$$

$$+ D_{17} \boldsymbol{k} \cdot \boldsymbol{Q} (\boldsymbol{k} \times \boldsymbol{P}) \cdot (\boldsymbol{\sigma}_{1} \times \boldsymbol{\sigma}_{2})$$
(2)

with k = p' - p and $Q = \frac{p' + p}{2}$, p and p' being the initial and final relative momenta, and $P = p_1 + p_2$ the total pair momentum. However, as it was pointed out in Ref. [22], only 12 independent LECs survive on shell and can thus be determined from 2N scattering data. This redundancy amounts to a unitary ambiguity, i.e. to the possibility of generating shifts of the LECs by unitary transforming the one-body kinetic energy

operator H_0 as

$$H_0 \to U^{\dagger} H_0 U.$$
 (3)

Here U is the most general unitary 2-body contact transformation depending on 5 arbitrary parameters α_i ,

$$U = \exp\left[\sum_{i=1}^{5} \alpha_i T_i\right],\tag{4}$$

and the independent generators T_i , which are given explicitly in Ref. [17, 23], induce a shift in the N3LO contact LECs, $D_i \to D_i + \delta D_i$. Specifically, the induced shifts for D_{16} and D_{17} are given by:

$$\delta D_{16} = -\frac{2}{m}\alpha_4,\tag{5}$$

$$\delta D_{17} = -\frac{4}{m}\alpha_3 - \frac{2}{m}\alpha_5. \tag{6}$$

At the same time, the unitary transformation, when applied to the LO 2N contact Hamiltonian, induces a shift of the LECs parametrizing the subleading three-body interaction E_i , as

$$V_{3N,\Lambda}^{(2)} = \sum_{ijk} \left[E_1 + E_2 \boldsymbol{\tau}_i \cdot \boldsymbol{\tau}_j + \left(E_3 + E_4 \boldsymbol{\tau}_i \cdot \boldsymbol{\tau}_j \right) \boldsymbol{\sigma}_i \cdot \boldsymbol{\sigma}_j \right]$$

$$\times \left[Z_{\Lambda}''(r_{ij}) + 2 \frac{Z_{\Lambda}'(r_{ij})}{r_{ij}} \right] Z_{\Lambda}(r_{ik})$$

$$+ \left(E_5 + E_6 \boldsymbol{\tau}_i \cdot \boldsymbol{\tau}_j \right) S_{ij} \left[Z_{\Lambda}''(r_{ij}) - \frac{Z_{\Lambda}'(r_{ij})}{r_{ij}} \right] Z_{\Lambda}(r_{ik})$$

$$+ \left(E_7 + E_8 \boldsymbol{\tau}_i \cdot \boldsymbol{\tau}_k \right) \left\{ \left(\mathbf{L} \cdot \mathbf{S} \right)_{ij}, \frac{Z_{\Lambda}'(r_{ij})}{r_{ij}} Z_{\Lambda}(r_{ik}) \right\}$$

$$+ \left[\left(E_9 + E_{10} \boldsymbol{\tau}_j \cdot \boldsymbol{\tau}_k \right) \boldsymbol{\sigma}_j \cdot \hat{\boldsymbol{r}}_{ij} \boldsymbol{\sigma}_k \cdot \hat{\boldsymbol{r}}_{ik} \right]$$

$$+ \left(E_{11} + E_{12} \boldsymbol{\tau}_j \cdot \boldsymbol{\tau}_k + E_{13} \boldsymbol{\tau}_i \cdot \boldsymbol{\tau}_j \right) \boldsymbol{\sigma}_k \cdot \hat{\boldsymbol{r}}_{ij} \boldsymbol{\sigma}_j \cdot \hat{\boldsymbol{r}}_{ik} \right]$$

$$\times Z_{\Lambda}'(r_{ij}) Z_{\Lambda}'(r_{ik}),$$

$$(7)$$

where S_{ij} and $(\mathbf{L} \cdot \mathbf{S})_{ij}$ are respectively the tensor and spin-orbit operators for particles i and j, and the profile functions

$$Z_{\Lambda}(r) = \int \frac{d\mathbf{p}}{(2\pi)^3} e^{i\mathbf{p}\cdot\mathbf{r}} F(\mathbf{p}^2; \Lambda), \tag{8}$$

representing the cutoff in coordinate space, chosen as

$$F(\mathbf{p}^2, \Lambda) = \exp\left[-\left(\frac{\mathbf{p}^2}{\Lambda^2}\right)^2\right]. \tag{9}$$

In the following the value $\Lambda = 500\,\mathrm{MeV}$ will be used. The explicit expression for the N4LO LECs shift δE_i of the three-body force can be found in Ref. [17] (see Eqs. (39)-(51)). The induced contributions δE_i are enhanced as compared to the genuine ones E_i , due to the presence of the nucleon mass factor, scaling as $m \sim O(\Lambda_\chi^2/p)$ (being Λ_χ hard or breakdown scale of the theory) in the Weinberg counting [2], which effectively promotes them to N3LO. In this work, the LECs E_i will be thought of as constituted only of the induced contributions of the P-dependent D_{16} and D_{17} . Thus, at N3LO the 3N contact interaction depends on two combinations of the 2N LECs D_i , appearing in Eqs. (5)-(6), which cannot be determined from 2N scattering data, but have to be fitted to experimental observables in A > 2 systems.

We investigate the sensitivity of polarization observables in low-energy N-d scattering to the two P-dependent N3LO LECs. The AV18 potential is used as a representative 2N interaction, and the meaning of the LECs C_S and C_T in this framework is found treating the LO contact pionless potential as a very low-energy representation of the AV18 potential, with a local cutoff introduced. The values of C_S and C_T are thus taken from a fit of the LO 2N contact interaction

$$V_{2N,\Lambda}^{(0)} = \left[C_S + C_T \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2\right] Z_{\Lambda}(r) \tag{10}$$

to the singlet and triplet n-p scattering lengths as predicted by the AV18 potential. In the above expression a local cutoff has been introduced as in Eq. (8). From this procedure we get

$$C_S = -66.53 \text{ GeV}^{-2}, \quad C_T = -3.47 \text{ GeV}^{-2}.$$
 (11)

The 3-body Schrödinger equation is solved as in Ref.[17] employing the Hyperspherical Harmonic (HH) method. Below the deuteron breakup threshold, the N-dscattering wave function is expressed as the sum of an internal Ψ_C and an asymptotic part Ψ_A

$$\Psi_{LSJJ_z} = \Psi_C + \Psi_A \,, \tag{12}$$

where the internal part Ψ_C is expanded on the HH basis as

$$\Psi_C = \sum_{\mu} c_{\mu} \Phi_{\mu}. \tag{13}$$

Here μ denotes all the quantum numbers required to define the basis element.

The asymptotic part describes the relative motion between the nucleon and the deuteron at large distances, involving regular (R) and irregular (I) solutions.

Denoting these solution $\Omega_L^{\lambda}SJJz$, with $\lambda=R,I$ respectively, and defining

$$\Omega_{LSJJ_z}^{\pm} = i\Omega_{LSJJ_z}^R \pm \Omega_{LSJJ_z}^I, \tag{14}$$

we have

$$\Psi_A = \Omega_{LSJJ_z}^- + \sum_{L'S'} \mathcal{S}_{LS,L'S'}^J(q) \Omega_{L'S'JJJ_z}^+.$$
 (15)

Here $S_{LS,L'S'}^J$ are the S-matrix elements and q is defined as the modulus of the N-d relative momentum. From the S-matrix it is possible to compute phase shifts and mixing angles, from which the scattering observables are calculated. The S-matrix in Eq. (15) and the coefficients c_{μ} in Eq. (13) are determined by the complex formulation of the Kohn variational principle [24]. This principle demands that a certain functional be stationary under variations of trial parameters, leading to a linear system whose solution provides the weights and coefficients.

The Hamiltonian is decomposed into H_{2N} (kinetic energy plus AV18 2N interaction with Coulomb potential) and $V_{3N,\Lambda}^{(2)}$ (containing 3N interaction induced by D_{16} and D_{17} terms). The linear system for coefficients involves the computation of matrix elements between HH basis elements and asymptotic functions.

A specific set of LECs allows the computation of the associated S-matrix for each J^{π} state using the Kohn variational principle, providing observables at a specific energy.

4 Fit results

The observables used in the fitting procedure include the p-d differential cross section, the two vector analyzing powers A_y and iT_{11} , the three tensor analyzing powers T_{20}, T_{21}, T_{22} and the doublet and quartet n-d scattering lengths, with the experimental values $^2a_{nd}=(0.645\pm0.003\pm0.007)$ fm [25] and $^4a_{nd}=(6.35\pm0.02)$ fm [26]. In particular we fit the experimental doublet and quartet n-d scattering lengths and the six p-d scattering observables at center-of-mass energy $E_{\rm cm}=2\,{\rm MeV}$ [27], amounting to 282 experimental data. In addition we also fix the $^3{\rm H}$ binding energy to 8.469 MeV. This value takes into account the contribution of the neutron-proton mass difference, which is not described in the HH method, amounting to $\sim 7\,{\rm keV}$, and additional amount of $\sim 6\,{\rm keV}$ from the truncation of the HH expansion.

In the case of the differential cross section, we introduce an overall normalization factor Z in the definition of χ^2 . Specifically,

$$\chi^2 = \sum_i \frac{\left(d_i^{\text{exp}}/Z - d_i^{\text{th}}\right)^2}{\left(\sigma_i^{\text{exp}}/Z\right)^2},\tag{16}$$

where Z is determined from the minimization condition:

$$Z = \frac{\sum_{i} d_{i}^{\text{exp}} d_{i}^{\text{th}} / \left(\sigma_{i}^{\text{exp}}\right)^{2}}{\sum_{i} \left(d_{i}^{\text{th}}\right)^{2} / \left(\sigma_{i}^{\text{exp}}\right)^{2}}.$$
(17)

Here, $d_i^{\exp/\text{th}}$ represents the experimental data points and their theoretical predictions, respectively, while σ_i^{\exp} is the experimental error. For other observables, we treat the normalization $Z=1.00\pm0.01$ as an additional experimental datum, considering the systematic uncertainty estimated as 1% according to Ref. [27].

The fitting procedure involves a global 2-parameter fit including only the P-dependent 2N interaction. We also perform 3-parameter fits including the LO 3N

Fitting procedure	2-param.	3-param.
$\chi^2/\mathrm{d.o.f.}$	2.1	1.9
e_0	-	0.459
$\tilde{lpha}_4 C_S$	1.751	1.894
$ ilde{lpha}_5 C_S$	-0.495	-1.175
$^2a_{nd}$ (fm)	0.573	0.599

Table 2 Results of the 2-parameters and 3-parameters fits, the latter one obtained including the leading order 3N contact interaction. Here the fitted parameters α_i and the corresponding values of the LO 3N contact LEC E_0 are dimensionless, i.e. $e_0 = E_0 F_\pi^4 \Lambda$, $\tilde{\alpha}_i = \alpha_i F_\pi^4 \Lambda^3$ with $F_\pi = 92.4$ MeV. In the last row we report the value obtained for the n-d doublet scattering length $^2a_{nd}$, which should be compared with the experimental value $^2a_{nd} = (0.645 \pm 0.003 \pm 0.007)$ fm [25].

contact interaction.

$$V_{3N,\Lambda}^{(0)} = E_0 \sum_{ijk} Z_{\Lambda}(r_{ij}) Z_{\Lambda}(r_{ik}), \tag{18}$$

where the LEC E_0 introduces further flexibility to the fit and it is mainly determined by the ${}^{3}\mathrm{H}$ binding energy.

We start the iterative minimization procedure by solving the scattering problem for an initial random set of α_4 and α_5 parameters, calculating the corresponding observables. Employing the POUNDerS algorithm [28], we repeat the process with different initial random α_i values, aiming to converge to the deepest minimum. The resulting $\chi^2/\text{d.o.f.}$ is $\sim 2.1(1.9)$ for the two(three)-parameter fits. The fitted parameters E_0 , α_4 and α_5 can be read from the first column of Table 2.

Figure 1 shows the best fit curves for various analyzing powers and observables, compared to predictions from the 2N AV18 potential and the addition of the 3N Urbana IX interaction. The effective N3LO 3N contact interaction induced by the D_{16} and D_{17} terms successfully addresses the A_y problem, and the description of the vector analyzing power iT_{11} is significantly improved. We also conclude by saying that the introduction into the fit of the LEC E_0 corresponding to the three-body contact force at leading order does not substantially change the description of the experimental data, except for the observable A_y , as can be seen in Figure 1.

5 Conclusions

In this analysis, we have explored the relativistic constraints on the $\mathcal{O}(p^4)$ 2N contact Lagrangian bringing out two P-dependent terms in the potential accompanied by two unconstrained LECs D_{16} and D_{17} . It should be emphasized that the above terms are not to be understood as relativistic corrections but for all intents are within the N3LO 2N contact potential. These LECs, whose effect vanishes in the 2N center-of-mass frame, can play a crucial role in larger nuclear systems. The unitary equivalence to 3N contact operators implies a connection between these LECs and off-shell effects.




Fig. 1 Proton and deuteron analyzing power and differential cross-section at $E_{\rm cm}=2$ MeV. The red lines result from a global 2-parameter fit, the black lines represent the 3-parameters fit including the E_0 term, the blue lines are the predictions from the 2N AV18 potential, while the green lines are the predictions including also the 3N Urbana IX interaction. Experimental data are from Ref. [27].

Using an hybrid approach where the three-nucleon interaction, parametrized by D_{16} and D_{17} LECs, is considered alongside the phenomenological AV18 2N potential, we fit experimental data on polarization observables in low-energy p-d scattering, specifically focusing on the A_y asymmetry. The results indicate that the inclusion of terms represented by D_{16} and D_{17} in the three-nucleon interaction is crucial for accurately reproducing experimental data, in low-energy proton-deuteron scattering. This implies that these LECs can be resolutive for the long-standing A_y puzzle. The D_{16} and D_{17} LECs on systems with A>2 should be further quantified in future investigations. Of course it will be undoubtedly intriguing to conduct a thorough reexamination of the aforementioned analysis within a fully consistent framework of ChEFT.

References

[1] S. Weinberg, Nuclear forces from chiral Lagrangians. Phys. Lett. B **251**, 288–292 (1990). https://doi.org/10.1016/0370-2693(90)90938-3

- [2] S. Weinberg, Effective chiral Lagrangians for nucleon pion interactions and nuclear forces. Nucl. Phys. B 363, 3–18 (1991). https://doi.org/10.1016/0550-3213(91)90231-L
- [3] S. Weinberg, Three body interactions among nucleons and pions. Phys. Lett. B **295**, 114–121 (1992). https://doi.org/10.1016/0370-2693(92)90099-P. arXiv:hep-ph/9209257
- P.F. Bedaque, U. van Kolck, Effective field theory for few nucleon systems. Ann. Rev. Nucl. Part. Sci. 52, 339–396 (2002). https://doi.org/10.1146/annurev.nucl. 52.050102.090637. arXiv:nucl-th/0203055
- [5] E. Epelbaum, Few-nucleon forces and systems in chiral effective field theory. Prog. Part. Nucl. Phys. 57, 654-741 (2006). https://doi.org/10.1016/j.ppnp.2005.09. 002. arXiv:nucl-th/0509032
- [6] E. Epelbaum, H.W. Hammer, U.G. Meissner, Modern Theory of Nuclear Forces. Rev. Mod. Phys. 81, 1773–1825 (2009). https://doi.org/10.1103/RevModPhys. 81.1773. arXiv:0811.1338 [nucl-th]
- [7] R. Machleidt, D.R. Entem, Chiral effective field theory and nuclear forces.
 Phys. Rept. 503, 1–75 (2011). https://doi.org/10.1016/j.physrep.2011.02.001.
 arXiv:1105.2919 [nucl-th]
- [8] R.J. Furnstahl, D.R. Phillips, S. Wesolowski, A recipe for EFT uncertainty quantification in nuclear physics. J. Phys. G 42(3), 034028 (2015). https://doi.org/10.1088/0954-3899/42/3/034028. arXiv:1407.0657 [nucl-th]
- [9] E. Epelbaum, H. Krebs, U.G. Meißner, Improved chiral nucleon-nucleon potential up to next-to-next-to-leading order. The European Physical Journal A 51(5) (2015). https://doi.org/10.1140/epja/i2015-15053-8. URL https://doi.org/ 10.1140%2Fepja%2Fi2015-15053-8
- [10] R.J. Furnstahl, N. Klco, D.R. Phillips, S. Wesolowski, Quantifying truncation errors in effective field theory. Phys. Rev. C 92(2), 024005 (2015). https://doi. org/10.1103/PhysRevC.92.024005. arXiv:1506.01343 [nucl-th]
- [11] S. König, A. Ekström, K. Hebeler, D. Lee, A. Schwenk, Eigenvector Continuation as an Efficient and Accurate Emulator for Uncertainty Quantification. Phys. Lett. B 810, 135814 (2020). https://doi.org/10.1016/j.physletb.2020.135814. arXiv:1909.08446 [nucl-th]
- [12] L.L. Foldy, Relativistic particle systems with interactions. Phys. Rev. **122**, 275–288 (1961). https://doi.org/10.1103/PhysRev.122.275
- [13] R.A. Krajcik, L.L. Foldy, Relativistic center-of-mass variables for composite systems with arbitrary internal interactions. Phys. Rev. D **10**, 1777–1795 (1974).

- [14] Y. Xiao, L.S. Geng, X.L. Ren, Covariant chiral nucleon-nucleon contact Lagrangian up to order $\mathcal{O}(q^4)$. Phys. Rev. C **99**(2), 024004 (2019). https://doi.org/10.1103/PhysRevC.99.024004. arXiv:1812.03005 [nucl-th]
- [15] E. Filandri, L. Girlanda, Momentum dependent nucleon-nucleon contact interaction from a relativistic lagrangian. Phys. Lett. B 841, 137957 (2023). https://doi.org/10.1016/j.physletb.2023.137957. URL https://doi.org/10.1016 %2Fj.physletb.2023.137957
- [16] G.J. et al., Low-energy neutron-deuteron reactions with n 3 lo chiral forces. The European Physical Journal A **50**(11) (2014). https://doi.org/10.1140/epja/i2014-14177-7. URL https://doi.org/10.1140/epja/i2014-14177-7
- [17] L. Girlanda, E. Filandri, A. Kievsky, L.E. Marcucci, M. Viviani, Effect of the n3lo three-nucleon contact interaction on p-d scattering observables. Phys. Rev. C 107, L061001 (2023). https://doi.org/10.1103/PhysRevC.107.L061001. URL https://link.aps.org/doi/10.1103/PhysRevC.107.L061001
- [18] L. Girlanda, M. Viviani, Relativistic Covariance of the 2-Nucleon Contact Interactions. Few Body Syst. 49, 51–60 (2011). https://doi.org/10.1007/s00601-010-0185-6
- [19] S. Petschauer, N. Kaiser, Relativistic SU(3) chiral baryon-baryon Lagrangian up to order q^2 . Nucl. Phys. A **916**, 1–29 (2013). https://doi.org/10.1016/j.nuclphysa. 2013.07.010. arXiv:1305.3427 [nucl-th]
- [20] D. Entem, R. Machleidt, Accurate nucleon–nucleon potential based upon chiral perturbation theory. Phys. Lett. B 524(1-2), 93–98 (2002). https://doi.org/10.1016/s0370-2693(01)01363-6. URL https://doi.org/10.1016%2Fs0370-2693 %2801%2901363-6
- [21] D.R. Entem, R. Machleidt, Accurate charge-dependent nucleon-nucleon potential at fourth order of chiral perturbation theory. Phys. Rev. C **68**, 041001 (2003). https://doi.org/10.1103/PhysRevC.68.041001. URL https://link.aps.org/doi/10.1103/PhysRevC.68.041001
- [22] P. Reinert, H. Krebs, E. Epelbaum, Semilocal momentum-space regularized chiral two-nucleon potentials up to fifth order. The European Physical Journal A 54, 1–49 (2017)
- [23] L. Girlanda, A. Kievsky, L.E. Marcucci, M. Viviani, Unitary ambiguity of nn contact interactions and the 3n force. Physical Review C $\bf 102(6)$ (2020). https://doi.org/10.1103/physrevc.102.064003. URL https://doi.org/10.1103%2Fphysrevc. 102.064003

- [24] A. Kievsky, The complex kohn variational method applied to n-d scattering. Nuclear Physics A **624**(2), 125–139 (1997). https://doi.org/https://doi.org/10.1016/S0375-9474(97)81832-5. URL https://www.sciencedirect.com/science/article/pii/S0375947497818325
- [25] W. Dilg, L. Koester, W. Nistler, The neutron-deuteron scattering lengths. Phys. Lett. B **36**(3), 208–210 (1971). https://doi.org/https://doi.org/10.1016/0370-2693(71)90070-0. URL https://www.sciencedirect.com/science/article/pii/0370269371900700
- [26] K. Schoen, D.L. Jacobson, M. Arif, P.R. Huffman, T.C. Black, W.M. Snow, S.K. Lamoreaux, H. Kaiser, S.A. Werner, Precision neutron interferometric measurements and updated evaluations of the n-p and n-d coherent neutron scattering lengths. Phys. Rev. C **67**, 044005 (2003). https://doi.org/10.1103/PhysRevC.67. 044005. URL https://link.aps.org/doi/10.1103/PhysRevC.67.044005
- [27] S. Shimizu, K. Sagara, H. Nakamura, K. Maeda, T. Miwa, N. Nishimori, S. Ueno, T. Nakashima, S. Morinobu, Analyzing powers of p+d scattering below the deuteron breakup threshold. Phys. Rev. C 52, 1193–1202 (1995). https://doi.org/10.1103/PhysRevC.52.1193. URL https://link.aps.org/doi/10.1103/PhysRevC.52.1193
- [28] T. Munson, J. Sarich, S. Wild, S. Benson, L. McInnes, Tao 2.0 users manual technical report anl/mcs-tm-322 URL http://www.mcs.anl.gov/tao

