SUP-NORM BOUNDS FOR JACOBI CUSP FORMS

ANILATMAJA ARYASOMAYAJULA, JÜRG KRAMER, AND ANNA-MARIA VON PIPPICH

ABSTRACT. In this article, we give bounds for the natural invariant norm of cusp forms of real weight k and character χ for any cofinite Fuchsian subgroup $\Gamma \subset \mathrm{SL}_2(\mathbb{R})$. Using the representation of Jacobi cusp forms of integral weight k and index m for the modular group $\Gamma_0 = \mathrm{SL}_2(\mathbb{Z})$ as linear combinations of modular forms of weight $k - \frac{1}{2}$ for some congruence subgroup of Γ_0 (depending on m) and suitable Jacobi theta functions, we derive bounds for the natural invariant norm of these Jacobi cusp forms. More specifically, letting $J_{k,m}^{\mathrm{cusp}}(\Gamma_0)$ denote the complex vector space of Jacobi cusp forms under consideration and $\|\cdot\|_{\mathrm{Pet}}$ the pointwise Petersson norm on $J_{k,m}^{\mathrm{cusp}}(\Gamma_0)$, we prove that for given $\epsilon > 0$, the bound

$$\sup_{\tau,z)\in\mathbb{H}\times\mathbb{C}} \|f(\tau,z)\|_{\mathrm{Pet}} = O_{\epsilon}\left(k^{\frac{3}{4}}m^{\frac{3}{2}+\epsilon}\right)$$

holds for any $f \in J_{k,m}^{\text{cusp}}(\Gamma_0)$, which is normalized with respect to the Petersson inner product, where the implied constant depends only on the choice of $\epsilon > 0$.

1. INTRODUCTION

1.1. **Background.** In general, bounds for automorphic forms and for their Fourier coefficients represent an area of great interest in number theory. More specifically, we mention in this respect the results of [FJK16], where J. Friedman, J. Jorgenson, and J. Kramer obtained optimal sup-norm bounds on average for cusp forms of even weight k for any cofinite Fuchsian subgroup $\Gamma \subset SL_2(\mathbb{R})$. These bounds turn out to be uniform with respect to the subgroup Γ . Moreover, in [FJK19], effective versions for these sup-norm bounds are given. With regard to sup-norm bounds for individual Hecke eigenforms of large level, we mention, for example, the results by V. Blomer and R. Holowinsky in [BH11].

So far, less attention has been devoted to the study of sup-norm bounds for Jacobi cusp forms. The first comprehensive study of Jacobi forms was undertaken by M. Eichler and D. Zagier in [EZ85]. Subsequently, various authors have built on their work. In contrast to their analytical approach, a geometrical approach to the theory of Jacobi forms was given by J. Kramer in [Kr91].

Let k, m be positive integers. A Jacobi form of weight k and index m for the modular group $\Gamma_0 := \operatorname{SL}_2(\mathbb{Z})$ is a holomorphic function on the product $\mathbb{H} \times \mathbb{C}$ of the upper half-plane \mathbb{H} with the complex plane \mathbb{C} having a suitable transformation behaviour with respect to Γ_0 and vanishing "at infinity". We denote the complex vector space of Jacobi cusp forms of weight k and index m for Γ_0 by $J_{k,m}^{\operatorname{cusp}}(\Gamma_0)$. The pointwise Petersson norm of a Jacobi form $f \in J_{k,m}^{\operatorname{cusp}}(\Gamma_0)$ is then defined by

$$\|f(\tau,z)\|_{\operatorname{Pet}}^2 \coloneqq |f(\tau,z)|^2 \operatorname{Im}(\tau)^k e^{-\frac{4\pi m \operatorname{Im}(z)^2}{\operatorname{Im}(\tau)}} \qquad (\tau \in \mathbb{H}, \ z \in \mathbb{C}).$$

Let F be a Siegel cusp form of weight k for the Siegel modular group $\text{Sp}_4(\mathbb{Z})$, and let $\{f_m\}_{m\geq 1}$ be the set of Jacobi forms appearing in the Fourier–Jacobi expansion of F, i.e.,

Date: April 23, 2024.

²⁰¹⁰ Mathematics Subject Classification. 11F11, 11F12.

 $f_m \in J_{k,m}^{\text{cusp}}(\Gamma_0)$. Then, for any $\epsilon > 0$, W. Kohnen proved the following sup-norm bound for the pointwise Petersson norm of f_m in [Ko93]

(1)
$$\sup_{(\tau,z)\in\mathbb{H}\times\mathbb{C}} \|f_m(\tau,z)\|_{\mathrm{Pet}} = O_{F,\epsilon}\left(m^{\frac{k}{2}-\frac{2}{9}+\epsilon}\right),$$

where the implied constant depends on the Siegel cusp form F and on the choice of $\epsilon > 0$. Motivated by the Ramanujan–Petersson conjecture, W. Kohnen then conjectured the bound

$$\sup_{(\tau,z)\in\mathbb{H}\times\mathbb{C}}\|f_m(\tau,z)\|_{\mathrm{Pet}}=O_{F,\epsilon}\left(m^{\frac{k-1}{2}+\epsilon}\right),$$

where the implied constant depends on the Siegel cusp form F and on the choice of $\epsilon > 0$.

More recently, P. Anamby and S. Das established in [AD23] a general sup-norm bound for the pointwise Petersson norm of $f \in J_{k,m}^{\text{cusp}}(\Gamma_0)$, which is normalized with respect to the Petersson inner product, i.e., for which we have

$$\int_{\Gamma_0 \ltimes \mathbb{Z}^2 \setminus \mathbb{H} \times \mathbb{C}} \|f(\tau, z)\|_{\text{Pet}}^2 \frac{\mathrm{d}\xi \wedge \mathrm{d}\eta \wedge \mathrm{d}x \wedge \mathrm{d}y}{\eta^3} = 1 \qquad (\tau = \xi + i\eta, \, z = x + iy).$$

Their bound is (see Theorem 1.4 in [AD23])

(2)
$$\sup_{(\tau,z)\in\mathbb{H}\times\mathbb{C}} \|f(\tau,z)\|_{\mathrm{Pet}} = O_{\epsilon}(k\,m),$$

where the implied constant depends on the choice of $\epsilon > 0$.

1.2. Main results. The goal of this article is to provide new sup-norm bounds for the pointwise Petersson norm for Jacobi forms of integral weight k and integral index m for Γ_0 , which are normalized with respect to the Petersson inner product. The main result in this respect is given in Theorem 4.4 and states for $f \in J_{k,m}^{\text{cusp}}(\Gamma_0)$, which is normalized with respect to the Petersson inner product, that

(3)
$$\sup_{(\tau,z)\in\mathbb{H}\times\mathbb{C}} \|f(\tau,z)\|_{\mathrm{Pet}} = O_{\epsilon} \left(k^{\frac{3}{4}} m^{\frac{3}{2}+\epsilon}\right),$$

where the implied constant depends only on the choice of $\epsilon > 0$. For the proof, we essentially use the representation of the Jacobi cusp forms under consideration as linear combinations of modular forms of weight $k - \frac{1}{2}$ for some congruence subgroup of Γ_0 (depending on m) and suitable Jacobi theta functions; we then need to derive bounds for the pointwise Petersson norms of these functions to arrive at our result. Comparing our bound with the bound (2) by P. Anamby and S. Das, we realize an improvement with regard to the polynomial growth in k, while there is a price to be paid with regard to the polynomial growth in m.

In order to be able to derive our bound (3), we need sup-norm bounds for the pointwise Petersson norm of cusp forms of positive real weight k and character χ for any cofinite Fuchsian subgroup Γ of $SL_2(\mathbb{R})$. While such bounds could be derived from [FJK16] with some extra work, due to the lack of a precise reference, we provide new, alternative proofs for the results of [FJK16] applying to any positive real weight k and any character χ by using the Bergman kernel for the modular curve associated to Γ .

More specifically, given $\Gamma \subset SL_2(\mathbb{R})$ a Fuchsian subgroup, $k \in \mathbb{R}_{>0}$, and $\chi \colon \Gamma \to \mathbb{C}^{\times}$ a character, we let $S_{k,\chi}(\Gamma)$ denote the space of cusp forms of weight k and character χ for Γ . Denoting by d_k the dimension of $S_{k,\chi}(\Gamma)$ and letting $\{f_1, \ldots, f_{d_k}\}$ be an orthonormal basis

of $S_{k,\chi}(\Gamma)$ with respect to the Petersson inner product, the Bergman kernel associated to $S_{k,\chi}(\Gamma)$ is then defined by

$$B_{k,\chi}(\tau,\tau') := \sum_{j=1}^{d_k} f_j(\tau) \overline{f_j(\tau')};$$

it is straightforward that this definition does not depend on the choice of an orthonormal basis of $S_{k,\chi}(\Gamma)$. The pointwise Petersson norm of the Bergman kernel is defined by

$$||B_{k,\chi}(\tau,\tau')||_{\operatorname{Pet}} = |B_{k,\chi}(\tau,\tau')| \left(\operatorname{Im}(\tau)\operatorname{Im}(\tau')\right)^{\frac{\kappa}{2}},$$

which gives on the diagonal

(4)
$$||B_{k,\chi}(\tau,\tau)||_{\text{Pet}} = \sum_{j=1}^{d_k} ||f_j(\tau)||_{\text{Pet}}^2$$

As a second main result of this article, we establish in Theorem 3.3, assuming that $k \in \mathbb{R}_{\geq 5}$, for Γ being cocompact without elliptic elements the bound

$$\sup_{z \in \mathbb{H}} \|B_{k,\chi}(\tau,\tau)\|_{\text{Pet}} = O_{\Gamma}(k);$$

moreover, for Γ being cofinite, we give the bound

$$\sup_{z \in \mathbb{H}} \|B_{k,\chi}(\tau,\tau)\|_{\mathrm{Pet}} = O_{\Gamma}\left(k^{\frac{3}{2}}\right),$$

where the implied constants depend only on the Fuchsian subgroup Γ . Due to the relation (4), these results reprove the sup-norm bounds on average obtained in [FJK16], but now for any real weight $k \in \mathbb{R}_{\geq 5}$ and any character χ . Based on these results, we are then also able to prove the uniformity of the above bounds with respect to the subgroup Γ in Theorem 3.5.

1.3. **Outline.** Let us briefly outline the contents of this article. In the subsequent, second section we collect all the necessary prerequisites for the sequel of the paper. In particular, we introduce the definitions of cusp forms and Jacobi cusp forms together with their respective (pointwise) Petersson inner products. Furthermore, we define the Bergman kernel for modular curves and state its basic properties. We close the section by recalling asymptotics of Bergman kernels associated to tensor powers of holomorphic line bundles on compact complex Kähler manifolds due to [Bo96], which are crucial in the derivation of the bound (3).

The third section is devoted to the revisiting of the sup-norm bounds on average obtained in [FJK16], but now for any real weight $k \in \mathbb{R}_{\geq 5}$ and any character χ . Here, the proofs of Theorem 3.3 and Theorem 3.5 are provided.

In the fourth section, the bound (3) is proven in Theorem 4.4. In addition to some straightforward bounds established for the relative L^2 -norm of classical Jacobi theta functions, the above mentioned asymptotics of Bergman kernels associated to tensor powers of holomorphic line bundles corresponding to these Jacobi theta functions are crucial in the derivation of the proof of Theorem 4.4.

2. Preliminaries

2.1. Hyperbolic metric. Let $\mathbb{H} := \{\tau \in \mathbb{C} \mid \tau = \xi + i\eta, \eta > 0\}$ be the upper half-plane. We denote by $ds_{hyp}^2(\tau)$ the line element and by $\mu_{hyp}(\tau)$ the volume form corresponding to the hyperbolic metric on \mathbb{H} , which is compatible with the complex structure of \mathbb{H} and has constant curvature equal to -1. Locally on \mathbb{H} , we have

$$\mathrm{d}s_{\mathrm{hyp}}^2(\tau) = \frac{\mathrm{d}\xi^2 + \mathrm{d}\eta^2}{\eta^2}$$
 and $\mu_{\mathrm{hyp}}(\tau) = \frac{\mathrm{d}\xi \wedge \mathrm{d}\eta}{\eta^2}$

For $\tau, \tau' \in \mathbb{H}$, we let $\operatorname{dist}_{\operatorname{hyp}}(\tau, \tau')$ denote the hyperbolic distance between these two points. For later purposes, it is useful to introduce the displacement function

(5)
$$\sigma(\tau,\tau') \coloneqq \cosh^2\left(\frac{\operatorname{dist}_{\operatorname{hyp}}(\tau,\tau')}{2}\right) = \frac{|\tau-\overline{\tau}'|^2}{4\operatorname{Im}(\tau)\operatorname{Im}(\tau')}.$$

2.2. Quotient space. Let $\Gamma \subset SL_2(\mathbb{R})$ be a Fuchsian subgroup acting by fractional linear transformations on \mathbb{H} . Let X_{Γ} be the quotient space $\Gamma \setminus \mathbb{H}$ and g_{Γ} the genus of X_{Γ} . In the sequel, we identify X_{Γ} with a fundamental domain $\mathcal{F}_{\Gamma} \subset \mathbb{H}$ for the group Γ , which we assume to be closed and connected.

Denote by

$$\mathcal{P}_{\Gamma} = \{p_1, \ldots, p_s\}$$

the set of cusps of \mathcal{F}_{Γ} . Let $\sigma_{\mathcal{P},j} \in \mathrm{SL}_2(\mathbb{R})$ be a scaling matrix of the cusp p_j , that is, $p_j = \sigma_{\mathcal{P},j} i \infty$ with stabilizer subgroup Γ_{p_j} described as

(6)
$$\sigma_{\mathcal{P},j}^{-1}\Gamma_{p_j}\sigma_{\mathcal{P},j} = \begin{cases} \left\langle \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \right\rangle, & \text{if } -\text{id} \notin \Gamma, \\ \left\langle \pm \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \right\rangle, & \text{if } -\text{id} \in \Gamma, \end{cases} \quad (j = 1, \dots, s).$$

For Y > 0, we let $\mathcal{F}_j^Y \subset \mathcal{F}_{\Gamma}$ denote the neighborhood of the cusp p_j characterized by

$$\sigma_{\mathcal{P},j}^{-1}\mathcal{F}_{j}^{Y} = \{\tau = \xi + i\eta \in \mathbb{H} \mid -1/2 \le \xi \le 1/2, \eta \ge Y\} \qquad (j = 1, \dots, s).$$

With these notations, we define \mathcal{F}_Y to be the closure of the complement of the union $\mathcal{F}_1^Y \cup \ldots \cup \mathcal{F}_s^Y$ in \mathcal{F}_{Γ} , i.e.,

(7)
$$\mathcal{F}_Y \coloneqq \operatorname{cl}\bigl(\mathcal{F}_\Gamma \setminus \bigl(\mathcal{F}_1^Y \cup \ldots \cup \mathcal{F}_s^Y\bigr)\bigr)$$

which is compact; we note that $\mathcal{F}_Y = \mathcal{F}_{\Gamma}$, if Γ is cocompact. We choose $0 < m_Y < M_Y$ such that for all $\tau \in \mathcal{F}_Y$ the inequalities

$$m_Y \leq \operatorname{Im}(\sigma_{\mathcal{P},j}^{-1}\tau) \leq M_Y$$

hold for all j = 1, ..., s; we note that m_Y and M_Y depend on the choice of Y.

Denote by

$$\mathcal{E}_{\Gamma} = \{e_1, \ldots, e_t\}$$

the set of elliptic fixed points of \mathcal{F}_{Γ} . Let Γ_{e_j} and m_j denote the stabilizer subgroup and order of the elliptic fixed point e_j , respectively.

We denote the hyperbolic length of the shortest closed geodesic on X_{Γ} by ℓ_{Γ} . For a domain $D \subset \mathbb{H}$, we denote its hyperbolic diameter by $\operatorname{diam}_{\operatorname{hyp}}(D)$ and its hyperbolic volume by $\operatorname{vol}_{\operatorname{hyp}}(D)$. Finally, the injectivity radius r_{Γ} is defined by

(8)
$$r_{\Gamma} \coloneqq \inf \left\{ \operatorname{dist}_{\operatorname{hyp}}(\tau, \gamma \tau) \, \middle| \, \tau \in \mathcal{F}_{\Gamma}, \, \gamma \in \Gamma \setminus \left(\bigcup_{j=1}^{s} \Gamma_{p_{j}} \cup \bigcup_{j=1}^{t} \Gamma_{e_{j}} \right) \right\}.$$

We note that if X_{Γ} is compact without elliptic fixed points, i.e., $\mathcal{P}_{\Gamma} = \mathcal{E}_{\Gamma} = \emptyset$, then the injectivity radius r_{Γ} equals the length of the shortest closed geodesic ℓ_{Γ} of X_{Γ} .

2.3. Cusp forms and Bergman kernel. For $k \in \mathbb{R}_{>0}$ and a character $\chi: \Gamma \to \mathbb{C}^{\times}$, we let $S_{k,\chi}(\Gamma)$ denote the space of cusp forms of weight k and character χ for Γ , i. e., the space of holomorphic functions $f: \mathbb{H} \to \mathbb{C}$, which have the transformation behavior

$$f(\gamma z)(cz+d)^{-k} = \chi(\gamma)f(z)$$

for all $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$, and which vanish at all the cusps of \mathcal{F}_{Γ} . Given $f \in S_{k,\chi}(\Gamma)$, we define

$$\|f(\tau)\|_{\text{Pet}}^2 \coloneqq |f(\tau)|^2 \eta^k \qquad (\tau = \xi + i\eta),$$

which defines a Γ -invariant function on \mathbb{H} called the pointwise Petersson norm of f.

The space $S_{k,\chi}(\Gamma)$ is equipped with the Petersson inner product

(9)
$$\langle f_1, f_2 \rangle_{\text{Pet}} \coloneqq \int_{\mathcal{F}_{\Gamma}} f_1(\tau) \overline{f_2(\tau)} \eta^k \mu_{\text{hyp}}(\tau) \qquad (f_1, f_2 \in S_{k,\chi}(\Gamma)).$$

Let d_k denote the dimension of $S_{k,\chi}(\Gamma)$ and let $\{f_1, \ldots, f_{d_k}\}$ be an orthonormal basis of $S_{k,\chi}(\Gamma)$ with respect to the Petersson inner product. Then, the Bergman kernel associated to $S_{k,\chi}(\Gamma)$ is defined by

$$B_{k,\chi}(\tau,\tau') \coloneqq \sum_{j=1}^{d_k} f_j(\tau) \overline{f_j(\tau')}$$

It is obvious that this definition does not depend on the choice of an orthonormal basis of $S_{k,\chi}(\Gamma)$.

The Bergman kernel $B_{k,\chi}(\tau, \tau')$ is a holomorphic cusp form of weight k and character χ for Γ in the τ -variable, and an anti-holomorphic cusp form of weight k and character $\overline{\chi}$ for Γ in the τ' -variable. Hence, the pointwise Petersson norm of the Bergman kernel is given by

$$||B_{k,\chi}(\tau,\tau')||_{\operatorname{Pet}} = |B_{k,\chi}(\tau,\tau')| \, (\eta\eta')^{\frac{\kappa}{2}},$$

which is a Γ -invariant function on $\mathbb{H} \times \mathbb{H}$ with respect to both variables.

Moreover, $B_{k,\chi}(\tau,\tau')$ is the reproducing kernel for $S_{k,\chi}(\Gamma)$, i.e., we have

$$\int_{\mathcal{F}_{\Gamma}} B_{k,\chi}(\tau,\tau') f(\tau') \eta'^k \mu_{\text{hyp}}(\tau') = f(\tau) \qquad (\tau' = \xi' + i\eta')$$

for any $f \in S_{k,\chi}(\Gamma)$. Therefore, for $k \in \mathbb{R}_{>3}$, the Bergman kernel $B_{k,\chi}(\tau, \tau')$ can also be represented in the following form (see Proposition 1.3 on p. 77 in [Fr90])

(10)
$$B_{k,\chi}(\tau,\tau') = \frac{(2i)^k(k-1)}{4\pi} \sum_{\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma} \frac{1}{(\tau - \gamma \overline{\tau'})^k} \frac{1}{\chi(\gamma)(c\overline{\tau'} + d)^k}$$

Note that the formula for the Bergman kernel given in [Fr90] is missing a factor of $(2i)^k$.

2.4. Counting function. Given $\tau \in \mathbb{H}$ and $\rho \in \mathbb{R}_{\geq 0}$, we recall from [JL95] the counting function

$$N_{\Gamma}(\tau;\rho) \coloneqq |\mathcal{N}_{\Gamma}(\tau;\rho)|,$$

where

$$\mathcal{N}_{\Gamma}(\tau;\rho) \coloneqq \bigg\{ \gamma \in \Gamma \setminus \bigg(\bigcup_{j=1}^{s} \Gamma_{p_j} \cup \bigcup_{j=1}^{t} \Gamma_{e_j} \bigg) \, \bigg| \, \mathrm{dist}_{\mathrm{hyp}}(\tau,\gamma\tau) \le \rho \bigg\}.$$

Let now f be a positive, smooth, and decreasing function on $\mathbb{R}_{\geq 0}$. Then, adapting the arguments from [JL95] to Fuchsian subgroups of $\mathrm{SL}_2(\mathbb{R})$, we have for any $\tau \in \mathbb{H}$ and any $\delta \geq r_{\Gamma}/2$ the inequality

(11)
$$\int_{0}^{\infty} f(\rho) \, \mathrm{d}N_{\Gamma}(\tau;\rho) \leq \int_{0}^{\delta} f(\rho) \, \mathrm{d}N_{\Gamma}(\tau;\rho) + \frac{2|\mathrm{Cent}(\Gamma)| \cosh(r_{\Gamma}/4)}{\sinh(r_{\Gamma}/4)} \sinh(\delta) f(\delta) + \frac{|\mathrm{Cent}(\Gamma)|}{2\sinh^{2}(r_{\Gamma}/4)} \int_{\delta}^{\infty} f(\rho) \sinh(\rho + r_{\Gamma}/2) \, \mathrm{d}\rho;$$

here $\text{Cent}(\Gamma)$ denotes the center of Γ . Note that our definition (8) of injectivity radius differs from the one used in [JL95] by a factor of 2, and the inequality (11) takes this fact into account.

2.5. Jacobi forms. For $k, m \in \mathbb{N}$, we let $J_{k,m}^{\text{cusp}}(\Gamma_0)$ denote the space of Jacobi cusp forms of weight k and index m for $\Gamma_0 = \text{SL}_2(\mathbb{Z})$, i. e., the space of holomorphic functions $f : \mathbb{H} \times \mathbb{C} \to \mathbb{C}$, which have the transformation behaviour

$$f\left(\frac{a\tau+b}{c\tau+d},\frac{z+\lambda\tau+\mu}{c\tau+d}\right)(c\tau+d)^{-k}\exp\left(2\pi im\left(\lambda^2\tau+2\lambda z-\frac{c(z+\lambda\tau+\mu)^2}{c\tau+d}\right)\right)=f(\tau,z)$$

for all $\left[\begin{pmatrix} a & b \\ c & d \end{pmatrix}, (\lambda, \mu) \right] \in \Gamma_0 \ltimes \mathbb{Z}^2$, and which have a Fourier expansion of the form

$$f(\tau, z) = \sum_{\substack{n \in \mathbb{N}, r \in \mathbb{Z} \\ 4mn - r^2 > 0}} c(n, r) q^n \zeta^r \qquad (q = e^{2\pi i \tau}, \, \zeta = e^{2\pi i z}).$$

Given $f \in J_{k,m}^{\text{cusp}}(\Gamma_0)$, we define

$$||f(\tau, z)||_{\text{Pet}}^2 \coloneqq |f(\tau, z)|^2 \eta^k e^{-\frac{4\pi m y^2}{\eta}} \qquad (\tau = \xi + i\eta, \, z = x + iy),$$

which defines a $\Gamma_0 \ltimes \mathbb{Z}^2$ -invariant function on $\mathbb{H} \times \mathbb{C}$ called the pointwise Petersson norm of f.

Let \mathcal{D}_{Γ_0} denote a fundamental domain of the quotient space $Y_{\Gamma_0} \coloneqq \Gamma_0 \ltimes \mathbb{Z}^2 \setminus \mathbb{H} \times \mathbb{C}$, which is a 2-dimensional complex manifold. The space $J_{k,m}^{\text{cusp}}(\Gamma_0)$ is equipped with the Petersson inner product

(12)

$$\langle f_1, f_2 \rangle_{\text{Pet}} \coloneqq \int_{\mathcal{D}_{\Gamma_0}} f_1(\tau, z) \overline{f_2(\tau, z)} \, \eta^k \, e^{-\frac{4\pi m y^2}{\eta}} \, \frac{\mathrm{d}\xi \wedge \mathrm{d}\eta \wedge \mathrm{d}x \wedge \mathrm{d}y}{\eta^3} \qquad (f_1, f_2 \in J_{k,m}^{\mathrm{cusp}}(\Gamma_0)).$$

For $f \in J_{k,m}^{\text{cusp}}(\Gamma_0)$, one has the decomposition

(13)
$$f(\tau, z) = \sum_{\mu=0}^{2m-1} \varphi_{\mu}(\tau) \vartheta_{\mu,m}(\tau, z),$$

where the function φ_{μ} is a cusp form of weight $(k - \frac{1}{2})$ for the finite index subgroup $\Gamma_1 \coloneqq \Gamma_0(4m)$ of Γ_0 , and $\vartheta_{\mu,m}$ is the Jacobi theta function

(14)
$$\vartheta_{\mu,m}(\tau,z) \coloneqq \sum_{n \in \mathbb{Z}} e^{2\pi i m \tau \left(n - \frac{\mu}{2m}\right)^2 + 2\pi i z (2mn - \mu)}$$

As we will see below, the theta functions $\vartheta_{\mu,m}$ ($\mu = 0, \ldots, 2m-1$) arise for fixed $\tau \in \mathbb{H}$ as global sections of a suitable line bundle on the elliptic curve associated to τ . In fact, it is shown in Theorem 5.1 of [EZ85] that the decomposition (14) gives rise to the isomorphism

$$J_{k,m}^{\mathrm{cusp}}(\Gamma_0) \cong \mathcal{V}_{k-\frac{1}{2}}(\Gamma_0)$$

where $\mathcal{V}_{k-\frac{1}{2}}(\Gamma_0)$ denotes the complex vector space of vector-valued cusp forms of weight $(k-\frac{1}{2})$ with suitable transformation behaviour with respect to Γ_0 .

Let now

$$f_1(\tau, z) = \sum_{\mu=0}^{2m-1} \varphi_{\mu,1}(\tau) \vartheta_{\mu,m}(\tau, z) \quad \text{and} \quad f_2(\tau, z) = \sum_{\mu=0}^{2m-1} \varphi_{\mu,2}(\tau) \vartheta_{\mu,m}(\tau, z)$$

be two Jacobi cusp forms of weight k and index m for Γ_0 . Then, the decomoposition (13) gives rise to the equality (see Theorem 5.3 in [EZ85])

(15)
$$\langle f_1, f_2 \rangle_{\text{Pet}} = \frac{1}{\sqrt{4m}} \int_{\mathcal{F}_{\Gamma_0}} \sum_{\mu=0}^{2m-1} \varphi_{\mu,1}(\tau) \overline{\varphi_{\mu,2}(\tau)} \eta^{k-\frac{1}{2}} \frac{\mathrm{d}\xi \wedge \mathrm{d}\eta}{\eta^2},$$

where we recall that \mathcal{F}_{Γ_0} is a fundamental domain for the quotient space $X_{\Gamma_0} = \Gamma_0 \setminus \mathbb{H}$.

For a fixed $\tau = \xi + i\eta \in \mathbb{H}$, consider the elliptic curve $E_{\tau} \coloneqq \mathbb{C}/\Lambda_{\tau}$ with $\Lambda_{\tau} \coloneqq \mathbb{Z} \oplus \tau \mathbb{Z}$. Let O_{τ} denote the identity element of E_{τ} , when considered as an abelian group with \oplus_{τ} denoting the group operation, and let

$$[2]: E_{\tau} \longrightarrow E_{\tau}$$

be multiplication by 2, given by the assignment $z \mapsto 2z := z \oplus_{\tau} z$, which is an isogeny of degree 4. Let \mathcal{M}_{τ} be the line bundle associated to the divisor O_{τ} . Then, the theorem of the cube gives the isomorphism

$$[2]^*\mathcal{M}_{\tau}\cong\mathcal{M}_{\tau}^{\otimes 4}.$$

We then put $\mathcal{L}_{\tau} := \mathcal{M}_{\tau}^{\otimes 2}$ and find that the theta functions $\vartheta_{\mu,m}$ ($\mu = 0, \ldots, 2m - 1$) arise as global holomorphic sections of the line bundle $\mathcal{L}_{\tau}^{\otimes m}$.

The pointwise norm of $\vartheta_{\mu,m} \in H^0(E_{\tau}, \mathcal{L}_{\tau}^{\otimes m})$ at the point $z = x + iy \in E_{\tau}$ (identifying E_{τ} with its universal cover \mathbb{C}) is given by the following formula

(16)
$$\|\vartheta_{\mu,m}(\tau,z)\|_{\mathcal{L}^{\otimes m}_{\tau}}^{2} \coloneqq |\vartheta_{\mu,m}(\tau,z)|^{2} \sqrt{\eta} e^{-\frac{4\pi m y^{2}}{\eta}}.$$

Let μ_{eucl} denote the Euclidean metric on E_{τ} ; at the point $z = x + iy \in E_{\tau}$, it is given by the formula

(17)
$$\mu_{\text{eucl}}(z) = \frac{i}{2} \cdot \frac{\mathrm{d}z \wedge \mathrm{d}\overline{z}}{\eta} = \frac{\mathrm{d}x \wedge \mathrm{d}y}{\eta}$$

Furthermore, the L²-norm of $\vartheta_{\mu,m} \in H^0(E_\tau, \mathcal{L}_\tau^{\otimes m})$ is given by the following formula

$$\begin{aligned} \|\vartheta_{\mu,m}(\tau,\cdot)\|_{\mathrm{L}^{2},\mathcal{L}^{\otimes m}_{\tau}}^{2} &= \int_{E_{\tau}} \|\vartheta_{\mu,m}(\tau,z)\|_{\mathcal{L}^{\otimes m}_{\tau}}^{2} \mu_{\mathrm{eucl}}(z) \\ &= \int_{0}^{\eta} \int_{0}^{1} |\vartheta_{\mu,m}(\tau,z)|^{2} \sqrt{\eta} \, e^{-\frac{4\pi m y^{2}}{\eta}} \, \frac{\mathrm{d}x \wedge \mathrm{d}y}{\eta} \end{aligned}$$

2.6. Asymptotics for Bergman kernels. In this subsection, we recall asymptotics of Bergman kernels associated to tensor powers of holomorphic line bundles on compact complex Kähler manifolds, which are used in section 4 to derive bounds for theta functions.

Let (M, ω) be a compact complex Kähler manifold of dimension n with positive closed (1, 1)-form ω . Let \mathcal{L} be a positive hermitian holomorphic line bundle on M and let $H^0(M, \mathcal{L}^{\otimes m})$ denote the vector space of global holomorphic sections its m-th tensor power $\mathcal{L}^{\otimes m}$ for $m \in \mathbb{Z}_{\geq 1}$. Let $|\cdot|_{\mathcal{L}^{\otimes m}}$ and $\langle \cdot, \cdot \rangle_{\mathrm{L}^2, \mathcal{L}^{\otimes m}}$ denote the pointwise hermitian metric and the L^2 -inner product on $H^0(M, \mathcal{L}^{\otimes m})$, respectively.

Let $\{s_j\}$ denote an orthonormal basis of $H^0(M, \mathcal{L}^{\otimes m})$ with respect to the L²-inner product. For any $z \in M$, the function

$$B_{\mathcal{L}^{\otimes m}}(z) \coloneqq \sum_{j} |s_j(z)|^2_{\mathcal{L}^{\otimes m}}$$

is called the Bergman kernel associated to the line bundle $\mathcal{L}^{\otimes m}$. We note that the above definition is independent on the choice of an orthonormal basis of $H^0(M, \mathcal{L}^{\otimes m})$.

Let

(18)
$$c_1(\mathcal{L}, |\cdot|_{\mathcal{L}})(z) \coloneqq -\frac{i}{2\pi} \partial_z \partial_{\overline{z}} \log |s(z)|_{\mathcal{L}}^2$$

denote the curvature form of the line bundle \mathcal{L} at the point $z \in M$, where s is any meromorphic section of \mathcal{L} . At any $z \in M$, there exists a coordinate chart around the point z such that

$$\omega(z) = \sum_{j=1}^{n} \frac{i}{2} \cdot \mathrm{d}z_{j} \wedge \mathrm{d}\overline{z}_{j} \quad \text{and} \quad \mathrm{c}_{1}(\mathcal{L}, |\cdot|_{\mathcal{L}})(z) = \sum_{j=1}^{n} \frac{i}{2} \cdot \alpha_{j} \cdot \mathrm{d}z_{j} \wedge \mathrm{d}\overline{z}_{j}.$$

The complex numbers $\alpha_1, \ldots, \alpha_n$ are called the eigenvalues of the curvature form $c_1(\mathcal{L}, |\cdot|_{\mathcal{L}})(z)$ at the point $z \in M$. We set

$$\det_{\omega} \big(c_1(\mathcal{L}, |\cdot|_{\mathcal{L}})(z) \big) \coloneqq \prod_{j=1}^n \alpha_j.$$

Since the line bundle \mathcal{L} is positive, we have $\alpha_j > 0$ for $j = 1, \ldots, n$. Finally, we recall from Theorem 2.1 in [Bo96] the bound

(19)
$$B_{\mathcal{L}^{\otimes m}}(z) = \det_{\omega} \big(c_1(\mathcal{L}, |\cdot|_{\mathcal{L}})(z) \big) m^n + O(m^{n-1}),$$

provided that \mathcal{L} is a positive line bundle for any $z \in M$.

3. Sup-norm bounds for cusp forms revisited

Refining arguments of [AM17] and [AM18], we first derive bounds for the Bergman kernel along the diagonal.

Proposition 3.1. With notations as above, let $\Gamma \subset SL_2(\mathbb{R})$ be a cocompact Fuchsian subgroup without elliptic elements. Then, for $k \in \mathbb{R}_{\geq 5}$ and $\tau \in \mathbb{H}$, we have the bound

$$\|B_{k,\chi}(\tau,\tau)\|_{\text{Pet}} \le \frac{k-1}{2\pi} + \frac{3(k-1)}{\pi \cosh^{k-4}(r_{\Gamma}/4)} \left(1 + \frac{1}{\sinh^2(r_{\Gamma}/4)}\right).$$

Proof. Letting $k \in \mathbb{R}_{\geq 5}$ and considering the Bergman kernel (10) on the diagonal, we derive by means of relation (5) the bound

$$||B_{k,\chi}(\tau,\tau)||_{\text{Pet}} = \frac{2^k (k-1)}{4\pi} \bigg| \sum_{\gamma \in \binom{a \ b}{c \ d} \in \Gamma} \frac{1}{(\tau - \gamma \overline{\tau})^k} \frac{1}{\chi(\gamma)(c\overline{\tau} + d)^k} \bigg| \operatorname{Im}(\tau)^k$$
$$\leq \frac{k-1}{4\pi} \sum_{\gamma \in \Gamma} \left(\frac{4 \operatorname{Im}(\tau) \operatorname{Im}(\gamma \tau)}{|\tau - \gamma \overline{\tau}|^2} \right)^{k/2} = \frac{k-1}{4\pi} \sum_{\gamma \in \Gamma} \frac{1}{\cosh^k(\operatorname{dist}_{\operatorname{hyp}}(\tau, \gamma \tau)/2)}$$
$$= \frac{k-1}{4\pi} \bigg(|\operatorname{Cent}(\Gamma)| + \sum_{\gamma \in \Gamma \setminus \operatorname{Cent}(\Gamma)} \frac{1}{\cosh^k(\operatorname{dist}_{\operatorname{hyp}}(\tau, \gamma \tau)/2)} \bigg).$$

Substituting $\delta = r_{\Gamma}/2$ in inequality (11) and using the fact that $|\text{Cent}(\Gamma)| \leq 2$, we derive

(21)
$$\sum_{\gamma \in \Gamma \setminus \operatorname{Cent}(\Gamma)} \frac{1}{\cosh^{k}(\operatorname{dist}_{\operatorname{hyp}}(\tau, \gamma \tau)/2)} \leq \int_{0}^{r_{\Gamma}/2} \frac{\mathrm{d}N_{\Gamma}(\tau; \rho)}{\cosh^{k}(\rho/2)} + \frac{8}{\cosh^{k-2}(r_{\Gamma}/4)} + \frac{1}{\sinh^{2}(r_{\Gamma}/4)} \int_{r_{\Gamma}/2}^{\infty} \frac{\sinh(\rho + r_{\Gamma}/2)}{\cosh^{k}(\rho/2)} \,\mathrm{d}\rho.$$

From the defining equation (8) of the injectivity radius r_{Γ} , we find for the first term of (21) that

(22)
$$\int_{0}^{r_{\Gamma}/2} \frac{\mathrm{d}N_{\Gamma}(\tau;\rho)}{\cosh^{k}(\rho/2)} = 0.$$

(20)

With regard to the third term of (21), we recall the bound (12) in [AM17], which states for any $k \in \mathbb{R}_{\geq 5}$ and any $\delta \geq 0$ (note that we have replaced 2k by k) that

$$\frac{1}{\sinh^2(r_{\Gamma}/4)} \int_{\delta}^{\infty} \frac{\sinh(\rho + r_{\Gamma}/2)}{\cosh^k(\rho/2)} d\rho$$
(23) $\leq \frac{4}{(k-2)\cosh^{k-2}(\delta/2)} \left(2 + \frac{1}{\sinh^2(r_{\Gamma}/4)}\right) + \frac{8}{(k-4)\cosh^{k-4}(\delta/2)} \cdot \frac{1}{\sinh^2(r_{\Gamma}/4)}.$

From the elementary inequality $\cosh^{k-4}(r_{\Gamma}/4) \leq \cosh^{k-2}(r_{\Gamma}/4)$ and recalling that $k \in \mathbb{R}_{\geq 5}$, we now derive from (23) with $\delta = r_{\Gamma}/2$ the bound

$$\frac{1}{\sinh^{2}(r_{\Gamma}/4)} \int_{r_{\Gamma}/2}^{\infty} \frac{\sinh(\rho + r_{\Gamma}/2)}{\cosh^{k}(\rho/2)} d\rho \\
\leq \frac{4}{(k-2)\cosh^{k-2}(r_{\Gamma}/4)} \left(2 + \frac{1}{\sinh^{2}(r_{\Gamma}/4)}\right) + \frac{8}{(k-4)\cosh^{k-4}(r_{\Gamma}/4)} \cdot \frac{1}{\sinh^{2}(r_{\Gamma}/4)} \\
(24) \leq \frac{4}{\cosh^{k-4}(r_{\Gamma}/4)} \left(1 + \frac{1}{\sinh^{2}(r_{\Gamma}/4)}\right) + \frac{8}{(k-4)\cosh^{k-4}(r_{\Gamma}/4)} \cdot \frac{1}{\sinh^{2}(r_{\Gamma}/4)}.$$

Combining the bounds (20), (21) with (22), (24), and using the fact that $k \in \mathbb{R}_{\geq 5}$, completes the proof of the proposition.

Proposition 3.2. With notations as above, let $\Gamma \subset SL_2(\mathbb{R})$ be a cofinite Fuchsian subgroup. Then, for $k \in \mathbb{R}_{\geq 5}$ and $\tau \in \mathbb{H}$, we have the bound

$$\|B_{k,\chi}(\tau,\tau)\|_{\text{Pet}} \leq \frac{k-1}{2\pi} + \frac{3(k-1)}{\pi \cosh^{k-4}(r_{\Gamma}/4)} \left(1 + \frac{1}{\sinh^{2}(r_{\Gamma}/4)}\right) \\ + \frac{k-1}{4\pi} \sum_{e_{j} \in \mathcal{E}_{\Gamma}} (m_{j}-1) + \frac{2(k-1)}{\sqrt{\pi}} \cdot \frac{\Gamma((k-1)/2)}{\Gamma(k/2)} \sum_{p_{j} \in \mathcal{P}_{\Gamma}} \operatorname{Im}(\sigma_{\mathcal{P},j}^{-1}\tau),$$

where $\sigma_{\mathcal{P},j}$ is the scaling matrix associated to the cusp $p_j \in \mathcal{P}_{\Gamma}$ defined in (6).

Proof. For $k \in \mathbb{R}_{\geq 5}$ and $\tau \in \mathbb{H}$, using the bound (20) and the fact that $|\text{Cent}(\Gamma)| \leq 2$, we derive

$$||B_{k,\chi}(\tau,\tau)||_{\text{Pet}} \leq \frac{k-1}{2\pi} + \frac{k-1}{4\pi} \sum_{\gamma \in \Gamma \setminus \left(\bigcup_{e_j \in \mathcal{E}_{\Gamma}} \Gamma_{e_j} \cup \bigcup_{p_j \in \mathcal{P}_{\Gamma}} \Gamma_{p_j}\right)} \frac{1}{\cosh^k (\operatorname{dist}_{\operatorname{hyp}}(\tau,\gamma\tau)/2)} + \frac{k-1}{4\pi} \sum_{e_j \in \mathcal{E}_{\Gamma}} \sum_{\gamma \in \Gamma_{e_j} \setminus \operatorname{Cent}(\Gamma)} \frac{1}{\cosh^k (\operatorname{dist}_{\operatorname{hyp}}(\tau,\gamma\tau)/2)} + \frac{k-1}{4\pi} \sum_{p_j \in \mathcal{P}_{\Gamma}} \sum_{\gamma \in \Gamma_{p_j} \setminus \operatorname{Cent}(\Gamma)} \frac{1}{\cosh^k (\operatorname{dist}_{\operatorname{hyp}}(\tau,\gamma\tau)/2)}.$$
(25)

Adapting our arguments from Proposition 3.1 to the second summand on the right-hand side of (25), we arrive at the bound

(26)
$$\frac{k-1}{4\pi} \sum_{\gamma \in \Gamma \setminus \left(\bigcup_{e_j \in \mathcal{E}_{\Gamma}} \Gamma_{e_j} \cup \bigcup_{p_j \in \mathcal{P}_{\Gamma}} \Gamma_{p_j}\right)} \frac{1}{\cosh^k (\operatorname{dist}_{\operatorname{hyp}}(\tau, \gamma \tau)/2)} \leq \frac{3(k-1)}{\pi \cosh^{k-4}(r_{\Gamma}/4)} \left(1 + \frac{1}{\sinh^2(r_{\Gamma}/4)}\right).$$

For the third term on the right-hand side of (25), we trivially have the bound

(27)
$$\frac{k-1}{4\pi} \sum_{e_j \in \mathcal{E}_{\Gamma}} \sum_{\gamma \in \Gamma_{e_j} \setminus \text{Cent}(\Gamma)} \frac{1}{\cosh^k (\text{dist}_{\text{hyp}}(\tau, \gamma \tau)/2)} \le \frac{k-1}{4\pi} \sum_{e_j \in \mathcal{E}_{\Gamma}} (m_j - 1)$$

From the definition of the scaling matrix (6) and using the fact that $|\text{Cent}(\Gamma)| \leq 2$, we find

(28)
$$\frac{k-1}{4\pi} \sum_{p_j \in \mathcal{P}_{\Gamma}} \sum_{\gamma \in \Gamma_{p_j} \setminus \operatorname{Cent}(\Gamma)} \frac{1}{\cosh^k (\operatorname{dist}_{\operatorname{hyp}}(\tau, \gamma \tau)/2)} \leq \frac{k-1}{2\pi} \sum_{p_j \in \mathcal{P}_{\Gamma}} \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{1}{\cosh^k (\operatorname{dist}_{\operatorname{hyp}}(\sigma_{\mathcal{P},j}^{-1}\tau, \sigma_{\mathcal{P},j}^{-1}\tau + n)/2)}.$$

We now recall the bound (18) in [AM17], which gives for $k \in \mathbb{R}_{\geq 5}$, $p_j \in \mathcal{P}_{\Gamma}$, and $\tau, \tau' \in \mathbb{H}$ (note that we have replaced 2k by k) the bound

(29)
$$\frac{k-1}{2\pi} \sum_{p_j \in \mathcal{P}_{\Gamma}} \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{1}{\cosh^k (\operatorname{dist}_{\operatorname{hyp}}(\sigma_{\mathcal{P},j}^{-1}\tau, \sigma_{\mathcal{P},j}^{-1}\tau' + n)/2)} \\ \leq \frac{k-1}{\sqrt{\pi}} \cdot \frac{\Gamma((k-1)/2)}{\Gamma(k/2)} \sum_{p_j \in \mathcal{P}_{\Gamma}} \frac{\left(4\operatorname{Im}(\sigma_{\mathcal{P},j}^{-1}\tau)\operatorname{Im}(\sigma_{\mathcal{P},j}^{-1}\tau')\right)^{k/2}}{\left(\operatorname{Im}(\sigma_{\mathcal{P},j}^{-1}\tau) + \operatorname{Im}(\sigma_{\mathcal{P},j}^{-1}\tau')\right)^{k-1}}.$$

Substituting $\tau = \tau'$ in (29) and combining it with (28), we arrive for the fourth term on the right-hand side of (25) at the bound

(30)
$$\frac{k-1}{4\pi} \sum_{p_j \in \mathcal{P}_{\Gamma}} \sum_{\gamma \in \Gamma_{p_j} \setminus \text{Cent}(\Gamma)} \frac{1}{\cosh^k(\text{dist}_{\text{hyp}}(\tau, \gamma \tau)/2)} \\ \leq \frac{2(k-1)}{\sqrt{\pi}} \cdot \frac{\Gamma((k-1)/2)}{\Gamma(k/2)} \sum_{p_j \in \mathcal{P}_{\Gamma}} \text{Im}(\sigma_{\mathcal{P},j}^{-1}\tau).$$

Combining the bounds (26), (27), and (30) with (25) completes the proof of the proposition. \Box

Theorem 3.3. With notations as above, let $\Gamma \subset SL_2(\mathbb{R})$ be a cofinite Fuchsian subgroup and $k \in \mathbb{R}_{\geq 5}$. Then, if Γ is cocompact without elliptic elements, we have the bound

(31)
$$\sup_{\tau \in \mathbb{H}} \|B_{k,\chi}(\tau,\tau)\|_{\mathrm{Pet}} = O_{\Gamma}(k).$$

Moreover, if Γ is cofinite, we have the bound

(32)
$$\sup_{\tau \in \mathbb{H}} \|B_{k,\chi}(\tau,\tau)\|_{\text{Pet}} = O_{\Gamma}\left(k^{\frac{3}{2}}\right)$$

The implied constants in the bounds (31) and (32) depend only on the Fuchsian subgroup Γ .

Proof. When Γ is cocompact without elliptic elements, the claimed bound (31) follows directly from Proposition 3.1.

Let next Γ be a cofinite Fuchsian subgroup. From the proof of Theorem 6.1 in [FJK16], it follows that

(33)
$$\sup_{\tau \in \mathbb{H}} \|B_{k,\chi}(\tau,\tau)\|_{\operatorname{Pet}} = \sup_{\substack{\tau \in \partial \mathcal{F}_Y \\ Y = k/(2\pi)}} \|B_{k,\chi}(\tau,\tau)\|_{\operatorname{Pet}},$$

where $\partial \mathcal{F}_Y$ denotes the boundary of the truncated fundamental domain \mathcal{F}_Y defined in (7). Combining Proposition 3.2 with (33) and employing the fact that

$$\frac{\Gamma((k-1)/2)}{\Gamma(k/2)} = O\left(\frac{1}{\sqrt{k}}\right),\,$$

we derive

$$\sup_{\tau \in \mathbb{H}} \|B_{k,\chi}(\tau,\tau)\|_{\text{Pet}} \leq \frac{k-1}{2\pi} + \frac{3(k-1)}{\pi \cosh^{k-4}(r_{\Gamma}/4)} \left(1 + \frac{1}{\sinh^{2}(r_{\Gamma}/4)}\right) + (k-1)C_{\Gamma,\text{ell}} + k^{\frac{3}{2}}C_{\Gamma,\text{par}}$$

for some positive constants $C_{\Gamma,\text{ell}}, C_{\Gamma,\text{par}}$, which depend only on the Fuchsian subgroup Γ . This completes the proof of the theorem.

Corollary 3.4. With notations as above, let $\Gamma \subset SL_2(\mathbb{R})$ be a cofinite Fuchsian subgroup. For $k \in \mathbb{R}_{\geq 5}$, let $f \in S_{k,\chi}(\Gamma)$ be a cusp form, which is normalized with respect to the Petersson inner product (9). If Γ is cocompact without elliptic elements, we have the bound

(34)
$$\sup_{\tau \in \mathbb{H}} \|f(\tau)\|_{\text{Pet}}^2 = O_{\Gamma}(k).$$

Moreover, if Γ is cofinite, we have the bound

(35)
$$\sup_{\tau \in \mathbb{H}} \|f(\tau)\|_{\text{Pet}}^2 = O_{\Gamma}\left(k^{\frac{3}{2}}\right)$$

The implied constants in the bounds (34) and (35) depend only on the Fuchsian subgroup Γ .

Proof. Choose an orthonormal basis $\{f_1 = f, \ldots, f_{d_k}\}$ of $S_{k,\chi}(\Gamma)$ with respect to the Petersson inner product (9). For $\tau \in \mathbb{H}$, we then have the bound

$$||f(\tau)||_{\text{Pet}}^2 \le ||B_{k,\chi}(\tau,\tau)||_{\text{Pet}}$$

The proof of the corollary now immediately follows from Theorem 3.3.

Theorem 3.5. With notations as above, let $\Gamma_0 \subset SL_2(\mathbb{R})$ be a fixed cofinite Fuchsian subgroup and let $\Gamma \subseteq \Gamma_0$ be a finite index subgroup of Γ_0 . For $k \in \mathbb{R}_{\geq 5}$, let $f \in S_{k,\chi}(\Gamma)$ be a cusp form, which is normalized with respect to the Petersson inner product (9). If Γ_0 is cocompact without elliptic elements, we have the bound

(36)
$$\sup_{\tau \in \mathbb{H}} \|f(\tau)\|_{\operatorname{Pet}}^2 = O_{\Gamma_0}(k).$$

Moreover, if Γ_0 is cofinite, we have the bound

(37)
$$\sup_{\tau \in \mathbb{H}} \|f(\tau)\|_{\text{Pet}}^2 = O_{\Gamma_0}\left(k^{\frac{3}{2}}\right).$$

The implied constants in the bounds (36) and (37) depend only on the Fuchsian subgroup Γ_0 .

Proof. Choose an orthonormal basis $\{f_1 = f, \ldots, f_{d_k}\}$ of $S_{k,\chi}(\Gamma)$ with respect to the Petersson inner product (9).

Let now Γ_0 be a cocompact Fuchsian subgroup without elliptic elements. From the proof of Proposition 3.1, we derive the bound

$$\begin{split} \sup_{\tau \in \mathbb{H}} \|f(\tau)\|_{\text{Pet}}^2 &\leq \sup_{\tau \in \mathbb{H}} \|B_{k,\chi}(\tau,\tau)\|_{\text{Pet}} \\ &\leq \frac{k-1}{4\pi} \sup_{\tau \in \mathbb{H}} \sum_{\gamma \in \Gamma} \frac{1}{\cosh^k(\text{dist}_{\text{hyp}}(\tau,\gamma\tau)/2)} \\ &\leq \frac{k-1}{4\pi} \sup_{\tau \in \mathbb{H}} \sum_{\gamma \in \Gamma_0} \frac{1}{\cosh^k(\text{dist}_{\text{hyp}}(\tau,\gamma\tau)/2)} = O_{\Gamma_0}(k), \end{split}$$

which completes the proof of the theorem in the case that Γ_0 is cocompact without elliptic elements.

Let next Γ_0 be a cofinite Fuchsian subgroup. Given Y > 0, we recall from (7) the fundamental domain decomposition

$$\mathcal{F}_{\Gamma} = \mathcal{F}_{Y} \cup \left(\mathcal{F}_{1}^{Y} \cup \ldots \cup \mathcal{F}_{s}^{Y}\right),$$

where \mathcal{F}_Y is a compact subset of \mathcal{F}_{Γ} and the \mathcal{F}_j^Y 's are neighborhoods of the cusps $p_j \in \mathcal{P}_{\Gamma}$ $(j = 1, \ldots, s)$. Choosing Y large enough, we can assume without loss of generality in the sequel that the neighborhoods \mathcal{F}_j^Y are pairwise disjoint. We now first provide a bound for the pointwise Petersson norm of f, when τ ranges across the compact set \mathcal{F}_Y , and subsequently we compute bounds for the pointwise Petersson norm of f, when τ ranges across the neighborhoods \mathcal{F}_j^Y of the cusps for fixed, large enough Y. Adapting arguments from the proof of Proposition 3.2, we obtain the bound

$$\sup_{\tau \in \mathcal{F}_{Y}} \|B_{k,\chi}(\tau,\tau)\|_{\text{Pet}} \leq \frac{k-1}{4\pi} \sup_{\tau \in \mathcal{F}_{Y}} \sum_{\gamma \in \Gamma} \frac{1}{\cosh^{k}(\operatorname{dist}_{\operatorname{hyp}}(\tau,\gamma\tau)/2)} \leq \frac{k-1}{4\pi} \sup_{\tau \in \mathcal{F}_{Y}} \sum_{\gamma \in \Gamma_{0}} \frac{1}{\cosh^{k}(\operatorname{dist}_{\operatorname{hyp}}(\tau,\gamma\tau)/2)} \leq \frac{k-1}{2\pi} + \frac{3(k-1)}{\pi \cosh^{k-4}(r_{\Gamma_{0},Y}/4)} \left(1 + \frac{1}{\sinh^{2}(r_{\Gamma_{0},Y}/4)}\right) + \frac{k-1}{4\pi} \sum_{e_{j} \in \mathcal{E}_{\Gamma_{0}}} (m_{e_{j}} - 1),$$

where

$$r_{\Gamma_0,Y} = \inf \left\{ \operatorname{dist}_{\operatorname{hyp}}(\tau,\gamma\tau) \, \middle| \, \tau \in \mathcal{F}_Y, \, \gamma \in \Gamma_0 \setminus \bigcup_{e_j \in \mathcal{E}_{\Gamma_0}} \Gamma_{0,e_j} \right\} > 0.$$

From this, we immediately conclude that

(39)
$$\sup_{\tau\in\mathcal{F}_Y} \|f(\tau)\|_{\operatorname{Pet}}^2 \leq \sup_{\tau\in\mathcal{F}_Y} \|B_{k,\chi}(\tau,\tau)\|_{\operatorname{Pet}} = O_{\Gamma_0,Y}(k),$$

where the implied constant depends on Γ_0 and the choice of Y.

We are left to provide bounds for the pointwise Petersson norm of f, when τ ranges across the neighborhoods \mathcal{F}_j^Y of the cusps. For this, we will have to distinguish between the two cases $Y > \frac{k}{2\pi}$ and $Y < \frac{k}{2\pi}$. Without loss of generality, we can assume that j = 1, when $p_1 \in \mathcal{P}_{\Gamma}$ is the cusp at infinity for Γ lying above the cusp p at infinity for Γ_0 with ramification index $[\Gamma_{0,p} : \Gamma_{p_1}]$ (note that $\Gamma_{0,p}$ denotes the stabilizer subgroup of p in Γ_0). With the above notations, we obtain the inclusion $\Gamma \setminus \Gamma_{p_1} \subseteq \Gamma_0 \setminus \Gamma_{0,p}$.

We first treat the case $Y > \frac{k}{2\pi}$, which implies that $\mathcal{F}_1^Y \subset \mathcal{F}_1^{k/(2\pi)}$. Arguing as in the proof of Theorem 6.1 in [FJK16], we deduce, recalling the inclusion $\Gamma \setminus \Gamma_{p_1} \subseteq \Gamma_0 \setminus \Gamma_{0,p}$, that

$$\sup_{\tau \in \mathcal{F}_{1}^{Y}} \|B_{k,\chi}(\tau,\tau)\|_{\operatorname{Pet}} \leq \sup_{\tau \in \mathcal{F}_{1}^{k/(2\pi)}} \|B_{k,\chi}(\tau,\tau)\|_{\operatorname{Pet}} \leq \sup_{\tau \in \partial \mathcal{F}_{1}^{k/(2\pi)}} \|B_{k,\chi}(\tau,\tau)\|_{\operatorname{Pet}} \\
\leq \frac{k-1}{4\pi} \left(\sup_{\tau \in \partial \mathcal{F}_{1}^{k/(2\pi)}} \sum_{\gamma \in \Gamma \setminus \Gamma_{p_{1}}} \frac{1}{\cosh^{k}(\operatorname{dist}_{\operatorname{hyp}}(\tau,\gamma\tau)/2)} + \sup_{\tau \in \partial \mathcal{F}_{1}^{k/(2\pi)}} \sum_{\gamma \in \Gamma_{p_{1}}} \frac{1}{\cosh^{k}(\operatorname{dist}_{\operatorname{hyp}}(\tau,\gamma\tau)/2)} \right) \\
\leq \frac{k-1}{4\pi} \left(\sup_{\tau \in \partial \mathcal{F}_{1}^{k/(2\pi)}} \sum_{\gamma \in \Gamma_{0} \setminus \Gamma_{0,p}} \frac{1}{\cosh^{k}(\operatorname{dist}_{\operatorname{hyp}}(\tau,\gamma\tau)/2)} + \sup_{\tau \in \partial \mathcal{F}_{1}^{k/(2\pi)}} \sum_{\gamma \in \Gamma_{p_{1}}} \frac{1}{\cosh^{k}(\operatorname{dist}_{\operatorname{hyp}}(\tau,\gamma\tau)/2)} \right) \\$$
(40)

Arguments similar to the ones used to derive the bound (38), lead for the first term of (40) to the bound

$$\frac{k-1}{4\pi} \sup_{\tau \in \partial \mathcal{F}_{1}^{k/(2\pi)}} \sum_{\gamma \in \Gamma_{0} \setminus \Gamma_{0,p}} \frac{1}{\cosh^{k}(\operatorname{dist}_{\operatorname{hyp}}(\tau, \gamma \tau)/2)} \\ \leq \frac{k-1}{2\pi} + \frac{3(k-1)}{\pi \cosh^{k-4}(r_{\Gamma_{0},k/(2\pi)}/4)} \left(1 + \frac{1}{\sinh^{2}(r_{\Gamma_{0},k/(2\pi)}/4)}\right) + \frac{k-1}{4\pi} \sum_{e_{j} \in \mathcal{E}_{\Gamma_{0}}} (m_{e_{j}} - 1),$$

where

$$r_{\Gamma_{0,k/(2\pi)}} = \inf \left\{ \operatorname{dist}_{\operatorname{hyp}}(\tau, \gamma \tau) \, \middle| \, \tau \in \partial \mathcal{F}_{1}^{k/(2\pi)}, \, \gamma \in \Gamma_{0} \setminus \left(\Gamma_{0,p} \cup \bigcup_{e_{j} \in \mathcal{E}_{\Gamma_{0}}} \Gamma_{0,e_{j}} \right) \right\} > 0$$

Since it is easy to see that

$$\frac{1}{\sinh^2(r_{\Gamma_0,k/(2\pi)})} = O_{\Gamma_0}(1),$$

we arrive for the first term of (40) at the bound

(41)
$$\frac{k-1}{4\pi} \sup_{\tau \in \partial \mathcal{F}_1^{k/(2\pi)}} \sum_{\gamma \in \Gamma_0 \setminus \Gamma_{0,p}} \frac{1}{\cosh^k (\operatorname{dist}_{\operatorname{hyp}}(\tau, \gamma \tau)/2)} = O_{\Gamma_0}(k).$$

Using the same arguments as in the proof of Proposition 3.2, we derive for the second term of (40) the bound

(42)
$$\frac{k-1}{4\pi} \sup_{\tau \in \partial \mathcal{F}_1^{k/(2\pi)}} \sum_{\gamma \in \Gamma_{p_1}} \frac{1}{\cosh^k (\operatorname{dist}_{\operatorname{hyp}}(\tau, \gamma \tau)/2)} = O(k^{\frac{3}{2}}).$$

By means of (40), we thus deduce from (41) and (42) in the case $Y > \frac{k}{2\pi}$ the bound

(43)
$$\sup_{\tau \in \mathcal{F}_1^Y} \|f(\tau)\|_{\text{Pet}}^2 \le \sup_{\tau \in \mathcal{F}_1^Y} \|B_{k,\chi}(\tau,\tau)\|_{\text{Pet}} = O_{\Gamma_0}(k^{\frac{3}{2}}).$$

We finally turn to the case $Y < \frac{k}{2\pi}$, which implies that $\mathcal{F}_1^{k/(2\pi)} \subset \mathcal{F}_1^Y$. Here we find, arguing as in the preceding case that

(44)

$$\sup_{\tau \in \mathcal{F}_{1}^{Y}} \|B_{k,\chi}(\tau,\tau)\|_{\operatorname{Pet}} \leq \sup_{\tau \in \mathcal{F}_{1}^{Y} \setminus \mathcal{F}_{1}^{k/(2\pi)}} \|B_{k,\chi}(\tau,\tau)\|_{\operatorname{Pet}} \leq \frac{k-1}{4\pi} \sup_{\tau \in \mathcal{F}_{1}^{Y} \setminus \mathcal{F}_{1}^{k/(2\pi)}} \sum_{\gamma \in \Gamma_{0} \setminus \Gamma_{0,p}} \frac{1}{\cosh^{k}(\operatorname{dist}_{\operatorname{hyp}}(\tau,\gamma\tau)/2)} + \frac{k-1}{4\pi} \sup_{\tau \in \mathcal{F}_{1}^{Y} \setminus \mathcal{F}_{1}^{k/(2\pi)}} \sum_{\gamma \in \Gamma_{p_{1}}} \frac{1}{\cosh^{k}(\operatorname{dist}_{\operatorname{hyp}}(\tau,\gamma\tau)/2)}.$$

As before, we now obtain the bounds

$$\frac{k-1}{4\pi} \sup_{\tau \in \mathcal{F}_{1}^{Y} \setminus \mathcal{F}_{1}^{k/(2\pi)}} \sum_{\gamma \in \Gamma_{0} \setminus \Gamma_{0,p}} \frac{1}{\cosh^{k}(\operatorname{dist}_{\operatorname{hyp}}(\tau, \gamma \tau)/2)} \le \frac{k-1}{2\pi} + \frac{3(k-1)}{\pi \cosh^{k-4}(r_{\Gamma_{0},Y}'/4)} \left(1 + \frac{1}{\sinh^{2}(r_{\Gamma_{0},Y}'/4)}\right) + \frac{k-1}{4\pi} \sum_{e_{j} \in \mathcal{E}_{\Gamma_{0}}} (m_{e_{j}} - 1)$$

 $(45) \qquad = O_{\Gamma_0,Y}(k),$

noting that

$$r_{\Gamma_{0},Y}' = \inf \left\{ \operatorname{dist}_{\operatorname{hyp}}(\tau,\gamma\tau) \, \middle| \, \tau \in \mathcal{F}_{1}^{Y}, \, \gamma \in \Gamma_{0} \setminus \left(\Gamma_{0,p} \cup \bigcup_{e_{j} \in \mathcal{E}_{\Gamma_{0}}} \Gamma_{0,e_{j}} \right) \right\} > 0,$$

as well as the bound

(46)
$$\frac{k-1}{4\pi} \sup_{\tau \in \mathcal{F}_1^Y \setminus \mathcal{F}_1^{k/(2\pi)}} \sum_{\gamma \in \Gamma_{p_1}} \frac{1}{\cosh^k (\operatorname{dist}_{\operatorname{hyp}}(\tau, \gamma \tau)/2)} = O\left(k^{\frac{3}{2}}\right)$$

By means of (44), we thus deduce from (45) and (46) in the case $Y < \frac{k}{2\pi}$ the bound

(47)
$$\sup_{\tau \in \mathcal{F}_{1}^{Y}} \|f(\tau)\|_{\text{Pet}}^{2} \leq \sup_{\tau \in \mathcal{F}_{1}^{Y}} \|B_{k,\chi}(\tau,\tau)\|_{\text{Pet}} = O_{\Gamma_{0},Y}(k^{\frac{3}{2}}).$$

Since Y has been fixed, the claim of the theorem follows from (39), (43), and (47). \Box

Remark 3.6. If $f \in S_{k,\chi}(\Gamma)$ is not a Hecke eigenform, then there is no evidence from the literature to suggest that the estimates (34) and (35) can be improved. Thus the estimates (34) and (35) are expected to be optimal.

4. Sup-norm bounds for Jacobi Cusp forms

For $k \in \mathbb{Z}_{\geq 5}$ and $m \in \mathbb{Z}_{\geq 1}$, let $f \in J_{k,m}^{cusp}(\Gamma_0)$ be a Jacobi cusp form of weight k and index m for the full modular group $\Gamma_0 = \mathrm{SL}_2(\mathbb{Z})$, which is normalized with respect to the Petersson inner product defined by (12). We now aim at bounding the quantity

$$\sup_{(\tau,z)\in\mathbb{H}\times\mathbb{C}}\|f(\tau,z)\|_{\mathrm{Pet}}.$$

Recall from (13) that we have the decomposition

$$f(\tau, z) = \sum_{\mu=0}^{2m-1} \varphi_{\mu}(\tau) \vartheta_{\mu,m}(\tau, z),$$

where the functions φ_{μ} are cusp forms of weight $(k - \frac{1}{2})$ with respect to the finite index subgroup $\Gamma_1 = \Gamma_0(4m)$ of Γ_0 and the theta functions $\vartheta_{\mu,m}$ are defined in (14).

Proposition 4.1. With notations as above, given $\epsilon > 0$, we have the bound

$$\sup_{(\tau,z)\in\mathbb{H}\times\mathbb{C}}\|f(\tau,z)\|_{\mathrm{Pet}}^2 = O_\epsilon\left(k^{\frac{3}{2}}m^{\frac{5}{2}+\epsilon}\|\vartheta_m\|_{\mathcal{L}^{\otimes m}_\tau}^2\right),$$

where

$$\|\vartheta_m\|_{\mathcal{L}^{\otimes m}_{\tau}}^2 \coloneqq \sup_{\substack{(\tau,z) \in \mathbb{H} \times \mathbb{C} \\ 0 \le \mu \le 2m-1}} \|\vartheta_{\mu,m}(\tau,z)\|_{\mathcal{L}^{\otimes m}_{\tau}}^2,$$

and the implied constant depends only on the choice of ϵ .

Proof. Substituting the decomposition (13) of the Jacobi form $f(\tau, z)$ into its pointwise Petersson norm, we compute

$$\begin{split} \|f(\tau,z)\|_{\text{Pet}}^{2} &= \left|\sum_{\mu=0}^{2m-1} \varphi_{\mu}(\tau) \vartheta_{\mu,m}(\tau,z)\right|^{2} \eta^{k} e^{-\frac{4\pi m y^{2}}{\eta}} \\ &= \sum_{\mu=0}^{2m-1} \sum_{\mu'=0}^{2m-1} \varphi_{\mu}(\tau) \vartheta_{\mu,m}(\tau,z) \overline{\varphi_{\mu'}(\tau)} \overline{\vartheta_{\mu',m}(\tau,z)} \eta^{k} e^{-\frac{4\pi m y^{2}}{\eta}} \\ &\leq \sum_{\mu=0}^{2m-1} \sum_{\mu'=0}^{2m-1} |\varphi_{\mu}(\tau) \vartheta_{\mu,m}(\tau,z)| \left|\varphi_{\mu'}(\tau) \vartheta_{\mu',m}(\tau,z)\right| \eta^{k} e^{-\frac{4\pi m y^{2}}{\eta}} \\ &\leq \frac{1}{2} \sum_{\mu=0}^{2m-1} \sum_{\mu'=0}^{2m-1} \left(|\varphi_{\mu}(\tau) \vartheta_{\mu,m}(\tau,z)|^{2} + |\varphi_{\mu'}(\tau) \vartheta_{\mu',m}(\tau,z)|^{2}\right) \eta^{k} e^{-\frac{4\pi m y^{2}}{\eta}}. \end{split}$$

From this we immediately derive

$$\begin{split} \|f(\tau,z)\|_{\text{Pet}}^{2} &\leq \frac{4m}{2} \sum_{\mu=0}^{2m-1} |\varphi_{\mu}(\tau)|^{2} \eta^{k-1/2} \cdot |\vartheta_{\mu,m}(\tau,z)|^{2} \eta^{1/2} e^{-\frac{4\pi m y^{2}}{\eta}} \\ &\leq 2m \sum_{\mu=0}^{2m-1} \|\varphi_{\mu}(\tau)\|_{\text{Pet}}^{2} \cdot \|\vartheta_{m}(\tau,z)\|_{\mathcal{L}_{\tau}^{\otimes m}}^{2} \\ &\leq 2m \|\vartheta_{m}\|_{\mathcal{L}_{\tau}^{\otimes m}}^{2} \sum_{\mu=0}^{2m-1} \|\varphi_{\mu}(\tau)\|_{\text{Pet}}^{2}. \end{split}$$

Since f is normalized with respect to the Petersson inner product, we have from (15) that

(49)
$$\langle f, f \rangle_{\text{Pet}} = \frac{1}{\sqrt{4m} \left[\Gamma_0 : \Gamma_1 \right]} \sum_{\mu=0}^{2m-1} \langle \varphi_\mu, \varphi_\mu \rangle_{\text{Pet}} = 1.$$

Combining (49) with Theorem 3.5, we have the bound

(50)
$$\sum_{\mu=0}^{2m-1} \|\varphi_{\mu}(\tau)\|_{\operatorname{Pet}}^{2} = \sum_{\mu=0}^{2m-1} \langle \varphi_{\mu}, \varphi_{\mu} \rangle_{\operatorname{Pet}} \frac{\|\varphi_{\mu}(\tau)\|_{\operatorname{Pet}}^{2}}{\langle \varphi_{\mu}, \varphi_{\mu} \rangle_{\operatorname{Pet}}} \le C \sqrt{4m} \left[\Gamma_{0} : \Gamma_{1}\right] k^{\frac{3}{2}},$$

where C is a positive constant depending on Γ_0 . Finally, recalling that

(51)
$$[\Gamma_0:\Gamma_1] = [\operatorname{SL}_2(\mathbb{Z}):\Gamma_0(4m)] = O_\epsilon(m^{1+\epsilon}),$$

where the implied constant depends only on the choice of $\epsilon > 0$, the claim follows combining the bounds (48), (50), and (51).

Proposition 4.2. With notations as above, we have for fixed $\tau \in \mathbb{H}$ and $0 \le \mu \le 2m - 1$ the bound

$$\sup_{z\in\mathbb{C}} \|\vartheta_{\mu,m}(\tau,z)\|_{\mathcal{L}^{\otimes m}_{\tau}}^{2} = O\big(m \,\|\vartheta_{\mu,m}(\tau,\cdot)\|_{\mathrm{L}^{2},\,\mathcal{L}^{\otimes m}_{\tau}}^{2}\big).$$

Proof. Let $B_{\mathcal{L}_{\tau}^{\otimes m}}$ denote the Bergman kernel associated to the space of global holomorphic sections $H^0(E_{\tau}, \mathcal{L}_{\tau}^{\otimes m})$ of the line bundle $\mathcal{L}_{\tau}^{\otimes m}$ on the elliptic curve E_{τ} equipped with the Kähler form $\omega = \mu_{\text{eucl}}$ (see formula (17)). From the bound (19), we then derive for $0 \leq \mu \leq 2m - 1$ that

$$B_{\mathcal{L}_{\tau}^{\otimes m}}(z) = \det_{\omega} \left(c_1(\mathcal{L}_{\tau}, \|\cdot\|_{\mathcal{L}_{\tau}})(z) \right) m + O(1),$$

where $c_1(\mathcal{L}_{\tau}, \|\cdot\|_{\mathcal{L}_{\tau}})(z)$ is the curvature form of \mathcal{L}_{τ} at a point $z \in E_{\tau}$. Using equations (16) and (18), the curvature form $c_1(\mathcal{L}_{\tau}, \|\cdot\|_{\mathcal{L}_{\tau}})(z)$ is given by the formula (identifying E_{τ} with its universal cover \mathbb{C} and writing z = x + iy)

$$c_1(\mathcal{L}_{\tau}, \|\cdot\|_{\mathcal{L}_{\tau}})(z) = -\frac{i}{2\pi} \partial_z \partial_{\overline{z}} \log \|\vartheta_{\mu,1}(\tau, z)\|_{\mathcal{L}_{\tau}}^2$$
$$= -\frac{i}{2\pi} \partial_z \partial_{\overline{z}} \left(-\frac{4\pi y^2}{\eta}\right) = 2\mu_{\text{eucl}}(z) = 2\omega(z).$$

Finally, observing that we have by the definition of the Bergman kernel $B_{\mathcal{L}_{\tau}^{\otimes m}}$ that

$$\frac{\|\vartheta_{\mu,m}(\tau,z)\|_{\mathcal{L}^{\otimes m}_{\tau}}^2}{\|\vartheta_{\mu,m}(\tau,\cdot)\|_{\mathrm{L}^2,\,\mathcal{L}^{\otimes m}_{\tau}}^2} \le B_{\mathcal{L}^{\otimes m}_{\tau}}(z),$$

the claim follows.

(48)

Proposition 4.3. With notations as above, for $0 \le \mu \le 2m - 1$, we have the bound

$$\sup_{\tau \in \mathbb{H}} \|\vartheta_{\mu,m}(\tau,\cdot)\|_{\mathrm{L}^{2},\mathcal{L}_{\tau}^{\otimes m}}^{2} = O(1/\sqrt{m}).$$

Proof. Let $\tau = \xi + i\eta \in \mathbb{H}$, $z = x + iy \in \mathbb{C}$, and $0 \le \mu \le 2m - 1$. For $\tau \in \mathbb{H}$, we need to bound the quantity

(52)
$$\begin{aligned} \|\vartheta_{\mu,m}(\tau,\cdot)\|_{\mathrm{L}^{2},\mathcal{L}_{\tau}^{\otimes m}}^{2} &= \int_{E_{\tau}}^{\pi} \|\vartheta_{\mu,m}(\tau,z)\|_{\mathcal{L}_{\tau}^{\otimes m}}^{2} \mu_{\mathrm{eucl}}(z) \\ &= \int_{0}^{\eta} \int_{0}^{1} |\vartheta_{\mu,m}(\tau,z)|^{2} \sqrt{\eta} \, e^{-\frac{4\pi m y^{2}}{\eta}} \, \frac{\mathrm{d}x \wedge \mathrm{d}y}{\eta}, \end{aligned}$$

where we have used (16) and (17). Recalling (14) and setting $a(n) \coloneqq n - \mu/(2m)$ as well as $b(n) \coloneqq 2mn - \mu$ in (52), we then compute

(53)
$$\|\vartheta_{\mu,m}(\tau,\cdot)\|_{L^{2},\mathcal{L}_{\tau}^{\otimes m}}^{2} = \int_{0}^{\eta} \sum_{n \in \mathbb{Z}} \sum_{n' \in \mathbb{Z}} e^{2\pi i m \tau a(n)^{2}} \overline{e^{2\pi i m \tau a(n')^{2}}} \left(\int_{0}^{1} e^{2\pi i z b(n)} \overline{e^{2\pi i z b(n')}} \, \mathrm{d}x\right) e^{-\frac{4\pi m y^{2}}{\eta}} \frac{\mathrm{d}y}{\sqrt{\eta}}$$

Since

$$\int_{0}^{1} e^{2\pi i z b(n)} \overline{e^{2\pi i z b(n')}} \, \mathrm{d}x = e^{-2\pi y(b(n) + b(n'))} \int_{0}^{1} e^{2\pi i x(b(n) - b(n'))} \, \mathrm{d}x = \begin{cases} e^{-4\pi y b(n)}, & \text{if } n = n', \\ 0, & \text{else,} \end{cases}$$

we arrive at

(54)
$$\sum_{n \in \mathbb{Z}} \sum_{n' \in \mathbb{Z}} e^{2\pi i m \tau a(n)^2} \overline{e^{2\pi i m \tau a(n')^2}} \int_{0}^{1} e^{2\pi i z b(n)} \overline{e^{2\pi i z b(n')}} \, \mathrm{d}x$$
$$= \sum_{n \in \mathbb{Z}} e^{-4\pi m \eta \left(n - \frac{\mu}{2m}\right)^2 - 4\pi y(2mn - \mu)}.$$

Substituting (54) into (53) and using an integral test, we find the bound

(55)
$$\|\vartheta_{\mu,m}(\tau,\cdot)\|_{L^{2},\mathcal{L}_{\tau}^{\otimes m}}^{2} = \sum_{n\in\mathbb{Z}}\int_{0}^{\eta} e^{-4\pi m\eta \left(n-\frac{\mu}{2m}\right)^{2}-4\pi y(2mn-\mu)} e^{-\frac{4\pi my^{2}}{\eta}} \frac{\mathrm{d}y}{\sqrt{\eta}}$$
$$\leq \int_{0}^{\eta}\int_{-\infty}^{\infty} e^{-4\pi m\eta \left(\nu-\frac{\mu}{2m}\right)^{2}-4\pi y(2m\nu-\mu)-\frac{4\pi my^{2}}{\eta}} \,\mathrm{d}\nu \frac{\mathrm{d}y}{\sqrt{\eta}}$$

(56)
$$+ \int_{0}^{\infty} e^{-\frac{\pi\eta\mu^2}{m} + 4\pi y\mu - \frac{4\pi my^2}{\eta}} \frac{\mathrm{d}y}{\sqrt{\eta}}.$$

Now, we rewrite the exponent of the integrand in (55) in the form

$$4\pi m\eta \left(\nu - \frac{\mu}{2m}\right)^2 + 4\pi y (2m\nu - \mu) + \frac{4\pi my^2}{\eta}$$
$$= \left(\sqrt{\frac{\pi\eta}{m}} (2m\nu - \mu)\right)^2 + 4\pi y (2m\nu - \mu) + \left(2\sqrt{\frac{\pi m}{\eta}}y\right)^2$$
$$= \left(\sqrt{\frac{\pi\eta}{m}} (2m\nu - \mu) + 2\sqrt{\frac{\pi m}{\eta}}y\right)^2.$$

Substituting

$$\rho \coloneqq \sqrt{\frac{\pi\eta}{m}}(2m\nu - \mu) + 2\sqrt{\frac{\pi m}{\eta}}y$$

into (55), we obtain for the inner integral

$$\int_{-\infty}^{\infty} e^{-4\pi m\eta \left(\nu - \frac{\mu}{2m}\right)^2 - 4\pi y(2m\nu - \mu) - \frac{4\pi my^2}{\eta}} \,\mathrm{d}\nu = \frac{1}{2\sqrt{\pi m\eta}} \int_{-\infty}^{\infty} e^{-\rho^2} \,\mathrm{d}\rho = \frac{1}{2\sqrt{m}} \frac{1}{\sqrt{\eta}}$$

From this, we compute the double integral (55) as

(57)
$$\int_{0}^{\eta} \int_{-\infty}^{\infty} e^{-4\pi m\eta \left(\nu - \frac{\mu}{2m}\right)^2 - 4\pi y(2m\nu - \mu) - \frac{4\pi my^2}{\eta}} \, \mathrm{d}\nu \, \frac{\mathrm{d}y}{\sqrt{\eta}} = \frac{1}{2\sqrt{m}} \frac{1}{\sqrt{\eta}} \int_{0}^{\eta} \frac{\mathrm{d}y}{\sqrt{\eta}} = \frac{1}{2\sqrt{m}} \frac{1}{\sqrt{\eta}} \frac{1}{\sqrt{\eta}} \frac{\mathrm{d}y}{\sqrt{\eta}} = \frac{1}{2\sqrt{\eta}} \frac{1}{\sqrt{\eta}} \frac{\mathrm{d}y}{\sqrt{\eta}} = \frac{1}{2\sqrt{m}} \frac{1}{\sqrt{\eta}} \frac{1}{\sqrt{\eta}} \frac{\mathrm{d}y}{\sqrt{\eta}} = \frac{1}{2\sqrt{m}} \frac{1}{\sqrt{\eta}} \frac{1}{\sqrt{\eta}} \frac{\mathrm{d}y}{\sqrt{\eta}} = \frac{1}{2\sqrt{m}} \frac{1}{\sqrt{\eta}} \frac{\mathrm{d}y}{\sqrt{\eta}} = \frac{1}{2\sqrt{m}} \frac{1}{\sqrt{\eta}} \frac{1}{\sqrt{\eta}} \frac{\mathrm{d}y}{\sqrt{\eta}} = \frac{1}{2\sqrt{m}} \frac{1}{\sqrt{\eta}} \frac{\mathrm{d}y}{\sqrt{\eta}} = \frac{1}{2\sqrt{m}} \frac{1}{\sqrt{\eta}} \frac{1}{\sqrt{$$

For the integral (56), we find in a similar way

(58)
$$\int_{0}^{\eta} e^{-\frac{\pi\eta\mu^{2}}{m}+4\pi y\mu-\frac{4\pi my^{2}}{\eta}} \frac{\mathrm{d}y}{\sqrt{\eta}} \leq \int_{-\infty}^{\infty} e^{-\left(-\sqrt{\frac{\pi\eta}{m}}\mu+2\sqrt{\frac{\pi m}{\eta}}y\right)^{2}} \frac{\mathrm{d}y}{\sqrt{\eta}} = \frac{1}{2\sqrt{m}}.$$

Adding up the bounds (57) and (58), yields

$$\|\vartheta_{\mu,m}(\tau,\cdot)\|_{\mathrm{L}^{2},\,\mathcal{L}_{\tau}^{\otimes m}}^{2} \leq \frac{1}{2\sqrt{m}} \int_{0}^{\eta} \frac{\mathrm{d}y}{\eta} = \frac{1}{\sqrt{m}},$$

which proves the claim.

Theorem 4.4. For $k \in \mathbb{Z}_{\geq 5}$ and $m \in \mathbb{Z}_{\geq 1}$, let $f \in J_{k,m}^{\text{cusp}}(\Gamma_0)$ be a Jacobi cusp form of weight k and index m for the full modular group $\Gamma_0 = \text{SL}_2(\mathbb{Z})$, which is normalized with respect to the Petersson inner product. Then, we have the bound

(59)
$$\sup_{(\tau,z)\in\mathbb{H}\times\mathbb{C}} \|f(\tau,z)\|_{\mathrm{Pet}}^2 = O_{\epsilon}\left(k^{\frac{3}{2}}m^{3+\epsilon}\right),$$

where the implied constant depends only on the choice of $\epsilon > 0$.

Proof. The proof of the theorem follows from combining Propositions 4.1-4.3.

Remark 4.5. The bound (59) is polynomial in k and m, and thus improves W. Kohnen's bound (1), which is exponential in k. Comparing our bound with the one obtained by P. Anamby and S. Das in [AD23], there is an improvement with regard to the polynomial growth in k, while the polynomial growth in m is slightly worse.

Acknowledgements

The first-named author expresses his gratitude to the Departments of Mathematics of Technical University of Darmstadt and of Humboldt University of Berlin for their hospitality, where this article was realized. The first-named author also acknowledges support from the INSPIRE research grant DST/INSPIRE/04/2015/002263 and the SERB Matrics grant MTR/2018/000636. The first-named author also would like to thank Prof. D. S. Nagaraj for helpful discussions. The second-named author acknowledges support from the DFG Cluster of Excellence MATH+. The third-named author acknowledges support from the LOEWE research unit Uniformized Structures in Arithmetic and Geometry. Lastly, all the three authors would like to acknowledge the hospitality of the International Centre for Theoretical Sciences (ICTS) during a visit for participating in the program Algebraic and Analytic Aspects of Automorphic Forms (Code: ICTS/aforms2019/02), where this project was initiated.

References

- [AD23] P. Anamby and S. Das, Jacobi forms, Saito-Kurokawa lifts, their pullbacks and sup-norms on average, Res. Math. Sci. 10, 14 (2023).
- [AM17] A. Aryasomayajula and P. Majumder, Off-diagonal estimates of the Bergman kernel on hyperbolic Riemann surfaces of finite volume, Proc. Amer. Math. Soc. 146 (2018), 4009–4020.
- [AM18] A. Aryasomayajula and P. Majumder, Off-diagonal estimates of the Bergman kernel on hyperbolic Riemann surfaces of finite volume-II, Ann. Fac. Sci. Toulouse Math. 29 (2020), 795–804.
- [BH11] V. Blomer and R. Holowinsky, Bounding sup-norms of cusp forms of large level, Invent. Math. 179 (2010), 645–681.
- [Bo96] T. Bouche, Asymptotic results for hermitian line bundles over complex manifolds: The heat kernel approach, In: Higher-dimensional complex varieties, 67–81, de Gruyter, Berlin, 1996.
- [EZ85] M. Eichler and D. Zagier, The theory of Jacobi cusp forms, Progress in Math. 55, Birkhäuser, 1985.
- [FJK16] J. Friedman, J. Jorgenson, and J. Kramer, Uniform sup-norm bounds on average for cusp forms of higher weights, Arbeitstagung Bonn 2013, 127–154, Progr. Math. 319, Birkhäuser/Springer, Cham, 2016.
- [FJK19] J. Friedman, J. Jorgenson, and J. Kramer, Effective sup-norm bounds on average for cusp forms of even weight, Trans. Amer. Math. Soc. 372 (2019), 7735–7766.
- [JL95] J. Jorgenson and R. Lundelius, Convergence of the heat kernel and the resolvent kernel on degenerating hyperbolic Riemann surfaces of finite volume, Quaestiones Math. 18 (1995), 345– 363.
- [Ko93] W. Kohnen, Estimates for Fourier coefficients of Siegel cusp forms of degree two, Compositio Math. 87 (1993), 231–240.
- [Kr91] J. Kramer, A geometrical approach to the theory of Jacobi cusp forms, Compositio Math. 79 (1991), 1–19.

Indian Institute of Science Education and Research, Tirupati, Karakambadi Road, Mangalam (P.O.) Tirupati-517507, Andhra Pradesh, India

Email address: anilatmaja@gmail.com

Institut für Mathematik, Humboldt-Universität zu Berlin, Unter den Linden 6, D-10099 Berlin, Germany

Email address: kramer@math.hu-berlin.de

FACHBEREICH MATHEMATIK UND STATISTIK, UNIVERSITÄT KONSTANZ, UNIVERSITÄTSSTRASSE 10, D-78464 KONSTANZ, GERMANY

Email address: anna.pippich@uni-konstanz.de