

Non-trivial fixed point of a ψ_d^4 fermionic theory, II. Anomalous exponent and scaling operators

Alessandro Giuliani¹, Vieri Mastropietro², Slava Rychkov³, Giuseppe Scola⁴

April 24, 2024

Abstract

We consider the Renormalization Group (RG) fixed-point theory associated with a fermionic ψ_d^4 model in $d = 1, 2, 3$ with fractional kinetic term, whose scaling dimension is fixed so that the quartic interaction is weakly relevant in the RG sense. The model is defined in terms of a Grassmann functional integral with interaction V^* , solving a fixed-point RG equation in the presence of external fields, and a fixed ultraviolet cutoff. We define and construct the field and density scale-invariant response functions, and prove that the critical exponent of the former is the naive one, while that of the latter is anomalous and analytic. We construct the corresponding (almost-)scaling operators, whose two point correlations are scale-invariant up to a remainder term, which decays like a stretched exponential at distances larger than the inverse of the ultraviolet cutoff. Our proof is based on constructive RG methods and, specifically, on a convergent tree expansion for the generating function of correlations, which generalizes the approach developed by three of the authors in a previous publication [1].

Contents

1	Introduction	2
2	The model and the main results	5
2.1	Generating function of the scale-invariant response functions	5
2.2	The RG map	7
2.2.1	The fixed point potential in the non-interacting case	9
2.3	Scaling dimension, relevant and irrelevant operators	11
2.4	Trimmed sequences	12
2.5	The fixed point potential	14
2.6	Almost-scaling operators	14
2.7	Main results	15
2.8	Structure of the proof	16

¹Università degli Studi Roma Tre, Dipartimento di Matematica e Fisica, L.go S. L. Murialdo 1, 00146 Roma (Italy), alessandro.giuliani@uniroma3.it

²Università degli Studi di Milano, Dipartimento di Matematica, Via C. Saldini 50, 20133 Milano (Italy), vieri.mastropietro@unimi.it

³Institut des Hautes Études Scientifiques, 35 Rte de Chartres, 91440 Bures-sur-Yvette (France), slava@ihes.fr

⁴Scuola Internazionale Superiore di Studi Avanzati, Mathematics Area, Via Bonomea 265, 34136 Trieste (Italy), giuseppescola89@gmail.com

3 Renormalization map and fixed point equation	17
3.1 Integrating out	17
3.2 Trimming	18
3.2.1 Case $(n, m, l) = (1, 0, 1)$	19
3.2.2 Case $(n, m, l) = (0, 1, 2)$	19
3.3 Fixed point equation	20
4 Solution to the FPE via the tree expansion	21
4.1 The FPE for $\ell \in L'$	22
4.2 The FPE for the local terms. Analyticity of the scaling exponents.	27
4.2.1 The scaling exponent Δ_1	28
4.2.2 The scaling exponent Δ_2	29
5 Pointwise bounds on the response functions	30
6 Stretched exponential decay of the correction terms $\mathcal{E}_1, \mathcal{E}_2$	40
A First order computation of ζ_2	49
B Proof of (2.43) and (2.44)	50
Acknowledgements	54

1 Introduction

Since the seminal works of Kadanoff, Wilson, Fisher and others in the 1970s, renormalization group (RG) fixed points and renormalization group flows connecting them are among central objects of study of theoretical physics, with many experimental applications. Only a small part of conjectures and intuitions accumulated about RG fixed points in theoretical physics have been put on rigorous mathematical footing. Pioneering results in this area were obtained by Gawedzki and Kupiainen, who in [2] showed non-perturbative stability of the Gaussian fixed point of a scalar field in four dimensions ($d = 4$), while in [3] they could follow, non-perturbatively, renormalization group flow of the Gross-Neveu two-dimensional model from the Gaussian fixed point of the Dirac field at short distances to some intermediate distance scale where the four-fermion coupling is still small. Later on [4] they constructed a non-perturbative interacting fixed point in a fractional Gross-Neveu model, which has the fermion propagator $\not{k}/k^2$ replaced by $\not{k}/|k|^{2-\epsilon}$. This ϵ serves as a small parameter which is somewhat similar to the Wilson-Fisher deviation from the integer number of dimensions. Fractional Gross-Neveu model instead lives in the integer number of dimension, but has long-ranged interactions. Analogous fractional model can be considered also for a scalar bosonic field, where they should describe critical points of the long range Ising model [5, 6, 7]. Fixed points in fractional bosonic models were also rigorously constructed [8, 9, 10].

The present paper is the second one in the series started by our recent work [1]. In [1], three of us considered a fermionic model closely related to the fractional Gross-Neveu model of Gawedzki and Kupiainen. In our fractional symplectic fermion model, the fields are spinless fermions ψ_a in d dimensions ($d = 1, 2, 3$) with propagator $\Omega_{ab}/|k|^{d/2+\epsilon}$ where Ω is a symplectic $N \times N$ matrix, and N is assumed even. The model is symmetric with respect to the symplectic group $Sp(N)$

and the bare potential includes the invariant mass and the quartic coupling terms. In this model, Ref. [1] demonstrated the existence of an interacting fixed point for any ϵ sufficiently small, which could be positive or negative, or in fact complex. One of the main results of Ref. [1] is that the fixed point potential depends analytically on ϵ in a small disk $|\epsilon| < \epsilon_0$. This interesting analyticity property should be a general feature of fixed points of fermionic models, but it was overlooked in prior literature. For bosonic models no such property holds, as they are well defined only for a positive quartic coupling. For fermionic models there is no such contradiction, since the quartic interaction may have any sign for fermions.

As discussed in [1, Section 8], the fractional symplectic fermion model of [1] is an excellent theoretical laboratory to study various aspects of the RG. In longer term perspective, one would also like to test the predictions of Conformal Field Theory (CFT) for the fixed point theory of the RG map, and possibly prove them starting from a microscopic model. We refer here, in particular, to conformal invariance of the correlation functions of suitable local operators $\mathcal{O}_i$ and the validity of the Operator Product Expansion (OPE), which are among the basic CFT axioms, see e.g. [11].

The focus of [1] was on the construction of the fixed point potential, which is not directly observable. In this second paper of the series we continue the study of the same model, making contact with physically interesting quantities such as critical exponents and correlation functions. Our goals will be twofold. On the one hand, we will show that the exponents in our model are analytic in a small disk $|\epsilon| < \epsilon_0$. This should be contrasted with the critical exponents of the fractional ϕ_d^4 theory [8, 9, 10], whose expansion in ϵ is expected to be asymptotic and conjectured to be Borel summable; note, however, that their regularity properties have not been investigated yet: current results are limited to the lowest order computation of a few non-trivial, anomalous, exponents [10].¹

On the other hand, we will make progress on clarifying the rigorous meaning of the so called scaling operators, formally defined as densities of eigenstates of the renormalization group transformation linearized around the fixed point. These operators are sometimes also called eigenoperators, see e.g. [21], Eq. (3.10). Scaling operators are used in the theoretical physics literature since the 1970s, but to our knowledge it is the first time they appear in a rigorous mathematical study of a non-integrable interacting field theory. In the special context of Liouville field theory, a full probabilistic construction of scale-invariant correlation functions has been achieved in a remarkable series of papers [22, 23, 24]. However, integrability of Liouville theory plays a key role in the construction of its correlation functions, and it is important to introduce more robust methods for a microscopic computation of scaling operators. Scaling operators are so called because they are believed to have a well-defined scaling dimension and exactly scale-invariant correlation functions. This is important for connecting the RG framework to another framework for understanding the critical state, the one based on the CFT axioms (or bootstrap axioms) in d dimensions. An outline of CFT/bootstrap axioms can be found e.g. in [11] (in physics language) or in [25, 26] (in a more mathematical language).

¹Other previous results on critical exponents of interacting field theories in the mathematical physics literature include the following. In the context of models with marginally irrelevant interactions, like ϕ_4^4 theories, the large-distance asymptotic behavior of correlation functions is characterized by logarithmic corrections with specific, fractional, exponents, computed in [12, 13, 14]; similar logarithmic corrections with fractional exponents are expected to characterize the short-distance asymptotic behavior of fermionic models with marginally relevant interactions, like the Gross-Neveu model in $d = 2$: its exponents can be computed via a generalization of [3, 15, 16], even though they have not been explicitly worked out. In the context of 2D models with ‘marginally marginal’ interactions, i.e., marginal interactions with asymptotically vanishing beta function [17, 18], the exponents displayed by the large-distance behavior of correlation functions are analytic in the interaction strength in a small disk in the complex plane; universality for these exponents have been proved, as well, in the sense that extended scaling relations are verified [19, 20].

The very first assumption of CFT is that there is a basis of local operators of the theory given by operators of well-defined scaling dimension, i.e. whose correlation functions are exact power laws. For the CFT and RG frameworks to be consistent, basis operators of CFT should be the same as the scaling operators of RG. In this paper, we focus on the fixed point (FP) theory defined in terms of a FP potential in the presence of external source fields and a reference Gaussian part with *fixed ultraviolet cutoff*. While it is standard to consider the generating function in the presence of linearly-coupled external fields, a novelty of our approach is to find a FP potential which remains exactly invariant after an RG step (and not just at the linear level), after an appropriate rescaling of the external source fields. As usual, correlation functions of our FP theory are obtained by first differentiating with respect to the external source fields and then by setting them to zero. In this setting, we have not been able to find exact scaling operators, and we actually believe that they do not exist, due to the presence of a short-distance cutoff. However, we save the day by identifying almost-scaling operators, whose correlation functions are scale-invariant up to remainder terms decreasing as a stretched exponential at large distances – that is our second main result. This is to be contrasted with the behavior of correlation functions of generic (non-scaling) local operators which are expected to decay like power laws at large distances, up to remainder terms decaying also like power laws, albeit with a larger power than the leading term. It appears plausible that generic local operators should be expressible as (infinite) linear combinations of almost-scaling operators, each of which contributes to the correlation function with a power law behavior with a different critical exponents. Such an expansion in operators with larger and larger scaling dimension is standard in physics considerations of critical phenomena, see e.g. [21, Eq. (3.11)] or [27, Eq. (3.7)]. We expect that our techniques will allow us to rigorously justify such expansion, but this remains to be done.

We stress that in our paper we choose to work with a fixed ultraviolet cutoff. This is essential to observe power law sub-leading corrections and, therefore, to be able to distinguish non-scaling from almost-scaling operators. Of course, for a critical theory, as the one we are looking at here, there is always an easy way to send the ultraviolet cutoff to infinity, by rescaling distances while simultaneously rescaling the fields appropriately, and taking the limit of the correlation functions we compute. When doing this, any sub-leading correction, being it power law or stretched exponential, would point-wise tend to zero. So we don't do this in our work.

Outline. The paper is structured as follows: In Section 2 we define the model and state our main results, summarized in Theorems 2.3 and 2.4. In Section 3 we describe in detail the RG map and derive the corresponding FP equation in the presence of the external source fields. In Section 4 we solve the FP equation for the potential with external source fields via a tree expansion, and prove its absolute convergence in a weighted L^1 norm. In Section 5 we prove the convergence of the tree expansion in a mixed L^1/L^∞ norm, required for the control of the correlation functions at fixed positions, and we conclude the proof of Theorem 2.3. In Section 6 we adapt the discussion of the previous section to the estimate of the remainder terms in the correlation functions of the almost-scaling operators and conclude the proof of Theorem 2.4. The lowest order computation of the anomalous critical exponent and of the two point function are collected in two appendices.

Many aspects of the proof are generalizations of the methods introduced and discussed in detail in [1]: therefore, we will often refer the reader to specific sections of our previous paper and emphasize the analogies and novelties of the present work compared to [1].

2 The model and the main results

2.1 Generating function of the scale-invariant response functions

Let us recall that in [1] we constructed the non-trivial FP potential $H = H^*$ for a d -dimensional Grassmann theory with partition function $\int d\mu_{\leq 0}(\psi) e^{H(\psi)}$, where ψ_a is an N -component (with $N \geq 4$ even and different from 8) Grassmann field in $\mathcal{V} \subset \mathbb{R}^d$ with $d = 1, 2, 3$ and $\mathcal{V}$ a cube whose side is eventually sent to ∞ , and $d\mu_{\leq 0}$ (denoted $d\mu_P$ in [1]) is a reference Gaussian integration characterized by the following two-point function, or *propagator*²:

$$\begin{aligned} G_{ab}^{(\leq 0)}(x-y) &= \int d\mu_{\leq 0}(\psi) \psi_a(x) \psi_b(y) = \Omega_{ab} \int \frac{d^d k}{(2\pi)^d} \frac{\chi(k)}{|k|^{d/2+\epsilon}} e^{ik \cdot (x-y)} \\ &\equiv \Omega_{ab} P_{\leq 0}(x-y). \end{aligned} \quad (2.1)$$

Here Ω is the standard $N \times N$ symplectic matrix in block-diagonal form (see [1, Eq.(1.3)]), and the cutoff function χ is radial, monotone decreasing in the radial direction, and such that

$$\chi(k) = \begin{cases} 1, & (|k| \leq 1/2) \\ 0, & (|k| \geq 1). \end{cases} \quad (2.2)$$

As in [1], we assume that χ belongs to the Gevrey class G^s , $s > 1$ [1, App. A]. The FP potential constructed in [1] reads:

$$H^*(\psi) = \nu^* \int d^d x \psi^2(x) + \lambda^* \int d^d x \psi^4(x) + H_{\text{IRR}}^*(\psi), \quad (2.3)$$

where ν^*, λ^* are non-zero real-analytic functions of ϵ , of order ϵ , for ϵ sufficiently small, $\psi^2(x) := \sum_{a,b=1}^N \Omega_{ab} \psi_a(x) \psi_b(x)$, $\psi^4(x) := (\psi^2(x))^2$, and H_{IRR}^* is an infinite sum of irrelevant terms, whose kernels are non-vanishing real-analytic functions of ϵ (see [1] for more precise claims).

In this paper, we consider the FP theory in the presence of external source fields and the corresponding generating function for correlations, formally defined as

$$W^*(\phi, J) = \log \frac{\int d\mu_{\leq 0}(\psi) e^{V^*(\phi, J, \psi)}}{\int d\mu_{\leq 0}(\psi) e^{H^*(\psi)}}. \quad (2.4)$$

Here V^* is a solution of the FP equation $V^* = RV^*$, with R the RG map, of the form

$$V^*(\phi, J, \psi) = H^*(\psi) + (\phi, \psi) + (J, \psi^2) + \mathcal{R}^*(\phi, J, \psi) + \mathcal{S}^*(\phi, J), \quad (2.5)$$

where (ϕ, ψ) and (J, ψ^2) are shorthand notations for $\sum_{a=1}^N \int d^d x \phi_a(x) \psi_a(x)$ and $\int d^d x J(x) \psi^2(x)$, respectively, $\mathcal{R}^*(\phi, J, \psi)$ is a sum of irrelevant terms depending explicitly both on (ϕ, J) and on

²In finite volume $\mathcal{V}$ the symbol $\int \frac{d^d k}{(2\pi)^d}$ should be understood a shorthand notation for the corresponding Riemann sum of discrete allowed momenta. As discussed in [1, Appendix H], the RG map leading to the definition of H^* can be defined in finite volume and proved to converge strongly (as an operator acting on the Banach space of interactions defined in [1, Section 4]) to a well-defined infinite volume counterpart. An analogous discussion can be repeated in the generalized setting considered in this paper, where external fields and correlation functions are considered. In the following we will always write all the relevant equations directly in the infinite volume setting and leave the details of the proof of convergence of the finite volume theory to its thermodynamic limit to the reader.

ψ , i.e., $\mathcal{R}^*(\phi, J, 0) = \mathcal{R}^*(0, 0, \psi) = 0$, and $\mathcal{S}^*(\phi, J)$ is an external potential, depending only upon the external fields (ϕ, J) , such that $\mathcal{S}^*(0, 0) = 0$. The precise definition of the RG map and of the irrelevant part of V^* will be given shortly, see Definition 2.2 below. Note also that, a priori, the right side of (2.4) may be plagued by infrared divergences. Therefore, (2.4) should be interpreted as:

$$\begin{aligned} W^*(\phi, J) &:= \lim_{h \rightarrow -\infty} W_{[h, 0]}^*(\phi, J) \\ W_{[h, 0]}^*(\phi, J) &:= \log \frac{\int d\mu_{[h, 0]}(\psi) e^{V^*(\phi, J, \psi)}}{\int d\mu_{[h, 0]}(\psi) e^{H^*(\psi)}}, \end{aligned} \quad (2.6)$$

where $d\mu_{[h, 0]}$ is the Grassmann Gaussian integration with propagator³

$$\begin{aligned} G_{ab}^{[h, 0]}(x - y) &= \Omega_{ab} \int \frac{d^d k}{(2\pi)^d} \frac{\chi(k) - \chi(\gamma^{-h} k)}{|k|^{d/2+\epsilon}} e^{ik \cdot (x-y)} \\ &\equiv \Omega_{ab} P_{[h, 0]}(x - y), \end{aligned}$$

and $\gamma > 1$ is an arbitrarily chosen rescaling parameter, which is fixed once and for all.

By differentiating W^* with respect to (w.r.t.) the external source fields, we define the field-field and density-density *response functions*:

$$\mathcal{G}_{ab}^*(x) := \frac{\delta^2 W^*(\phi, 0)}{\delta \phi_a(x) \delta \phi_b(0)} \Big|_{\phi=0}, \quad \mathcal{F}^*(x) := \frac{\delta^2 W^*(0, J)}{\delta J(x) \delta J(0)} \Big|_{J=0}, \quad (2.7)$$

which are the central objects of interest of this paper. These response functions are expected⁴ to have the same large distance asymptotic behavior, up to a finite multiplicative renormalization, as the two-point functions

$$\langle \psi_a(x) \psi_b(0) \rangle_{H^*}, \quad \langle \psi^2(x); \psi^2(0) \rangle_{H^*}, \quad (2.8)$$

respectively, where $\langle \cdots \rangle_{H^*} := \lim_{h \rightarrow -\infty} \frac{\int d\mu_{[h, 0]}(\psi) e^{H^*(\psi)} \cdots}{\int d\mu_{[h, 0]}(\psi) e^{H^*(\psi)}}$ (in the second expression, the semicolon indicates truncated, or connected, expectation, i.e., $\langle A; B \rangle_{H^*} := \langle AB \rangle_{H^*} - \langle A \rangle_{H^*} \langle B \rangle_{H^*}$); in particular, they are expected to define the same critical exponents. However, contrary to the two-point functions in (2.8), the response functions $\mathcal{G}^*(x)$ and $\mathcal{F}^*(x)$ are *scale-invariant*, see Theorem 2.3 below. This may look surprising, at first sight, due to the presence of a fixed ultraviolet cutoff in our theory; but it turns out that the very definition of fixed point potential V^* fixes the irrelevant terms $\mathcal{R}^*$ and the external potential $\mathcal{S}^*$ in a special way, so that the resulting response functions are, in fact, scale invariant. Even in the non-interacting theory corresponding to the Gaussian fixed point, the mechanism fixing these additional terms is not completely trivial: it is in fact instructive, as a preliminary exercise, to compute V^* and $\mathcal{G}^*, \mathcal{F}^*$ in the non-interacting case, see Section 2.2.1.

³More generally, for any pair of integers $h_1 < h_2$, we let $G_{ab}^{[h_1, h_2]}(x) := \Omega_{ab} \int \frac{d^d k}{(2\pi)^d} \frac{\chi(\gamma^{-h_2} k) - \chi(\gamma^{-h_1} k)}{|k|^{d/2+\epsilon}} e^{ik \cdot x} \equiv \Omega_{ab} P_{[h_1, h_2]}(x)$. We also denote $G_{ab}^{(\leq h)} := \lim_{h_1 \rightarrow -\infty} G_{ab}^{[h_1, h]}(x) \equiv \Omega_{ab} P_{\leq h}(x)$ and $G_{ab}^{(\geq h)} := \lim_{h_2 \rightarrow +\infty} G_{ab}^{[h, h_2]}(x) \equiv \Omega_{ab} P_{\geq h}(x)$. If $h_2 = h_1 + 1 \equiv h$, we denote the *single-scale* propagator by $g_{ab}^{(h)}(x) := G_{ab}^{[h-1, h]}(x)$.

⁴In this paper, we won't discuss how to compute the two-point functions and how to prove that they have the same asymptotic behavior as the response functions: this follows from a stability analysis of the RG flow in the vicinity of the fixed point potential, which is deferred to a third paper in this series. In such a third paper, we will also prove the independence of the fixed point potential from the scale parameter γ entering the definition of the RG map, see next section, and, most importantly, the universality of the critical exponents with respect to the choice of the ultraviolet cutoff function χ in (2.2).

2.2 The RG map

In order to clarify the notion of FP potential V^* in (2.6), we need, first of all, to specify the action of the RG map R . Before doing this formally, let us remind the reader the logic behind the definition of the R , which maps a potential V to a new potential V' as follows (this discussion parallels the one in Section 2.1 of [1]): decomposing

$$G^{(\leq 0)}(x) = G^{(\leq -1)}(x) + g^{(0)}(x), \quad (2.9)$$

see footnote 3 for the definitions of $G^{(\leq -1)}$ and $g^{(0)}$, we write ('addition principle for Gaussian integrals'):

$$\frac{1}{\mathcal{N}_{\leq 0}} \int d\mu_{\leq 0}(\psi) e^{V(\phi, J, \psi)} F(\psi) = \frac{1}{\mathcal{N}_{\leq 0}} \int d\mu_{\leq -1}(\psi_\gamma) \int d\mu_0(\psi') e^{V(\phi, J, \psi_\gamma + \psi')} F(\psi_\gamma + \psi'), \quad (2.10)$$

where $\mathcal{N}_{\leq 0} = \int d\mu_{\leq 0}(\psi) e^{V(0, 0, \psi)}$ is a normalization constant, and $d\mu_{\leq -1}$ (resp. μ_0) is the Grassmann Gaussian integration with propagator $G^{(\leq -1)}$ (resp. $g^{(0)}$). If $F(\psi)$ is a 'large scale observable', i.e., if it does not depend on the Fourier modes $\hat{\psi}(k)$ with k in the support of $\hat{g}^{(0)}(k)$, then

$$\begin{aligned} \frac{1}{\mathcal{N}_{\leq 0}} \int d\mu_{\leq 0}(\psi) e^{V(\phi, J, \psi)} F(\psi) &= \frac{1}{\mathcal{N}_{\leq 0}} \int d\mu_{\leq -1}(\psi_\gamma) F(\psi_\gamma) \int d\mu_0(\psi') e^{V(\phi, J, \psi_\gamma + \psi')} \\ &= \frac{\mathcal{N}_0}{\mathcal{N}_{\leq 0}} \int d\mu_{\leq -1}(\psi_\gamma) e^{V_{\text{eff}}(\phi, J, \psi_\gamma)} F(\psi_\gamma), \end{aligned} \quad (2.11)$$

where $\mathcal{N}_0 := \int d\mu_0(\psi) e^{V(0, 0, \psi)}$ and $V_{\text{eff}}(\phi, J, \psi_\gamma) = \log(\mathcal{N}_0^{-1} \int d\mu_0(\psi') e^{V(\phi, J, \psi_\gamma + \psi')})$. Thanks to the scaling property $G^{(\leq -1)}(x) = \gamma^{-2[\psi]} G^{(\leq 0)}(x/\gamma)$, with $[\psi] = \frac{d}{4} - \frac{\epsilon}{2}$ (see [1, eqs.(2.14),(2.15)]), the random field ψ_γ has the same distribution as $\gamma^{-[\psi]} \psi(\cdot/\gamma) \equiv D\psi$, with D the dilatation operator, ψ a random field with distribution $d\mu_{\leq 0}$, so that

$$(2.11) = \frac{1}{\mathcal{N}'_{\leq 0}} \int d\mu_{\leq 0}(\psi) e^{V_{\text{eff}}(\phi, J, D\psi)} F(D\psi),$$

where $\mathcal{N}'_{\leq 0} := \int d\mu_{\leq 0}(\psi) e^{V_{\text{eff}}(0, 0, D\psi)} \equiv \mathcal{N}_{\leq 0}/\mathcal{N}_0$. Now, if V has the same structure as (2.5), i.e.,

$$V(\phi, J, \psi) = H^*(\psi) + (\phi, \psi) + (J, \psi^2) + \mathcal{R}(\phi, J, \psi) + \mathcal{S}(\phi, J), \quad (2.12)$$

with $\mathcal{R}$ and $\mathcal{S}$ two potentials in an $O(\epsilon)$ neighborhood of the non-interacting fixed point discussed in Section 2.2.1 below (with $\mathcal{R}$ a sum of irrelevant terms, in the sense discussed in Section 2.3 below, and $\mathcal{S}$ an external potential, depending only upon the external fields ϕ, J), then it turns out, see Section 3, that

$$V_{\text{eff}}(\phi, J, D\psi) = H^*(\psi) + \tilde{Z}_1(\phi, D\psi) + \tilde{Z}_2(J, (D\psi)^2) + \tilde{\mathcal{R}}(\phi, J, D\psi) + \tilde{\mathcal{S}}(\phi, J), \quad (2.13)$$

with $\tilde{Z}_1 = 1 + O(\epsilon)$, $\tilde{Z}_2 = 1 + O(\epsilon)$ two constants produced by the integration of the fluctuation field ψ' , and $\tilde{\mathcal{R}}, \tilde{\mathcal{S}}$ two new potentials that are, again, $O(\epsilon)$ -close to the non-interacting fixed point, with $\tilde{\mathcal{R}}$ a sum of irrelevant terms. In view of (2.13), it is natural to rescale the external fields in a way similar to ψ , in such a way to recast the terms involving the external fields in a form as close as possible to the one in which they originally appeared in V : therefore, we

let $\phi \equiv D\phi'$ and $J \equiv DJ'$, with D the dilatation operator, acting on the external fields as: $D\phi(x) = \gamma^{-[\phi]}\phi(x/\gamma)$ and $DJ(x) = \gamma^{-[J]}J(x/\gamma)$, with the scaling dimensions $[\phi], [J]$ to be fixed appropriately, in a way explained shortly. In terms of these definitions: $(\phi, D\psi) = (D\phi', D\psi) = \sum_{a=1}^N \gamma^{-[\phi]-[\psi]} \int d^d x \phi'_a(x/\gamma) \psi_a(x/\gamma) = \gamma^{d-[\phi]-[\psi]}(\phi', \psi)$, and similarly for $(J, (D\psi)^2)$. Therefore,

$$\begin{aligned} V_{\text{eff}}(\phi, J, D\psi) &= V_{\text{eff}}(D\phi', DJ', D\psi) \\ &= H^*(\psi) + \tilde{Z}_1 \gamma^{d-[\phi]-[\psi]}(\phi', \psi) + \tilde{Z}_2 \gamma^{d-[J]-2[\psi]}(J', \psi^2) + \tilde{\mathcal{R}}(D\phi', DJ', D\psi) + \tilde{\mathcal{S}}(D\phi', DJ') \\ &\equiv V'(\phi', J', \psi). \end{aligned} \tag{2.14}$$

By definition, V' is the image of V under the RG map, i.e., $V' = RV$; more precisely, V' is *equivalent*, in the sense of [1, Section 5.2.1], to RV : the missing ingredient, in the previous informal discussion, is the action of the trimming map T , which is an operator, equivalent to the identity, which isolates the relevant terms from the irrelevant ones, see Section 3.2 below.

Note that the RG map depends, among other things, upon the choice of the scaling dimensions $[\phi]$ and $[J]$. In order to fix these dimensions, we use the requirement that, at the fixed point, $Z_1 \gamma^{d-[\phi]-[\psi]} = Z_2 \gamma^{d-[J]-2[\psi]} = 1$, with $Z_1 = \tilde{Z}_1^*$ and $Z_2 = \tilde{Z}_2^*$ the constants as in (2.13)-(2.14) associated with the fixed point potential V^* . In this way, the fixed point equation (FPE) defines an unambiguous criterium for assigning a scaling dimension to the external fields, which has a simple relation with the critical exponents of the corresponding response functions (cf. (2.16) with (2.41)); once that $[\phi]$ and $[J]$ are fixed, the FPE becomes an equation for $\mathcal{R}^*, \mathcal{S}^*$, of the form: $\tilde{\mathcal{R}}^*(D\phi', DJ', D\psi) = \mathcal{R}^*(\phi', J', \psi)$, $\tilde{\mathcal{S}}^*(D\phi', DJ') = \mathcal{S}^*(\phi', J')$. The uniqueness of the solution for the FPE for $[\phi], [J], \mathcal{R}^*, \mathcal{S}^*$, proved below, see Theorem 2.3, and the fact that the critical exponent of the two-point correlations (2.8) are the same as those of the response functions $\mathcal{G}^*, \mathcal{F}^*$ associated with V^* , proved in the next paper of this series, confirms the physical significance of the scaling dimensions $[\phi]$ and $[J]$, and of the fixed point potential V^* .

Given these premises, let us define the RG map R a bit more formally as follows:

$$R := D^{-(h-1)} T S_h D^h \equiv D T S_0$$

(independence from $h \in \mathbb{Z}$ is a consequence of the definitions of S_h, T, D below), where:

- S_h is the integration on scale $h \leq 0$, i.e.,

$$S_h V(\phi, J, \psi) = \log \frac{\int d\mu_h(\psi') e^{V(\phi, J, \psi' + \psi)}}{\int d\mu_h(\psi') e^{V(0, 0, \psi')}}, \tag{2.15}$$

where $d\mu_h$ is the Grassmann Gaussian integration with propagator $g_{ab}^{(h)}(x)$, see definition in footnote 3;

- T is the trimming map, defined in detail in Section 3.2 below, which is *equivalent to the identity*: when acting on a potential V it produces an equivalent potential $V' \sim V$, in the sense of [1, Sect.5.2.1], of the form $V' = TV = \mathcal{L}V + \mathcal{I}V$; here $\mathcal{L}V$ is a sum of finitely many *local* relevant and marginal terms (belonging to a finite dimensional subspace of the Banach space of potentials), and $\mathcal{I}V$ is an equivalent rewriting of $TV - \mathcal{L}V$ as an infinite linear combination of non-local, *irrelevant*, monomials in $\phi, J, \psi, \partial\psi$;

- D is the dilatation, acting on $V(\phi, J, \psi)$ as $DV(\phi, J, \psi) := V(D\phi, DJ, D\psi)$, with $D\psi(x) = \gamma^{-[\psi]}\psi(x/\gamma)$ (recall: $[\psi] = d/4 - \epsilon/2$, see [1, eqs.(2.14),(2.15)]), while $D\phi(x) = \gamma^{-[\phi]}\phi(x/\gamma)$, $DJ(x) = \gamma^{-[J]}J(x/\gamma)$, where $[\phi] = d - [\psi] + O(\epsilon)$ and $[J] = d - 2[\psi] + O(\epsilon)$ are two analytic functions of ϵ , for ϵ sufficiently small, to be determined in such a way that V^* , of the form described in Definition 2.2 below, is a solution of the FP equation $V^* = RV^*$. A simple argument, spelled out in Section 2.8 below⁵ shows that $2(d - [\phi])$ and $2(d - [J])$ are the critical exponents of the response functions $\mathcal{G}^*$ and $\mathcal{F}^*$, respectively: therefore, denoting by $2\Delta_1$ and $2\Delta_2$ these critical exponents, we rewrite the action of the dilatation operator on the external fields in terms of $\Delta_1 := d - [\phi]$, $\Delta_2 := d - [J]$:

$$D\phi(x) = \gamma^{-d+\Delta_1}\phi(x/\gamma), \quad DJ(x) = \gamma^{-d+\Delta_2}J(x/\gamma); \quad (2.16)$$

we will see that $\Delta_1 = [\psi]$ and $\Delta_2 = 2[\psi] + \eta_2(\epsilon)$, with $\eta_2(\epsilon) = 2\epsilon\frac{N-2}{N-8} + O(\epsilon^2)$.

2.2.1 The fixed point potential in the non-interacting case

Before introducing the notion of relevant and irrelevant operators, and giving a more formal definition of fixed point potential V^* , it is instructive to discuss the ‘Gaussian’ case corresponding, in the absence of external fields, to the trivial fixed point $H^*(\psi) = 0$. Even in this case, the fixed point potential in the presence of the external fields, to be denoted by V_0^* , is not trivial, and its structure is crucial to guarantee that the response functions $\mathcal{G}^*$ and $\mathcal{F}^*$ are scale-invariant, notwithstanding the presence of an ultraviolet cutoff. For $H^*(\psi) = 0$, we let $\Delta_1 = [\psi]$ and $\Delta_2 = 2[\psi]$ in (2.16), so that the action of the dilatation operator reduces to:

$$D\psi(x) = \gamma^{-\frac{d}{4}+\frac{\epsilon}{2}}\psi(x/\gamma), \quad D\phi(x) = \gamma^{-\frac{3}{4}d-\frac{\epsilon}{2}}\phi(x/\gamma), \quad DJ(x) = \gamma^{-\frac{d}{2}-\epsilon}J(x/\gamma). \quad (2.17)$$

The non-interacting fixed point potential V_0^* we consider is the one obtained as the image of the ‘naive’ potential $V_0(\phi, J, \psi) := (\phi, \psi) + (J, \psi^2)$ under the iterated action of the RG map DS_0 , in the limit as the number of iterations goes to infinity:

$$V_0^*(\phi, J, \psi) = \lim_{n \rightarrow \infty} (DS_0)^n V_0(\phi, J, \psi). \quad (2.18)$$

The action of S_0 on a potential V returns (see [1, Section 5.1]):

$$S_0 V(\phi, J, \psi) = \sum_{s \geq 1} \frac{1}{s!} \underbrace{\langle V(\phi, J, \psi + \cdot); \dots; V(\phi, J, \psi + \cdot) \rangle_0^{(0)}}_{s \text{ times}},$$

where $\langle A_1; \dots; A_s \rangle_0^{(0)}$ denotes connected expectation of order s with respect to (w.r.t.) the Gaussian integration $d\mu_0$ with propagator $g^{(0)}$ (the lower label ‘0’ indicates that the expectation is computed w.r.t. a Gaussian, non-interacting, measure, while the upper label ‘(0)’ refers to the scale of the propagator $g^{(0)}$). Using the fact that the naive potential $V_0(\phi, J, \psi)$ is invariant under the action of D , one finds that (2.18) can be re-expressed as

$$V_0^*(\phi, J, \psi) = S_{\geq 1} V_0(\phi, J, \psi), \quad (2.19)$$

⁵See in particular (2.52). Note that the discussion in Section 2.8 is written in terms of $\Delta_1 = d - [\phi]$ and $\Delta_2 = d - [J]$ rather than in terms of $[\phi], [J]$, but this is just a matter of notation.

where

$$S_{\geq 1}V(\phi, J, \psi) = \sum_{s \geq 1} \frac{1}{s!} \underbrace{\langle V(\phi, J, \psi + \cdot); \cdots; V(\phi, J, \psi + \cdot) \rangle_0}_{s \text{ times}}^{(\geq 1)}, \quad (2.20)$$

and $\langle A_1; \dots; A_s \rangle_0^{(\geq 1)}$ denotes connected expectation of order s w.r.t. the Gaussian integration $d\mu_{\geq 1}$ with propagator $G^{(\geq 1)}$, see footnote 3. As well known (see also [1, Appendix D.2]), connected expectations w.r.t. a Gaussian Grassmann measure can be expressed in terms of connected Feynman diagrams associated with the corresponding propagator. Using this fact, (2.20) implies that the fixed point potential in (2.18) is $V_0^*(\phi, J, \psi) = (\phi, \psi) + (J, \psi^2) + \mathcal{R}_0^*(\phi, J, \psi) + \mathcal{S}_0^*(\phi, J)$, where, if the two terms (ϕ, ψ) and (J, ψ^2) in V_0 are graphically represented as in Figure 1, then $\mathcal{R}_0^*$ and $\mathcal{S}_0^*$




Figure 1: The graphical representations of the two vertices (ϕ, ψ) and (J, ψ^2) , respectively.

are graphically represented as in Figure 2, with the solid lines representing the propagator $G^{(\geq 1)}$. In formulae, recalling the definition of $P_{>1}$ in footnote 3:

$$\mathcal{R}_0^*(\phi, J, \psi) =$$

$$\sum_{n \geq 1} \text{---} \bullet_{y_0} \text{---} \overset{\geq 1}{\text{wavy}} \bullet_{y_1} \text{---} \overset{\geq 1}{\text{wavy}} \bullet_{y_2} \text{---} \cdots \text{---} \bullet_{y_n} \text{---} + \sum_{n \geq 2} \text{---} \overset{\geq 1}{\text{wavy}} \bullet_{y_1} \text{---} \overset{\geq 1}{\text{wavy}} \bullet_{y_2} \text{---} \cdots \text{---} \bullet_{y_n} \text{---},$$

$$\mathcal{S}_0^*(\phi, J) =$$

$$\sum_{n \geq 0} \text{---} \bullet_{y_0} \text{---} \overset{\geq 1}{\text{wavy}} \bullet_{y_1} \text{---} \cdots \text{---} \overset{\geq 1}{\text{wavy}} \bullet_{y_n} \text{---} \text{---} \bullet_{y_{n+1}} \text{---} + \sum_{n \geq 1} \text{---} \overset{\geq 1}{\text{wavy}} \bullet_{y_0} \text{---} \overset{\geq 1}{\text{wavy}} \bullet_{y_1} \text{---} \cdots \text{---} \overset{\geq 1}{\text{wavy}} \bullet_{y_{n-1}} \text{---} \text{---} \bullet_{y_n} \text{---} \overset{\geq 1}{\text{wavy}},$$

Figure 2: The graphical representations of $\mathcal{R}_0^*$ and $\mathcal{S}_0^*$, respectively.

$$\begin{aligned}
\mathcal{R}_0^*(\phi, J, \psi) &= \sum_{m \geq 1} (-2)^m \sum_{a=1}^N \int d^d y_0 \cdots d^d y_m \phi_a(y_0) P_{\geq 1}(y_0 - y_1) J(y_1) P_{\geq 1}(y_1 - y_2) \cdots J(y_m) \psi_a(y_m) \\
&\quad + \sum_{m \geq 2} (-2)^{m-1} \sum_{a,b=1}^N \Omega_{ab} \int d^d y_1 \cdots d^d y_m \psi_a(y_1) J(y_1) P_{\geq 1}(y_1 - y_2) \cdots J(y_m) \psi_b(y_m) \\
\mathcal{S}_0^*(\phi, J) &= \sum_{m \geq 0} (-2)^{m-1} \sum_{a,b=1}^N \Omega_{ab} \int d^d y_0 \cdots d^d y_{m+1} \phi_a(y_0) P_{\geq 1}(y_0 - y_1) \cdots J(y_m) P_{\geq 1}(y_m - y_{m+1}) \phi_b(y_{m+1}) \\
&\quad + \sum_{m \geq 1} (-2)^{m-1} \frac{N}{m} \int d^d y_1 \cdots d^d y_m J(y_1) P_{\geq 1}(y_1 - y_2) \cdots J(y_m) P_{\geq 1}(y_m - y_1)
\end{aligned} \tag{2.21}$$

Therefore, the non-interacting fixed point generating function is:

$$W_0^*(\phi, J) = \mathcal{S}_0^*(\phi, J) + \log \int d\mu_{\leq 0}(\psi) e^{(\phi, \psi) + (J, \psi^2) + \mathcal{R}_0^*(\phi, J, \psi)}, \tag{2.22}$$

from which a straightforward computation shows that

$$\begin{aligned}
\mathcal{G}_{0;ab}^*(x) &:= \frac{\delta^2 W_0^*(\phi, 0)}{\delta \phi_a(x) \delta \phi_b(0)} \Big|_{\phi=0} = \Omega_{ab} [P_{\leq 0}(x) + P_{\geq 1}(x)], \\
\mathcal{F}_0^*(x) &:= \frac{\delta^2 W_0^*(0, J)}{\delta J(x) \delta J(0)} \Big|_{J=0} = -2N [P_{\leq 0}^2(x) + 2P_{\leq 0}(x)P_{\geq 1}(x) + P_{\geq 1}^2(x)],
\end{aligned} \tag{2.23}$$

which are scale invariant, as anticipated above.

2.3 Scaling dimension, relevant and irrelevant operators

Let us consider a monomial in ϕ, J, ψ of the form⁶

$$\begin{aligned}
M(\phi, J, \psi) &= \\
&= \iiint d\mathbf{x} d\mathbf{y} d\mathbf{z} \phi_{a_1}(x_1) \cdots \phi_{a_n}(x_n) J(y_1) \cdots J(y_m) \partial_{\mu_1}^{p_1} \psi_{b_1}(z_1) \cdots \partial_{\mu_l}^{p_l} \psi_{b_l}(z_l) [H_{n,m,l,\mathbf{p}}(\mathbf{x}, \mathbf{y}, \mathbf{z})]_{\mathbf{a}, \mu, \mathbf{b}}
\end{aligned} \tag{2.24}$$

with $\mathbf{x} = (x_1, \dots, x_n)$, etc., $p_i \in \{0, 1\}$, $\mu_i \in \{0, 1, \dots, d\}$ ⁷, $n + l$ even, $H_{n,m,l,\mathbf{p}}$ translationally invariant and of finite L^1 norm, i.e.,

$$\|H_{n,m,l,\mathbf{p}}\| := \max_{\mathbf{a}, \mu, \mathbf{b}} \iiint^* d\mathbf{x} d\mathbf{y} d\mathbf{z} | [H_{n,m,l,\mathbf{p}}(\mathbf{x}, \mathbf{y}, \mathbf{z})]_{\mathbf{a}, \mu, \mathbf{b}} | < +\infty \tag{2.25}$$

⁶For later reference, and in order to make contact with the notation used in [1, Section 4.1], given the n -ples $\mathbf{a}, \mathbf{x}$, the m -ple $\mathbf{y}$ and the l -ples $\mathbf{p}, \mu, \mathbf{z}$, we shall also denote $[H_{n,m,l,\mathbf{p}}(\mathbf{x}, \mathbf{y}, \mathbf{z})]_{\mathbf{a}, \mu, \mathbf{b}}$ by $H(\mathbf{a}, \mathbf{x}, \mathbf{y}, \mathbf{B}, \mathbf{z})$, with $\mathbf{B} = ((p_1, \mu_1, b_1), \dots, (p_l, \mu_l, b_l))$. Moreover, we let $\Phi(\mathbf{a}, \mathbf{x}) := \phi_{a_1}(x_1) \cdots \phi_{a_n}(x_n)$, $J(\mathbf{y}) := J(y_1) \cdots J(y_m)$ and $\Psi(\mathbf{B}, \mathbf{z}) := \partial_{\mu_1}^{p_1} \psi_{b_1}(z_1) \cdots \partial_{\mu_l}^{p_l} \psi_{b_l}(z_l)$. Without loss of generality, we can assume, and will do so from now on, that $H(\mathbf{a}, \mathbf{x}, \mathbf{y}, \mathbf{B}, \mathbf{z})$ is: anti-symmetric under simultaneous permutations of the elements of $\mathbf{a}, \mathbf{x}$, symmetric under permutations of the elements of $\mathbf{y}$; anti-symmetric under simultaneous permutations of the elements of $\mathbf{B}, \mathbf{z}$.

⁷If $p_i = 1$, then $\mu_i \in \{1, \dots, d\}$. If $p_i = 0$, the index μ_i needs not to be specified, but, conventionally, in that case we set $\mu_i = 0$. If $\mathbf{p} = \mathbf{0}$ we shall often drop the label μ .

(here the $*$ on the integral indicates that we are *not* integrating over one of the elements of $(\mathbf{x}, \mathbf{y}, \mathbf{z})$, say x_1). The action of D on $M(\phi, J, \psi)$ is defined as $DM(\phi, J, \psi) := M(D\phi, DJ, D\psi)$. After a change of variables, this can be rewritten as:

$$DM(\phi, J, \psi) = \iiint d\mathbf{x} d\mathbf{y} d\mathbf{z} \phi_{a_1}(x_1) \cdots \phi_{a_n}(x_n) J(y_1) \cdots J(y_m) \partial_{\mu_1}^{p_1} \psi_{b_1}(z_1) \cdots \partial_{\mu_l}^{p_l} \psi_{b_l}(z_l) [DH_{n,m,l,\mathbf{p}}(\mathbf{x}, \mathbf{y}, \mathbf{z})]_{\mathbf{a}, \boldsymbol{\mu}, \mathbf{b}} \quad (2.26)$$

where

$$DH_{n,m,l,\mathbf{p}}(\mathbf{x}, \mathbf{y}, \mathbf{z}) := \gamma^{\delta_{sc}(n,m,l,\mathbf{p})} H_{n,m,l,\mathbf{p}}(\gamma\mathbf{x}, \gamma\mathbf{y}, \gamma\mathbf{z}), \quad (2.27)$$

with

$$\delta_{sc}(n, m, l, \mathbf{p}) := d(n + m + l) - n(d - \Delta_1) - m(d - \Delta_2) - l[\psi] - \|\mathbf{p}\|_1, \quad (2.28)$$

defines the action of the dilatation operator D on the kernels. Note also that the L^1 norm of $H_{n,m,l,\mathbf{p}}$ rescales as follows under the action of D :

$$\|DH_{n,m,l,\mathbf{p}}\| = \gamma^{D_{sc}(n,m,l,\mathbf{p})} \|H_{n,m,l,\mathbf{p}}\|, \quad (2.29)$$

where

$$\begin{aligned} D_{sc}(n, m, l, \mathbf{p}) &:= \delta_{sc}(n, m, l, \mathbf{p}) - d(n + m + l - 1) \\ &= d - n(d - \Delta_1) - m(d - \Delta_2) - l[\psi] - \|\mathbf{p}\|_1 \end{aligned} \quad (2.30)$$

is the *scaling dimension* of $H_{n,m,l,\mathbf{p}}$. Depending on whether $D_{sc}(n, m, l, \mathbf{p})$ is positive, vanishing or negative, we say that the corresponding monomial is relevant, marginal or irrelevant. Note that, anticipating the fact that $\Delta_1 = [\psi]$ and $\Delta_2 = 2[\psi] + O(\epsilon)$, the only relevant or marginal terms⁸ with $l > 0$ are those with $(n, m, l, \mathbf{p}) = (0, 0, 2, \mathbf{0})$, $(0, 0, 4, \mathbf{0})$, $(1, 0, 1, \mathbf{0})$, $(0, 1, 2, \mathbf{0})$, and $(0, 0, 2, \mathbf{p})$ with $\|\mathbf{p}\|_1 = 1$, for which we have $D_{sc}(0, 0, 2, \mathbf{0}) = d/2 + \epsilon$, $D_{sc}(0, 0, 4, \mathbf{0}) = 2\epsilon$, $D_{sc}(1, 0, 1, \mathbf{0}) = \Delta_1 - [\psi]$, $D_{sc}(0, 1, 2, \mathbf{0}) = \Delta_2 - 2[\psi]$, and $D_{sc}(0, 0, 2, (1, 0)) = D_{sc}(0, 0, 2, (0, 1)) = d/2 + \epsilon - 1$.

2.4 Trimmed sequences

A potential $H(\phi, J, \psi)$ is an infinite sum of monomials, of the form

$$H(\phi, J, \psi) = \sum_{(n,m,l,\mathbf{p}) \in L} \sum_{\mathbf{a}, \boldsymbol{\mu}, \mathbf{b}} \iiint d\mathbf{x} d\mathbf{y} d\mathbf{z} \Phi(\mathbf{a}, \mathbf{x}) J(\mathbf{y}) \Psi(\mathbf{B}, \mathbf{z}) [H_{n,m,l,\mathbf{p}}(\mathbf{x}, \mathbf{y}, \mathbf{z})]_{\mathbf{a}, \boldsymbol{\mu}, \mathbf{b}} \quad (2.31)$$

where

$$L := \{(n, m, l, \mathbf{p}) : \mathbf{p} = (p_1, \dots, p_l) \text{ with } p_i \in \{0, 1\}, \ n, m, l, |\mathbf{p}| \geq 0, \ n + l \text{ even}, \ n + m + l \geq 1\}, \quad (2.32)$$

⁸To be precise, the relevance or irrelevance of the term with $(n, m, l, \mathbf{p}) = (0, 1, 2, \mathbf{0})$ depends upon the sign of $\eta_2 = \Delta_2 - 2[\psi]$. In any case, even if irrelevant, this term is barely so, with a scaling dimension of order $O(\epsilon)$. For this reason, it is convenient to treat it differently from the irrelevant terms with scaling dimensions well separated from 0, uniformly in ϵ . Moreover, the terms with $(0, 0, 2, \mathbf{p})$ with $\|\mathbf{p}\|_1 = 1$ are relevant only if $d = 2, 3$: in $d = 1$ they are irrelevant, with scaling dimension equal to $-1/2 + \epsilon$; however, the treatment of these terms in $d = 1$ does not present any significant simplification, as compared to the cases $d = 2, 3$. For this reason, and for uniformity of notation, we shall treat them exactly in the same way for $d = 1$ and for $d = 2, 3$, notwithstanding this difference: in particular, the notion of of trimmed sequence and of ‘trimming’ given below, which defines the way in which we ‘localize’ and ‘interpolate’ the relevant terms, is the same in $d = 1, 2, 3$.

and, given $(n, m, l, \mathbf{p}) \in L$, the sums over $\mathbf{a}$, $\boldsymbol{\mu}$ and $\mathbf{b}$ run over the sets $\{1, \dots, N\}^n$, $\{0, 1\}^l$ and $\{1, \dots, N\}^l$, respectively; moreover, we used the notations of footnote 6, with $\mathbf{B} = ((p_1, \mu_1, b_1), \dots, (p_l, \mu_l, b_l))$. The (formal) infinite sum in (2.31) should be thought of as a way of representing the collection of kernels $H_{n,m,l,\mathbf{p}}$. Therefore, in the following, we shall equivalently write a potential H either as in (2.31) or as

$$H = \{H_\ell\}_{\ell \in L}. \quad (2.33)$$

We shall say that a sequence $\{H_\ell\}_{\ell \in L}$, with translationally invariant kernels H_ℓ of finite L^1 norm, is a *trimmed sequence* if the relevant or marginal kernels with $\ell = (0, 0, 2, \mathbf{0})$, $(0, 0, 4, \mathbf{0})$, $(1, 0, 1, \mathbf{0})$, $(0, 1, 2, \mathbf{0})$, and $(0, 0, 2, \mathbf{p})$ with $\|\mathbf{p}\|_1 = 1$ have the following local structure:

$$\begin{aligned} [H_{0,0,2,\mathbf{0}}(\mathbf{z})]_{\mathbf{b}} &= c_1 \delta(z_1 - z_2) \Omega_{b_1 b_2}, \\ [H_{0,0,2,\mathbf{p}}(\mathbf{z})]_{\boldsymbol{\mu}, \mathbf{b}} &= 0, \quad \forall \mathbf{p} : \|\mathbf{p}\|_1 = 1, \\ [H_{0,0,4,\mathbf{0}}(\mathbf{z})]_{\mathbf{b}} &= c_2 \delta(z_1 - z_2) \delta(z_1 - z_3) \delta(z_1 - z_4) q_{b_1 b_2 b_3 b_4}, \\ [H_{1,0,1,\mathbf{0}}(x_1, z_1)]_{a_1, b_1} &= c_3 \delta(x_1 - z_1) \delta_{a_1, b_1}, \\ [H_{0,1,2,\mathbf{0}}(y_1, \mathbf{z})]_{\mathbf{b}} &= c_4 \delta(y_1 - z_1) \delta(z_1 - z_2) \Omega_{b_1 b_2}, \end{aligned} \quad (2.34)$$

where c_1, c_2, c_3, c_4 are constants, and $q_{abce} = \Omega_{ab} \Omega_{ce} - \Omega_{ac} \Omega_{be} + \Omega_{ae} \Omega_{bc}$ is the totally anti-symmetric tensor, as in [1, Eq.(4.4)].

As discussed in Section 3, the RG transformation R is a map from the space of trimmed sequences of kernels to itself.

Remark 2.1. The non-interacting fixed point potential $V_0^*(\phi, J, \psi) = (\phi, \psi) + (J, \psi^2) + \mathcal{R}_0^*(\phi, J, \psi) + \mathcal{S}_0^*(\phi, J)$, constructed in Section 2.2.1, can be written as $V_0^* = \{V_{0;\ell}^*\}_{\ell \in L}$, where the only non-zero kernels in the sequence are the following:

$$\begin{aligned} [V_{0;(1,m,1,\mathbf{0})}(x_1, \mathbf{y}, z_1)]_{a_1, b_1} \Big|_{\epsilon=0} &= (-2)^m \delta_{a_1, b_1} \delta(y_m - z_1) \prod_{i=0}^{m-1} P_{\geq 1}(y_i - y_{i+1}), & m \geq 0 \\ [V_{0;(0,m,2,\mathbf{0})}(\mathbf{y}, \mathbf{z})]_{b_1, b_2} \Big|_{\epsilon=0} &= (-2)^{m-1} \Omega_{b_1, b_2} \delta(z_1 - y_1) \delta(y_m - z_2) \prod_{i=1}^{m-1} P_{\geq 1}(y_i - y_{i+1}), & m \geq 1 \\ [V_{0;(2,m,0,\emptyset)}(\mathbf{x}, \mathbf{y})]_{a_1, a_2} \Big|_{\epsilon=0} &= (-2)^{m-1} \Omega_{a_1 a_2} \prod_{i=0}^m P_{\geq 1}(y_i - y_{i+1}), & m \geq 0 \\ V_{0;(0,m,0,\emptyset)}^*(\mathbf{y}) \Big|_{\epsilon=0} &= (-2)^{m-1} \frac{N}{m} \prod_{i=1}^m P_{\geq 1}(y_i - y_{i+1}), & m \geq 1 \end{aligned} \quad (2.35)$$

where: $P_{\geq 1}$ was defined in footnote 3 above; in the first two lines we dropped the $\boldsymbol{\mu}$ labels; in the first line, y_0 should be interpreted as x_1 , and, for $m = 0$, the right side should be interpreted as that of the third line of (2.34) with $c_3 = 1$; for $m = 1$, the right side of the second line should be interpreted as that of the fourth line of (2.34) with $c_4 = 1$; in the third line, y_0 and y_{m+1} should be interpreted as x_1 and x_2 , respectively; in the fourth line, y_{m+1} should be interpreted as y_1 .

In particular, V_0^* is associated with a *trimmed sequence*, such that $c_1 = c_2 = 0$ and $c_3 = c_4 = 1$.

2.5 The fixed point potential

We are now in good position to define more precisely the fixed point potential $V^*(\phi, J, \psi)$ that we construct in this paper.

Definition 2.2. $V^* = \{V_\ell^*\}_{\ell \in L}$ is the solution of the fixed point equation $V^* = RV^*$ in the space of trimmed sequences, such that: (1) $V_{0,0,l,\mathbf{p}}^* = H_{l,\mathbf{p}}^*$, with $H_{l,\mathbf{p}}^*$ the kernels of the FP potential constructed in [1]; (2) for $\ell = (1, 0, 1, \mathbf{0})$, $(0, 1, 2, \mathbf{0})$, V_ℓ^* is as in (2.34) with $c_3 = c_4 = 1$, i.e., the local terms (ϕ, ψ) and (J, ψ^2) in V^* have pre-factor equal to 1; (3) the scaling exponents Δ_1, Δ_2 entering the definition of D and, therefore, of R , are analytic in ϵ for ϵ sufficiently small, and such that $\Delta_1 - [\psi] = O(\epsilon)$, $\Delta_2 - 2[\psi] = O(\epsilon)$; (4) the kernels V_ℓ^* with $\ell \in L \setminus \{(1, 0, 1, \mathbf{0}), (0, 1, 2, \mathbf{0})\}$ are analytic in ϵ for ϵ sufficiently small and, at $\epsilon = 0$, satisfy $V_\ell^*|_{\epsilon=0} = V_{0;\ell}^*$, with $V_{0;\ell}^*$ as in (2.34) for $\ell \in \{(1, m, 1, \mathbf{0})\}_{m \geq 1} \cup \{(0, m, 2, \mathbf{0})\}_{m \geq 2} \cup \{(2, m, 0, \emptyset)\}_{m \geq 0} \cup \{(0, m, 0, \emptyset)\}_{m \geq 1}$, and equal to zero otherwise.

Note that the existence and uniqueness of the FP potential V^* specified in Definition 2.2 is part of the results proved in this paper. As anticipated above, we will write

$$V^*(\phi, J, \psi) = H^*(\psi) + (\phi, \psi) + (J, \psi^2) + \mathcal{R}^*(\phi, J, \psi) + \mathcal{S}^*(\phi, J), \quad (2.36)$$

where, letting⁹

$$L_{\text{ext},0} := \{(n, m, 0, \emptyset) \in L\}, \quad L_{\text{ext},f} := \{(n, m, l, \mathbf{p}) \in L : n + m > 0, l > 0\}, \quad (2.37)$$

and $L'_{\text{ext},f} := L_{\text{ext},f} \setminus \{(1, 0, 1, \mathbf{0}), (0, 1, 2, \mathbf{0})\}$, we let: $\mathcal{R}^* = \{V_\ell^*\}_{\ell \in L'_{\text{ext},f}}$ and $\mathcal{S}^* = \{V_\ell^*\}_{\ell \in L_{\text{ext},0}}$.

2.6 Almost-scaling operators

The response functions are not correlation functions of any local operators built out the field ψ . We can however identify two quasi-local operators $\mathcal{O}^{(1)}$ and $\mathcal{O}^{(2)}$, whose two point functions will end up being the same as $\mathcal{G}^*$ and $\mathcal{F}^*$, up to corrections that decay faster than any power at large distances. Because of these correction terms, we will call these operators *almost-scaling operators*. We do not expect that scaling operators with exactly scale invariant two point functions exist in our theory, due to the presence of the fixed ultraviolet cutoff $\chi(k)$.

In order to identify $\mathcal{O}^{(1)}$ and $\mathcal{O}^{(2)}$, let us expand $V^*(\phi, J, \psi)$ as

$$V^*(\phi, J, \psi) = H^*(\psi) + (\phi, \mathcal{O}^{(1)}) + (J, \mathcal{O}^{(2)}) + \mathcal{Q}^*(\phi, J, \psi) + \mathcal{S}^*(\phi, J), \quad (2.38)$$

where $\mathcal{Q}^*$ is the part of $\mathcal{R}^*$ that is at least quadratic in (ϕ, J) , and $\mathcal{S}^*$ is the same as in (2.36). In other words, $\mathcal{O}_a^{(1)}(x) = \frac{\delta V^*(\phi, J, \psi)}{\delta \phi_a(x)}|_{\phi=J=0}$, and similarly for $\mathcal{O}^{(2)}(x)$. From the definition of response functions, we find

$$\begin{aligned} \mathcal{G}_{ab}^*(x) &= \langle \mathcal{O}_a^{(1)}(x) \mathcal{O}_b^{(1)}(0) \rangle_{H^*} + [\mathcal{E}_1(x)]_{ab}, \\ \mathcal{F}^*(x) &= \langle \mathcal{O}^{(2)}(x); \mathcal{O}^{(2)}(0) \rangle_{H^*} + \mathcal{E}_2(x), \end{aligned} \quad (2.39)$$

⁹The rationale behind the labels ext, 0, and f in (2.37) is the following: ‘ext’ stands for ‘external field’, it refers to the fact that $n + m > 0$, i.e., there is at least one ‘external field’ ϕ or J field; ‘ f ’ stands for ‘fluctuation field’, it refers to the fact that $l > 0$, i.e., there is at least one ‘fluctuation field’ ψ ; similarly, the label ‘0’ in the first definition indicates that $l = 0$, i.e., there is no fluctuation field.

where the correction terms $\mathcal{E}_1, \mathcal{E}_2$ are given by

$$\begin{aligned} [\mathcal{E}_1(x)]_{ab} &:= \frac{\delta^2 \mathcal{S}^*(\phi, 0)}{\delta \phi_a(0) \delta \phi_b(x)} \Big|_{\phi=0} + \left\langle \frac{\delta^2 \mathcal{Q}^*(\phi, 0, \psi)}{\delta \phi_a(0) \delta \phi_b(x)} \Big|_{\phi=0} \right\rangle_{H^*}, \\ \mathcal{E}_2(x) &:= \frac{\delta^2 \mathcal{S}^*(0, J)}{\delta J(0) \delta J(x)} \Big|_{J=0} + \left\langle \frac{\delta^2 \mathcal{Q}^*(0, J, \psi)}{\delta J(0) \delta J(x)} \Big|_{J=0} \right\rangle_{H^*}. \end{aligned} \quad (2.40)$$

2.7 Main results

Theorem 2.3 (Analyticity of the anomalous critical exponent and discrete scale invariance). *There exists $\epsilon_0 > 0$ such that, for $\epsilon \in B_{\epsilon_0}(0) := \{\epsilon \in \mathbb{C} : |\epsilon| < \epsilon_0\}$, there is a unique potential V^* as in Definition 2.2, analytic in ϵ in $B_{\epsilon_0}(0)$. Moreover, the response functions $\mathcal{G}^*$ and $\mathcal{F}^*$ defined in (2.7) are analytic in ϵ in the same domain, and behave under rescaling $x \rightarrow \rho x$, with $\rho = \gamma^k$, $k \in \mathbb{Z}$, as*

$$\mathcal{G}_{ab}^*(\rho x) = \rho^{-2\Delta_1} \mathcal{G}_{ab}^*(x), \quad \mathcal{F}^*(\rho x) = \rho^{-2\Delta_2} \mathcal{F}^*(x), \quad (2.41)$$

with $\Delta_1 = [\psi]$ and $\Delta_2 = 2[\psi] + \eta_2(\epsilon)$, where $\eta_2(\epsilon)$ is analytic in ϵ for $|\epsilon| < \epsilon_0$, with

$$\eta_2(\epsilon) = 2\epsilon \frac{N-2}{N-8} + O(\epsilon^2). \quad (2.42)$$

We actually expect that $\mathcal{G}^*$ and $\mathcal{F}^*$ are exactly scale invariant, which means that Eq.(2.41) should be true for any $\rho > 0$: this will be proved in the third paper of this series, see also the recent paper [28] for a proof of full scale invariance with a different choice of the cutoff function. In this paper, we limit ourselves to show, by an explicit lowest-order computation (see Appendix B), that the dominant contributions in ϵ to $\mathcal{G}^*(x)$ and $\mathcal{F}^*(x)$ are exactly scale-invariant terms, namely $\mathcal{G}^*(x) = \mathcal{G}_0^*(x) + \mathcal{G}_{\text{h.o.}}^*(x)$, where, in the sense of distributions [29],

$$\mathcal{G}_{0,ab}^*(x) = \Omega_{ab} \int \frac{d^d k}{|k|^{d/2+\epsilon}} e^{ikx} \equiv \Omega_{ab} C_0 |x|^{-2\Delta_1} \quad (2.43)$$

and the higher order correction $\mathcal{G}_{\text{h.o.}}^*$ satisfies $|\mathcal{G}_{\text{h.o.}}^*(x)| \leq C\epsilon |x|^{-2\Delta_1}$. Similarly, $\mathcal{F}^*(x) = \mathcal{F}_0^*(x) + \mathcal{F}_{\text{h.o.}}^*(x)$, where

$$\mathcal{F}_0^*(x) = -2N \int \frac{d^d k}{(2\pi)^d |k|^{d/2+\epsilon}} \int \frac{d^d p}{(2\pi)^d |p|^{d/2+\epsilon-2\eta_2}} e^{i(k+p)x} \equiv C'_0 |x|^{-2\Delta_2}, \quad (2.44)$$

and $|\mathcal{F}_{\text{h.o.}}^*(x)| \leq C'\epsilon |x|^{-2\Delta_2}$.

Theorem 2.4 (Almost-scaling operators). *There exist $C_i, c_i > 0$, $i = 1, 2$ such that $\mathcal{E}_1(x)$ and $\mathcal{E}_2(x)$ are analytic in ϵ in the same domain $B_{\epsilon_0}(0)$ as in Theorem 2.3, and bounded as follows:*

$$|[\mathcal{E}_1(x)]_{ab}| \leq C_1 e^{-c_1 |x|^\sigma} (\min\{1, |x|\})^{-2\Delta_1}, \quad |\mathcal{E}_2(x)| \leq C_2 e^{-c_2 |x|^\sigma} (\min\{1, |x|\})^{-2\Delta_2}, \quad (2.45)$$

with $\sigma = 1/s$ and s the order of the Gevrey class G^s which the UV cutoff function χ belongs to.

Putting together Eqs. (2.39) with Theorems 2.3 and 2.4, we conclude that the two point functions $\langle \mathcal{O}_a^{(1)} \mathcal{O}_b^{(1)} \rangle_{H^*}$ and $\langle \mathcal{O}^{(2)}; \mathcal{O}^{(2)} \rangle_{H^*}$ are discrete scale invariant up to stretched-exponentially small corrections at large distances, i.e. that $\mathcal{O}^{(1)}$ and $\mathcal{O}^{(2)}$ are almost-scaling operators. As mentioned, we cannot hope for anything better than that.

Remark 2.5. In the non-interacting case discussed in Section 2.2.1, the stretched exponential decay of the correction terms, denoted by $\mathcal{E}_{0,1}, \mathcal{E}_{0,2}$ (the ‘0’ label recalling the absence of interaction), is immediate. In fact, rewriting the non-interacting fixed point potential V_0^* as in (2.38), we see that $\mathcal{O}_a^{(1)}(x) = \psi_a(x)$ and $\mathcal{O}^{(2)}(x) = \psi^2(x)$. Therefore, for $H^* = 0$, the two-point functions $\langle \mathcal{O}_a^{(1)}(x) \mathcal{O}_b^{(1)}(0) \rangle_{H^*}$ and $\langle \mathcal{O}^{(2)}(x); \mathcal{O}^{(2)}(0) \rangle_{H^*}$ reduce to $\langle \psi_a(x) \psi_b(0) \rangle_0 = \Omega_{ab} P_{\leq 1}(x)$ and $\langle \psi^2(x); \psi^2(0) \rangle_0 = -2N P_{\leq 0}^2(x)$, respectively. Therefore, from (2.23) and the non-interacting analogue of (2.39), that is

$$\begin{aligned} \mathcal{G}_{0;ab}^*(x) &= \langle \psi_a(x) \psi_b(0) \rangle_0 + [\mathcal{E}_{0,1}(x)]_{ab}, \\ \mathcal{F}_0^*(x) &= \langle \psi^2(x); \psi^2(0) \rangle_0 + \mathcal{E}_{0,2}(x), \end{aligned} \quad (2.46)$$

we see that $\mathcal{E}_{0,1}, \mathcal{E}_{0,2}$ are, explicitly:

$$\begin{aligned} [\mathcal{E}_{0,1}(x)]_{ab} &= \Omega_{ab} P_{\geq 1}(x) \\ \mathcal{E}_{0,2} &= -2N [2P_{\leq 0}(x)P_{\geq 1}(x) + P_{\geq 1}^2(x)], \end{aligned} \quad (2.47)$$

which can be easily shown to satisfy the decay bounds in (2.45).

2.8 Structure of the proof

The proofs of the Theorem 2.3 and 2.4 are based on the following considerations. First of all, from (2.6) we have:

$$W^*(\phi, J) = \lim_{h \rightarrow -\infty} (TS_{h+1} \cdots TS_0 V^*)(\phi, J, 0). \quad (2.48)$$

Now, recalling that the RG map $R = D^{-(h-1)} TS_h D^h \equiv DTS_0$, for all $h \leq 0$, we see that (2.48) can be written as:

$$W^*(\phi, J) = \lim_{h \rightarrow -\infty} D^h (R^{|h|} V^*)(\phi, J, 0) = \lim_{h \rightarrow -\infty} D^h V^*(\phi, J, 0) \quad (2.49)$$

where in the last equality we used that V^* is fixed point for the RG map, i.e. $RV^* = V^*$. Denote by $V_{n,m,l,p}^*$ the kernels of V^* , in the sense of (2.24). Using the definitions (2.7), and recalling the action (2.27) of D on the kernels, we find, letting $\mathbf{x} = (x, 0)$,

$$\begin{aligned} \mathcal{G}^*(x) &= 2 \lim_{h \rightarrow -\infty} D^h V_{2,0,0,\emptyset}^*(\mathbf{x}) \\ &= 2 \lim_{h \rightarrow -\infty} \gamma^{2h\Delta_1} V_{2,0,0,\emptyset}^*(\gamma^h \mathbf{x}), \end{aligned} \quad (2.50)$$

and, letting $\mathbf{y} = (y, 0)$,

$$\begin{aligned} \mathcal{F}^*(y) &= 2 \lim_{h \rightarrow -\infty} D^h V_{0,2,0,\emptyset}^*(\mathbf{y}) \\ &= 2 \lim_{h \rightarrow -\infty} \gamma^{2h\Delta_2} V_{0,2,0,\emptyset}^*(\gamma^h \mathbf{y}). \end{aligned} \quad (2.51)$$

After having defined, in Section 3, the fixed point equation for V^* , in Section 4 we will show that the kernels $V_{n,m,l,p}^*$, and in particular $V_{2,0,0,\emptyset}^*, V_{0,2,0,\emptyset}^*$, can be written in terms of a tree expansion that is absolutely convergent in a weighted L^1 norm. As a consequence, these kernels, as well as the scaling exponents Δ_1, Δ_2 , turn out to be analytic in ϵ for ϵ small enough. In particular $\Delta_1 = [\psi]$

and $\Delta_2 = 2[\psi] + \eta_2(\epsilon)$, with η_2 as in (2.42) (the explicit computation of the first order contribution is deferred to Appendix A).

In Section 5, we will prove the pointwise existence of the limits in (2.50) and (2.51), as well as the analyticity of the limiting functions. These will require to control the absolute convergence of the tree expansion for the kernels of the fixed point potential in a pointwise norm. The ideas here generalize, and adapt to the present context, those used in [18, 30, 31, 32] to estimate the n -point correlation functions. Once the pointwise existence of the limits in (2.50)-(2.51) is established, the scale invariant property (2.41) is an immediate corollary: in fact, from (2.50) one finds, renaming $h + k \equiv h'$,

$$\begin{aligned} \mathcal{G}^*(\gamma^k x) &= 2 \lim_{h \rightarrow -\infty} \gamma^{2h\Delta_1} V_{2,0,0,\emptyset}^*(\gamma^{h+k} x) \\ &= 2\gamma^{-2k\Delta_1} \lim_{h' \rightarrow -\infty} \gamma^{2h'\Delta_1} V_{2,0,0,\emptyset}^*(\gamma^{h'} x) \equiv \gamma^{-2k\Delta_1} \mathcal{G}^*(x), \end{aligned} \quad (2.52)$$

provided the limit exists, and similarly for $\mathcal{F}^*$, thus concluding the proof of Theorem 2.3.

Finally, in Section 6, we derive a tree expansion for the correction terms $\mathcal{E}_1, \mathcal{E}_2$, and show that its convergence in a pointwise norm introduced in Section 5 implies Theorem 2.4.

3 Renormalization map and fixed point equation

In this section we describe more in detail the RG transformation introduced in Section 2.2. Moreover, we write down the fixed point equation for V^* , including the equations for the scaling exponents Δ_1, Δ_2 . The discussion follows the analogous one in [1, Section 5], which we refer to for additional details.

3.1 Integrating out

Consider a potential $H(\phi, J, \psi) = \{H_\ell\}_{\ell \in L}$, whose sequence of kernels is trimmed, in the sense of Section 2.4. Let us discuss the effect of integrating out the fluctuation field on scale 0 from $H(\phi, J, \psi)$:

$$H_{\text{eff}}(\phi, J, \psi) = S_0 H(\phi, J, \psi), \quad (3.1)$$

with S_0 the integrating-out map on scale 0 defined in (2.15). In analogy with [1, Eq.(5.3)], using the notation introduced in footnote 6, the kernels of the effective interaction can be written as:

$$H_{\text{eff}}(\mathbf{a}, \mathbf{x}, \mathbf{y}, \mathbf{B}, \mathbf{z}_{\mathbf{B}}) = \mathcal{P} \sum_{s \geq 1} \frac{1}{s!} \sum_{\substack{\mathbf{B}_1, \dots, \mathbf{B}_s \\ \sum_i \mathbf{B}_i = \mathbf{B}}} \sum_{\substack{\mathbf{A}_1, \dots, \mathbf{A}_s \\ \mathbf{A}_i \supset \mathbf{B}_i}} \sum_{\substack{\mathbf{a}_1, \dots, \mathbf{a}_s \\ \mathbf{x}_1, \dots, \mathbf{x}_s \\ \mathbf{y}_1, \dots, \mathbf{y}_s}} (-)^{\#} \int d\mathbf{z}_{\bar{\mathbf{B}}} \mathcal{C}(\mathbf{z}_{\bar{\mathbf{B}}}) \prod_{i=1}^s H(\mathbf{a}_i, \mathbf{x}_i, \mathbf{y}_i, \mathbf{A}_i, \mathbf{z}_{\mathbf{A}_i}) \quad (3.2)$$

where: $\mathcal{P}$ is the operator that anti-symmetrizes under simultaneous permutations of the elements of $\mathbf{a}$ and $\mathbf{x}$, symmetrizes over permutations of the elements of $\mathbf{y}$, and anti-symmetrizes under simultaneous permutations of the elements of $\mathbf{B}$ and $\mathbf{z}_{\mathbf{B}}$ (which generalizes the antisymmetrization operator $\mathcal{A}$ of [1, Eq.(5.3)]); the sums over $\mathbf{B}_i$ and $\mathbf{A}_i$ must be interpreted as described after [1, Eq.(5.3)]; the sum over $\mathbf{a}_1, \dots, \mathbf{a}_s$ is over all ways to represent $\mathbf{a}$ as a concatenation $\mathbf{a}_1 + \dots + \mathbf{a}_s$, and similarly for the sums over $\mathbf{x}_1, \dots, \mathbf{x}_s$ and over $\mathbf{y}_1, \dots, \mathbf{y}_s$. Finally, letting, as in [1, Eq.(5.3)], $\bar{\mathbf{B}}_i = \mathbf{A}_i \setminus \mathbf{B}_i$ and $\bar{\mathbf{B}} = \bar{\mathbf{B}}_1 + \dots + \bar{\mathbf{B}}_s$, we denote $\mathcal{C}(\mathbf{z}_{\bar{\mathbf{B}}}) := \langle \Psi'(\bar{\mathbf{B}}_1, \mathbf{z}_{\bar{\mathbf{B}}_1}); \dots; \Psi'(\bar{\mathbf{B}}_s, \mathbf{z}_{\bar{\mathbf{B}}_s}) \rangle_0^{(0)}$ where $\Psi'(\bar{\mathbf{B}}_i, \mathbf{z}_{\bar{\mathbf{B}}_i})$ is interpreted as in footnote 6, and $\langle \dots \rangle_0^{(0)}$ denotes the expectation w.r.t. $d\mu_0(\psi')$.

We compactly rewrite (3.2) as

$$(H_{\text{eff}})_\ell = \sum_{s \geq 1} \sum_{(\ell_i)_{i=1}^s} S_\ell^{\ell_1, \dots, \ell_s}(H_{\ell_1}, \dots, H_{\ell_s}), \quad (3.3)$$

where ℓ belongs to the label set L in (2.32), while $\ell_1, \dots, \ell_s$ belong to the subset of L ‘with at least one fluctuation field label’, i.e., to

$$L_f = \{(n, m, l, \mathbf{p}) \in L : l > 0\}, \quad (3.4)$$

the label f standing for ‘fluctuation field’.

3.2 Trimming

Even if the input potential $H = \{H_\ell\}_{\ell \in L}$ is trimmed, the output H_{eff} will not in general be so. However, we can act on H_{eff} with a linear operator, equivalent to the identity in the sense of [1, Section 5.2.1], called the trimming operator T , which returns an equivalent trimmed potential. Denoting equivalence between potentials by the symbol $\sim$, the action of the trimming operator will allow us to rewrite

$$H_{\text{eff}} \sim \mathcal{L}H_{\text{eff}} + \mathcal{I}H_{\text{eff}}, \quad (3.5)$$

with $\mathcal{L}H_{\text{eff}}$ the local, relevant, part of H_{eff} , and $\mathcal{I}H_{\text{eff}}$ the non-local, irrelevant, part of H_{eff} . The action of T on the kernels with $n = m = 0$ has already been described in [1, Sect.5.2 and App.C] and won’t be repeated here. The action of T on the kernels with $n + m > 0$ is non-trivial iff (n, m, l) equals $(1, 0, 1)$ or $(0, 1, 2)$; in the complementary case, we let $(\mathcal{I}H_{\text{eff}})_{n,m,l,\mathbf{p}} = (H_{\text{eff}})_{n,m,l,\mathbf{p}}$ and $(\mathcal{L}H_{\text{eff}})_{n,m,l,\mathbf{p}} = 0$.

Let us now consider the cases $(n, m, l) = (1, 0, 1)$, $(0, 1, 2)$. As in the case without source fields, trimming involves localization and interpolation: localization extracts the local parts of $(H_{\text{eff}})_{1,0,1,0}$ and $(H_{\text{eff}})_{0,1,2,0}$ (cf. with [1, Eq.(5.12)]), as follows:

$$(\mathcal{L}H_{\text{eff}})_{1,0,1,0} = T_{1,0,1,0}^{1,0,1,0}(H_{\text{eff}})_{1,0,1,0}, \quad (\mathcal{L}H_{\text{eff}})_{0,1,2,0} = T_{0,1,2,0}^{0,1,2,0}(H_{\text{eff}})_{0,1,2,0}, \quad (3.6)$$

(the operators $T_{1,0,1,0}^{1,0,1,0}$ and $T_{0,1,2,0}^{0,1,2,0}$ are defined in the following subsections), and we let

$$(\mathcal{L}H_{\text{eff}})_{1,0,1,\mathbf{p}} = (\mathcal{L}H_{\text{eff}})_{0,1,2,\mathbf{p}} = 0 \quad \text{for } \mathbf{p} \neq \mathbf{0}.$$

On the other hand, letting $P_1 = \{(0), (1)\}$ and $P_2 = \{(0, 0), (1, 0), (0, 1), (1, 1)\}$, interpolation rearranges the difference between $\{(H_{\text{eff}})_{1,0,1,\mathbf{p}}\}_{\mathbf{p} \in P_1}$ (resp. $\{(H_{\text{eff}})_{0,1,2,\mathbf{p}}\}_{\mathbf{p} \in P_2}$) and its local part $\{(\mathcal{L}H_{\text{eff}})_{1,0,1,\mathbf{p}}\}_{\mathbf{p} \in P_1}$ (resp. $\{(\mathcal{L}H_{\text{eff}})_{0,1,2,\mathbf{p}}\}_{\mathbf{p} \in P_2}$) in such a way to equivalently rewrite it as $\{(\mathcal{I}H_{\text{eff}})_{1,0,1,\mathbf{p}}\}_{\mathbf{p} \in P_1}$ (resp. $\{(\mathcal{I}H_{\text{eff}})_{0,1,2,\mathbf{p}}\}_{\mathbf{p} \in P_2}$) with $(\mathcal{I}H_{\text{eff}})_{1,0,1,0} = (\mathcal{I}H_{\text{eff}})_{0,1,2,0} = 0$. More precisely, we let

$$\begin{aligned} (\mathcal{I}H_{\text{eff}})_{1,0,1,\mathbf{p}} &= \begin{cases} 0 & \text{if } \mathbf{p} = \mathbf{0} \\ (H_{\text{eff}})_{1,0,1,\mathbf{p}} + T_{1,0,1,\mathbf{p}}^{1,0,1,0}(H_{\text{eff}})_{1,0,1,0} & \text{if } \|\mathbf{p}\|_1 = 1, \end{cases} \\ (\mathcal{I}H_{\text{eff}})_{0,1,2,\mathbf{p}} &= \begin{cases} 0 & \text{if } \mathbf{p} = \mathbf{0} \\ (H_{\text{eff}})_{0,1,2,\mathbf{p}} + T_{0,1,2,\mathbf{p}}^{0,1,2,0}(H_{\text{eff}})_{0,1,2,0} & \text{if } \|\mathbf{p}\|_1 = 1 \\ (H_{\text{eff}})_{0,1,2,\mathbf{p}} & \text{if } \|\mathbf{p}\|_1 = 2. \end{cases} \end{aligned} \quad (3.7)$$

Identifying the kernels with the corresponding Grassmann monomials as in [1, Eq.(C.1)], the manipulations equivalent to the identity leading to the definitions of $T_{1,0,1,\mathbf{p}}^{1,0,1,0}$, $T_{0,1,2,\mathbf{p}}^{0,1,2,0}$ are described in the following subsections.

3.2.1 Case $(n, m, l) = (1, 0, 1)$

We will use the following interpolation identity (in most cases, we drop and leave implicit the component indices of the Grassmann fields; moreover, summation over repeated indices is understood), which is the same as [1, Eq.(5.8)]:

$$\psi(y) = \psi(x) + (y - x)_\mu \int_0^1 ds \partial_\mu \psi(x + s(y - x)). \quad (3.8)$$

This is used to split a non-local but relevant term into a local relevant term plus an irrelevant one (for the definition of relevant and irrelevant, see after (2.30)). Hence considering $\int d^d x d^d z \phi(x) G(x, z) \psi(z)$ with G translationally invariant, playing the role of $(H_{\text{eff}})_{1,0,1,0}$, we get:

$$\begin{aligned} \int d^d x d^d z \phi(x) G(x, z) \psi(z) &= \int d^d x \phi(x) \hat{G}(0) \psi(x) \\ &+ \int d^d x d^d z \phi(x) G(x, z) (z - x)_\mu \int_0^1 ds \partial_\mu \psi(x + s(z - x)) \\ &\equiv \int d^d x \phi(x) \hat{G}(0) \psi(x) + \int d^d x d^d z' \phi(x) G^\mu(x, z') \partial_\mu \psi(z'), \end{aligned} \quad (3.9)$$

with $\hat{G}(0) := \int d^d z G(x, z)$ and

$$G^\mu(x, z) = \int_0^1 \frac{ds}{s^{d+1}} G(x, x + (z - x)/s) (z - x)_\mu. \quad (3.10)$$

In light of this,

$$(T_{1,0,1,0}^{1,0,1,0} G)(x, z) = \hat{G}(0) \delta(x - z) \quad (3.11)$$

and, for $\mathbf{p} = (1)$,

$$[(T_{1,0,1,(1)}^{1,0,1,0} G)(x, z)]_\mu = G^\mu(x, z) \quad (3.12)$$

By the same considerations as in [1, Appendix C], we have that $T_{1,0,1,\mathbf{p}}^{1,0,1,0}$ satisfies the following norm bounds:

$$\|T_{1,0,1,\mathbf{p}}^{1,0,1,0} G\| \leq \|G\|, \quad \|T_{1,0,1,(1)}^{1,0,1,0} G\| \leq \max_{a,b} \int d^d z |[G(0, z)]_{a,b}| |z| \quad (3.13)$$

and similarly for the weighted L^1 norms to be used below.

3.2.2 Case $(n, m, l) = (0, 1, 2)$

Consider now $\int d^d y d^d z_1 d^d z_2 J(y) \psi(z_1) F(y, \mathbf{z}) \psi(z_2)$ with F translationally invariant, playing the role of $(H_{\text{eff}})_{(0,1,2,0)}$. Proceeding analogously we find:

$$\begin{aligned} &\int d^d y d\mathbf{z} J(y) \psi(z_1) F(y, \mathbf{z}) \psi(z_2) \\ &= \int d^d y d\mathbf{z} J(y) \left[\psi(y) F(y, \mathbf{z}) \psi(y) + (z_1 - y)_\mu \int_0^1 ds \partial_\mu \psi(y + s(z_1 - y)) F(y, \mathbf{z}) \psi(y + s(z_2 - y)) \right. \\ &\quad \left. + (z_2 - y)_\mu \int_0^1 ds \psi(y + s(z_1 - y)) F(y, \mathbf{z}) \partial_\mu \psi(y + s(z_2 - y)) \right] \\ &\equiv \sum_{\mathbf{p}: \|\mathbf{p}\|_1 \leq 1} \int d^d y d\mathbf{z}' J(y) \partial_{\mu_1}^{p_1} \psi(z'_1) [F^{\mathbf{p}}(y, \mathbf{z}')]_{\mu} \partial_{\mu_2}^{p_2} \psi(z'_2), \end{aligned} \quad (3.14)$$

with $F^{\mathbf{0}}(y, \mathbf{z}) = \delta(z_1 - y)\delta(z_2 - y) \int d^{2d} \mathbf{z}' F(x, \mathbf{z}')$,

$$F^{(1,0)}(y, \mathbf{z}) = (z_1 - y)_\mu \int_0^1 \frac{ds}{s^{2d+1}} F(y, y + (z_1 - y)/s, y + (z_2 - y)/s), \quad (3.15)$$

and similarly for $F^{(0,1)}$. Then, identifying $\mathbf{p}$ with the pair (p_1, p_2) :

$$\begin{aligned} (T_{0,1,2,\mathbf{0}}^{0,1,2,\mathbf{0}} F)(y, \mathbf{z}) &= \delta(z_1 - y)\delta(z_2 - y) \int d^{2d} \mathbf{z}' F(0, \mathbf{z}'), \\ (T_{0,1,2,\mathbf{p}}^{0,1,2,\mathbf{0}} F)(y, \mathbf{z}) &= F^{\mathbf{p}}(y, \mathbf{z}), \quad \text{if } \|\mathbf{p}\|_1 = 1. \end{aligned} \quad (3.16)$$

In analogy with (3.13), we have: $\|T_{0,1,2,\mathbf{0}}^{0,1,2,\mathbf{0}} F\| \leq \|F\|$, while, if $\mathbf{p} = (1, 0)$,

$$\|T_{0,1,2,(1,0)}^{0,1,2,\mathbf{0}} F\| \leq \max_{a,b} \int dz |[F(0, \mathbf{z})]_{a,b}| |z_1|, \quad (3.17)$$

and similarly for $\mathbf{p} = (0, 1)$.

3.3 Fixed point equation

We are now ready to write the definition of the renormalization map $H \rightarrow RH = DTS_0 H$ component-wise. Recalling (3.1)-(3.3), the definition of the trimming map in the previous subsection, and the definition of the dilatation D in (2.27), for any $\ell \in L$, we can write

$$(RH)_\ell = \sum_{s \geq 1} \sum_{(\ell_i)_{i=1}^s} R_\ell^{\ell_1, \dots, \ell_s}(H_{\ell_1}, \dots, H_{\ell_s}), \quad (3.18)$$

where the sum runs over s -ples of labels in the set L_f defined in (3.4) and, letting $\mathfrak{L} = \{(0, 0, 2), (0, 0, 4), (1, 0, 1), (0, 1, 2)\}$,

$$R_\ell^{\ell_1, \dots, \ell_s} = D \begin{cases} S_\ell^{\ell_1, \dots, \ell_s}, & \text{if } \ell = (n, m, l, \mathbf{p}) \text{ with } (n, m, l) \notin \mathfrak{L}, \\ T_{n,m,l,\mathbf{0}}^{n,m,l,\mathbf{0}} S_{(n,m,l,\mathbf{0})}^{\ell_1, \dots, \ell_s}, & \text{if } \ell = (n, m, l, \mathbf{0}) \text{ with } (n, m, l) \in \mathfrak{L}, \\ \sum_{\mathbf{p}'} T_{n,m,l,\mathbf{p}}^{n,m,l,\mathbf{p}'} S_{(n,m,l,\mathbf{p}')}^{\ell_1, \dots, \ell_s}, & \text{if } \ell = (n, m, l, \mathbf{p}) \text{ with } (n, m, l) \in \mathfrak{L} \text{ and } \mathbf{p} \neq \mathbf{0}, \end{cases} \quad (3.19)$$

and in the second and third lines the operator $T_{n,m,l,\mathbf{p}}^{n,m,l,\mathbf{p}'}$ is defined as follows: it is the identity, if $\mathbf{p} = \mathbf{p}' \neq \mathbf{0}$; it is the one defined in subsections 3.2.1-3.2.2 above, if $(n, m, l) = (1, 0, 1), (0, 1, 2)$ and either $\mathbf{p} = \mathbf{p}' = \mathbf{0}$ or $\mathbf{p}' = \mathbf{0}$, $\|\mathbf{p}\|_1 = 1$; it is given by [1, eq.s (5.12), (5.13), (5.14)] if $(n, m, l) = (0, 0, 2)$ and either $\mathbf{p} = \mathbf{p}' = \mathbf{0}$ or $\|\mathbf{p}'\|_1 < \|\mathbf{p}\|_1 = 2$, or if $(n, m, l) = (0, 0, 4)$ and either $\mathbf{p} = \mathbf{p}' = \mathbf{0}$ or $\mathbf{p}' = \mathbf{0}$, $\|\mathbf{p}\|_1 = 1$; it vanishes, otherwise.

In view of these definitions, and recalling the fact that we look for a fixed point potential with pre-factor in front of the local terms (ϕ, ψ) and (J, ψ^2) in V^* equal to 1 (see item (2) of Definition

2.2), the fixed point equation (FPE) for the ‘local’ components $\ell \in \{(n, m, l, \mathbf{0})\}_{(n, m, l) \in \mathcal{L}}$ reads:

$$\begin{aligned}
\nu &= \gamma^{d/2+\varepsilon} \nu + \sum_{s \geq 1} \sum_{(\ell_i)_{i=1}^s}^* R_{(0,0,2,\mathbf{0})}^{\ell_1, \dots, \ell_s}(H_{\ell_1}, \dots, H_{\ell_s}) \\
\lambda &= \gamma^{2\varepsilon} \lambda + \sum_{s \geq 1} \sum_{(\ell_i)_{i=1}^s}^* R_{(0,0,4,\mathbf{0})}^{\ell_1, \dots, \ell_s}(H_{\ell_1}, \dots, H_{\ell_s}) \\
1 &= \gamma^{\Delta_1 - [\psi]} + \sum_{s \geq 1} \sum_{(\ell_i)_{i=1}^s}^* R_{(1,0,1,\mathbf{0})}^{\ell_1, \dots, \ell_s}(H_{\ell_1}, \dots, H_{\ell_s}) \equiv \gamma^{\Delta_1 - [\psi]}(1 + \zeta_1) \\
1 &= \gamma^{\Delta_2 - 2[\psi]} + \sum_{s \geq 1} \sum_{(\ell_i)_{i=1}^s}^* R_{(0,1,2,\mathbf{0})}^{\ell_1, \dots, \ell_s}(H_{\ell_1}, \dots, H_{\ell_s}) \equiv \gamma^{\Delta_2 - 2[\psi]}(1 + \zeta_2)
\end{aligned} \tag{3.20}$$

where, again, the labels ℓ_i in the sums in the right hand sides are summed over L_f , and the * indicates the constraint that the term with $s = 1$ and $\ell_1 = \ell$ should be excluded from the corresponding sums. The FPE for the components $(0, 0, 2, \mathbf{p})$ with $\|\mathbf{p}\|_1 = 1$ is by construction trivial, $0 = 0$, while the FPE for all the other components reads:

$$H_\ell = \sum_{s \geq 1} \sum_{(\ell_i)_{i=1}^s}^* R_\ell^{\ell_1, \dots, \ell_s}(H_{\ell_1}, \dots, H_{\ell_s}). \tag{3.21}$$

Note that the FPE for the components $\ell = (n, m, l, \mathbf{p})$ with $n = m = 0$ is the same as the one studied and solved in [1], the solution being the sequence of kernels $H_{l, \mathbf{p}}^*$ constructed there. In particular, the first two components of (3.20) are solved by the fixed point values λ^*, ν^* of λ, ν computed in [1], and proved there to be analytic in ϵ for ϵ sufficiently small. So, from now on, we will set $\lambda = \lambda^*$ and $\nu = \nu^*$. Moreover, we will prove below that ζ_1 and ζ_2 are sums of convergent series in λ, ν , for ϵ sufficiently small and this implies that they are themselves analytic in ϵ for ϵ small. Letting $Z_1 := 1 + \zeta_1$ and $Z_2 := 1 + \zeta_2$, the FPE requires

$$\Delta_1 = [\psi] - \log_\gamma Z_1, \quad \Delta_2 = 2[\psi] - \log_\gamma Z_2 \equiv 2[\psi] + \eta_2. \tag{3.22}$$

We will see in Section 4.2.1 that $\zeta_1 = 0$, so that $\Delta_1 = [\psi]$. Moreover, in Section 4.2.2 and Appendix A, we will show that

$$Z_2 = 1 - \frac{2(N-2)}{N-8} \epsilon \log \gamma + O(\epsilon^2), \tag{3.23}$$

which gives $\eta_2 = 2\epsilon \frac{N-2}{N-8} + O(\epsilon^2)$.

4 Solution to the FPE via the tree expansion

Let us now discuss how to determine the solution to the FPE introduced in the previous section, satisfying the properties spelled out in Definition 2.2. We use an analogue of the tree expansion discussed in [1, App.J], which provides an explicit analytic solution. Our construction automatically proves the uniqueness of the solution within the class of analytic potentials with a prescribed form at $\epsilon = 0$. Uniqueness of the solution in the larger Banach space of potentials that are close enough to the prescribed unperturbed potential in the appropriate weighted L^1 norm, could be proved via

a contraction argument similar to the one underlying the proof of [1, Key Lemma], but we will not belabor the details of this proof here.

As already mentioned above, we fix $\lambda = \lambda^*$, $\nu = \nu^*$, the fixed point values of the quartic and quadratic interactions computed in [1], once and for all. We also recall that our ansatz for the fixed point potential V^* requires that the local terms (ϕ, ψ) and (J, ψ^2) have pre-factor equal to 1, see condition (2) in Definition 2.2, and that the components $(0, 0, 2, \mathbf{p})$ with $\|\mathbf{p}\|_1 = 1$ are zero, by the requirement that $\{V_\ell^*\}_{\ell \in L}$ is trimmed, see the second line of (2.34). We first describe how to solve the FPE for the remaining components, i.e., those in

$$L' := L \setminus \left(\{(n, m, l, \mathbf{0})\}_{(n, m, l) \in \mathcal{L}} \cup \{(0, 0, 2, \mathbf{p})\}_{\|\mathbf{p}\|_1 = 1} \right), \quad (4.1)$$

and then we will discuss the local components $\ell = (1, 0, 1, \mathbf{0}), (0, 1, 2, \mathbf{0})$.

4.1 The FPE for $\ell \in L'$

The FPEs for the components $\ell \in L'$ are those in (3.21). We proceed in a way similar to [1, Appendix J]. For any $\ell \in L'$, we isolate from the right side of the FPE the term with $s = 1$ and $\ell_1 = \ell$, i.e. the term DH_ℓ , move it to the left side, and multiply both sides by $(1 - D)^{-1}$. The resulting equation takes the form

$$H_\ell = \sum_{s \geq 1} \sum_{(\ell_i)_{i=1}^s}^* (1 - D)^{-1} R_\ell^{\ell_1, \dots, \ell_s} (H_{\ell_1}, \dots, H_{\ell_s}) \quad (4.2)$$

where we recall that the second sum runs over s -ples of labels in L_f , see (3.4), and $*$ denotes the constraint that, if $s = 1$, then $\ell_1 \neq \ell$, while $R_\ell^{\ell_1, \dots, \ell_s}$ is defined in (3.19). As shown below, the (diagonal in ℓ) operator $(1 - D)$ is invertible in L^1 on all the components $\ell \in L'$: this is immediate for the components such that the scaling dimension at $\epsilon = 0$ is different from zero, $D_{\text{sc}}(\ell)|_{\epsilon=0} \neq 0$, see (2.30), that is, for all the indices in L' but $(0, 2, 0, \emptyset)$. In order for $(1 - D)$ to be invertible on this component as well, we need extra information about Δ_2 , besides knowing that $\Delta_2 = 2[\psi] + O(\epsilon)$, as we are assuming (see condition (3) in Definition 2.2). Anticipating the fact that $\Delta_2 = 2[\psi] + 2\epsilon \frac{N-2}{N-8} + O(\epsilon^2)$, the desired invertibility for $\epsilon \neq 0$ follows, because this explicit expression for Δ_2 implies that $D_{\text{sc}}(0, 2, 0, \emptyset) = 2\epsilon \frac{N+4}{N-8} + O(\epsilon^2)$, which is different from zero for $\epsilon \neq 0$.

We look for a solution in the form of a sum over rooted trees,

$$V_\ell^* = \sum_{\tau} H_\ell[\tau], \quad (4.3)$$

The value $H_\ell[\tau]$ is fixed so that the following recursive equation is satisfied:

$$H_\ell[\tau] = \sum_{(\ell_i)_{i=1}^{s_{v_0}}}^* (1 - D)^{-1} R_\ell^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_n}[\tau_{s_{v_0}}]). \quad (4.4)$$

where $\tau_1, \dots, \tau_{s_{v_0}}$ are the subtrees of τ rooted in the vertices $v_1, \dots, v_{s_{v_0}}$ that are ‘children’ of the root vertex v_0 of τ . The rooted trees (with root v_0) involved in the sum (4.3) have the structure exemplified in Fig.3.

The iterative application of (4.4) leads to a representation of the tree value $H_\ell[\tau]$ in terms of an iterated action of the operator $(1 - D)^{-1} R_{\ell_v}^{\ell_{v_1}, \dots, \ell_{s_v}}$, one per vertex v , summed over the labels




Figure 3: An example of a tree rooted in v_0 , contributing to the sum in (4.3). Here $s_{v_0} = 3$ and the 3 children of v_0 are v_1, v_2, v_3 .

ℓ_v, ℓ_{v_i} (here the labels ℓ_{v_i} at exponent refer to the vertices $v_1, \dots, v_{s_v}$, which are the children of v on τ ; the label ℓ_{v_0} associated with the root v_0 is the only one that is kept fixed, all the others are summed over). The labels ℓ_v associated with vertices $v \neq v_0$ that are not endpoints are summed over $L'_f := L_f \cap L'$, while, if v is an endpoint, then ℓ_v takes one of the values in $\{(n, m, l, \mathbf{0})\}_{(n, m, l) \in \mathcal{L}}$, depending on the nature of the endpoint, as graphically described in Fig.4.




Figure 4: The four types of endpoints and the corresponding ℓ labels.

In the evaluation of the tree value $H_\ell[\tau]$, each of these endpoints is associated with the kernel of the corresponding ‘interaction vertex’, see Fig.5.




Figure 5: The four ‘interaction vertices’, graphically representing the contributions associated with the four types of endpoints depicted in Fig.4.

For example, for the tree τ represented in Fig.3, $s_{v_0} = 3$ and the three subtrees τ_1, τ_2, τ_3 ‘exiting’ from the root v_0 are those represented in Fig.6 (note that τ_2 is ‘trivial’, in that it consists of a single vertex, which is both the root and the endpoint of τ_2).

We now intend to use the formula for the tree values described above in order to derive norm bounds on $H_\ell[\tau]$ and to prove that the tree expansion for H_ℓ is absolutely convergent in the appropriate norms. We start by discussing bounds for the following **weighted L_1 norm** of $H_\ell[\tau]$, generalizing the definition in (2.25). Recall that $H_\ell[\tau](\mathbf{x}, \mathbf{y}, \mathbf{z})$ with $\ell = (n, m, l, \mathbf{p})$ must be understood as a tensor-valued function of components $[H_\ell[\tau](\mathbf{x}, \mathbf{y}, \mathbf{z})]_{\mathbf{a}, \mathbf{\mu}, \mathbf{b}}$, as in (2.24). For notational convenience, in some of the equations below, we will equivalently rewrite these components as




Figure 6

$H_\ell[\tau](\mathbf{a}, \mathbf{x}, \mathbf{y}, \mathbf{B}, \mathbf{z})$, with $\mathbf{B} = ((p_1, \mu_1, b_1), \dots, (p_l, \mu_l, b_l))$, as in footnote 6 (the label ℓ attached to $H_\ell[\tau](\mathbf{a}, \mathbf{x}, \mathbf{y}, \mathbf{B}, \mathbf{z})$ is actually redundant, but we will keep it for clarity); given such a $\mathbf{B}$, we shall write $\mathbf{p}(\mathbf{B}) = \mathbf{p}$. In analogy with (2.25), we let

$$\|H_\ell[\tau]\|_w := \max_{\substack{\mathbf{a}, \mathbf{B}: \\ \mathbf{p}(\mathbf{B}) = \mathbf{p}}} \iiint^* d\mathbf{x} d\mathbf{y} d\mathbf{z} |H_\ell[\tau](\mathbf{a}, \mathbf{x}, \mathbf{y}, \mathbf{B}, \mathbf{z})| w(\mathbf{x}, \mathbf{y}, \mathbf{z}) \quad (4.5)$$

where

$$w(\mathbf{x}, \mathbf{y}, \mathbf{z}) := e^{\bar{C}(St(\mathbf{x}, \mathbf{y}, \mathbf{z})/\gamma)^\sigma}, \quad (4.6)$$

with: $St(\mathbf{x}, \mathbf{y}, \mathbf{z})$ the Steiner diameter, or ‘tree distance’, of $(\mathbf{x}, \mathbf{y}, \mathbf{z})$, see [1, footnote 19]; $\bar{C} \equiv \frac{1}{2}C_{\chi^2}$ with C_{χ^2} the positive constant in (4.14) below; and $\sigma = 1/s$ with s the Gevrey regularity of χ , see the line after (2.2).

In order to recursively estimate the norm (4.5) of $H_\ell[\tau]$ via (4.4), note that, from (3.2) and (3.19), if $\ell = (n, m, l, \mathbf{p}) \in L'$ with $(n, m, l) \notin \mathcal{L}$, we can write

$$\begin{aligned} & H_\ell[\tau](\mathbf{a}, \mathbf{x}, \mathbf{y}, \mathbf{B}, \mathbf{z}_{\mathbf{B}}) \\ &= \frac{D}{1-D} \mathcal{P} \frac{1}{s_{v_0}!} \sum_{\substack{\mathbf{B}_1, \dots, \mathbf{B}_{s_{v_0}} \\ \sum \mathbf{B}_i = \mathbf{B}}} \sum_{\substack{\mathbf{A}_1, \dots, \mathbf{A}_{s_{v_0}} \\ \mathbf{A}_i \supset \mathbf{B}_i}} \sum_{\substack{\mathbf{a}_1, \dots, \mathbf{a}_{s_{v_0}} \\ \mathbf{x}_1, \dots, \mathbf{x}_{s_{v_0}} \\ \mathbf{y}_1, \dots, \mathbf{y}_{s_{v_0}}}} (-)^\sharp \int d\mathbf{z}_{\bar{\mathbf{B}}} \mathcal{C}(\mathbf{z}_{\bar{\mathbf{B}}}) \prod_{i=1}^{s_{v_0}} H_{\ell_i}[\tau_i](\mathbf{a}_i, \mathbf{x}_i, \mathbf{y}_i, \mathbf{A}_i, \mathbf{z}_{\mathbf{A}_i}), \end{aligned} \quad (4.7)$$

where $\ell_i \equiv \ell_{v_i}$, and we used that for these components the trimming map is the identity. If, instead, $\ell = (n, m, l, \mathbf{p}) \in L'$ with $(n, m, l) \in \mathcal{L}$ and $\mathbf{p} \neq 0$, then trimming acts non-trivially, and the FPE reads:

$$\begin{aligned} H_{n,m,l,\mathbf{p}}[\tau](\mathbf{a}, \mathbf{x}, \mathbf{y}, \mathbf{B}, \mathbf{z}_{\mathbf{B}}) &= \frac{D}{1-D} \sum_{\mathbf{p}'} T_{n,m,l,\mathbf{p}}^{n,m,l,\mathbf{p}'} \mathcal{P} \frac{1}{s_{v_0}!} \sum_{\substack{\mathbf{B}_1, \dots, \mathbf{B}_{s_{v_0}} \\ \sum \mathbf{B}_i = \mathbf{B}'}} \sum_{\substack{\mathbf{A}_1, \dots, \mathbf{A}_{s_{v_0}} \\ \mathbf{A}_i \supset \mathbf{B}_i}} \sum_{\substack{\mathbf{a}_1, \dots, \mathbf{a}_{s_{v_0}} \\ \mathbf{x}_1, \dots, \mathbf{x}_{s_{v_0}} \\ \mathbf{y}_1, \dots, \mathbf{y}_{s_{v_0}}}} \times \\ &\times (-)^\sharp \int d\mathbf{z}_{\bar{\mathbf{B}}} \mathcal{C}(\mathbf{z}_{\bar{\mathbf{B}}}) \prod_{i=1}^{s_{v_0}} H_{\ell_i}[\tau_i](\mathbf{a}_i, \mathbf{x}_i, \mathbf{y}_i, \mathbf{A}_i, \mathbf{z}_{\mathbf{A}_i}), \end{aligned} \quad (4.8)$$

where, given $\mathbf{B} = ((p_1, \mu_1, b_1), \dots, (p_l, \mu_l, b_l))$ and an l -ple $\mathbf{p}'$ such that $T_{n,m,l,\mathbf{p}}^{n,m,l,\mathbf{p}'} \neq 0$, the set $\mathbf{B}'$ appearing in the right side in the condition $\sum \mathbf{B}_i = \mathbf{B}'$ is equal to: $\mathbf{B}$ if $\mathbf{p}' = \mathbf{p}$; $\mathbf{B}_0 := ((0, 0, b_1), \dots, (0, 0, b_l))$ if $\mathbf{p}' = \mathbf{0}$; $\mathbf{B}_{(1,0)} = ((1, \mu_1, b_1), (0, 0, b_2))$ if $l = 2$, $\mathbf{p} = (1, 1)$ and $\mathbf{p}' = (1, 0)$;

and $\mathbf{B}_{(0,1)} = ((0, 0, b_1), (1, \mu_2, b_2))$ if $l = 2$, $\mathbf{p} = (1, 1)$ and $\mathbf{p}' = (0, 1)$ (these two last cases are the only ones where $\mathbf{p}' \neq \mathbf{0}, \mathbf{p}$).

Now, in order to bound the right sides of (4.7) and (4.8) we recall a few basic bounds from [1] (or their analogues adapted to the present more general context). First of all, as in [1, (5.37)], for $\ell \in L'$,

$$\|D(H_\ell[\tau])\|_w \leq \gamma^{D_{\text{sc}}(\ell)} \|H_\ell[\tau]\|_{w(\cdot/\gamma)} \leq \gamma^{D_{\text{sc}}(\ell)} \|H_\ell[\tau]\|_w, \quad (4.9)$$

with $D_{\text{sc}}(\ell)$ as in (2.30). Note that, assuming Δ_1 and Δ_2 to be ϵ -close to $[\psi]$ and $2[\psi]$, respectively, with ϵ small enough, then $\min_{\ell \in L' \setminus \{(0,2,0,0)\}} |D_{\text{sc}}(\ell)| = \min\{1, d/2\} + O(\epsilon)$. Therefore, we also have that, for $\ell \in L' \setminus \{(0, 2, 0, \emptyset)\}$,

$$\|(1 - D)^{-1}(H_\ell[\tau])\|_w \leq d_\gamma \|H_\ell[\tau]\|_w, \quad (4.10)$$

with $d_\gamma = \max_{\ell \in L' \setminus \{(0,2,0,0)\}} |1 - \gamma^{D_{\text{sc}}(\ell)}|^{-1}$, which is finite and bounded from above, uniformly in ϵ . On the other hand, recalling that $\Delta_2 = 2[\psi] + \eta_2$ with $[\psi] = d/4 - \epsilon/2$, as in (3.22),

$$\|(1 - D)^{-1}(H_{(0,2,0,0)}[\tau])\|_w \leq \alpha_\gamma \|H_{(0,2,0,0)}[\tau]\|_w, \quad (4.11)$$

where $\alpha_\gamma := |1 - \gamma^{-2\epsilon+2\eta_2}|^{-1}$, which is finite and of order ϵ^{-1} iff $\epsilon \neq 0$ and $\eta_2 \neq \epsilon$. From the first order computation of η_2 in Appendix A, which implies $\eta_2 = 2\epsilon \frac{N-2}{N-8} + O(\epsilon^2)$, see (2.42), we see that this condition is always verified for $\epsilon \neq 0$ small enough.

Concerning the action of the trimming operator, for any $\ell = (n, m, l, \mathbf{p}) \in L'$ and $\mathbf{p}' \neq \mathbf{p}$ such that $T_{n,m,l,\mathbf{p}}^{n,m,l,\mathbf{p}'}$ does not vanish, we have the analogue of [1, (5.43)]:

$$\|T_{n,m,l,\mathbf{p}}^{n,m,l,\mathbf{p}'} H_{(n,m,l,\mathbf{p}')} \|_{w(\cdot/\gamma)} \leq C_R \gamma^{\|\mathbf{p}-\mathbf{p}'\|_1} \|H_{(n,m,l,\mathbf{p}')} \|_w. \quad (4.12)$$

For the estimate of $S_\ell^{\ell_1, \dots, \ell_s}$, with $\ell \in L'$, $\ell_1, \dots, \ell_s \in L_f$, we proceed as follows. From [1, Eq.(5.39)], we see that $\mathcal{C}(\mathbf{z})$ satisfies:

$$|\mathcal{C}(\mathbf{z})| \leq C_{GH}^s \sum_{\mathcal{T}} \prod_{(z,z') \in \mathcal{T}} M(z - z') \quad (4.13)$$

where C_{GH} constant given by the Gram-Hadamard bound [1, Lemma D.2] and M as in [1, Eq.(4.15)], i.e., such that:

$$|g_{a,b}^{(0)}(x)|, |\partial_\nu g_{a,b}^{(0)}(x)|, |\partial_{\mu,\nu} g_{a,b}^{(0)}(x)| \leq M(x) \equiv C_{\chi^1} e^{-C_{\chi^2}|x/\gamma|^\sigma}, \quad (4.14)$$

where C_{χ^1}, C_{χ^2} are constants depending on χ but independent of γ and $\sigma = 1/s \in (0, 1)$, with s the Gevrey regularity of χ , see the line after (2.2).

Using these estimates, and proceeding as in [1, Sect. 5.6] and [1, App.E], we get, letting $\ell = (n, m, l, \mathbf{p})$ and $\ell_i = (n_i, m_i, l_i, \mathbf{p}_i)$:

$$\|S_\ell^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}])\|_w \leq C_\gamma^{s_{v_0}-1} C_0^{\sum_{i=1}^{s_{v_0}} l_i - l} N_\ell^{\ell_1, \dots, \ell_{s_{v_0}}} \prod_{i=1}^{s_{v_0}} \|H_{\ell_i}[\tau_i]\|_w, \quad (4.15)$$

where $N_\ell^{\ell_1, \dots, \ell_{s_{v_0}}}$ is the number of ways in which $\ell = (n, m, l, \mathbf{p})$ can be realized, given $\ell_1, \dots, \ell_{s_{v_0}}$, via the action of $S_\ell^{\ell_1, \dots, \ell_{s_{v_0}}}$ on $(H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}])$, which is such that

$$\sum_{\mathbf{p}} N_{(n,m,l,\mathbf{p})}^{\ell_1, \dots, \ell_{s_{v_0}}} \leq \binom{\sum_{i=1}^{s_{v_0}} l_i}{l}. \quad (4.16)$$

Moreover, $C_\gamma = N^2 d^2 \|M\|_w = \text{Cost} \cdot \gamma^d$ and C_0 a constant independent of γ , see [1, Remark J.1]. Recall also that $S_\ell^{\ell_1, \dots, \ell_{s_{v_0}}}$ is non-zero only if $\sum_{i=1}^{s_{v_0}} l_i \geq l + 2(s_{v_0} - 1)$, $\sum_{i=1}^{s_{v_0}} n_i = n$ and $\sum_{i=1}^{s_{v_0}} m_i = m$.

Plugging eqs.(4.9)–(4.12) and (4.15) into (4.4), with $R_\ell^{\ell_1, \dots, \ell_{s_{v_0}}}$ defined as in (3.19), we obtain:

$$\begin{aligned} \|H_\ell[\tau]\|_w &\leq \sum_{(\ell_i)_{i=1}^{s_{v_0}}}^* \left\| \frac{D}{1-D} T S_\ell^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}]) \right\|_w \\ &\leq (\alpha_\gamma/d_\gamma)^{\mathbb{1}(\ell=(0,2,0,\emptyset))} \sum_{(\ell_i)_{i=1}^{s_{v_0}}}^* d_\gamma \gamma^{D_{\text{sc}}(\ell)} C_R \gamma^2 C_\gamma^{s_{v_0}-1} C_0^{\sum_{i=1}^{s_{v_0}} l_i - l} \mathcal{N}_\ell^{\ell_1, \dots, \ell_{s_{v_0}}} \prod_{i=1}^{s_{v_0}} \|H_{\ell_i}[\tau_i]\|_w, \end{aligned} \quad (4.17)$$

where $\mathcal{N}_\ell^{\ell_1, \dots, \ell_{s_{v_0}}}$ denotes the number of ways in which $\ell = (n, m, l, \mathbf{p})$ can be realized, given $\ell_1, \dots, \ell_{s_{v_0}}$, via the action of $R_\ell^{\ell_1, \dots, \ell_{s_{v_0}}}$ on $(H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}])$. Recalling the definition of $R_\ell^{\ell_1, \dots, \ell_s}$ in (3.19), and in the particular the definition of the components $T_{n,m,l,\mathbf{p}'}^{n,m,l,\mathbf{p}'}$ of the trimming operator given right below (3.19), as well as the bound on $\mathcal{N}_\ell^{\ell_1, \dots, \ell_{s_{v_0}}}$ stated right after (4.15), we find:

$$\begin{aligned} \sum_{\mathbf{p}}^* \mathcal{N}_{(n,m,l,\mathbf{p})}^{\ell_1, \dots, \ell_{s_{v_0}}} &\leq \binom{\sum_{i=1}^{s_{v_0}} l_i}{l} \cdot \begin{cases} 1 & \text{if } (n, m, l) \notin \mathfrak{L} \\ 4 & \text{if } (n, m, l) \in \mathfrak{L} \end{cases} \\ &\leq 4 \binom{\sum_{i=1}^{s_{v_0}} l_i}{l}, \end{aligned} \quad (4.18)$$

where $\sum_{\mathbf{p}}^*$ denotes the sum over the $\mathbf{p}$'s such that $(n, m, l, \mathbf{p}) \in L'$, and 4 is the maximum number of different $\mathbf{p}$'s for which, given $(n, m, l) \in \mathfrak{L}$ and $\mathbf{p}' \in \{0, 1\}^l$, the tuple $(n, m, l, \mathbf{p})$ is in L' and the operator $T_{n,m,l,\mathbf{p}}^{n,m,l,\mathbf{p}'}$ is non zero (such maximum is realized for $(n, m, l) = (0, 0, 4)$ and $\mathbf{p}' = \mathbf{0}$; in this case, the different $\mathbf{p}$'s with the stated property are: $(1, 0, 0, 0)$, $(0, 1, 0, 0)$, $(0, 0, 1, 0)$, $(0, 0, 0, 1)$).

By using iteratively the estimates above, and recalling that $|\lambda|, |\nu| \leq K\epsilon$ for some $K > 0$, we find

$$\begin{aligned} \|H_\ell[\tau]\|_w &\leq (\alpha_\gamma/d_\gamma)^{\mathbb{1}(\ell=(0,2,0,\emptyset))} \sum_{\{\ell_v\}} \left(\prod_{v \text{ not e.p.}} d_\gamma C_\gamma^{s_v-1} \gamma^{D_{\text{sc}}(\ell_v)} C_R \gamma^2 C_0^{\sum_{i=1}^{s_v} l_{v_i} - l_v} \mathcal{N}_{\ell_v}^{\ell_{v_1}, \dots, \ell_{v_{s_v}}} \right) \\ &\quad \cdot \left(\prod_{v \text{ e.p.}} (K\epsilon)^{\delta_{n_v+m_v, 0}} \right) \end{aligned} \quad (4.19)$$

where the sum over $\{\ell_v\}$ in the right hand side runs over $L'_f = L' \cap L_f$ for each ℓ_v associated with one the vertices of the tree other than the root v_0 and the endpoints; moreover, we denoted $\ell_v = (n_v, m_v, l_v, \mathbf{p}_v)$, and v_i is the i -th child of v . Let us note that, thanks to the definition of the trimming map, $D_{\text{sc}}(\ell_v) \leq -\frac{l_v}{12} < 0$ for all $\ell_v \in L'_f$. If we now first sum over the choices of $\mathbf{p}_v$, given

(n_v, m_v, l_v) , for all the vertices v of τ other than the root v_0 and the endpoints, using (4.18) we get:

$$\begin{aligned} \|H_\ell[\tau]\|_w &\leq (\alpha_\gamma/d_\gamma)^{\mathbb{1}(\ell=(0,2,0,\emptyset))} \gamma^{D_{\text{sc}}(\ell)+l/12} \sum_{\{l_v\}} \left(\prod_{v \text{ not e.p.}} K' C_\gamma^{s_v-1} \gamma^{-l_v/12} C_0^{\sum_{i=1}^{s_v} l_{v_i}-l_v} \binom{\sum_{i=1}^s l_{v_i}}{l_v} \right) \\ &\quad \cdot \left(\prod_{v \text{ e.p.}} (K\epsilon)^{\delta_{n_v+m_v,0}} \right) \end{aligned} \quad (4.20)$$

where: $\ell \equiv \ell_{v_0} = (n, m, l, \mathbf{p})$; the sum over $\{l_v\}$ runs over the positive integers, $l_v \geq 1$, for all the vertices v of τ other than the root v_0 and the endpoints; and $K' = 4d_\gamma C_R \gamma^2$. Now, letting $n_{\text{e.p.}}[\tau]$ be the number of endpoints of τ , we have that:

1. the number of vertices of τ that are not endpoints is $\leq 2n_{\text{e.p.}}[\tau]$ (see [1, eq.(J.10)]);
2. $\prod_{v \text{ not e.p.}} C_\gamma^{s_v-1} = C_\gamma^{n_{\text{e.p.}}[\tau]-1}$;
3. $\prod_{v \text{ not e.p.}} C_0^{\sum_{i=1}^{s_v} l_{v_i}-l_v} = C_0^{-l} \prod_{v \text{ e.p.}} C_0^{l_v} \leq C_0^{4n_{\text{e.p.}}[\tau]-l}$;
4. $\sum_{\{l_v\}} \prod_{v \text{ not e.p.}} \gamma^{-l_v/12} \binom{\sum_{i=1}^s l_{v_i}}{l_v} \leq (1 - \gamma^{-1/12})^{-4n_{\text{e.p.}}[\tau]}$ (see [33, Appendix A.6.1]).

Therefore, for any $\gamma > 1$,

$$\|H_\ell[\tau]\|_w \leq (\alpha_\gamma/d_\gamma)^{\mathbb{1}(\ell=(0,2,0,\emptyset))} \gamma^{D_{\text{sc}}(\ell)} C_\gamma^{-1} (\gamma^{\frac{1}{12}}/C_0)^l (K\epsilon)^{-(n+m)} \left[\left(\frac{C_0}{1 - \gamma^{-\frac{1}{12}}} \right)^4 (K')^2 C_\gamma K\epsilon \right]^{n_{\text{e.p.}}[\tau]}. \quad (4.21)$$

In view of (4.21), recalling that the number of the trees with k endpoints is less than 4^k , see e.g. [33, Lemma A.1], the sum over trees in the right hand side of (4.3) converges absolutely for any $\gamma > 1$ and ϵ small enough in the weighted L^1 norm (4.5).

Remark 4.1. Note that (4.21) is increasing in C_0 , so that, if desired, for any prescribed $\rho \geq 1$, we can make the factor $\gamma^{D_{\text{sc}}(n,m,l,\mathbf{p})} (\gamma^{\frac{1}{12}}/C_0)^l$ in the right and side of (4.21) smaller than $C_{n,m} \rho^{-l}$ for some $C_{n,m} > 0$, possibly at the cost of increasing C_0 , thus getting:

$$\|H_{n,m,l,\mathbf{p}}[\tau]\|_w \leq (\alpha_\gamma/d_\gamma)^{\mathbb{1}(\ell=(0,2,0,\emptyset))} C_{n,m} \rho^{-l} (C\rho\epsilon)^{n_{\text{e.p.}}[\tau]} (K\epsilon)^{-n-m}, \quad (4.22)$$

for any $\rho \geq 1$ and some ρ -independent constants $C_{n,m}, C, K > 0$.

4.2 The FPE for the local terms. Analyticity of the scaling exponents.

Consider now the components of the FPE associated with the local parts of the effective potential, (3.20). We focus on the last two components of the equation, the first two having been discussed and solved in [1]. Using the definition of D and the second line of (3.19), we see that in those components we can replace $\gamma^{[\psi]-\Delta_1} R_{(1,0,1,\mathbf{0})}^{\ell_1, \dots, \ell_s}(H_{\ell_1}, \dots, H_{\ell_s})$ by $T_{1,0,1,\mathbf{0}}^{1,0,1,\mathbf{0}} S_{(1,0,1,\mathbf{0})}^{\ell_1, \dots, \ell_s}(H_{\ell_1}, \dots, H_{\ell_s})$, and similarly for the component with $\ell = (0, 1, 2, \mathbf{0})$, thus getting that, at the fixed point

$$\begin{aligned} \zeta_1 &= \sum_{s \geq 1} \sum_{(\ell_i)_{i=1}^s}^* T_{1,0,1,\mathbf{0}}^{1,0,1,\mathbf{0}} S_{\ell}^{\ell_1, \dots, \ell_s}(V_{\ell_1}^*, \dots, V_{\ell_s}^*), \\ \zeta_2 &= \sum_{s \geq 1} \sum_{(\ell_i)_{i=1}^s}^* T_{0,1,2,\mathbf{0}}^{0,1,2,\mathbf{0}} S_{\ell}^{\ell_1, \dots, \ell_s}(V_{\ell_1}^*, \dots, V_{\ell_s}^*). \end{aligned} \quad (4.23)$$

To get, out of this, a representation of ζ_1, ζ_2 in terms of a convergent tree expansion, we insert the rewriting (4.3) in the right sides of (4.23), so that

$$\zeta_1 = \sum_{\tau} \tilde{H}_{1,0,1,\mathbf{0}}[\tau], \quad \zeta_2 = \sum_{\tau} \tilde{H}_{0,1,2,\mathbf{0}}[\tau], \quad (4.24)$$

where the tree values $\tilde{H}_{\ell}[\tau]$ with $\ell = (1, 0, 1, \mathbf{0}), (0, 1, 2, \mathbf{0})$ are defined essentially in the same way as in the previous subsection, with the only difference that the root vertex v_0 is associated with the action of an operator $T_{\ell}^{\ell_1, \dots, \ell_{s_{v_0}}}$ rather than $(1 - D)^{-1} R_{\ell}^{\ell_1, \dots, \ell_{s_{v_0}}}$. In other words, for $\ell = (1, 0, 1, \mathbf{0}), (0, 1, 2, \mathbf{0})$, we have $\tilde{H}_{\ell}[\tau] = \sum_{(\ell_i)_{i=1}^{s_{v_0}}} T_{\ell}^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_n}[\tau_{s_{v_0}}])$ where, recalling that $\ell_1, \dots, \ell_{s_{v_0}} \in L'_f$, the values $H_{\ell_i}[\tau_i]$ for $i \in \{1, \dots, s_{v_0}\}$ have been constructed and bounded in the previous subsection. On the other hand, recalling that the weighted L^1 norm of T_{ℓ}^{ℓ} is bounded by 1, we find that for $\ell = (1, 0, 1, \mathbf{0}), (0, 1, 2, \mathbf{0})$ the norms $\|\tilde{H}_{\ell}[\tau]\|_w$ are bounded in a way analogous to (4.21), namely:

$$\|\tilde{H}_{\ell}[\tau]\|_w \leq C'_{\gamma} (K\epsilon)^{-1} (K''\epsilon)^{n_{\text{e.p.}}[\tau]}, \quad (4.25)$$

where we can choose $C'_{\gamma} = C_{\gamma}^{-1} (\gamma^{\frac{1}{12}} / C_0)^2$ and $K'' = \left(\frac{C_0}{1 - \gamma^{-\frac{1}{12}}} \right)^4 (K')^2 C_{\gamma} K$. Absolute summability over τ follows by the same considerations after (4.21). In conclusion, both ζ_1 and ζ_2 are expressed in terms of absolutely convergent tree expansions. Recalling that, for $\epsilon_0, \delta_0 > 0$ small enough:

- $\lambda = \lambda^*(\epsilon)$ and $\nu = \nu^*(\epsilon)$ are analytic functions of ϵ of order ϵ , for $|\epsilon| < \epsilon_0$;
- the single-scale propagator $g^{(0)}$ depends analytically upon ϵ for $|\epsilon| < \epsilon_0$; therefore, the connected expectation $\mathcal{C}(z)$ in (3.2) (see [1, Appendix D] for the explicit representation of $\mathcal{C}(z)$ in terms of $g^{(0)}$), which enters the definition of the ‘integrating out’ map S_0 and, as a consequence, of the tree values themselves, is analytic in ϵ in the same domain, as well;
- the dilatation operator D in (2.27), which enters the definition of the tree values, is analytic in the scaling exponents Δ_1 and Δ_2 , for $|\Delta_1 - [\psi]| < \delta_0$ and $|\Delta_2 - 2[\psi]| < \delta_0$;

all the tree values $H_{\ell}[\tau]$ are analytic in $(\epsilon, \Delta_1, \Delta_2)$ in the domain

$$\mathcal{D}_0 := \{(\epsilon, \Delta_1, \Delta_2) \in \mathbb{C}^3 : |\epsilon| < \epsilon_0, |\Delta_1 - [\psi]| < \delta_0, |\Delta_2 - 2[\psi]| < \delta_0\}.$$

Therefore, by absolute convergence of the tree expansion, uniform in $\mathcal{D}_0$, analyticity of the sums in the right sides of (4.23) follows in the same domain, by Weierstrass’ theorem on the uniform convergence of sequences of analytic functions. We shall then write:

$$\zeta_1 = F_1(\epsilon, \Delta_1, \Delta_2), \quad \zeta_2 = F_2(\epsilon, \Delta_1, \Delta_2), \quad (4.26)$$

with F_1, F_2 analytic in $\mathcal{D}_0$.

4.2.1 The scaling exponent Δ_1

From the previous considerations and inspection of perturbation theory, it follows that $\zeta_1 = 0$. In fact, in view of the convergence of the tree expansion $\sum_{\tau} \tilde{H}_{1,0,1,\mathbf{0}}[\tau] \equiv F_1(\epsilon, \Delta_1, \Delta_2)$, in order to prove that $\zeta_1 = 0$ it is enough to prove that any tree contributing to F_1 has vanishing value. Note that the trees contributing to F_1 have one ‘white square’ endpoint (i.e., the first in the second line

of Fig.4, corresponding to the third interaction vertex in Fig.5) and $k \geq 1$ additional endpoints of type ν or λ (i.e., those in the first line of Fig.4, corresponding to the first two interaction vertices in Fig.5). It is straightforward to check that, for any such τ , by applying the definition of tree value, $[\tilde{H}_{1,0,1,0}[\tau](x_1, z_1)]_{a_1, b_1}$ is local, i.e., it is equal to $\delta_{x_1, z_1} F_{a_1, b_1}(\tau)$ for some $F_{a, b}(\tau)$ that is a (in general infinite, absolutely convergent) linear combination of terms of the following form:

$$\sum_{b_2} \int g_{a, b_2}^{(h)}(x_1 - z_2) f_{b_2, b}(z_2) dz_2, \quad (4.27)$$

for appropriate functions $f_{b', b}$ (with the correct $Sp(N)$ invariance properties, such that (4.27) is in fact proportional to $\delta_{a, b}$ and independent of a). For example, it is instructive to check that the sum of the values of the trees with one white square endpoint and one additional endpoint, either of type ν or λ , is

$$\sum_{b_2} \int \left[2\nu g_{a, b_2}^{(0)}(x_1 - z_2) \Omega_{b_2, b} + 4\lambda \sum_{b_3, b_4} q_{b_2 b_3 b_4 b} \sum_{h \geq 0} \gamma^{(d + \Delta_1 - 5[\psi])h} g_{ab_2}^{(h)}(x_1 - z_2) g_{b_3 b_4}^{(0)}(0) \right] dz_2, \quad (4.28)$$

with q_{abcd} the totally antisymmetric tensor defined after (2.34). Now, the key remark is that (the summand over b_2 in) (4.27) is proportional to $\hat{g}_{a, b_2}^{(h)}(0)$, which is zero, because the support of the Fourier transform of $g^{(h)}$ does not contain the origin. Therefore, as anticipated above, all the contributions to F_1 vanishing, thus implying that $\zeta_1 = 0$ and, therefore, recalling (3.22), $\Delta_1 = [\psi]$.

Remark 4.2. The considerations above, leading to the conclusion that the scaling exponent of the external field ϕ , coupled linearly to the fluctuation field ψ , has scaling dimension $[\phi] = d - [\psi]$, is a general fact, valid for any infrared, critical, even, interacting theory, treatable perturbatively close to a Gaussian fixed point: it implies that, in general, the critical exponent describing the asymptotic, large distance, polynomial decay of the interacting two point function $\langle \psi(x) \psi(y) \rangle$ is the same as the one associated with the Gaussian part of the infrared RG fixed point. This is true, in particular, in theories where the fluctuation field ψ has an infrared anomalous scaling dimension (i.e., $[\psi]$ differs from the naive scaling dimension associated with the bare propagator), as it is the case, for example, for models in the ‘Luttinger liquid’ universality class, in their fermionic formulation (i.e., models admitting a large distance effective description in terms of 2D spinless fermions with quartic interaction), see [33].

4.2.2 The scaling exponent Δ_2

Contrary to the case of ζ_1 , the right hand side of the equation for ζ_2 does not vanish. Recalling that Δ_2 is related to ζ_2 via (3.22), we rewrite the second equation of (4.26) as

$$\zeta_2 = f_2(\epsilon, \zeta_2), \quad (4.29)$$

where $f_2(\epsilon, \zeta_2) := F_2(\epsilon, [\psi], 2[\psi] - \log_\gamma(1 + \zeta_2))$ is analytic in (ϵ, ζ_2) in a small complex neighborhood of the origin. An explicit computation shows that $f_2(\epsilon, \zeta_2) = -2\epsilon^{\frac{N-2}{N-8}} \log \gamma + O(\epsilon^2, \epsilon \zeta_2)$, see Appendix A. Therefore, applying the analytic implicit function theorem, see e.g. [34, Section 5.11], it follows that (4.29) admits a unique analytic solution $\zeta_2(\epsilon)$ in a neighborhood of the origin, satisfying $\zeta_2(\epsilon) = -2\epsilon^{\frac{N-2}{N-8}} \log \gamma + O(\epsilon^2)$.

This concludes the proof of the part of the statement of Theorem 2.3 concerning the analyticity of V^* and of the scaling exponents, as well as the facts that $\Delta_1 = [\psi]$ and $\Delta_2 = 2[\psi] + \eta_2$, with η_2 as in (2.42).

5 Pointwise bounds on the response functions

In this section, we prove the following result on the pointwise convergence of the limits in (2.50)-(2.51). Recall that the scaling exponents satisfy $\Delta_1 = [\psi]$ and $\Delta_2 = 2[\psi] + \eta_2$, with η_2 as in (2.42), as proved in the previous section. We make use of the same notations and conventions on trees, tree values, components of the renormalization map, etc., as in the previous two sections. Moreover, whenever possible, we will keep the flavor indices implicit (e.g., in the first line of (5.1) below we drop the a, b indices labelling $\mathcal{G}^*$, which should be then thought of as an $N \times N$ anti-symmetric matrix).

Proposition 5.1. *There exists $\epsilon_0 > 0$ such that the limits in (2.50)-(2.51) exist and are analytic in $|\epsilon| < \epsilon_0$, and, letting $\mathbf{x} = (x, 0)$ and $\mathbf{y} = (y, 0)$, they can be explicitly written as*

$$\begin{aligned}\mathcal{G}^*(x) &= 2 \sum_{\tau} \sum_{h \in \mathbb{Z}} \gamma^{2h\Delta_1} \sum_{(\ell_i)_{i=1}^{s_{v_0}}}^* S_{2,0,0,\emptyset}^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}])(\gamma^h \mathbf{x}), \\ \mathcal{F}^*(y) &= 2 \sum_{\tau} \sum_{h \in \mathbb{Z}} \gamma^{2h\Delta_2} \sum_{(\ell_i)_{i=1}^{s_{v_0}}}^* S_{0,2,0,\emptyset}^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}])(\gamma^h \mathbf{y}),\end{aligned}\tag{5.1}$$

with the understanding that the sums in the right hand sides are absolutely summable in h and τ . Moreover, for any $\alpha > 0$ small enough, there exists $C_\alpha > 0$ such that

$$\begin{aligned}|2\gamma^{2h\Delta_1} V_{2,0,0,\emptyset}^*(\gamma^h \mathbf{x}) - \mathcal{G}^*(x)| &\leq \frac{C_\alpha}{|x|^{2\Delta_1}} \left(\min\{1, \gamma^h |x|\} \right)^{2[\psi]-\alpha}, \\ |2\gamma^{2h\Delta_2} V_{0,2,0,\emptyset}^*(\gamma^h \mathbf{y}) - \mathcal{F}^*(y)| &\leq \frac{C_\alpha}{|y|^{2\Delta_2}} \left(\min\{1, \gamma^h |y|\} \right)^{2[\psi]-\alpha}.\end{aligned}\tag{5.2}$$

As proved in (2.52), Proposition 5.1 implies the scale invariance property (2.41). Therefore, in view of the fact that the other statements of Theorem 2.3 have already been proved above, see the comment at the end of Section 4, Proposition 5.1 implies Theorem 2.3.

Proof of Proposition 5.1. Using the tree representation (4.3), we write:

$$\begin{aligned}\gamma^{2h\Delta_1} V_{2,0,0,\emptyset}^*(\gamma^h \mathbf{x}) &= \gamma^{2h\Delta_1} \sum_{\tau} H_{2,0,0,\emptyset}[\tau](\gamma^h \mathbf{x}), \\ \gamma^{2h\Delta_2} V_{0,2,0,\emptyset}^*(\gamma^h \mathbf{y}) &= \gamma^{2h\Delta_2} \sum_{\tau} H_{0,2,0,\emptyset}[\tau](\gamma^h \mathbf{y}),\end{aligned}\tag{5.3}$$

with $H_\ell[\tau]$ recursively defined as in (4.4). We emphasize that the trees τ contributing to the sum in the first line of (5.3) have two ‘white square’ endpoints, and those contributing to the second line have two ‘white rombus’ endpoints, see Fig.4. Using (3.19), the identity $(1 - D)^{-1} D = \sum_{k \geq 1} D^k$ with D as in (2.27), the definition of δ_{sc} in (2.28) (from which $\delta_{\text{sc}}(2, 0, 0, \emptyset) = 2\Delta_1$ and $\delta_{\text{sc}}(0, 2, 0, \emptyset) = 2\Delta_2$), and the fact that for $\ell = (2, 0, 0, \emptyset), (0, 2, 0, \emptyset)$ the trimming map is the identity, we get (after

renaming $h + k \equiv h'$):

$$\begin{aligned}\gamma^{2h\Delta_1} H_{2,0,0,\emptyset}[\tau](\gamma^h \mathbf{x}) &= \sum_{h' > h} \gamma^{2h'\Delta_1} \sum_{(\ell_i)_{i=1}^{s_{v_0}}}^* S_{2,0,0,\emptyset}^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}])(\gamma^{h'} \mathbf{x}), \\ \gamma^{2h\Delta_2} H_{0,2,0,\emptyset}[\tau](\gamma^h \mathbf{y}) &= \sum_{h' > h} \gamma^{2h'\Delta_2} \sum_{(\ell_i)_{i=1}^{s_{v_0}}}^* S_{0,2,0,\emptyset}^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}])(\gamma^{h'} \mathbf{y}).\end{aligned}\tag{5.4}$$

If we multiply by 2 both sides, sum over τ and take $h \rightarrow -\infty$, we obtain the representations (5.1), provided that the sums in the right hand sides are absolutely summable in τ and h .

The right hand sides of (5.4) can be bounded via the following lemma that, for later purposes, is formulated in greater generality than required for the moment. In order to state the lemma, we define a mixed L^1/L^∞ norm: using the same notations as in the definition of the weighted L^1 norm (4.5), we let, for $\ell = (n, m, l, \mathbf{p}) \in L$,

$$\|H_\ell[\tau](\mathbf{x}, \mathbf{y})\| := \max_{\mathbf{a}, \mathbf{B}} \int d\mathbf{z}_\mathbf{B} |H_\ell[\tau](\mathbf{a}, \mathbf{x}, \mathbf{y}, \mathbf{B}, \mathbf{z}_\mathbf{B})| \tag{5.5}$$

with the understanding that, if $\ell = (n, m, 0, \emptyset)$, then $\|H_{n,m,0,\emptyset}[\tau](\mathbf{x}, \mathbf{y})\|$ should be interpreted as being equal to $\max_{\mathbf{a}} |H_{n,m,0,\emptyset}[\tau](\mathbf{a}, \mathbf{x}, \mathbf{y})|$.

Lemma 5.2. *Consider a tree τ contributing to one of the sums in the right sides of (5.3), denote by $\tau_1, \dots, \tau_{s_{v_0}}$ its subtrees rooted in v_0 , and by $n_{e.p.}[\tau]$ be the number of its endpoints. Let $\mathbf{x} = (x, 0)$ and $\mathbf{y} = (y, 0)$. For any $\alpha > 0$ sufficiently small, there exists $C = C(\alpha) > 0$ such that, for any $k \in \mathbb{Z}$, any $l \geq 0$ even and any $\mathbf{p} \in \{0, 1\}^l$,*

$$\begin{aligned}\sum_{(\ell_i)_{i=1}^{s_{v_0}}}^* \left\| D^k S_{2,0,l,\mathbf{p}}^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}])(\mathbf{x}) \right\| \\ \leq C(C\epsilon)^{n_{e.p.}[\tau]-2} \gamma^{k(2\Delta_1-l[\psi]-\|\mathbf{p}\|_1)} e^{-\frac{C}{2}(\gamma^{k-1}|x|)^\sigma} (\min\{1, \gamma^k|x|\})^{-\alpha},\end{aligned}\tag{5.6}$$

with D the dilatation operator and $\bar{C}$ the same constant as in (4.6), and

$$\begin{aligned}\sum_{(\ell_i)_{i=1}^{s_{v_0}}}^* \left\| D^k S_{0,2,l,\mathbf{p}}^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}])(\mathbf{y}) \right\| \\ \leq C(C\epsilon)^{n_{e.p.}[\tau]-2} \gamma^{k(2\Delta_2-l[\psi]-\|\mathbf{p}\|_1)} e^{-\frac{\bar{C}}{2}(\gamma^{k-1}|y|)^\sigma} (\min\{1, \gamma^k|y|\})^{-2\Delta_2+2[\psi]-\alpha}.\end{aligned}\tag{5.7}$$

Assuming the validity of this lemma, the proof of Proposition 5.1 goes as follows. Let us focus, e.g., on the component with $\ell = (2, 0, 0, \emptyset)$, the case $\ell = (0, 2, 0, \emptyset)$ being analogous. Plugging (5.6) with $k = h'$, $l = 0$ and $\mathbf{p} = \emptyset$ in the right hand side of the first line of (5.4), recalling that $D^{h'} S_{2,0,l,\mathbf{p}}^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}])(\mathbf{x})$ is just a rewriting of $\gamma^{2h'\Delta_1} S_{2,0,l,\mathbf{p}}^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}])(\gamma^{h'} \mathbf{x})$, we find:

$$\begin{aligned}\gamma^{2h\Delta_1} |H_{2,0,0,\emptyset}[\tau](\gamma^h \mathbf{x})| &\leq C(C\epsilon)^{n_{e.p.}[\tau]-2} |x|^{-2\Delta_1} \\ &\cdot \sum_{h' > h} e^{-\frac{C}{2}(\gamma^{h'-1}|x|)^\sigma} \max\{(\gamma^{h'}|x|)^{2\Delta_1}, (\gamma^{h'}|x|)^{2\Delta_1-\alpha}\}.\end{aligned}\tag{5.8}$$

Now note that, for any $\beta, \kappa > 0$, letting $h_x := \lfloor \log_\gamma |x|^{-1} \rfloor$, $(\gamma^{h'} |x|)^\beta e^{-\kappa(\gamma^{h'} |x|)^\sigma}$ is bounded from above by $\gamma^{\beta(h'-h_x)} e^{-\kappa\gamma^{\sigma(h'-h_x-1)}}$, which is summable in h' over $\mathbb{Z}$. Therefore,

$$\sum_{h' > h} (\gamma^{h'} |x|)^\beta e^{-\kappa(\gamma^{h'} |x|)^\sigma} \leq C_{\beta, \kappa} := \sum_{h \in \mathbb{Z}} \gamma^{\beta h} e^{-\kappa\gamma^{\sigma(h-1)}}, \quad (5.9)$$

uniformly in h and $|x|$. For later purpose, let us also observe that

$$\begin{aligned} \sum_{h' \leq h} (\gamma^{h'} |x|)^\beta e^{-\kappa(\gamma^{h'} |x|)^\sigma} &\leq \sum_{h' \leq h} \gamma^{\beta(h'-h_x)} e^{-\kappa\gamma^{\sigma(h'-h_x-1)}} \\ &\leq \begin{cases} C_{\beta, \kappa} & \text{if } h > h_x \\ \gamma^{\beta(h-h_x)} / (1 - \gamma^{-\beta}) & \text{if } h \leq h_x \end{cases} \leq C'_{\beta, \kappa} (\min\{1, \gamma^h |x|\})^\beta, \end{aligned} \quad (5.10)$$

for a suitable $C'_{\beta, \kappa} > 0$. Using (5.9) in (5.8), we find that, for any $\alpha > 0$ small enough,

$$\gamma^{2h\Delta_1} |H_{(2,0,0,\emptyset)}[\tau](\gamma^h \mathbf{x})| \leq C'(C\epsilon)^{n_{e.p}[\tau]-2} |x|^{-2\Delta_1}, \quad (5.11)$$

for some $C' > 0$, uniformly in h . Since the right hand side of this inequality is summable over τ , the sum being bounded by $(\text{const.})|x|^{-2\Delta_1}$, this proves the absolute convergence of the sum in the right hand side of the first line of (5.1) and, therefore, as already observed after (5.4), it implies the very validity of the first line of (5.1).

Concerning the difference $2\gamma^{2h\Delta_1} V_{2,0,0,\emptyset}^*(\gamma^h \mathbf{x}) - \mathcal{G}^*(x)$, using (5.1), (5.3), (5.4) and (5.6) with $k = h'$, $l = 0$ and $\mathbf{p} = \emptyset$, we find:

$$\begin{aligned} &|2\gamma^{2h\Delta_1} V_{2,0,0,\emptyset}^*(\gamma^h \mathbf{x}) - \mathcal{G}^*(x)| \\ &\leq 2 \sum_{\tau} \sum_{h' \leq h} \gamma^{2h'\Delta_1} \sum_{(\ell_i)_{i=1}^{s_{v_0}}}^* |S_{2,0,0,\emptyset}^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}]) (\gamma^{h'} \mathbf{x})| \\ &\leq 2C|x|^{-2\Delta_1} \sum_{\tau} (C\epsilon)^{n_{e.p}[\tau]-2} \sum_{h' \leq h} e^{-\frac{C}{2}(\gamma^{h'-1}|x|)^\sigma} \max\{(\gamma^{h'} |x|)^{2\Delta_1}, (\gamma^{h'} |x|)^{2\Delta_1-\alpha}\}. \end{aligned} \quad (5.12)$$

Now, using (5.10), we find that the sum over $h' \leq h$ in the last line is bounded from above by $(\text{const.})(\min\{1, \gamma^h |x|\})^{2\Delta_1-\alpha}$. Moreover, recalling that the number of trees with k endpoints is smaller than 4^k , see [33, Lemma A.1], and noting that the trees contributing to the sum in the last line have at least two endpoints (because τ has at least two ‘white square’ endpoints), we see that $(C_\alpha \epsilon)^{n_{e.p}[\tau]-2}$ is summable over τ , and the sum is bounded by a positive constant independent of ϵ . In conclusion,

$$|2\gamma^{2h\Delta_1} V_{2,0,0,\emptyset}^*(\gamma^h \mathbf{x}) - \mathcal{G}^*(x)| \leq C'|x|^{-2\Delta_1} (\min\{1, \gamma^h |x|\})^{2\Delta_1-\alpha},$$

which is the desired estimate in the first line of (5.2), up to a redefinition of C_α .

The proof of the second line of (5.1) and of (5.2) is completely analogous and left to the reader. $\square$

We are left with proving Lemma 5.2.

Proof of Lemma 5.2. We focus on the proof of (5.6), the one of (5.7) being analogous (we will make a few comments at the end on the minor differences between the two cases). We recall that $\mathbf{x} = (x, 0)$.

In order to obtain a point-wise bound on the left hand side of (5.6), we intend to apply iteratively the definition of tree value, similarly to what we did in Section 4. We recall that the trees τ contributing to the sum in the first line of (5.3) have two ‘white square’ endpoints, which will be denoted v_1^* and v_2^* . Note that the coordinates associated with these endpoints are fixed, i.e., not integrated out in the computation of the tree value: this implies that, for the purpose of recursively deriving bounds on the values of the subtrees of τ , we must be careful in proceeding slightly differently, depending on whether the subtree under consideration contains both v_1^* and v_2^* , or just one of them, or none.

We define v_{12}^* to be the rightmost vertex of τ that is an ancestor both of v_1^* and of v_2^* , and we let n_τ be the length of the path connecting the root vertex v_0 with v_{12}^* , see Fig.8 below (where $n \equiv n_\tau$).

Let us first discuss the case that $n_\tau = 0$, i.e., v_1^* and v_2^* belong to two different subtrees of τ rooted in two distinct children vertices of $v_0 \equiv v_{12}^*$, called v_1 and v_2 , respectively; the two subtrees rooted in v_1 and v_2 will be denoted by τ_1 and τ_2 , respectively, see Figure 7.




Figure 7: Proof of Lemma 5.2, the case with $n_\tau = 0$. The two subtrees rooted in v_1 and v_2 are denoted by τ_1 and τ_2 .

In this case, recalling the definition (5.5) and proceeding as in the proof of (4.15), we get:

$$\begin{aligned}
& \left\| D^k S_{2,0,l,\mathbf{p}}^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}])(\mathbf{x}) \right\| e^{\bar{C}(\gamma^{k-1}|x|)^\sigma} \\
&= \gamma^{k\delta_{sc}(2,0,l,\mathbf{p})} \gamma^{-kld} \left\| S_{2,0,l,\mathbf{p}}^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}])(\gamma^k \mathbf{x}) \right\| e^{\bar{C}(\gamma^{k-1}|x|)^\sigma} \\
&\leq \frac{\gamma^{k(2\Delta_1 - l[\psi] - \|\mathbf{p}\|_1)}}{s_{v_0}!} C_0^{\sum_{i=1}^{s_{v_0}} l_i - l} N_{(2,0,l,\mathbf{p})}^{\ell_1, \dots, \ell_{s_{v_0}}} \left(\sum_{\mathcal{T}} \prod_{\substack{(z,z') \in \mathcal{T} \\ (z,z') \neq \ell_0}} N^2 d^2 \|M\|_w \right). \quad (5.13) \\
&\cdot \sup_{x'} \{ N^2 d^2 M(x') e^{\bar{C}(|x'|/\gamma)^\sigma} \} \prod_{i=1}^{s_{v_0}} \|H_{\ell_i}[\tau_i]\|_w,
\end{aligned}$$

where $\bar{C}$ is the same as in (4.6), $\delta_{sc}(2,0,l,\mathbf{p})$ is given by (2.28), and γ^{-kld} comes from the rescaling of the $\mathbf{z}$ variable in $\|\cdot\|$. The summation in the right hand side of (5.13) is over the anchored trees described in [1, Appendix D.4] (see also (4.13)), ℓ_0 is an element of $\mathcal{T}$, chosen arbitrarily among

those in the path from the group of points associated with τ_1 and the one associated with τ_2 (say, the one closest to the group associated with τ_1), and the factor $\sup_{x'} \{M(x')e^{C(|x'|/\gamma)^\sigma}\}$ comes from the factor M associated with ℓ_0 . Now, recalling that $M(x) = C_{\chi_1} e^{-C_{\chi_2}|x/\gamma|^\sigma}$ and $\bar{C} \equiv \frac{1}{2}C_{\chi_2}$, bounding the sum over $\mathcal{T}$ as in [1, Appendix D.5], we find¹⁰

$$\begin{aligned} & \sum_{(\ell_i)_{i=1}^{s_{v_0}}}^* \left\| D^k S_{2,0,l,\mathbf{p}}^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}])(\mathbf{x}) \right\| \\ & \leq N^2 d^2 C_{\chi_1} \gamma^{k(2\Delta_1 - l[\psi] - \|\mathbf{p}\|_1)} e^{-\bar{C}(\gamma^{k-1}|x|)^\sigma} \sum_{(\ell_i)_{i=1}^{s_{v_0}}}^* C_\gamma^{s_{v_0}-2} (4C_0)^{\sum_{i=1}^{s_{v_0}} l_i - l} N_{(2,0,l,\mathbf{p})}^{\ell_1, \dots, \ell_{s_{v_0}}} \prod_{i=1}^{s_{v_0}} \|H_{\ell_i}[\tau_i]\|_w. \end{aligned} \quad (5.14)$$

with $C_\gamma = N^2 d^2 \|M\|_w$. Now, in the right side of (5.14) we bound $N_{(2,0,l,\mathbf{p})}^{\ell_1, \dots, \ell_{s_{v_0}}}$ from above by $(\sum_{i=1}^{s_{v_0}} l_i)$, see (4.16), and we bound $\|H_{\ell_i}[\tau_i]\|_w \leq C(\rho')^{-l_i} (C\epsilon)^{n_{\text{e.p.}}[\tau_i] - n_i}$ via (4.22); here n_i is the first component of $\ell_i = (n_i, m_i, l_i, \mathbf{p}_i)$ (by construction $n_1 = n_2 = 1$ and $n_i = 0$ for $i > 2$) and C is a ρ' -dependent constant; therefore, choosing $\rho' \geq 1$ sufficiently large, we can easily sum over $(\ell_i)_{i=1}^{s_{v_0}}$, thus obtaining (noting that $s_{v_0} \leq n_{\text{e.p.}}[\tau]$):

$$\begin{aligned} & \sum_{(\ell_i)_{i=1}^{s_{v_0}}}^* \left\| D^k S_{2,0,l,\mathbf{p}}^{\ell_1, \dots, \ell_{s_{v_0}}} (H_{\ell_1}[\tau_1], \dots, H_{\ell_{s_{v_0}}}[\tau_{s_{v_0}}])(\mathbf{x}) \right\| \\ & \leq C' (C'\epsilon)^{n_{\text{e.p.}}[\tau] - 2} (4C_0)^{-l} \gamma^{k(2\Delta_1 - l[\psi] - \|\mathbf{p}\|_1)} e^{-\bar{C}(\gamma^{k-1}|x|)^\sigma}, \end{aligned} \quad (5.15)$$

for some $C' > 0$, which proves (5.6) for $n_\tau = 0$ (simply because $1 \leq (\min\{1, \gamma^k|x|\})^{-\alpha}$ for $\alpha > 0$), provided we choose $C_0 \geq 1/4$ (the freedom to choose C_0 as large as desired follows, as already observed after (5.13), from the monotonicity of the right hand side of (5.15) on the choice of C_0 : recall, in fact, that C' is proportional to C_0).

Let us discuss next the case $n_\tau \geq 1$, in which v_1^* and v_2^* both belong to a common subtree among those rooted in the vertices immediately following v_0 on τ , say to τ_1 . In this case, letting $n \equiv n_\tau$, we denote by $v_0, v_1, \dots, v_n \equiv v_{12}^*$ the vertices of τ in the path from v_0 to v_{12}^* , as shown in Fig. 8. For later reference, we introduce (and partly recall) the following notations and definitions. For any vertex v of τ , we denote by $\ell_v = (n_v, m_v, l_v, \mathbf{p}_v)$ the components of the label associated with v : if $v = v_0$, then $\ell_{v_0} \equiv (2, 0, l, \mathbf{p})$; if $v \neq v_0$ is not an endpoint, then ℓ_v is summed over L'_f , while, if v is an endpoint, ℓ_v takes one of the values in $\{(n, m, l, \mathbf{0})\}_{(n,m,l) \in \mathcal{L}}$, depending on the nature of the endpoint, see Fig. 4. Moreover, for any vertex v of τ that is not an endpoint, we denote by

¹⁰The bound (5.14) is similar to (4.17) for the same value of $\ell = (2, 0, l, \mathbf{p})$ (for which the trimming operator T in the first line of (4.17) acts as the identity and, therefore, $\mathcal{N}_\ell^{\ell_1, \dots, \ell_s} = N_\ell^{\ell_1, \dots, \ell_s}$), the most relevant differences being that: (1) the operator $\sum_{h \geq 1} D^h = \frac{D}{1-D}$ in the first line of (4.17) is replaced by the action of D^k in the left hand side of (5.14); (2) in the left hand side of (5.14) we are not integrating over the coordinate x , contrary to the case of (4.17). Correspondingly, the proof of (5.14) is similar to that (4.17), and the details are left to the reader. Let us just mention that the two differences (1) and (2) mentioned above explain the presence of the decay factor $e^{-\bar{C}(\gamma^{k-1}|x|)^\sigma}$ in (5.14), as well as that of the dimensional factor $\gamma^{k[D_{\text{sc}}(2,0,l,\mathbf{p})+d]} = \gamma^{k(2\Delta_1 - l[\psi] - \|\mathbf{p}\|_1)}$ instead of $d_\gamma \gamma^{D_{\text{sc}}(2,0,l,\mathbf{p})}$ (roughly speaking, the fact that in (5.14) we have ‘one integration less’, in combination with the presence of D^k , comes with a dimensional factor γ^{kd} ; on the other hand, the factor $d_\gamma \gamma^{D_{\text{sc}}(2,0,l,\mathbf{p})}$ in (4.17) is a norm bound on $D(1-D)^{-1}$).




Figure 8: Proof of Lemma 5.2, the case with $n = n_\tau \geq 1$.

$v_i(v)$ the i -th child of v and, if τ_v is a tree rooted in v , $\tau_i(v)$ is the subtree of τ_v rooted in $v_i(v)$. Correspondingly, we let $\ell_i(v) \equiv \ell_{v_i(v)}$ and denote $\ell_i(v) = (n_i(v), m_i(v), l_i(v), \mathbf{p}_i(v))$. Note that, for $v_0, v_1, \dots, v_n$ in Figure 8, we have $\ell_{v_i} = (2, 0, l_{v_i}, \mathbf{p}_{v_i})$, where, for any $1 \leq i \leq n$, l_{v_i} is summed over the even, positive, integers.

We will iteratively use the following basic estimates:

1. For the vertices v_j , $j = 0, \dots, n-1$ in Fig.8, we use:

$$\begin{aligned}
& \left\| S_{\ell_{v_j}}^{\ell_1(v_j), \dots, \ell_{s_{v_j}}(v_j)} (H_{\ell_1(v_j)}[\tau_1(v_j)], \dots, H_{\ell_{s_{v_j}}(v_j)}[\tau_{s_{v_j}}(v_j)])(\mathbf{x}) \right\| \\
& \leq \frac{1}{s_{v_j}!} C_0^{\sum_{i=1}^{s_{v_j}} l_i(v_j) - l_{v_j}} N_{\ell_{v_j}}^{\ell_1(v_j), \dots, \ell_{s_{v_j}}(v_j)} \left(\sum_{\mathcal{T}} \prod_{(z, z') \in \mathcal{T}} N^2 d^2 \|M\|_1 \right) \\
& \quad \cdot \left(\prod_{i=2}^{s_{v_j}} \|H_{\ell_i(v_j)}[\tau_i(v_j)]\|_w \right) \|H_{\ell_1(v_j)}[\tau_1(v_j)](\mathbf{x})\|,
\end{aligned} \tag{5.16}$$

which is proved in the same way as (4.15) or (5.13). Bounding once again the sum over $\mathcal{T}$ as in [1, Appendix D.5], we further obtain

$$\begin{aligned}
& \left\| S_{\ell_{v_j}}^{\ell_1(v_j), \dots, \ell_{s_{v_j}}(v_j)} (H_{\ell_1(v_j)}[\tau_1(v_j)], \dots, H_{\ell_{s_{v_j}}(v_j)}[\tau_{s_{v_j}}(v_j)])(\mathbf{x}) \right\| \\
& \leq C_\gamma^{s_{v_j}-1} (4C_0)^{\sum_{i=1}^{s_{v_j}} l_i(v_j) - l_{v_j}} N_{\ell_{v_j}}^{\ell_1(v_j), \dots, \ell_{s_{v_j}}(v_j)} \left(\prod_{i=2}^{s_{v_j}} \|H_{\ell_i(v_j)}[\tau_i(v_j)]\|_w \right) \|H_{\ell_1(v_j)}[\tau_1(v_j)](\mathbf{x})\|,
\end{aligned} \tag{5.17}$$

with $C_\gamma = N^2 d^2 \|M\|_w$.

2. Using (4.4), with $R_\ell^{\ell_1, \dots, \ell_{s_{v_0}}} = D S_\ell^{\ell_1, \dots, \ell_{s_{v_0}}}$ for any ℓ of the form $\ell = (2, 0, l, \mathbf{p})$, see (3.19), we rewrite and bound each of the factors $\|H_{\ell_{v_i}}[\tau_{v_i}](\mathbf{x})\|$ in the right hand side of (5.16) (note

that, for $j = 0, \dots, n-1$, $\ell_1(v_j) = \ell_{v_i}$ and $\tau_1(v_j) = \tau_{v_i}$, with $i = j+1 = 1, \dots, n$ as follows:

$$\begin{aligned}
& \left\| H_{\ell_{v_i}}[\tau_{v_i}](\mathbf{x}) \right\| \\
& \leq \sum_{k_{v_i} \geq 1} \sum_{(\ell_j(v_i))_{j=1}^{s_{v_i}}}^* \left\| D^{k_{v_i}} S_{\ell_{v_i}}^{\ell_1(v_i), \dots, \ell_{s_{v_i}}(v_i)} (H_{\ell_1(v_i)}[\tau_1(v_i)], \dots, H_{\ell_{s_{v_i}}(v_i)}[\tau_{s_{v_i}}(v_i)])(\mathbf{x}) \right\| \\
& = \sum_{k_{v_i} \geq 1} \gamma^{k_{v_i} \delta_{sc}(2, 0, l_{v_i}, \mathbf{p}_{v_i})} \gamma^{-k_{v_i} dl_{v_i}} \cdot \\
& \quad \cdot \sum_{(\ell_j(v_i))_{j=1}^{s_{v_i}}}^* \left\| S_{\ell_{v_i}}^{\ell_1(v_i), \dots, \ell_{s_{v_i}}(v_i)} (H_{\ell_1(v_i)}[\tau_1(v_i)], \dots, H_{\ell_{s_{v_i}}(v_i)}[\tau_{s_{v_i}}(v_i)])(\gamma^{k_{v_i}} \mathbf{x}) \right\|,
\end{aligned} \tag{5.18}$$

where we used that $D(1-D)^{-1} = \sum_{k_{v_i} \geq 1} D^{k_{v_i}}$ together with (2.27), and $\gamma^{-k_{v_i} dl_{v_i}}$ comes from the rescaling of the $\mathbf{z}$ variables within the definition of the $\|\cdot\|$ norm.

Let us now explain how to apply these estimates for bounding the left hand side of (5.6): using once again the definition of D in (2.27), the fact that $\delta_{sc}(2, 0, l, \mathbf{p}) - ld = 2\Delta_1 - l[\psi] - \|\mathbf{p}\|_1$, and the bound (5.17) with $j = 0$ (note that $\ell_{v_0} = (2, 0, l, \mathbf{p})$ and that the labels $\ell_1, \dots, \ell_{s_{v_0}}$ in (5.6) must be identified with $\ell_1(v_0), \dots, \ell_{s_{v_0}}(v_0)$ in (5.17)), we get

$$\begin{aligned}
& \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}}^* \left\| D^k S_{2, 0, l, \mathbf{p}}^{\ell_1(v_0), \dots, \ell_{s_{v_0}}(v_0)} (H_{\ell_1(v_0)}[\tau_1(v_0)], \dots, H_{\ell_{s_{v_0}}(v_0)}[\tau_{s_{v_0}}(v_0)])(\mathbf{x}) \right\| \\
& \leq \gamma^{k(2\Delta_1 - l[\psi] - \|\mathbf{p}\|_1)} \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}}^* C_\gamma^{s_{v_0}-1} (4C_0)^{\sum_{i=1}^{s_{v_0}} l_i(v_0) - l_{v_0}} N_{\ell_{v_0}}^{\ell_1(v_0), \dots, \ell_{s_{v_0}}(v_0)} \\
& \quad \cdot \left(\prod_{i=2}^{s_{v_0}} \|H_{\ell_i(v_0)}[\tau_i(v_0)]\|_w \right) \left\| H_{\ell_1(v_0)}[\tau_1(v_0)](\gamma^k \mathbf{x}) \right\|.
\end{aligned} \tag{5.19}$$

In order to bound the last factor in the last line, we apply (5.18) with $i = 1$ and $v_1 \equiv v_1(v_0)$, thus getting:

$$\begin{aligned}
& \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}}^* \left\| D^k S_{2, 0, l, \mathbf{p}}^{\ell_1(v_0), \dots, \ell_{s_{v_0}}(v_0)} (H_{\ell_1(v_0)}[\tau_1(v_0)], \dots, H_{\ell_{s_{v_0}}(v_0)}[\tau_{s_{v_0}}(v_0)])(\mathbf{x}) \right\| \\
& \leq \gamma^{k(2\Delta_1 - l[\psi] - \|\mathbf{p}\|_1)} \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}}^* C_\gamma^{s_{v_0}-1} (4C_0)^{\sum_{i=1}^{s_{v_0}} l_i(v_0) - l_{v_0}} N_{\ell_{v_0}}^{\ell_1(v_0), \dots, \ell_{s_{v_0}}(v_0)} \\
& \quad \cdot \left(\prod_{i=2}^{s_{v_0}} \|H_{\ell_i(v_0)}[\tau_i(v_0)]\|_w \right) \sum_{k_{v_1} \geq 1} \gamma^{k_{v_1}(2\Delta_1 - l_{v_1}[\psi] - \|\mathbf{p}_{v_1}\|_1)} \\
& \quad \cdot \sum_{(\ell_i(v_1))_{i=1}^{s_{v_1}}}^* \left\| S_{\ell_{v_1}}^{\ell_1(v_1), \dots, \ell_{s_{v_1}}(v_1)} (H_{\ell_1(v_1)}[\tau_1(v_1)], \dots, H_{\ell_{s_{v_1}}(v_1)}[\tau_{s_{v_1}}(v_1)])(\gamma^{k+k_{v_1}} \mathbf{x}) \right\|,
\end{aligned} \tag{5.20}$$

where we used once again that $\delta_{sc}(2, 0, l_{v_1}, \mathbf{p}_{v_1}) - dl_{v_1} = 2\Delta_1 - l_{v_1}[\psi] - \|\mathbf{p}_{v_1}\|_1$.

In order to bound the last line of (5.20), we iteratively apply (5.17) and (5.18) $n-1$ more times, write at each step $\delta_{sc}(2, 0, l_{v_i}, \mathbf{p}_{v_i}) - dl_{v_i}$ as $2\Delta_1 - l_{v_i}[\psi] - \|\mathbf{p}_{v_i}\|_1$, thus finding:

$$\begin{aligned}
& \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}} \left\| D^k S_{2,0,l,\mathbf{p}}^{\ell_1(v_0), \dots, \ell_{s_{v_0}}(v_0)} (H_{\ell_1(v_0)}[\tau_1(v_0)], \dots, H_{\ell_{s_{v_0}}(v_0)}[\tau_{s_{v_0}}(v_0)])(\mathbf{x}) \right\| \\
& \leq \gamma^{k(2\Delta_1 - l[\psi] - \|\mathbf{p}\|_1)} \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}} \dots \sum_{(\ell_i(v_{n-1}))_{i=1}^{s_{v_{n-1}}}} \sum_{k_{v_1}, \dots, k_{v_n} \geq 1} \left(\prod_{i=1}^n \gamma^{k_{v_i}(2\Delta_1 - l_{v_i}[\psi] - \|\mathbf{p}_{v_i}\|_1)} \right) \\
& \cdot \left(\prod_{i=0}^{n-1} C_{\gamma}^{s_{v_i}-1} (4C_0)^{\sum_{j=1}^{s_{v_i}} l_j(v_i) - l_{v_i}} N_{\ell_{v_i}}^{\ell_1(v_i), \dots, \ell_{s_{v_i}}(v_i)} \prod_{j=2}^{s_{v_i}} \|H_{\ell_{v_j}(v_i)}[\tau_{v_j}(v_i)]\|_w \right) \\
& \cdot \sum_{(\ell_i(v_n))_{i=1}^{s_{v_n}}} \left\| S_{\ell_{v_n}}^{\ell_1(v_n), \dots, \ell_{s_{v_n}}(v_n)} (H_{\ell_1(v_n)}[\tau_1(v_n)], \dots, H_{\ell_{s_{v_n}}(v_n)}[\tau_{s_{v_n}}(v_n)])(\gamma^{k+k_{v_1}+\dots+k_{v_n}} \mathbf{x}) \right\|.
\end{aligned} \tag{5.21}$$

In order to bound the last line, recalling that v_1^* and v_2^* belong to two different subtrees of τ_{v_n} , we use the analogue of Eqs.(5.13)–(5.14), which is proved in the same way:

$$\begin{aligned}
& \sum_{(\ell_i(v_n))_{i=1}^{s_{v_n}}} \left\| S_{\ell_{v_n}}^{\ell_1(v_n), \dots, \ell_{s_{v_n}}(v_n)} (H_{\ell_1(v_n)}[\tau_1(v_n)], \dots, H_{\ell_{s_{v_n}}(v_n)}[\tau_{s_{v_n}}(v_n)])(\gamma^{k+k_{v_1}+\dots+k_{v_n}} \mathbf{x}) \right\| \\
& \leq N^2 d^2 C_{\chi_1} e^{-\bar{C}(\gamma^{k+k_{v_1}+\dots+k_{v_n}-1}|x|)^{\sigma}} \\
& \cdot \sum_{(\ell_i(v_n))_{i=1}^{s_{v_n}}} C_{\gamma}^{s_{v_n}-2} (4C_0)^{\sum_{i=1}^{s_{v_n}} l_i(v_n) - l_{v_n}} N_{\ell_{v_n}}^{\ell_1(v_n), \dots, \ell_{s_{v_n}}(v_n)} \prod_{i=1}^{s_{v_n}} \|H_{\ell_i(v_n)}[\tau_i(v_n)]\|_w.
\end{aligned} \tag{5.22}$$

Plugging (5.22) in (5.21), and bounding each factor $\|H_{\ell_v}[\tau_v]\|_w$ as in (4.22) (with ρ replaced by ρ'), we get:

$$\begin{aligned}
& \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}} \left\| D^k S_{2,0,l,\mathbf{p}}^{\ell_1(v_0), \dots, \ell_{s_{v_0}}(v_0)} (H_{\ell_1(v_0)}[\tau_1(v_0)], \dots, H_{\ell_{s_{v_0}}(v_0)}[\tau_{s_{v_0}}(v_0)])(\mathbf{x}) \right\| \\
& \leq C(C'\rho'\epsilon)^{n_{e.p.}[\tau]-2} (4C_0)^{-l} \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}} \dots \sum_{(\ell_i(v_n))_{i=1}^{s_{v_n}}} \sum_{k_{v_1}, \dots, k_{v_n} \geq 1} e^{-\bar{C}(\gamma^{k+\sum_{i=1}^n k_{v_i}-1}|x|)^{\sigma}} \\
& \cdot N_{\ell_{v_0}}^{\ell_1(v_0), \dots, \ell_{s_{v_0}}(v_0)} \gamma^{k(2\Delta_1 - l[\psi] - \|\mathbf{p}\|_1)} \left(\prod_{i=1}^n N_{\ell_{v_i}}^{\ell_1(v_i), \dots, \ell_{s_{v_i}}(v_i)} \gamma^{k_{v_i}(2\Delta_1 - l_{v_i}[\psi] - \|\mathbf{p}_{v_i}\|_1)} \right) \\
& \cdot \left(\prod_{i=0}^{n-1} \prod_{j=2}^{s_{v_i}} \left(\frac{\rho'}{4C_0} \right)^{-l_j(v_i)} \right) \left(\prod_{j=1}^{s_{v_n}} \left(\frac{\rho'}{4C_0} \right)^{-l_j(v_n)} \right)
\end{aligned} \tag{5.23}$$

for some $C, C' > 0$. We are left with sums over $\ell_{v_1} = (l_{v_1}, \mathbf{p}_{v_1}), \dots, \ell_{v_n} = (l_{v_n}, \mathbf{p}_{v_n})$, and over $(\ell_i(v_0))_{i=2}^{s_{v_0}}, \dots, (\ell_i(v_{n-1}))_{i=2}^{s_{v_{n-1}}}, (\ell_i(v_n))_{i=1}^{s_{v_n}}$. We first sum over $\mathbf{p}_{v_1}, \dots, \mathbf{p}_{v_n}$ and, using (4.16), we

get:

$$\begin{aligned}
& \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}} \left\| D^k S_{2,0,l,\mathbf{p}}^{\ell_1(v_0),\dots,\ell_{s_{v_0}}(v_0)} (H_{\ell_1(v_0)}[\tau_1(v_0)], \dots, H_{\ell_{s_{v_0}}(v_0)}[\tau_{s_{v_0}}(v_0)])(\mathbf{x}) \right\| \\
& \leq C(C'\rho'\epsilon)^{n_{e.p.}[\tau]-2} (4C_0)^{-l} \gamma^{k(2\Delta_1-l[\psi]-\|\mathbf{p}\|_1)} \sum_{(\ell_i(v_0))_{i=2}^{s_{v_0}}} \dots \sum_{(\ell_i(v_{n-1}))_{i=2}^{s_{v_{n-1}}}} \sum_{(\ell_i(v_n))_{i=1}^{s_{v_n}}} \cdot \\
& \cdot \left(\prod_{i=0}^{n-1} \left(\prod_{j=2}^{s_{v_i}} \left(\frac{\rho'}{4C_0} \right)^{-l_j(v_i)} \right) \right) \left(\prod_{j=1}^{s_{v_n}} \left(\frac{\rho'}{4C_0} \right)^{-l_j(v_n)} \right) \sum_{k_{v_1}, \dots, k_{v_n} \geq 1} e^{-\bar{C}(\gamma^{k+\sum_{i=1}^n k_{v_i}-1}|x|)^\sigma} \cdot \\
& \cdot \sum_{l_{v_1}, \dots, l_{v_n} \geq 2} \binom{\sum_{j=1}^{s_{v_0}} l_j(v_0)}{l} \left(\prod_{i=1}^n \binom{\sum_{j=1}^{s_{v_i}} l_j(v_i)}{l_{v_i}} \right) \gamma^{k_{v_i}(2\Delta_1-l_{v_i}[\psi])},
\end{aligned} \tag{5.24}$$

where, if $s_{v_{i_0}} = 1$ for some $i_0 \in \{0, \dots, n-1\}$, then $\sum_{(\ell_i(v_{i_0}))_{i=2}^{s_{v_{i_0}}}} \left(\prod_{j=2}^{s_{v_{i_0}}} \left(\frac{\rho'}{4C_0} \right)^{-l_j(v_{i_0})} \right)$ should be interpreted as being equal to 1. In the last line, we bound $\left(\prod_{j=1}^{s_{v_0}} l_j(v_0) \right)$ from above by $2^{\sum_{j=1}^{s_{v_0}} l_j(v_0)}$, and, recalling that $\Delta_1 = [\psi]$ and $k_{v_i} \geq 1$, we rewrite and bound: $\gamma^{k_{v_i}(2\Delta_1-l_{v_i}[\psi])} = \gamma^{-k_{v_i}(l_{v_i}-2)[\psi]} \leq \gamma^{-(l_{v_i}-2)[\psi]}$, so that, proceeding as in [33, Appendix A.6.1], the sum over $l_{v_1}, \dots, l_{v_n}$ in the last line can be bounded as follows:

$$\begin{aligned}
& 2^{\sum_{j=2}^{s_{v_0}} l_j(v_0)} \sum_{l_{v_1}, \dots, l_{v_n} \geq 2} 2^{l_{v_1}} \left(\prod_{i=1}^n \binom{\sum_{j=1}^{s_{v_i}} l_j(v_i)}{l_{v_i}} \right) \gamma^{k_{v_i}(2\Delta_1-l_{v_i}[\psi])} \\
& \leq \gamma^{2[\psi]n} 2^{\sum_{j=2}^{s_{v_0}} l_j(v_0)} \sum_{l_{v_1}, \dots, l_{v_n} \geq 0} 2^{l_{v_1}} \left(\prod_{i=1}^n \binom{\sum_{j=1}^{s_{v_i}} l_j(v_i)}{l_{v_i}} \right) \gamma^{-l_{v_i}[\psi]} \\
& \leq \gamma^{2[\psi]n} 2^{\sum_{j=2}^{s_{v_0}} l_j(v_0)} (1 + 2\gamma^{-[\psi]})^{\sum_{j=2}^{s_{v_1}} l_j(v_1)} \dots \left(\sum_{m=0}^{n-2} \gamma^{-m[\psi]} + 2\gamma^{-(n-1)[\psi]} \right)^{\sum_{j=2}^{s_{v_{n-1}}} l_j(v_{n-1})} \cdot \\
& \cdot \left(\sum_{m=0}^{n-1} \gamma^{-m[\psi]} + 2\gamma^{-n[\psi]} \right)^{\sum_{j=1}^{s_{v_n}} l_j(v_n)} \\
& \leq \gamma^{2[\psi]n} \left(\frac{2}{1 - \gamma^{-[\psi]}} \right)^{\sum_{i=0}^{n-1} \sum_{j=2}^{s_{v_i}} l_j(v_i) + \sum_{j=1}^{s_{v_n}} l_j(v_n)},
\end{aligned} \tag{5.25}$$

where in the last step we bounded $\max\{2, \max_{n \geq 1} \{\sum_{m=0}^{n-1} \gamma^{-m[\psi]} + 2\gamma^{-n[\psi]}\}\}$ by $2/(1 - \gamma^{-[\psi]})$.

Inserting this into (5.24) and noting that $1 \leq n \leq 2n_{e.p.}[\tau]$ (the upper bound on n following

from [1, Eq.(J.10)], so that the factor $\gamma^{2[\psi]n}$ can be re-absorbed in a re-definition of C, C' , we find:

$$\begin{aligned}
& \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}}^* \left\| D^k S_{2,0,l,\mathbf{p}}^{\ell_1(v_0), \dots, \ell_{s_{v_0}}(v_0)} (H_{\ell_1(v_0)}[\tau_1(v_0)], \dots, H_{\ell_{s_{v_0}}(v_0)}[\tau_{s_{v_0}}(v_0)])(\mathbf{x}) \right\| \\
& \leq C(C'\rho'\epsilon)^{n_{e.p.}[\tau]-2} (4C_0)^{-l} \gamma^{k(2\Delta_1-l[\psi]-\|\mathbf{p}\|_1)} \sum_{(\ell_i(v_0))_{i=2}^{s_{v_0}}}^* \dots \sum_{(\ell_i(v_{n-1}))_{i=2}^{s_{v_{n-1}}}}^* \sum_{(\ell_i(v_n))_{i=1}^{s_{v_n}}}^* \cdot \\
& \cdot \left(\prod_{i=0}^{n-1} \left(\prod_{j=2}^{s_{v_i}} \left(\frac{\rho'(1-\gamma^{-[\psi]})}{8C_0} \right)^{-l_j(v_i)} \right) \right) \left(\prod_{j=1}^{s_{v_n}} \left(\frac{\rho'(1-\gamma^{-[\psi]})}{8C_0} \right)^{-l_j(v_n)} \right) \sum_{k_{v_1}, \dots, k_{v_n} \geq 1} e^{-\bar{C}(\gamma^{k+\sum_{i=1}^n k_{v_i}-1}|x|)^\sigma}.
\end{aligned} \tag{5.26}$$

up to a possible re-definition of C, C' . As above, if $s_{v_{i_0}} = 1$ for some $i_0 \in \{0, \dots, n-1\}$, then $\sum_{(\ell_i(v_{i_0}))_{i=2}^{s_{v_{i_0}}}}^* \left(\prod_{j=2}^{s_{v_{i_0}}} \left(\frac{\rho'(1-\gamma^{-[\psi]})}{8C_0} \right)^{-l_j(v_{i_0})} \right)$ should be interpreted as being equal to 1. Choosing ρ' larger than $8C_0(1-\gamma^{-[\psi]})^{-1}$, the sums

$$\sum_{(\ell_i(v_0))_{i=2}^{s_{v_0}}}^* \dots \sum_{(\ell_i(v_{n-1}))_{i=2}^{s_{v_{n-1}}}}^* \sum_{(\ell_i(v_n))_{i=1}^{s_{v_n}}}^* \left(\prod_{i=0}^{n-1} \prod_{j=2}^{s_{v_i}} \left(\frac{\rho'(1-\gamma^{-[\psi]})}{8C_0} \right)^{-l_j(v_i)} \right) \left(\prod_{j=1}^{s_{v_n}} \left(\frac{\rho'(1-\gamma^{-[\psi]})}{8C_0} \right)^{-l_j(v_n)} \right)$$

are absolutely summable and bounded from above by $(\text{const.})^{n_{e.p.}[\tau]}$, which can be also re-absorbed into a re-definition of C, C' . In conclusion, for some $C'' > 0$,

$$\begin{aligned}
& \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}}^* \left\| D^k S_{2,0,l,\mathbf{p}}^{\ell_1(v_0), \dots, \ell_{s_{v_0}}(v_0)} (H_{\ell_1(v_0)}[\tau_1(v_0)], \dots, H_{\ell_{s_{v_0}}(v_0)}[\tau_{s_{v_0}}(v_0)])(\mathbf{x}) \right\| \\
& \leq C''(C''\epsilon)^{n_{e.p.}[\tau]-2} (4C_0)^{-l} \gamma^{k(2\Delta_1-l[\psi]-\|\mathbf{p}\|_1)} \sum_{k_{v_1}, \dots, k_{v_n} \geq 1} e^{-\bar{C}(\gamma^{k+\sum_{i=1}^n k_{v_i}-1}|x|)^\sigma} \\
& = C'''(C''\epsilon)^{n_{e.p.}[\tau]-2} (4C_0)^{-l} \gamma^{k(2\Delta_1-l[\psi]-\|\mathbf{p}\|_1)} \sum_{\bar{k} \geq n} \binom{\bar{k}-1}{n-1} e^{-\bar{C}(\gamma^{k+\bar{k}-1}|x|)^\sigma}.
\end{aligned} \tag{5.27}$$

In order to bound the sum over $\bar{k}$, we use that, for any $\alpha > 0$,

$$\binom{\bar{k}-1}{n-1} \leq \frac{(\bar{k}-1)^{n-1}}{(n-1)!} \leq (\alpha \log \gamma)^{1-n} \gamma^{(\bar{k}-1)\alpha},$$

so that

$$\begin{aligned}
& \sum_{\bar{k} \geq n} \binom{\bar{k}-1}{n-1} e^{-\bar{C}(\gamma^{k+\bar{k}-1}|x|)^\sigma} \leq (\alpha \log \gamma)^{1-n} e^{-\frac{\bar{C}}{2}(\gamma^{k+n-1}|x|)^\sigma} \sum_{\bar{k} \geq n} \gamma^{(\bar{k}-1)\alpha} e^{-\frac{\bar{C}}{2}(\gamma^{k+\bar{k}-1}|x|)^\sigma} \\
& \leq (\alpha \log \gamma)^{1-n} e^{-\frac{\bar{C}}{2}(\gamma^{k+n-1}|x|)^\sigma} C_{\alpha, \bar{C}/2} (\min\{1, \gamma^k|x|\})^{-\alpha},
\end{aligned} \tag{5.28}$$

where, in the last inequality, letting $C_{\alpha, \bar{C}/2}$ as in (5.9), we bounded $\sum_{\bar{k} \geq n} \gamma^{(\bar{k}-1)\alpha} e^{-\frac{\bar{C}}{2}(\gamma^{k+\bar{k}-1}|x|)^\sigma}$ from above by:

$$\begin{cases} \sum_{\bar{k} \geq n} \gamma^{(\bar{k}-1)\alpha} e^{-\frac{\bar{C}}{2}(\gamma^{k+\bar{k}-1})^\sigma} \leq \gamma^{-\alpha} C_{\alpha, \bar{C}/2}, & \text{if } \gamma^k |x| \geq 1, \\ (\gamma^k |x|)^{-\alpha} \sum_{\bar{k} \geq n} \gamma^{(k+\bar{k}-h_x-1)\alpha} e^{-\frac{\bar{C}}{2}(\gamma^{k+\bar{k}-h_x-2})^\sigma} \leq (\gamma^k |x|)^{-\alpha} C_{\alpha, \bar{C}/2}, & \text{if } \gamma^k |x| < 1, \end{cases} \quad (5.29)$$

where h_x was defined before (5.9). Plugging (5.28) in (5.27), and noting that $1 \leq n \leq 2n_{e.p}[\tau]$ (the upper bound on n following from [1, Eq.(J.10)]), we finally obtain the desired bound, (5.6), provided that C_0 is chosen larger than $1/4$ (once again, we have the freedom to choose C_0 as large as desired, due to the monotonicity of the right hand side of (5.27) on C_0 : recall, in fact, that C'' is proportional to C_0).

The proof of (5.7) is completely analogous, the only difference being that x must be replaced by y and Δ_1 by Δ_2 . Note that, by changing Δ_1 in Δ_2 , we get an extra factor $\gamma^{(2\Delta_2-2[\psi]) \sum_{i=1}^n k_{v_i}}$ in the analogues of the right hand side of (5.25) and of the following equations; in particular, it also reflects into the analogue of (5.27), which reads:

$$\begin{aligned} & \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}} \left\| S_{0,2,l,\mathbf{p}}^{\ell_1(v_0), \dots, \ell_{s_{v_0}}(v_0)} (H_{\ell_1(v_0)}[\tau_1(v_0)], \dots, H_{\ell_{s_{v_0}}(v_0)}[\tau_{s_{v_0}}(v_0)])(\mathbf{y}) \right\| \\ & \leq C''' (C'' \epsilon)^{n_{e.p}[\tau]-2} (4C_0)^{-l} \gamma^{k(2\Delta_2-l[\psi]-\|\mathbf{p}\|_1)} \sum_{\bar{k} \geq n} \binom{\bar{k}-1}{n-1} \gamma^{\bar{k}(2\Delta_2-2[\psi])} e^{-\bar{C}(\gamma^{k+\bar{k}-1}|y|)^\sigma}. \end{aligned} \quad (5.30)$$

Bounding the sum over $\bar{k}$ as explained after (5.27) and choosing C_0 larger than $1/4$ implies the desired estimate, (5.7). $\square$

6 Stretched exponential decay of the correction terms $\mathcal{E}_1, \mathcal{E}_2$

In this section we prove Theorem 2.4. We focus on the bound on $\mathcal{E}_1$, the bound on $\mathcal{E}_2$ being completely analogous (and, therefore, left to the reader). From the definition of $\mathcal{E}_1$ in (2.40), letting $\mathbf{x} = (x, 0)$, we have:

$$[\mathcal{E}_1(x)]_{a,b} = 2[V_{2,0,0,\emptyset}^*(\mathbf{x})]_{a,b} + [\mathfrak{E}_1(x)]_{a,b}, \quad (6.1)$$

where $[\mathfrak{E}_1(x)]_{a,b} = \langle \frac{\delta^2 \mathcal{Q}^*(\phi, 0, \psi)}{\delta \phi_b(0) \delta \phi_a(x)} |_{\phi=0} \rangle_{H^*}$ with $\mathcal{Q}^*(\phi, J, \psi)$ defined in (2.38) and following line, i.e.,

$$\mathcal{Q}^* = \{V_\ell^*\}_{\ell \in L_f(2,0)} \quad (6.2)$$

(here, letting $L(2,0) := \{(n, m, l, \mathbf{p}) \in L : (n, m) = (2, 0)\}$, we defined $L_f(2,0) := L(2,0) \setminus \{(2, 0, 0, \emptyset)\}$). The first term in the right hand side of (6.1) has already been analyzed in the previous sections, and proved to be analytic in ϵ for ϵ small. Moreover, as we shall see shortly, the tree representation (5.3) and the bounds derived in the preceding section on the tree values $H_{2,0,0,\emptyset}[\tau](\mathbf{x})$ readily imply the stretched exponential decay of $V_{2,0,0,\emptyset}^*(\mathbf{x})$. Concerning the second term, we will prove below that it can be computed via a convergent expansion in which the kernels $V_{2,0,l,\mathbf{p}}^*$ of $\mathcal{Q}^*$ are ‘contracted’ with those of H^* . Not surprisingly, $\mathfrak{E}_1(x)$ can be expressed once again in terms of a tree expansion, slightly different from the one of the preceding sections, in that it includes only a subset of the trees contributing to the scale-invariant, fixed-point, potential studied above. As we shall see, the trees contributing to $\mathfrak{E}_1$ are characterized by a constraint on the ‘scale

indices' (called $h, k_1, k_2, \dots$ in the following), implying the desired stretched exponential decay. In order to simplify the analysis and to rely on the bounds derived in the preceding sections as much as possible, we shall define the trees contributing to $\mathfrak{E}_1$ in terms of 'dressed endpoints' (rather than of the 'bare endpoints' in Figure 4): these are the big white and black endpoints of, e.g., Figure 9 below, representing the kernels $V_{2,0,l,\mathbf{p}}^*$ and $H_{l,\mathbf{p}}^*$, respectively. In order to estimate the values of the trees contributing to $\mathfrak{E}_1$, we shall use the bounds on $V_{2,0,l,\mathbf{p}}^*$ and $H_{l,\mathbf{p}}^*$ derived above and in [1], without re-expanding from scratch these kernel as sums over trees once again.

Let us consider the first term in the right hand side of (6.1). As mentioned above, we already proved in the previous sections that it is analytic in ϵ for ϵ small. Moreover, using the first identity in (5.3) and the bound (5.8) with $h = 0$, we have (dropping as usual the component labels):

$$\begin{aligned} |V_{2,0,0,\emptyset}^*(\mathbf{x})| &\leq C_\alpha |x|^{-2\Delta_1} \sum_{\tau} (C_\alpha \epsilon)^{n_{\text{e.p.}}[\tau]-2} \sum_{k \geq 1} e^{-\frac{\tilde{C}}{2}(\gamma^{k-1}|x|)^\sigma} \max\{(\gamma^k|x|)^{2\Delta_1}, (\gamma^k|x|)^{2\Delta_1-\alpha}\} \\ &\leq C'_\alpha |x|^{-2\Delta_1} e^{-\frac{\tilde{C}}{4}(|x|/\gamma)^\sigma} \sum_{k \geq 1} e^{-\frac{\tilde{C}}{4}(\gamma^{k-1}|x|)^\sigma} \max\{(\gamma^k|x|)^{2\Delta_1}, (\gamma^k|x|)^{2\Delta_1-\alpha}\}, \end{aligned} \quad (6.3)$$

where, in passing from the first to the second line, we performed the sum over τ of $(C_\alpha \epsilon)^{n_{\text{e.p.}}[\tau]-2}$ (which is summable, and whose sum is bounded by an ϵ -independent constant), and we extracted part of the exponential factor from the sum over k . Now, if $|x| \geq 1$, the summand in the sum over k can be bounded from above by $e^{-\frac{\tilde{C}}{4}\gamma^{\sigma(k-1)}} \gamma^{2k\Delta_1} |x|^{2\Delta_1}$, so that the right hand side of (6.3) can be bounded from above by $(\text{const.})e^{-\frac{\tilde{C}}{4}(|x|/\gamma)^\sigma}$, with $(\text{const.}) = C'_\alpha \sum_{k \geq 1} e^{-\frac{\tilde{C}}{4}\gamma^{\sigma(k-1)}} \gamma^{2k\Delta_1}$. On the other hand, if $|x| \leq 1$, the sum over k in the right hand side of (6.3) can be bounded via (5.9), so that the right hand side of (6.3) can be bounded from above by $(\text{const.})|x|^{-2\Delta_1} e^{-\frac{\tilde{C}}{4}(|x|/\gamma)^\sigma}$. Putting the two cases together, we find

$$|V_{2,0,0,\emptyset}^*(\mathbf{x})| \leq C e^{-\frac{\tilde{C}}{4}(|x|/\gamma)^\sigma} (\min\{1, |x|\})^{-2\Delta_1}. \quad (6.4)$$

Note that the right hand side has the same form as that of the first inequality in (2.45). Therefore, in order to complete the proof of Theorem 2.4, we need to prove a comparable bound on $\mathfrak{E}_1(x)$, as well as its analyticity.

For this purpose, we derive a tree expansion for $\mathfrak{E}_1(x)$. We write (keeping, once again, the flavor indices implicit):

$$\mathfrak{E}_1(x) = \frac{\delta^2 \mathcal{Q}_{\text{eff}}(\phi, 0)}{\delta\phi(0)\delta\phi(x)} \Big|_{\phi=0}, \quad (6.5)$$

where, in analogy with (2.6) and (2.49),

$$\begin{aligned} \mathcal{Q}_{\text{eff}}(\phi, J) &= \lim_{h \rightarrow -\infty} \log \frac{\int d\mu_{[h,0]}(\psi) e^{H^*(\psi) + \mathcal{Q}^*(\phi, J, \psi)}}{\int d\mu_{[h,0]}(\psi) e^{H^*(\psi)}} \\ &\equiv \lim_{h \rightarrow -\infty} D^h v^{(h)}(\phi, J, 0), \end{aligned} \quad (6.6)$$

with

$$v^{(h)}(\phi, J, \psi) := R^{|h|}(H^* + \mathcal{Q}^*)(\phi, J, \psi) - H^*(\psi), \quad (6.7)$$

and $\mathcal{Q}^*$ as in (6.2). Denoting by $v_\ell^{(h)}$ with $\ell \in L$ the kernels of $v^{(h)}$ (which are by construction non-vanishing only for indices $\ell = (n, m, l, \mathbf{p})$ such that $n + m \geq 2$), we can then rewrite

$$\begin{aligned}\mathfrak{E}_1(x) &= 2 \lim_{h \rightarrow -\infty} D^h v_{2,0,0,\emptyset}^{(h)}(\mathbf{x}) \\ &= 2 \lim_{h \rightarrow -\infty} \gamma^{2h\Delta_1} v_{2,0,0,\emptyset}^{(h)}(\gamma^h \mathbf{x}).\end{aligned}\tag{6.8}$$

In view of the definition (6.7) of $v^{(h)}$, of the fact that its kernels $v_\ell^{(h)}$ are non-vanishing only for indices $\ell = (n, m, l, \mathbf{p})$ such that $n + m \geq 2$, and of the representation (3.18) of the action of the RG map R in components, for any $\ell \in L(2, 0) = \{(n, m, l, \mathbf{p}) \in L : (n, m) = (2, 0)\}$, for any $h < 0$ we can write:

$$v_\ell^{(h)} = \sum_{s \geq 1} s \sum_{\ell_1 \in L_f(2,0)} \sum_{(\ell_i)_{i=2}^s} R_\ell^{\ell_1, \dots, \ell_s} (v_{\ell_1}^{(h+1)}, H_{\ell_2}^*, \dots, H_{\ell_s}^*),\tag{6.9}$$

where we recall that $L_f(2, 0) := L(2, 0) \setminus \{(2, 0, 0, \emptyset)\}$, and the third sum in the right hand side runs over the $(s-1)$ -ples of labels in $L(0, 0) := \{(n, m, l, \mathbf{p}) \in L : (n, m) = (0, 0)\}$ (note that for any such label, of the form $\ell = (0, 0, l, \mathbf{p})$, $V_{0,0,l,\mathbf{p}}^* \equiv H_{l,\mathbf{p}}^*$, with H^* the FP potential constructed in [1], see condition (1) in Definition 2.2; for this reason, with some abuse of notation, in (6.9) we wrote H_ℓ^* in place of V_ℓ^*). From (3.19) we see that, for any $\ell \in L(2, 0)$, $R_\ell^{\ell_1, \dots, \ell_s} = DS_\ell^{\ell_1, \dots, \ell_s}$, so that, isolating from the right hand side of (6.9) the contribution with $s = 1$ and $\ell_1 = \ell$, we can rewrite:

$$v_\ell^{(h)} = Dv_\ell^{(h+1)} + \sum_{s \geq 1} s \sum_{\ell_1 \in L_f(2,0)}^* \sum_{(\ell_i)_{i=2}^s} DS_\ell^{\ell_1, \dots, \ell_s} (v_{\ell_1}^{(h+1)}, H_{\ell_2}^*, \dots, H_{\ell_s}^*),\tag{6.10}$$

where the $*$ on the second sum in the right hand side indicates the constraint that, if $s = 1$, then $\ell_1 \neq \ell$. Note also that, by definition, for $h = -1$, the kernel $v_\ell^{(0)}$ in the right hand side is $v_\ell^{(0)} \equiv V_\ell^*$, for all $\ell \in L_f(2, 0)$, and zero otherwise. Therefore, for any $\ell \in L(2, 0)$,

$$v_\ell^{(-1)} = DV_\ell^* \mathbb{1}_{\ell \in L_f(2,0)} + \sum_{s \geq 1} s \sum_{\ell_1 \in L_f(2,0)}^* \sum_{(\ell_i)_{i=2}^s} DS_\ell^{\ell_1, \dots, \ell_s} (V_{\ell_1}^*, H_{\ell_2}^*, \dots, H_{\ell_s}^*).\tag{6.11}$$

If we graphically represent any kernel V_ℓ^* with $\ell \in L(2, 0)$ by a big white dot, the first term in the right hand side, whenever it is there (i.e., for $\ell \neq (2, 0, 0, \emptyset)$), can be represented as $D \bigcirc$, while the second as a sum over trees τ of length 1 (the length being the maximal number of branches along a path from the endpoints to the root) as in Fig.9. We correspondingly write:

$$v_\ell^{(-1)} = DV_\ell^* \mathbb{1}_{\ell \in L_f(2,0)} + \sum_{\tau \in \mathcal{T}_1} v_\ell[\tau],\tag{6.12}$$

where $\mathcal{T}_1$ is the family of trees of length 1 described in Figure 9 and in its caption, and $v_\ell[\tau]$ is its value: if τ has s_{v_0} endpoints, then $v_\ell[\tau] = s_{v_0} \sum_{\ell_1 \in L_f(2,0)}^* \sum_{(\ell_i)_{i=2}^{s_{v_0}}} DS_\ell^{\ell_1, \dots, \ell_{s_{v_0}}} (V_{\ell_1}^*, H_{\ell_2}^*, \dots, H_{\ell_{s_{v_0}}}^*)$.

We can easily iterate this procedure, via (6.10), thus ending up with the following tree representation for $v_\ell^{(h)}$:

$$v_\ell^{(h)} = D^{|h|} V_\ell^* \mathbb{1}_{\ell \in L_f(2,0)} + \sum_{\mathfrak{L}=1}^{|h|} \sum_{\substack{k_{\mathfrak{L}} \geq 0, \\ k_0, k_1, \dots, k_{\mathfrak{L}-1} \geq 1: \\ k_0 + \dots + k_{\mathfrak{L}} = |h|}} \sum_{\tau \in \mathcal{T}_{\mathfrak{L}}} v_\ell[\tau, \mathbf{k}],\tag{6.13}$$




Figure 9: A tree in $\mathcal{T}_1$, representing one of the contributions $v_\ell[\tau]$ in the right hand side of (6.11), with the action of the dilatation operator D associated with the root explicitly indicated. The elements of $\mathcal{T}_1$ consists of a root vertex v_0 , followed by $s_{v_0} \geq 1$ children: the first (i.e., the topmost) one, denoted v_1 , is a big white dot, representing $V_{\ell_1}^*$ (with ℓ_1 an index to be summed over $L_f(2,0)$, such that $\ell_1 \neq \ell$), while the others are all big-black-dot endpoints, representing $H_{\ell_i}^*$, $i = 2, \dots, s_{v_0}$, where ℓ_i are indices to be summed over $L(0,0)$.

where $\mathfrak{L}$ represents the length of the tree (i.e., the number of branches crossed by the straight path from the root v_0 to the big white end point, see Figure 10), $\mathcal{T}_{\mathfrak{L}}$ is the family of trees of length $\mathfrak{L}$ described in Figure 10 and in its caption, $\mathbf{k} := (k_0, k_1, \dots, k_{\mathfrak{L}})$ and $v_\ell[\tau, \mathbf{k}]^{11}$ is the value of the tree τ , in the presence of the action of the dilatations $D^{k_0}, D^{k_1}, \dots, D^{k_{\mathfrak{L}}}$ associated with the vertices $v_0, v_1, \dots, v_{\mathfrak{L}}$, as in Figure 10. Given $\tau \in \mathcal{T}_{\mathfrak{L}}$ with $\mathfrak{L} > 1$, letting $\tau_{v_1} \in \mathcal{T}_{\mathfrak{L}-1}$ be the subtree of τ




Figure 10: A tree in $\mathcal{T}_{\mathfrak{L}}$ with the action of the dilatations D^{k_i} associated with the vertices v_i , $i = 0, \dots, \mathfrak{L}$ explicitly indicated. For any $\mathfrak{L} > 1$, the elements of $\mathcal{T}_{\mathfrak{L}}$ are recursively characterized by the conditions that the root v_0 has $s_{v_0} \geq 1$ children, and that the first (i.e., the topmost) one, denoted v_1 , is the root of a tree in $\mathcal{T}_{\mathfrak{L}-1}$, while the other children are all big-black-dot endpoints. For $\mathfrak{L} = 1$, the set $\mathcal{T}_1$ was described in Fig.9. For any h such that $|h| \geq \mathfrak{L}$, the integers $k_0, \dots, k_{\mathfrak{L}}$ satisfy: $k_0, \dots, k_{\mathfrak{L}-1} \geq 1$, $k_{\mathfrak{L}} \geq 0$, and $k_0 + \dots + k_{\mathfrak{L}} = |h|$; in particular, for $\mathfrak{L} = 1$ and $h \leq -1$, the vertices v_0 and v_1 are associated with two dilatations operators D^{k_0} and D^{k_1} with $k_0 \geq 1$, $k_1 \geq 0$ and $k_0 + k_1 = |h|$: the case depicted in Fig.9 (where v_0 is associated with D , and v_1 with the identity) is a special case corresponding to $h = -1$.

rooted in v_1 , $v_\ell[\tau, (k_0, \dots, k_{\mathfrak{L}})]$, for $\ell \in L(2,0)$, is recursively defined as:

$$v_\ell[\tau, (k_0, k_1, \dots, k_{\mathfrak{L}})] = s_{v_0} \sum_{\ell_1 \in L_f(2,0)}^* \sum_{(\ell_i)_{i=2}^{s_{v_0}}} D^{k_0} S_{\ell}^{\ell_1, \dots, \ell_{s_{v_0}}} (v_{\ell_1}[\tau_{v_1}, (k_1, \dots, k_{\mathfrak{L}})], H_{\ell_2}^*, \dots, H_{\ell_{s_{v_0}}}^*), \quad (6.14)$$

¹¹Note that for $h = -1$ there is only one possible choice of $\mathbf{k} = (k_0, k_1)$ compatible with the constraints $k_0 \geq 1$, $k_1 \geq 0$, $k_0 + k_1 = 1$ indicated under the second sum in the right hand side of (6.13), that is $(k_0, k_1) = (1, 0)$: therefore, for $h = -1$, the value $v_\ell[\tau]$ in the right hand side of (6.12) is the same denoted by $v_\ell[\tau, (1, 0)]$ in (6.13).

where the sums over ℓ_1 and over $(\ell_i)_{i=1}^{s_{v_0}}$ must be interpreted as in (6.10), while, given $\tau \in \mathcal{T}_1$,

$$v_\ell[\tau, (k_0, k_1)] = s_{v_0} \sum_{\ell_1 \in L_f(2,0)}^* \sum_{(\ell_i)_{i=2}^{s_{v_0}}} D^{k_0} S_\ell^{\ell_1, \dots, \ell_{s_{v_0}}} (D^{k_1} V_{\ell_1}^*, H_{\ell_2}^*, \dots, H_{\ell_{s_{v_0}}}^*). \quad (6.15)$$

The validity of the tree representation (6.13) with the tree values defined above can be straightforwardly proved by induction, and is left to the reader.

We now use this tree representation to compute and bound the right hand side of (6.8). We have:

$$D^h v_{2,0,0,\emptyset}^{(h)}(\mathbf{x}) = \sum_{\mathfrak{L}=1}^{|h|} \sum_{\substack{k_{\mathfrak{L}} \geq 0, \\ k_0, k_1, \dots, k_{\mathfrak{L}-1} \geq 1: \\ k_0 + \dots + k_{\mathfrak{L}} = |h|}} \sum_{\tau \in \mathcal{T}_{\mathfrak{L}}} D^h v_{2,0,0,\emptyset}[\tau, \mathbf{k}](\mathbf{x}). \quad (6.16)$$

If $\tau \in \mathcal{T}_1$, using (6.15), the definition of D in (2.27), and the analogue of (5.17), we get that, for any $k_0 \geq 1, k_1 \geq 0$ such that $k_0 + k_1 = |h|$,

$$\begin{aligned} & |D^h v_{2,0,0,\emptyset}[\tau, (k_0, k_1)](\mathbf{x})| \\ & \leq s_{v_0} \sum_{\ell_1 \in L_f(2,0)} \sum_{(\ell_i)_{i=2}^{s_{v_0}}} \gamma^{2\Delta_1(h+k_0)} \left| S_{2,0,0,\emptyset}^{\ell_1, \dots, \ell_{s_{v_0}}} (D^{k_1} V_{\ell_1}^*, H_{\ell_2}^*, \dots, H_{\ell_{s_{v_0}}}^*) (\gamma^{h+k_0} \mathbf{x}) \right| \\ & \leq s_{v_0} \sum_{\ell_1 \in L_f(2,0)} \sum_{(\ell_i)_{i=2}^{s_{v_0}}} \gamma^{2\Delta_1(h+k_0)} C_\gamma^{s_{v_0}-1} (4C_0)^{\sum_{i=1}^{s_{v_0}} l_i} N_{(2,0,0,\emptyset)}^{\ell_1, \dots, \ell_{s_{v_0}}} \left(\prod_{i=2}^{s_{v_0}} \|H_{\ell_i}^*\|_w \right) \|D^{k_1} V_{\ell_1}^* (\gamma^{h+k_0} \mathbf{x})\|. \end{aligned} \quad (6.17)$$

Now, in the last line, for any $i = 2, \dots, s_{v_0}$, we use that

$$\|H_{\ell_i}^*\|_w \leq \sum_{\tau} \|H_{\ell_i}[\tau]\|_w \leq C \sum_{\tau} (C\epsilon)^{n_{\text{e.p.}}[\tau]} \leq C' (C'\epsilon)^{\max\{1, \frac{l_i}{2}-1\}}, \quad (6.18)$$

where, in the second inequality, we used (4.22) with $\rho = 1$ and the fact that the trees contributing to $H_{\ell_i}^*$, with $\ell_i = (0, 0, l_i, \mathbf{p}_i)$, necessarily have $n_{\text{e.p.}}[\tau] \geq \max\{1, \frac{l_i}{2} - 1\}$. Concerning the factor $\|D^{k_1} V_{\ell_1}^* (\gamma^{h+k_0} \mathbf{x})\|$, we rewrite it and bound it as follows: we use once again the tree representation (4.3) with $H_\ell[\tau]$ as in (4.4). Noting that for $\ell = (2, 0, l, \mathbf{p})$ the trimming operator acts as the identity, and rewriting $\frac{D}{1-D} = \sum_{k \geq 1} D^k$, with D as in (2.27), we find:

$$\|D^{k_1} V_{\ell_1}^* (\gamma^{h+k_0} \mathbf{x})\| \leq \sum_{k \geq 1} \sum_{\tau} \sum_{(\ell'_i)_{i=1}^{s_{v_0}}}^* \left\| D^{k+k_1} S_{\ell_1}^{\ell'_1, \dots, \ell'_{s_{v_0}}} (H_{\ell'_1}[\tau_1], \dots, H_{\ell'_{s_{v_0}}}[\tau_{s_{v_0}}]) (\gamma^{h+k_0} \mathbf{x}) \right\| \quad (6.19)$$

By Lemma 5.2, we thus obtain:

$$\begin{aligned} & \gamma^{2\Delta_1(h+k_0)} \|D^{k_1} V_{\ell_1}^* (\gamma^{h+k_0} \mathbf{x})\| \leq C \sum_{\tau} (C\epsilon)^{n_{\text{e.p.}}[\tau]-2} \sum_{k \geq 1} \gamma^{2\Delta_1(h+k_0+k+k_1)} \gamma^{-(k+k_1)(l_1[\psi] + \|\mathbf{p}_1\|_1)} \\ & \quad \cdot e^{-\frac{C}{2}(\gamma^{k+k_1+h+k_0-1}|x|)^\sigma} \left(\min\{1, \gamma^{k+k_1+h+k_0}|x|\} \right)^{-\alpha} \\ & \leq C' (C'\epsilon)^{\frac{l_1}{2}-1} \sum_{k \geq 1} \gamma^{2\Delta_1 k} \gamma^{-(k+k_1)(l_1[\psi] + \|\mathbf{p}_1\|_1)} e^{-\frac{C}{2}(\gamma^{k-1}|x|)^\sigma} \left(\min\{1, \gamma^k|x|\} \right)^{-\alpha}, \end{aligned} \quad (6.20)$$

where in the second inequality we performed the sum over τ (note that, for given $l_1 \geq 2$, $n_{\text{e.p.}}[\tau] - 2 \geq \frac{l_1}{2} - 1$, so that $\sum_{\tau} (C\epsilon)^{n_{\text{e.p.}}[\tau] - 2} \leq (\text{const.})(C'\epsilon)^{\frac{l_1}{2} - 1}$) and used the fact that $k_0 + k_1 + h = 0$. By the same considerations discussed after (6.3), performing the sum over $k \geq 1$ we get:

$$\gamma^{2\Delta_1(h+k_0)} \|D^{k_1} V_{\ell_1}^*(\gamma^{h+k_0} \mathbf{x})\| \leq C'(C'\epsilon)^{\frac{l_1}{2} - 1} \gamma^{-k_1(l_1[\psi] + \|\mathbf{p}_1\|_1)} e^{-\frac{\bar{C}}{4}(|x|/\gamma)^\sigma} (\min\{1, |x|\})^{-2\Delta_1}. \quad (6.21)$$

Plugging (6.18) and (6.21) into (6.17), and using that $N_{(2,0,0,\emptyset)}^{\ell_1, \dots, \ell_{s_{v_0}}} \leq 2^{\sum_{i=1}^{s_{v_0}} l_i}$, gives:

$$\begin{aligned} & |D^h v_{2,0,0,\emptyset}[\tau, (k_0, k_1)](\mathbf{x})| \\ & \leq s_{v_0} (C'')^{s_{v_0}} e^{-\frac{\bar{C}}{4}(|x|/\gamma)^\sigma} (\min\{1, |x|\})^{-2\Delta_1} \sum_{\ell_1 \in L_f(2,0)} (C''\epsilon)^{\frac{l_1}{2} - 1} \gamma^{-k_1(l_1[\psi] + \|\mathbf{p}_1\|_1)} \\ & \cdot \sum_{(\ell_i)_{i=2}^{s_{v_0}}} \prod_{i=2}^{s_{v_0}} (C''\epsilon)^{\max\{1, \frac{l_i}{2} - 1\}} \leq C''' (C'''\epsilon)^{s_{v_0} - 1} e^{-\frac{\bar{C}}{4}(|x|/\gamma)^\sigma} (\min\{1, |x|\})^{-2\Delta_1} \gamma^{-2k_1[\psi]}. \end{aligned} \quad (6.22)$$

Remark 6.1. From (6.20), it is clear that the constraint $k_0 + k_1 = |h|$ (or, in general, $k_0 + k_1 + \dots + k_{\mathfrak{L}} = |h|$ for trees $\tau \in \mathcal{T}_{\mathfrak{L}}$ with $\mathfrak{L} \geq 1$, to be discussed in the following) is essential in proving the stretched exponential decay in (6.22). This constraint, first appearing in (6.13), has its origin in the recursive equation (6.10), which is linear in $\{v_{\ell}^{(h')}\}_{\ell \in L_f(2,0)}^{h' \leq 0}$ (with $v_{\ell}^{(0)} \equiv V_{\ell}^*$); in turn, such linearity in $\{v_{\ell}^{(h')}\}_{\ell \in L_f(2,0)}^{h' \leq 0}$ originates from the fact that $\mathcal{Q}^*$ is quadratic in ϕ . In order to compare more closely the tree expansion for $D^h v_{2,0,0,\emptyset}^{(h)}$ in (6.13) with the one for $D^h V_{2,0,0,\emptyset}^*$ in the first line of (5.3) we can, if desired, re-expand the kernels $V_{\ell'}^*$ and $H_{\ell''}^*$, associated with the big white and black endpoints of the trees in $\mathcal{T}_{\mathfrak{L}}$ in terms of the tree expansions discussed in Section 5 (for $V_{\ell'}^*$) and in [1, Appendix J] (for $H_{\ell''}^*$): this would lead to a new tree expansion for $D^h v_{2,0,0,\emptyset}^{(h)}$ (more complex, but closer to the tree expansion of the previous section) in terms of trees with ‘bare endpoints’ as those in Figure 4. Such trees would be exactly of the same form as those contributing to $D^h V_{2,0,0,\emptyset}^*$, such as those in Figure 8, with the additional constraint on the scale labels $k_0 + k_1 + \dots + k_{\mathfrak{L}} = |h|$, which was absent in the discussion in Section 5; in fact, the analogue of the combination $h + k_0 + k_1$ in (6.20) and, more generally, of the combination $h + k_0 + k_1 + \dots + k_{\mathfrak{L}}$ appearing in the bounds below, is what in Section 5 was denoted $k + k_{v_1} + \dots + k_{v_n}$, see (5.22) and following equations; in those equations, the combination $k + \sum_{i=1}^n k_{v_i}$ could take arbitrarily negative values, and this was (rightly so) at the origin of the polynomial decay of the kernel of $D^h V_{2,0,0,\emptyset}^*$ in the limit $h \rightarrow -\infty$, proved in the previous section.

We now proceed in a similar fashion, in order to bound the general term in the right hand side of (6.16). We take $1 < \mathfrak{L} \leq |h|$, $\tau \in \mathcal{T}_{\mathfrak{L}}$, $\mathbf{k} = (k_0, k_1, \dots, k_{\mathfrak{L}})$ with $k_{\mathfrak{L}} \geq 0$, $k_0, k_1, \dots, k_{\mathfrak{L}-1} \geq 1$, and $h + k_0 + k_1 + \dots + k_{\mathfrak{L}} = 0$. Using the same notation introduced before (5.16), thanks to (6.14), we have:

$$D^h v_{2,0,0,\emptyset}[\tau, \mathbf{k}](\mathbf{x}) = s_{v_0} \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}}^* D^{h+k_0} S_{2,0,0,\emptyset}^{\ell_1(v_0), \dots, \ell_{s_{v_0}}(v_0)}(v_{\ell_{v_1}}[\tau_{v_1}, \mathbf{k}_{v_1}], H_{\ell_2(v_0)}^*, \dots, H_{\ell_{s_{v_0}}(v_0)}^*)(\mathbf{x}), \quad (6.23)$$

where $\sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}}^* \equiv \sum_{\ell_1(v_0) \in L_f(2,0)}^* \sum_{(\ell_i(v_0))_{i=2}^{s_{v_0}}}$, and $\mathbf{k}_{v_1} := (k_1, \dots, k_{\mathfrak{L}})$; moreover, we recall that

$\ell_{v_1} \equiv \ell_1(v_0)$. Using (the analogue of) (5.19) we get:

$$\begin{aligned}
|D^h v_{2,0,0,\emptyset}[\tau, \mathbf{k}](\mathbf{x})| &\leq \\
&\leq s_{v_0} \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}}^* \left| D^{h+k_0} S_{2,0,0,\emptyset}^{\ell_1(v_0), \dots, \ell_{s_{v_0}}(v_0)}(v_{\ell_{v_1}}[\tau_{v_1}, \mathbf{k}_{v_1}], H_{\ell_2(v_0)}^*, \dots, H_{\ell_{s_{v_0}}(v_0)}^*)(\mathbf{x}) \right| \\
&\leq \gamma^{2\Delta_1(h+k_0)} \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}}^* s_{v_0} C_\gamma^{s_{v_0}-1} (4C_0)^{\sum_{i=1}^{s_{v_0}} \ell_i(v_0)} N_{2,0,0,\emptyset}^{\ell_1(v_0), \dots, \ell_{s_{v_0}}(v_0)} \cdot \\
&\quad \cdot \left(\prod_{i=2}^{s_{v_0}} \|H_{\ell_i(v_0)}^*\|_w \right) \|v_{\ell_{v_1}}[\tau_{v_1}, \mathbf{k}_{v_1}](\gamma^{h+k_0} \mathbf{x})\|.
\end{aligned} \tag{6.24}$$

We now apply again the recursive definition of $v_{\ell_{v_1}}[\tau_{v_1}, \mathbf{k}_{v_1}]$ and iterate the same procedure until we reach $v_{\mathfrak{L}-1}$, thus getting:

$$\begin{aligned}
|D^h v_{2,0,0,\emptyset}[\tau, \mathbf{k}](\mathbf{x})| &\leq \gamma^{2\Delta_1 h} \sum_{(\ell_i(v_0))_{i=1}^{s_{v_0}}}^* \dots \sum_{(\ell_i(v_{\mathfrak{L}-1}))_{i=1}^{s_{v_{\mathfrak{L}-1}}}}^* \times \\
&\times \left(\prod_{j=0}^{\mathfrak{L}-1} s_{v_j} C_\gamma^{s_{v_j}-1} (4C_0)^{\sum_{i=1}^{s_{v_j}} \ell_i(v_j) - l_{v_j}} N_{\ell_{v_j}}^{\ell_1(v_j), \dots, \ell_{s_{v_j}}(v_j)} \gamma^{k_j(2\Delta_1 - l_{v_j}[\psi] - \|\mathbf{p}_{v_j}\|_1)} \right. \\
&\quad \cdot \left. \left(\prod_{i=2}^{s_{v_j}} \|H_{\ell_i(v_j)}^*\|_w \right) \right) \|D^{k_{\mathfrak{L}}} V_{\ell_{v_{\mathfrak{L}}}}^* (\gamma^{h+k_0+\dots+k_{\mathfrak{L}-1}} \mathbf{x})\|,
\end{aligned} \tag{6.25}$$

with $\ell_{v_0} = (n_{v_0}, m_{v_0}, l_{v_0}, \mathbf{p}_{v_0}) \equiv (2, 0, 0, \emptyset)$.

Remark 6.2. Every collection of labels $\{(\ell_i(v_j))_{i=1}^{s_{v_j}}\}_{j=0, \dots, \mathfrak{L}-1}$ contributing to the right hand side of (6.27) satisfies the following constraints (recall that we write $\ell_i(v_j) = (n_i(v_j), m_i(v_j), l_i(v_j), \mathbf{p}_i(v_j))$, and $l_1(v_j) \equiv l_{v_{j+1}}$): for any $j = 0, \dots, \mathfrak{L}-1$, if v_j is trivial, then $l_{v_j} \leq l_{v_{j+1}} - 2$ (with the understanding that $l_{v_0} \equiv 0$), while, if v_j is non-trivial, then $l_{v_j} \leq l_{v_{j+1}} + \sum_{i=2}^{s_{v_j}} l_i(v_j) - 2(s_{v_j} - 1)$. In the following, we use that the sums over $\{(\ell_i(v_j))_{i=1}^{s_{v_j}}\}_{j=0, \dots, \mathfrak{L}-1}$ are performed under these constraints.

Now, recalling that $h + k_0 + k_1 + \dots + k_{\mathfrak{L}} = 0$ and using the analogue of (6.21), we have

$$\begin{aligned}
&\gamma^{2\Delta_1(h+k_0+\dots+k_{\mathfrak{L}-1})} \|D^{k_{\mathfrak{L}}} V_{\ell_{v_{\mathfrak{L}}}}^* (\gamma^{h+k_0+\dots+k_{\mathfrak{L}-1}} \mathbf{x})\| \leq \\
&\leq C'(C'\epsilon)^{\frac{l_{v_{\mathfrak{L}}}}{2}-1} \gamma^{-k_{\mathfrak{L}}(l_{v_{\mathfrak{L}}}[\psi] + \|\mathbf{p}_{v_{\mathfrak{L}}}\|_1)} e^{-\frac{C}{4}(|x|/\gamma)^\sigma} (\min\{1, |x|\})^{-2\Delta_1}.
\end{aligned} \tag{6.26}$$

Inserting (6.18) and (6.26) into (6.25), summing over $\mathbf{p}_{v_1}, \dots, \mathbf{p}_{v_{\mathfrak{L}}}$ and using (4.16), implies:

$$\begin{aligned}
|D^h v_{2,0,0,\emptyset}[\tau, \mathbf{k}](\mathbf{x})| &\leq C^{n_{e.p.}[\tau]} e^{-\frac{\tilde{C}}{4}(|x|/\gamma)^\sigma} (\min\{1, |x|\})^{-2\Delta_1} \sum_{(\ell_i(v_0))_{i=2}^{s_{v_0}}} \cdots \sum_{(\ell_i(v_{\mathfrak{L}-1}))_{i=2}^{s_{v_{\mathfrak{L}-1}}}} \times \\
&\times \left(\prod_{j=0}^{\mathfrak{L}-1} \left(\prod_{i=2}^{s_{v_j}} (4C_0)^{l_i(v_j)} (C'\epsilon)^{\max\{1, \frac{l_i(v_j)}{2}-1\}} \right) \right) \sum_{l_{v_{\mathfrak{L}}}} (C'\epsilon)^{l_{v_{\mathfrak{L}}}/2-1} (8C_0)^{l_{v_{\mathfrak{L}}}} \gamma^{-k_{\mathfrak{L}} l_{v_{\mathfrak{L}}}[\psi]} \cdot \\
&\cdot \sum_{l_{v_1}, \dots, l_{v_{\mathfrak{L}-1}}} \left(\prod_{j=1}^{\mathfrak{L}-1} \left(\sum_{l_{v_j}}^{s_{v_j}} l_i(v_j) \right) \gamma^{-k_j l_{v_j}[\psi]} \right). \tag{6.27}
\end{aligned}$$

Now, in the last line, recalling that $l_{v_j} \geq 2$ and $k_j \geq 1$, we can bound from above each factor $\gamma^{-k_j l_{v_j}[\psi]}$ by $\gamma^{-2k_j[\psi]} \gamma^{-(l_{v_j}-2)[\psi]}$; next, by proceeding as in [33, Appendix A.6.1] and in (5.25), we obtain:

$$\begin{aligned}
\sum_{l_{v_1}, \dots, l_{v_{\mathfrak{L}-1}}} \left(\prod_{j=1}^{\mathfrak{L}-1} \left(\sum_{l_{v_j}}^{s_{v_j}} l_i(v_j) \right) \gamma^{-k_j l_{v_j}[\psi]} \right) &\leq \\
&\leq \gamma^{2[\psi](\mathfrak{L}-1)} \gamma^{-2[\psi](k_1 + \dots + k_{\mathfrak{L}-1})} (1 - \gamma^{-[\psi]})^{-\sum_{j=1}^{\mathfrak{L}-1} \sum_{i=2}^{s_{v_j}} l_i(v_j)} (1 - \gamma^{-[\psi]})^{-l_{v_{\mathfrak{L}}}}. \tag{6.28}
\end{aligned}$$

Hence, plugging (6.28) in (6.27), using $\gamma^{-k_{\mathfrak{L}} l_{v_{\mathfrak{L}}}[\psi]} \leq \gamma^{-2k_{\mathfrak{L}}[\psi]}$ and $\sum_{j=0}^{\mathfrak{L}-1} (s_{v_j} - 1) = n_{e.p.}[\tau] - 1$, we find

$$\begin{aligned}
|D^h v_{2,0,0,\emptyset}[\tau, \mathbf{k}](\mathbf{x})| &\leq (C'')^{n_{e.p.}[\tau]} e^{-\frac{\tilde{C}}{4}(|x|/\gamma)^\sigma} (\min\{1, |x|\})^{-2\Delta_1} \gamma^{2[\psi](\mathfrak{L}-1)} \gamma^{-2[\psi](k_1 + \dots + k_{\mathfrak{L}})} \cdot \\
&\cdot \sum_{(\ell_i(v_0))_{i=2}^{s_{v_0}}} \cdots \sum_{(\ell_i(v_{\mathfrak{L}-1}))_{i=2}^{s_{v_{\mathfrak{L}-1}}}} \left(\prod_{j=0}^{\mathfrak{L}-1} \left(\prod_{i=2}^{s_{v_j}} (C''\epsilon)^{\max\{1, \frac{l_i(v_j)}{2}-1\}} \right) \right) \sum_{l_{v_{\mathfrak{L}}}} (C''\epsilon)^{l_{v_{\mathfrak{L}}}/2-1} \tag{6.29}
\end{aligned}$$

for a suitable $C'' > 0$. In order to get the desired bound on $D^h v_{2,0,0,\emptyset}^{(h)}(\mathbf{x})$, uniformly as $h \rightarrow -\infty$ (see (6.8)), in view of (6.16), we still need to sum the right hand side of (6.29) over $\mathfrak{L}$, $\mathbf{k}$ and τ and show that the result of the sum is the same as the right hand side of (6.4) (up, possibly, to a redefinition of the constants).

Now, for the sums over $\mathbf{k}$, we simply use

$$\sum_{\substack{k_{\mathfrak{L}} \geq 0 \\ k_0, \dots, k_{\mathfrak{L}-1} \geq 1 \\ k_0 + \dots + k_{\mathfrak{L}} = |h|}} \gamma^{-2[\psi](k_1 + \dots + k_{\mathfrak{L}} - \mathfrak{L} + 1)} \leq (1 - \gamma^{-2[\psi]})^{\mathfrak{L}}, \tag{6.30}$$

so that (letting $\sum_{\mathbf{k}}$ be a shorthand notation for $\sum_{k_{\mathfrak{L}} \geq 0, k_0, \dots, k_{\mathfrak{L}-1} \geq 1}^{k_0 + \dots + k_{\mathfrak{L}} = |h|}$)

$$\begin{aligned}
\sum_{\mathbf{k}} |D^h v_{2,0,0,\emptyset}[\tau, \mathbf{k}](\mathbf{x})| &\leq C^{n_{e.p.}[\tau] + \mathfrak{L}} e^{-\frac{\tilde{C}}{4}(|x|/\gamma)^\sigma} (\min\{1, |x|\})^{-2\Delta_1} \cdot \\
&\cdot \sum_{(\ell_i(v_0))_{i=2}^{s_{v_0}}} \cdots \sum_{(\ell_i(v_{\mathfrak{L}-1}))_{i=2}^{s_{v_{\mathfrak{L}-1}}}} \left(\prod_{j=0}^{\mathfrak{L}-1} \left(\prod_{i=2}^{s_{v_j}} (C''\epsilon)^{\max\{1, \frac{l_i(v_j)}{2}-1\}} \right) \right) \sum_{l_{v_{\mathfrak{L}}}} (C''\epsilon)^{l_{v_{\mathfrak{L}}}/2-1} \tag{6.31}
\end{aligned}$$

up to a redefinition of C . If we now performed the sums in the second line using $(\ell_i(v_j))_{i=2}^{s_{v_j}}, l_{v_{\mathfrak{L}}}$, by forgetting all the constraints on the indices $(\ell_i(v_j))_{i=2}^{s_{v_j}}, l_{v_{\mathfrak{L}}}$ but $l_i(v_j) \geq 2$ and $l_{v_{\mathfrak{L}}} \geq 2$, we would find that the second line of (6.31) would be bounded by $C \prod_{j=0}^{\mathfrak{L}-1} (C\epsilon)^{s_{v_j}-1}$ for some $C > 0$. By re-plugging this back into (6.31), we would be led to a bound that is summable over $\tau \in \mathcal{T}_{\mathfrak{L}}$ (i.e., over $s_{v_j} \geq 1$ for $j = 0, \dots, \mathfrak{L}-1$), but not over $\mathfrak{L}$, and we would be in trouble. Therefore, we must take better advantage of the constraints spelled out in Remark 6.2, by proceeding as follows.

Given $\mathfrak{L} \geq 1$ and $\tau \in \mathcal{T}_{\mathfrak{L}}$, we consider the corresponding set of vertices $\{v_0, \dots, v_p\}$ as in Fig. 11, and rewrite it as the union of the set $V_t(\tau)$ of ‘trivial’ vertices (i.e., those with only one child, $s_{v_j} = 1$) and of the set $V_{nt}(\tau)$ of ‘non-trivial’ ones (i.e., those with $s_{v_j} \geq 2$). We let $p = p(\tau)$ be the cardinality of $V_{nt}(\tau)$ and, if $p \geq 1$, we denote $V_{nt}(\tau) \equiv \{v_{j_1}, \dots, v_{j_p}\}$, with $j_1 < \dots < j_p$. We shall think the tuple $(v_0, v_1, \dots, v_{\mathfrak{L}})$ as a concatenation of the tuples $(v_0, \dots, v_{j_1}), (v_{j_1+1}, \dots, v_{j_2}), \dots, (v_{j_p+1}, \dots, v_{\mathfrak{L}})$. We let $n_1 = j_1 + 1, n_2 = j_2 - j_1, \dots, n_{p+1} = \mathfrak{L} - j_p$ be the lengths of these tuples, and $s_1 = s_{v_{j_1}}, \dots, s_p = s_{v_{j_p}}$ the numbers of children of $v_{j_1}, \dots, v_{j_p}$. Note that the choice of the integers $p \geq 0, n_1, \dots, n_{p+1} \geq 1$ (with $n_1 \geq 2$ if $p = 0$) and $s_1, \dots, s_p \geq 2$ specifies uniquely the choice of $\mathfrak{L}$ and $\tau \in \mathcal{T}_{\mathfrak{L}}$. Therefore, we shall identify the double sum over $\mathfrak{L} \geq 1$ and $\tau \in \mathcal{T}_{\mathfrak{L}}$ with that over $p \geq 0, n_1, \dots, n_{p+1} \geq 1$ (with $n_1 \geq 2$ if $p = 0$) and $s_1, \dots, s_p \geq 2$.




Figure 11: A tree in $\mathcal{T}_{\mathfrak{L}}$ with $\mathfrak{L} = 9$. Its non trivial vertices are v_2 and v_6 , and $s_1 \equiv s_{v_2} = 4, s_2 \equiv s_{v_6} = 3$. The 10-ple $(v_0, \dots, v_9)$ is thought of as a concatenation of three tuples of length $n_1 = 3, n_2 = 4, n_3 = 3$, respectively, namely of $(v_0, v_1, v_2), (v_3, v_4, v_5, v_6)$ and (v_7, v_8, v_9) .

Let us now use the constraints spelled out in Remark 6.2, which we recall here for the reader's convenience: for $j = 0, \dots, \mathfrak{L}-1$, if v_j is trivial, then $l_{v_j} \leq l_{v_{j+1}} - 2$ (with $l_{v_0} \equiv 0$), while, if v_j is non-trivial, then $l_{v_j} \leq l_{v_{j+1}} + \sum_{i=2}^{s_{v_j}} l_i(v_{j_i}) - 2(s_{v_j} - 1)$. Therefore, if $p = 0$, then $l_{v_{\mathfrak{L}}} \geq 2(n_1 - 1)$; while, if $p \geq 1$:

$$\begin{aligned}
2(n_1 - 1) &\leq l_{v_{j_1}} \leq l_{v_{j_1+1}} + \sum_{i=2}^{s_1} l_i(v_{j_{i1}}) + 2(s_1 - 1) \\
l_{v_{j_1+1}} + 2(n_2 - 1) &\leq l_{v_{j_2}} \leq l_{v_{j_2+1}} + \sum_{i=2}^{s_2} l_i(v_{j_{i2}}) + 2(s_2 - 1) \\
&\vdots \\
l_{v_{j_{p-1}+1}} + 2(n_p - 1) &\leq l_{v_{j_p}} \leq l_{v_{j_p+1}} + \sum_{i=2}^{s_p} l_i(v_{j_{ip}}) + 2(s_p - 1) \\
l_{v_{j_p+1}} + 2(n_{p+1} - 1) &\leq l_{v_{\mathfrak{L}}},
\end{aligned} \tag{6.32}$$

which implies, in particular, that every collection of labels contributing to the right hand side of (6.31) satisfies

$$\sum_{i=2}^{s_1} l_i(v_{j_1}) + \cdots + \sum_{i=2}^{s_p} l_i(v_{j_p}) + l_{v_\Sigma} \geq 2(n_1 - 1) + \cdots + 2(n_{p+1} - 1) + 2(s_1 - 1) + \cdots + 2(s_p - 1). \quad (6.33)$$

Therefore, the right hand side of (6.31) can be bounded from above by

$$\begin{aligned} & C^{n_{e.p.}[\tau] + \mathfrak{L}} e^{-\frac{\tilde{C}}{4}(|x|/\gamma)^\sigma} (\min\{1, |x|\})^{-2\Delta_1} (\sqrt{\epsilon})^{[n_1 + \cdots + n_{p+1} - p - 2]_+} \\ & \cdot \sum_{(\ell_i(v_{j_1}))_{i=2}^{s_1}} \left(\prod_{i=2}^{s_1} (C'' \sqrt{\epsilon})^{\max\{1, \frac{l_i(v_{j_1})}{2} - 1\}} \right) \cdots \sum_{(\ell_i(v_{j_p}))_{i=2}^{s_p}} \left(\prod_{i=2}^{s_p} (C'' \sqrt{\epsilon})^{\max\{1, \frac{l_i(v_{j_p})}{2} - 1\}} \right) \sum_{l_{v_\Sigma}} (C'' \sqrt{\epsilon})^{\frac{l_{v_\Sigma}}{2} - 1} \end{aligned} \quad (6.34)$$

where $[\cdot]_+ := \max\{0, \cdot\}$ indicates the positive part. If we now perform the sums in the second line by forgetting all the constraints on the indices $(\ell_i(v_{j_k}))_{i=2}^{s_k}$, l_{v_Σ} , but $l_i(v_{j_k}) \geq 2$ and $l_{v_\Sigma} \geq 2$, we find that the second line of (6.34) is bounded by $C \prod_{k=1}^p (C \sqrt{\epsilon})^{s_k - 1}$ for some $C > 0$. In conclusion, plugging this back into (6.31), and recalling that $\sum_{k=1}^p (s_k - 1) = n_{e.p.}(\tau) - 1$ and $\sum_{k=1}^{p+1} n_k = \mathfrak{L} + 1$, we find, up to a re-definition of C :

$$\begin{aligned} & \sum_{\mathbf{k}} |D^h v_{2,0,0,\emptyset}[\tau, \mathbf{k}](\mathbf{x})| \leq \\ & \leq C^{n_1 + \cdots + n_{p+1}} (\sqrt{\epsilon})^{[n_1 + \cdots + n_{p+1} - p - 2]_+} (C \sqrt{\epsilon})^{\sum_{k=1}^p (s_k - 1)} e^{-\frac{\tilde{C}}{4}(|x|/\gamma)^\sigma} (\min\{1, |x|\})^{-2\Delta_1}, \end{aligned} \quad (6.35)$$

with the understanding that, if $p = 0$, the factor $(C \sqrt{\epsilon})^{\sum_{k=1}^p (s_k - 1)}$ should be interpreted as being equal to 1. Now, the right hand side of this equation is summable over $p \geq 0$, $n_1, \dots, n_{p+1} \geq 1$ (with $n_1 \geq 2$ if $p = 0$) and $s_1, \dots, s_p \geq 2$, the sum being bounded from above by $C e^{-\frac{\tilde{C}}{4}(|x|/\gamma)^\sigma} (\min\{1, |x|\})^{-2\Delta_1}$. In conclusion, in view of (6.16), we find

$$|D^h v_{2,0,0,\emptyset}^{(h)}(\mathbf{x})| \leq C e^{-\frac{\tilde{C}}{4}(|x|/\gamma)^\sigma} (\min\{1, |x|\})^{-2\Delta_1}, \quad (6.36)$$

as desired. Analyticity of $D^h v_{2,0,0,\emptyset}^{(h)}(\mathbf{x})$, uniformly in h , is a consequence of the absolute summability of its tree expansion, as well as of the uniform-in- h bounds that we just derived. A slight extension of the discussion above would also allow us to prove the existence of the limit in the right hand side of (6.8) and to derive explicit estimates on the speed of convergence to the limit. However, in order not to overwhelm an already lengthy discussion, we prefer not to discuss explicitly this point, which is left to the interested reader as a simple exercise. The bounds on $\mathcal{E}_2(y)$ and $\mathfrak{E}_2(y)$ are completely analogous to those discussed above; the minor changes (mostly notational) required for adapting the previous estimates to these functions are left to the reader. The proof of Theorem 2.4 is thus concluded.


A First order computation of ζ_2

In this appendix, we calculate the first order contribution in ϵ to $\zeta_2 = Z_2 - 1$, which implies (2.42) for η_2 . We recall that ζ_2 is expressed via a convergent tree expansion as in (4.24); isolating the first

order from the higher order contributions implies:

$$\zeta_2 = \tilde{H}_{0,1,2,\mathbf{0}}[\tau_0] + \tilde{H}_{0,1,2,\mathbf{0}}[\tau_1] + O(\epsilon^2), \quad (\text{A.1})$$

where τ_0 and τ_1 are the two trees in Fig.A.2 (the analogous trees with an endpoint of type ν replacing the endpoint of type λ give zero contribution).



$$(\text{A.2})$$

In order to compute $\tilde{H}_{0,1,2,\mathbf{0}}[\tau_0]$ and $\tilde{H}_{0,1,2,\mathbf{0}}[\tau_1]$, we recall that the kernel associated with the endpoint of type λ has the form in the third line of (2.34), with $c_2 = \frac{d}{3}$ and (see [1, (5.30), (G.7)]):

$$\begin{aligned} \lambda &= \lambda^* = \frac{-2\epsilon \log \gamma}{I_2} + O(\epsilon^2), \\ I_2 &= -4(N-8) \left[\frac{S_d}{(2\pi)^d} \log \gamma + O(\epsilon(\log \gamma)^2) \right] \end{aligned} \quad (\text{A.3})$$

(here S_d is the area of the unit sphere in $\mathbb{R}^d$). An application of the definition of tree value and a straightforward computation shows that:

$$\begin{aligned} \tilde{H}_{0,1,2,\mathbf{0}}[\tau_0] &= -4(N-2)\lambda^* \int \frac{d^d k}{(2\pi)^d} \frac{f_0(k)^2}{|k|^{d+2\epsilon}}, \\ \tilde{H}_{0,1,2,\mathbf{0}}[\tau_1] &= -8(N-2)\lambda^* \int \frac{d^d k}{(2\pi)^d} \frac{\gamma^{d+\Delta_2-6[\psi]} f_0(k) f_1(k)}{|k|^{d+2\epsilon}}, \end{aligned} \quad (\text{A.4})$$

where $f_h(k) := \chi(\gamma^{-h}k) - \chi(\gamma^{-h+1}k)$. Now, plugging these expressions in (A.1), using (A.3) and the fact that $d + \Delta_2 - 6[\psi] = 2\epsilon + \eta_2 = O(\epsilon)$, we find:

$$\zeta_2 = -2\epsilon \frac{N-2}{N-8} \frac{(2\pi)^d}{S_d} \int \frac{d^d k}{(2\pi)^d} \frac{f_0^2(k) + 2f_0(k)f_1(k)}{|k|^d} + O(\epsilon^2). \quad (\text{A.5})$$

A computation shows that $\int \frac{d^d k}{(2\pi)^d} \frac{f_0^2(k) + 2f_0(k)f_1(k)}{|k|^d} = \frac{S_d}{(2\pi)^d} \log \gamma$, which gives $\zeta_2 = -2\epsilon \frac{N-2}{N-8} \log \gamma + O(\epsilon^2)$, as already announced.

B Proof of (2.43) and (2.44)

Using (2.50)-(2.51) and (5.3), we write:

$$\begin{aligned} \mathcal{G}^*(x) &= 2 \lim_{h \rightarrow -\infty} \gamma^{2h\Delta_1} \sum_{\tau} H_{2,0,0,\emptyset}[\tau](\gamma^h \mathbf{x}), \\ \mathcal{F}^*(y) &= 2 \lim_{h \rightarrow -\infty} \gamma^{2h\Delta_2} \sum_{\tau} H_{0,2,0,\emptyset}[\tau](\gamma^h \mathbf{y}), \end{aligned} \quad (\text{B.1})$$

From Proposition 5.1 and its proof we know that the limits in the right hand sides exist, and that the contribution to $\mathcal{G}^*(x)$ (resp. $\mathcal{F}^*(y)$) from a given tree τ is bounded from above by


$C'(C\epsilon)^{n_{\text{e.p.}}[\tau]-2}|x|^{-2\Delta_1}$ (resp. $C'(C\epsilon)^{n_{\text{e.p.}}[\tau]-2}|y|^{-2\Delta_2}$), for some $C, C' > 0$, see (5.11). Therefore, using the fact that the number of trees with k endpoints is smaller than 4^k , we find that the sum of the contributions to $\mathcal{G}^*(x)$ (resp. $\mathcal{F}^*(y)$) from the trees with 3 or more endpoints is bounded from above by $C\epsilon|x|^{-2\Delta_1}$ (resp. $C\epsilon|y|^{-2\Delta_2}$). Therefore, in order to prove (2.43), it is sufficient to prove that the contribution to $\mathcal{G}^*(x)$ from the tree(s) with 2 endpoints is equal to the right hand side of (2.43) up, possibly, to an error term of the order $\epsilon|x|^{-2\Delta_1}$, and similarly for (2.44).

Dominant contribution to $\mathcal{G}^(x)$.* Direct inspection shows that there is only one tree with 2 endpoints contributing to $\mathcal{G}^*(x)$, which is the following:



whose contribution to $\mathcal{G}^*(x)$ is $\sum_{h' > h} 2^{2h'\Delta_1} g^{(0)}(\gamma^{h'}x)$, which implies (2.43).

Dominant contribution to $\mathcal{F}^(y)$.* Direct inspection shows that there are two trees with 2 endpoints contributing to $\mathcal{F}^*(y)$, which are the following:



whose total contribution to $\mathcal{F}^*(y)$ is (denoting by $\mathbf{p}_h(x)$ the scalar part of $g^{(h)}(x)$, i.e., $g_{ab}^{(h)}(x) \equiv \Omega_{ab}\mathbf{p}_h(x)$):

$$\begin{aligned} & -2N \lim_{h \rightarrow -\infty} \sum_{h' > h} \gamma^{h'(4[\psi]+2\eta_2)} \mathbf{p}_0(\gamma^{h'}y) \left[\mathbf{p}_0(\gamma^{h'}y) + 2 \sum_{k > 0} \gamma^{k(2[\psi]+2\eta_2)} \mathbf{p}_0(\gamma^{h'+k}y) \right] = \\ & = -2N \sum_{h' \in \mathbb{Z}} \mathbf{p}_{h'}(y) \left[\gamma^{2h'\eta_2} \mathbf{p}_{h'}(y) + 2 \sum_{h'' > h'} \gamma^{2h''\eta_2} \mathbf{p}_{h''}(y) \right]. \end{aligned} \quad (\text{B.4})$$

We let $f_h(k) := \chi_h(k) - \chi_{h-1}(k)$, with $\chi_h(k) := \chi(\gamma^{-h}k)$, and, after a relabelling of the scale indices, we rewrite (B.4) as:

$$\begin{aligned} & -2N \sum_{h' \in \mathbb{Z}} \gamma^{2h'\eta_2} \left[\mathbf{p}_{h'}^2(y) + 2 \sum_{h < h'} \mathbf{p}_h(y) \mathbf{p}_{h'}(y) \right] \\ & = -2N \sum_{h' \in \mathbb{Z}} \gamma^{2h'\eta_2} \int \frac{d^d k}{(2\pi)^d |k|^{d/2+\epsilon}} \int \frac{d^d p}{(2\pi)^d |p|^{d/2+\epsilon}} e^{i(k+p) \cdot y} f_{h'}(p) \left[f_{h'}(k) + 2 \sum_{h < h'} f_h(k) \right] \\ & = -2N \sum_{h' \in \mathbb{Z}} \gamma^{2h'\eta_2} \int \frac{d^d k}{(2\pi)^d |k|^{d/2+\epsilon}} \int \frac{d^d p}{(2\pi)^d |p|^{d/2+\epsilon}} e^{i(k+p) \cdot y} [\chi_{h'}(p) \chi_{h'}(k) - \chi_{h'-1}(p) \chi_{h'-1}(k)] \end{aligned} \quad (\text{B.5})$$

where we used:

$$f_{h'}(k) + 2 \sum_{h < h'} f_h(k) = \chi_{h'}(k) - \chi_{h'-1}(k) + 2\chi_{h'-1}(k) = \chi_{h'}(k) + \chi_{h'-1}(k). \quad (\text{B.6})$$

Now, noting that

$$(\gamma^{-h'}|p|)^{-2\eta_2} = 1 + \int_0^1 ds \frac{d}{ds} (\gamma^{-h'}|p|)^{-2s\eta_2} = 1 - 2\eta_2 \log(\gamma^{-h'}|p|) \int_0^1 ds (\gamma^{-h'}|p|)^{-2s\eta_2}, \quad (\text{B.7})$$

multiplying and dividing by $|p|^{-2\eta_2}$ in (B.5) we get:

$$\begin{aligned} (\text{B.5}) &= -2N \sum_{h' \in \mathbb{Z}} \int \frac{d^d k}{(2\pi)^d |k|^{d/2+\epsilon}} \int \frac{d^d p}{(2\pi)^d |p|^{d/2+\epsilon-2\eta_2}} e^{i(k+p) \cdot y} [\chi_{h'}(p) \chi_{h'}(k) - \chi_{h'-1}(p) \chi_{h'-1}(k)] \\ &\quad + 4\eta_2 N \sum_{h' \in \mathbb{Z}} \int \frac{d^d k}{(2\pi)^d |k|^{d/2+\epsilon}} \int \frac{d^d p}{(2\pi)^d |p|^{d/2+\epsilon-2\eta_2}} e^{i(k+p) \cdot y} \log(\gamma^{-h'}|p|) \int_0^1 ds (\gamma^{-h'}|p|)^{-2s\eta_2} \\ &\quad \cdot [\chi_{h'}(k) \chi_{h'}(p) - \chi_{h'-1}(k) \chi_{h'-1}(p)]. \end{aligned} \quad (\text{B.8})$$

We will prove below that the second addend in the right hand side can be bounded as:

$$\begin{aligned} 4|\eta_2|N \left| \sum_{h' \in \mathbb{Z}} \int \frac{d^d k}{(2\pi)^d |k|^{d/2+\epsilon}} \int \frac{d^d p}{(2\pi)^d |p|^{d/2+\epsilon-2\eta_2}} e^{i(k+p) \cdot y} \log(\gamma^{-h'}|p|) \int_0^1 ds (\gamma^{-h'}|p|)^{-2s\eta_2} \right. \\ \left. \cdot [\chi_{h'}(k) \chi_{h'}(p) - \chi_{h'-1}(k) \chi_{h'-1}(p)] \right| \leq C\epsilon |y|^{-2\Delta_2}, \end{aligned} \quad (\text{B.9})$$

that is, this term can be included in the error term $\mathcal{F}_{\text{h.o.}}^*(y)$ defined and bounded in the lines preceding and following (2.44). On the other hand, the first term in the right hand side of (B.8) can be rewritten as follows: recall that the sum over h' in $\mathbb{Z}$ should be interpreted as the limit as $h \rightarrow -\infty$ of the sum over $h' > h$; therefore, using the telescopic structure of the summand and, more specifically, the fact that $\lim_{h \rightarrow -\infty} \sum_{h' > h} [\chi_{h'}(p) \chi_{h'}(k) - \chi_{h'-1}(p) \chi_{h'-1}(k)] = \lim_{h \rightarrow -\infty} (1 - \chi_h(k) \chi_h(p)) = 1$ (where the identities are in the sense of distributions), we see that the first term in the right hand side of (B.8) is equal to

$$\mathcal{F}_0^*(y) = -2N \int \frac{d^d k}{(2\pi)^d |k|^{d/2+\epsilon}} \int \frac{d^d p}{(2\pi)^d |p|^{d/2+\epsilon-2\eta_2}} e^{i(k+p) \cdot y} = \frac{C'_0}{|y|^{d-2\epsilon+2\eta_2}} = \frac{C'_0}{|y|^{2\Delta_2}}, \quad (\text{B.10})$$

as desired.

We are left with proving (B.9). By adding and subtracting $\chi_{h'-1}(k) \chi_{h'}(p)$ to the term in square

brackets, we rewrite the second term in the right hand side of (B.8) as follows:

$$\begin{aligned}
& 4\eta_2 N \sum_{h' \in \mathbb{Z}} \int \frac{d^d k}{(2\pi)^d |k|^{d/2+\epsilon}} \int \frac{d^d p}{(2\pi)^d |p|^{d/2+\epsilon-2\eta_2}} e^{i(k+p) \cdot y} \log(\gamma^{-h'} |p|) \cdot \\
& \quad \cdot \int_0^1 ds (\gamma^{-h'} |p|)^{-2s\eta_2} [f_{h'}(k) \chi_{h'}(p) + \chi_{h'-1}(k) f_{h'}(p)] \\
& = 4\eta_2 N \sum_{h' \in \mathbb{Z}} \left(\frac{d^d k}{(2\pi)^d |k|^{d/2+\epsilon}} e^{ik \cdot y} f_{h'}(k) \right) \cdot \\
& \quad \cdot \left(\int \frac{d^d p}{(2\pi)^d |p|^{d/2+\epsilon-2\eta_2}} e^{ip \cdot y} \log(\gamma^{-h'} |p|) \int_0^1 ds (\gamma^{-h'} |p|)^{-2s\eta_2} \chi_{h'}(p) \right) \quad (\text{B.11}) \\
& + 4\eta_2 N \sum_{h' \in \mathbb{Z}} \left(\frac{d^d k}{(2\pi)^d |k|^{d/2+\epsilon}} e^{ik \cdot y} \chi_{h'-1}(k) \right) \cdot \\
& \quad \cdot \left(\int \frac{d^d p}{(2\pi)^d |p|^{d/2+\epsilon-2\eta_2}} e^{ip \cdot y} \log(\gamma^{-h'} |p|) \int_0^1 ds (\gamma^{-h'} |p|)^{-2s\eta_2} f_{h'}(p) \right) \\
& \equiv 4\eta_2 N \sum_{h' \in \mathbb{Z}} \int_0^1 ds \left[\mathfrak{p}_{h'}(y) \sum_{h \leq h'} \tilde{\mathfrak{p}}_{h;s,h'}(y) + P_{\leq h'-1}(y) \tilde{\mathfrak{p}}_{h';s,h'}(y) \right]
\end{aligned}$$

where we recall that $\mathfrak{p}_h(y)$ denotes the scalar part of $g^{(h)}(y)$ (i.e., $g_{ab}^{(h)}(y) \equiv \Omega_{ab} \mathfrak{p}_h(y)$) and, analogously, $P_{\leq h}(y)$ denotes the scalar part of $G^{(\leq h)}(y)$, see footnote 3; moreover,

$$\tilde{\mathfrak{p}}_{h;s,h'}(y) := \int \frac{d^d p}{(2\pi)^d |p|^{d/2+\epsilon-2\eta_2}} e^{ip \cdot y} \log(\gamma^{-h'} |p|) (\gamma^{-h'} |p|)^{-2s\eta_2} f_h(p). \quad (\text{B.12})$$

Now, recall that, by (4.14) and the definition of $\mathfrak{p}_h$,

$$|\mathfrak{p}_h(y)| \leq C_{\chi^1} \gamma^{h(d/2-\epsilon)} e^{-C_{\chi^2}(\gamma^{h-1}|y|)^\sigma}, \quad (\text{B.13})$$

whose proof is given in [1, Appendix A.2]. In order to get a bound on $\mathfrak{p}_{h;s,h'}(y)$ we proceed as follows: in the integral in the right hand side of (B.12), we rescale $p \rightarrow \gamma^h p$, thus getting:

$$\begin{aligned}
\tilde{\mathfrak{p}}_{h;s,h'}(y) & = \gamma^{h(d/2-\epsilon+2\eta_2)} \gamma^{(h'-h)2s\eta_2} \\
& \cdot \int \frac{d^d p}{(2\pi)^d |p|^{d/2+\epsilon+2(s-1)\eta_2}} e^{ip \cdot \gamma^h y} [\log |p| - (h' - h) \log \gamma] f_0(p) \quad (\text{B.14})
\end{aligned}$$

The integral in the second line can be bounded by proceeding exactly as in [1, Appendix A.2] (note, in particular, that $F(p) := \log |p|$ admits an analytic continuation into the polydisk centered at $p \neq 0$ of radius $R = \frac{1}{2} \max_i |p_i|$ such that the maximum of $|F(z)|$ in the polydisk is bounded by $C(1 + |F(p)|)$: this allows us to proceed as in [1, (A.15)] and following lines); we thus get, for some $C_1, C_2 > 0$:

$$|\tilde{\mathfrak{p}}_{h;s,h'}(y)| \leq C_1 (1 + h' - h) \gamma^{h(d/2-\epsilon+2\eta_2)} \gamma^{(h'-h)2s\eta_2} e^{-C_2(\gamma^{h-1}|y|)^\sigma}. \quad (\text{B.15})$$

We can now bound (B.11) via (B.13) and (B.15), thus getting, for some $C > 0$ and $\bar{C} = \min\{C_{\chi^2}, C_2\}$:

$$\begin{aligned}
& \left| 4\eta_2 N \sum_{h' \in \mathbb{Z}} \int_0^1 ds \left[\mathfrak{p}_{h'}(y) \sum_{h \leq h'} \tilde{\mathfrak{p}}_{h;s,h'}(y) + P_{\leq h'-1}(y) \tilde{\mathfrak{p}}_{h';s,h'}(y) \right] \right| \\
& \leq CN|\eta_2| \sum_{h' \in \mathbb{Z}} e^{-\bar{C}(\gamma^{h'-1}|y|)^\sigma} \sum_{h \leq h'} \left(\gamma^{h'(d/2-\epsilon)} \gamma^{h(d/2-\epsilon+2\eta_2)} (1+h'-h) \int_0^1 ds \gamma^{2s(h'-h)\eta_2} \right. \\
& \quad \left. + \gamma^{h(d/2-\epsilon)} \gamma^{h'(d/2-\epsilon+2\eta_2)} \right) \\
& \leq C'\epsilon \sum_{h' \in \mathbb{Z}} \gamma^{2h'\Delta_2} e^{-\bar{C}(\gamma^{h'-1}|y|)^\sigma} \leq C''\epsilon |y|^{-2\Delta_2}
\end{aligned} \tag{B.16}$$

where we used that $\eta_2 = O(\epsilon)$. This conclude the proof of (2.44).

Acknowledgements

We gratefully acknowledge support from: the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme, ERC CoG UniCoSM, grant agreement n.724939 (A.G.) and ERC StG MaMBoQ, n.80290 (G.S.); MUR, PRIN 2017 project MaQuMA cod. 2017ASFLJR (A.G., V.M., G.S.); MUR, PRIN 2022 project MaIQuFi cod. 20223J85K3; GNFM-INdAM Gruppo Nazionale per la Fisica Matematica.

References

- [1] A. Giuliani, V. Mastropietro, and S. Rychkov, “Gentle introduction to rigorous Renormalization Group: a worked fermionic example,” [JHEP 01 \(2021\) 026](#), [arXiv:2008.04361 \[hep-th\]](#).
- [2] K. Gawedzki and A. Kupiainen, “Massless lattice ϕ_4^4 theory: Rigorous control of a renormalizable asymptotically free model,” [Comm. Math. Phys. 99 no. 2, \(1985\) 197–252](#), <https://projecteuclid.org/euclid.cmp/1103942678>.
- [3] K. Gawedzki and A. Kupiainen, “Gross-Neveu Model Through Convergent Perturbation Expansions,” [Comm. Math. Phys. 102 \(1985\) 1](#), <https://projecteuclid.org/euclid.cmp/1104114300>.
- [4] K. Gawedzki and A. Kupiainen, “Renormalization of a Nonrenormalizable Quantum Field Theory,” [Nucl. Phys. B262 \(1985\) 33–48](#).
- [5] M. Fisher, S.-K. Ma, and B. Nickel, “Critical exponents for long-range interactions,” [Phys. Rev. Lett. \(1972\) 917–920](#).
- [6] J. Sak, “Recursion relations and fixed points for ferromagnets with long-range interactions,” [Phys. Rev. B 8 \(1973\) 281–285](#).

- [7] J. Sak, “Low-temperature renormalization group for ferromagnets with long-range interactions,” Phys. Rev. B **15** (1977) 4344–4347.
- [8] D. C. Brydges, P. K. Mitter, and B. Scoppola, “Critical $\Phi_{3,\epsilon}^4$,” Comm. Math. Phys. **240** (2003) 281–327, [arXiv:hep-th/0206040 \[hep-th\]](#).
- [9] A. Abdesselam, “A Complete Renormalization Group Trajectory Between Two Fixed Points,” Commun. Math. Phys. **276** (2007) 727–772, [arXiv:math-ph/0610018 \[math-ph\]](#).
- [10] G. Slade, “Critical exponents for long-range $O(n)$ models below the upper critical dimension,” Communications in Mathematical Physics **358** no. 1, (2018) 343–436, [arXiv:1611.06169 \[math-ph\]](#).
- [11] D. Poland, S. Rychkov, and A. Vichi, “The Conformal Bootstrap: Theory, Numerical Techniques, and Applications,” Rev. Mod. Phys. **91** (2019) 015002, [arXiv:1805.04405 \[hep-th\]](#).
- [12] R. Bauerschmidt, D. C. Brydges, and G. Slade, “Scaling limits and critical behaviour of the 4-dimensional n -component $|\phi|^4$ spin model,” J. Stat. Phys. **157** (2014) 692–742.
- [13] Bauerschmidt, R. and Slade, G. and Tomberg, A. and Wallace, B. C., “Finite-order correlation length for 4-dimensional weakly self-avoiding walk and $|\phi|^4$ spins,” Annales Henri Poincaré **18** (2017) 375–402.
- [14] G. Slade and A. Tomberg, “Critical correlation functions for the 4-dimensional weakly self-avoiding walk and n -component $|\phi|^4$ model,” Commun. Math. Phys. **342** (2016) 675–737.
- [15] Feldman, J. and Magnen, J. and Rivasseau, V. and Sénéor, R., “A renormalizable field theory: the massive Gross-Neveu model in two dimensions,” Comm. Math. Phys. **103** no. 1, (1986) 67–103.
- [16] Dimock, J. and Cheng Yuan, “Structural stability of the RG flow in the Gross-Neveu model,” [arXiv:2303.07916 \[math-ph\]](#).
- [17] G. Benfatto and V. Mastropietro, “Ward Identities and Chiral Anomaly in the Luttinger Liquid,” Comm. Math. Phys. **258** (2005) 609–655.
- [18] G. Benfatto, P. Falco, and V. Mastropietro, “Functional Integral Construction of the Massive Thirring model: Verification of Axioms and Massless Limit,” Comm. Math. Phys. **273** (2007) 67–118, [arXiv:0606177 \[hep-th\]](#).
- [19] G. Benfatto, P. Falco, and V. Mastropietro, “Extended Scaling Relations for Planar Lattice Models,” Comm. Math. Phys. **292** (2009) 569–605.
- [20] A. Giuliani, V. Mastropietro, and F. L. Toninelli, “Non-integrable dimers: Universal fluctuations of tilted height profiles,” Comm. Math. Phys. **377** (2020) 1883–1959.
- [21] F. Wegner, “The Critical State, General Aspects,” in Phase Transitions and Critical Phenomena, Vol.6, C. Domb and M. Green, eds., pp. 8–126. Academic Press, 1976.
- [22] A. Kupiainen, R. Rhodes, and V. Vargas, “Local Conformal Structure of Liouville Quantum Gravity,” Comm. Math. Phys. **371** (2019) 1005–1069.

- [23] A. Kupiainen, R. Rhodes, and V. Vargas, “Integrability of Liouville theory: proof of the DOZZ Formula,” Annals of Mathematics **191** no. 1, (2020) 81–166.
- [24] C. Guillarmou, A. Kupiainen, R. Rhodes, and V. Vargas, “Conformal bootstrap in Liouville Theory,” Acta Mathematica **to appear**, [arXiv:2005.11530 \[math probability\]](#).
- [25] P. Kravchuk, J. Qiao, and S. Rychkov, “Distributions in CFT. Part II. Minkowski space,” JHEP **08** (2021) 094, [arXiv:2104.02090 \[hep-th\]](#).
- [26] S. Rychkov, “3D Ising Model: a view from the Conformal Bootstrap Island,” Comptes Rendus Physique **21** no. 2, (2020) 185–198, [arXiv:2007.14315 \[math-ph\]](#).
- [27] A. Z. Patashinsky and V. N. Pokrovsky, Fluctuation Theory of Phase Transitions. Oxford, England: Pergamon Pr., 1979. 321 p.
- [28] R. L. Greenblatt, “Constructing a weakly-interacting fixed point of the Fermionic Polchinski equation,” [arXiv:2401.04462 \[math-ph\]](#).
- [29] G. Grubb, Distributions and operators, vol. 252. Springer Science & Business Media, 2008.
- [30] A. Giuliani, R. L. Greenblatt, and V. Mastropietro, “The scaling limit of the energy correlations in non integrable Ising models,” J. Math. Phys. **53** (2012) 095214, [arXiv:1204.4040 \[math-ph\]](#).
- [31] G. Antinucci, A. Giuliani, and R. L. Greenblatt, “Non-integrable Ising models in cylindrical geometry: Grassmann representation and infinite volume limit,” Annales Henri Poincaré **23** (2022) 1061–1139, [arXiv:2104.09328 \[math-ph\]](#).
- [32] G. Antinucci, A. Giuliani, and R. L. Greenblatt, “Energy correlations of non-integrable Ising models: The scaling limit in the cylinder,” Comm. Math. Phys. **397** (2023) 393–483, [arXiv:2006.04458 \[math-ph\]](#).
- [33] G. Gentile and V. Mastropietro, “Renormalization group for one-dimensional fermions. A review on mathematical results,” Phys. Rep. **352** (2001) 273–437.
- [34] G. Gallavotti, The elements of mechanics. Springer Science & Business Media, 1983.