

DENSITY OF SYSTOLES OF HYPERBOLIC MANIFOLDS

SAMI DOUBA AND JUNZHI HUANG

ABSTRACT. We show that for each $n \geq 2$, the systoles of closed hyperbolic n -manifolds form a dense subset of $(0, +\infty)$. We also show that for any $n \geq 2$ and any Salem number λ , there is a closed arithmetic hyperbolic n -manifold of systole $\log(\lambda)$. In particular, the Salem conjecture holds if and only if the systoles of closed arithmetic hyperbolic manifolds in some (any) dimension fail to be dense in $(0, +\infty)$.

The *systole* $\text{sys}(M)$ of a closed nonpositively curved Riemannian manifold M is the length of a shortest closed geodesic in M . It follows for instance from residual finiteness of surface groups that there is a sequence of closed hyperbolic surfaces with systole going to infinity. Since for any $\epsilon > 0$ and any closed surface M of negative Euler characteristic there is a hyperbolic metric on M of systole $< \epsilon$, continuity of the systole function on moduli space then yields for any $L > 0$ a closed hyperbolic surface of systole precisely L .

By Mostow rigidity [16], the set of systoles of closed hyperbolic n -manifolds is countable for $n \geq 3$. Nevertheless, we show the following.

Theorem 1. *For any $n \geq 2$, the set of systoles of closed hyperbolic n -manifolds is dense in $(0, +\infty)$.*

That 0 is an accumulation point of systoles of closed hyperbolic 3-manifolds follows for instance from Thurston’s hyperbolic Dehn filling theory (see [4, Sections E.5 and E.6]). The analogous fact for 4-manifolds was established by Agol [1], whose strategy was to “inbreed” arithmetic manifolds along totally geodesic hypersurfaces. This strategy was successfully extended to arbitrary dimensions independently by Belolipetsky–Thomson [3] and Bergeron–Haglund–Wise [6]. The present note is an elaboration on the utility of this inbreeding technique.

We also establish a variant of Theorem 1 for arithmetic hyperbolic manifolds. A *Salem number* is a real algebraic integer $\lambda > 1$ of degree ≥ 4 that is Galois-conjugate to λ^{-1} and all of whose remaining Galois conjugates lie on the unit circle.

Theorem 2. *Let $\lambda > 1$ be a Salem number. Then for any $n \geq 2$, there is a closed arithmetic hyperbolic n -manifold of systole precisely $\log(\lambda)$.*

The manifolds we exhibit in the proof of Theorem 2 are arithmetic of simplest type (the latter being the only arithmetic construction that applies in every dimension) and are moreover *classical* in the language of Emery–Ratcliffe–Tschantz [10]. For $n \geq 4$, it follows from Meyer’s theorem on quadratic forms and [10, Thm. 5.2] that the exponential lengths of closed geodesics in any classical simplest-type closed arithmetic hyperbolic n -manifold are Salem numbers.¹

¹Exponential geodesic lengths in closed arithmetic hyperbolic surfaces or 3-manifolds can be *quadratic* “Salem numbers”; the latter are sometimes included in the definition of a Salem number. Exponential geodesic lengths in odd-dimensional closed arithmetic hyperbolic orbifolds need not

Lehmer's conjecture for Salem numbers, or the *Salem conjecture* for short, asserts that the Salem numbers are bounded away from 1; see [13, Section 13] and [18, p. 31]. In light of the relationship between Salem numbers and lengths of geodesics in arithmetic hyperbolic manifolds, Theorem 2 yields the following reformulation of the Salem conjecture.

Corollary 2.1. *The Salem conjecture holds if and only if, for some, equivalently, any, $n \geq 2$, the set of systoles of closed arithmetic hyperbolic n -manifolds fails to be dense in $(0, +\infty)$.*

Proof. It is known generally that, given any irreducible symmetric space X of noncompact type, the Salem conjecture implies a uniform lower bound on systoles of closed arithmetic locally symmetric manifolds modeled on X ; see Frączyk–Pham [11]. On the other hand, since positive powers of Salem numbers remain Salem numbers, failure of the Salem conjecture would imply density of Salem numbers in $(1, +\infty)$, and hence density of systoles of closed arithmetic hyperbolic n -manifolds in $(0, +\infty)$ for each $n \geq 2$ by Theorem 2. \square

We motivate the proofs of Theorems 1 and 2 with the following remark.

Remark 3. We provide an argument along the lines of [1] that, given any $L > 0$, there is a closed hyperbolic surface of systole precisely L . Indeed, it suffices to find a closed hyperbolic surface M_L of systole $\geq L$ possessing two disjoint 2-sided simple closed geodesics Σ_1 and Σ_2 and an orthogeodesic segment ω joining the Σ_i of length $L/2$ and whose length is minimal among all orthogeodesic segments with endpoints on $\Sigma_1 \cup \Sigma_2$. Cutting M_L along the Σ_i and then doubling the resulting surface along its boundary yields a (possibly disconnected) closed hyperbolic surface of systole precisely L .

One can construct such a surface M_L as follows. Let $P \subset \mathbb{H}^2$ be a right-angled pentagon with an edge $\tilde{\omega}$ of length precisely $L/2$, and let $H_1, H_2 \subset \mathbb{H}^2$ be the walls of P adjacent to $\tilde{\omega}$. Let $\Gamma_P < \text{Isom}(\mathbb{H}^2)$ be the group generated by the reflections in the walls of P . We can find a larger right-angled convex polygon $Q \subset \mathbb{H}^2$ that is a union of finitely many Γ_P -translates of P such that the walls of Q that enter the $\frac{L}{2}$ -neighborhood of H_i are orthogonal to H_i for $i = 1, 2$ (this idea originates in work of Scott [19]; see also [2, Section 3.1]). By residual finiteness and virtual torsion-freeness of the group $\Gamma_Q < \text{Isom}(\mathbb{H}^2)$ generated by the reflections in the walls of Q , there is a finite-index subgroup $\Lambda < \Gamma_Q$ such that $\Lambda \backslash \mathbb{H}^2$ is a surface of systole $\geq L$. We may now take $M_L = \Lambda \backslash \mathbb{H}^2$, the Σ_i to be the projections to M_L of the H_i , and ω to be the projection of $\tilde{\omega}$.

Proof of Theorem 1. Given $L, \epsilon > 0$, we exhibit a closed hyperbolic n -manifold M with $|\text{sys}(M) - L| < \epsilon$. As in Remark 3, it suffices to find a closed hyperbolic n -manifold $M_{L,\epsilon}$ of systole $\geq L + \epsilon$ possessing two disjoint 2-sided properly embedded totally geodesic hypersurfaces Σ_1 and Σ_2 , and an orthogeodesic segment ω joining the Σ_i of length within $\epsilon/2$ from $L/2$ and whose length is minimal among all orthogeodesic segments with endpoints on $\Sigma_1 \cup \Sigma_2$. We can then cut $M_{L,\epsilon}$ along $\Sigma_1 \cup \Sigma_2$, select the component N containing ω of the resulting manifold with boundary, and define M to be the double of N . The double of ω is indeed a shortest closed geodesic in M of length within ϵ from L .

be Salem numbers even in this more general sense; see Lemma 4.10 and Theorem 4.11 in [17], as well as [10, Thm. 7.7].

To that end, let $k = \mathbb{Q}(\sqrt{2})$ and $\mathcal{O}_k = \mathbb{Z}[\sqrt{2}]$. Define f to be the quadratic form on \mathbb{R}^{n+1} given by

$$f(x_1, \dots, x_{n+1}) = x_1^2 + \dots + x_n^2 - \sqrt{2}x_{n+1}^2.$$

We identify the level set $\{x \in \mathbb{R}^{n+1} | f(x) = -1, x_{n+1} > 0\}$ with n -dimensional hyperbolic space \mathbb{H}^n and $O'(f; \mathbb{R})$ with $\text{Isom}(\mathbb{H}^n)$, where $O'(f; \mathbb{R})$ is the index-2 subgroup of $O(f; \mathbb{R})$ preserving \mathbb{H}^n . By the Borel–Harish-Chandra theorem [7], we have that $\Gamma := O'(f; \mathcal{O}_k)$ is a uniform arithmetic lattice of $\text{Isom}(\mathbb{H}^n)$.

Denote by H_1 the hyperplane $\{x_1 = 0\}$ in \mathbb{H}^n . Since $O'(f; k)$ is dense in $O'(f; \mathbb{R})$, there is an element $g \in O'(f; k)$ such that

$$\left| \text{dist}_{\mathbb{H}^n}(H_1, gH_1) - \frac{L}{2} \right| < \frac{\epsilon}{2}.$$

Let $\tilde{\omega}$ be the orthogeodesic segment in \mathbb{H}^n connecting H_1 and $H_2 := gH_1$. By the proof of Lemma 3.1 in Belolipetsky–Thomson [3], we may now pass to a non-zero ideal $I \subset \mathcal{O}_k$ such that $\text{dist}_{\mathbb{H}^n}(H_1, \gamma H_2) \geq \text{dist}_{\mathbb{H}^n}(H_1, H_2)$ for each γ in the principal congruence subgroup $\Gamma(I) < \Gamma$ of level I . Up to diminishing the ideal I , we can further assume that

- (1) the principal congruence subgroup $\Gamma(I)$ is torsion-free, so that $M_{L, \epsilon} := \Gamma(I) \backslash \mathbb{H}^n$ is a manifold;
- (2) we have $\text{sys}(M_{L, \epsilon}) \geq L + \epsilon$;
- (3) for $i = 1, 2$ and $\gamma \in \Gamma(I)$, either $\gamma H_i = H_i$, in which case γ furthermore preserves each side of H_i , or $\text{dist}_{\mathbb{H}^n}(H_i, \gamma H_i) \geq \text{dist}_{\mathbb{H}^n}(H_1, H_2)$; in particular, the projections Σ_i of the H_i to $M_{L, \epsilon}$ are 2-sided and properly *embedded*.

Note that item (2) can be arranged by residual finiteness of the ring \mathcal{O}_k , whereas item (3) can be ensured by the fact that the stabilizer in Γ of either side of H_i is an intersection of congruence subgroups of Γ for $i = 1, 2$; see [14] and [5, Lemme principal]. We now have that the Σ_i are disjoint and that any orthogeodesic segment in $M_{L, \epsilon}$ with endpoints on $\Sigma_1 \cup \Sigma_2$ has length at least that of the projection ω of $\tilde{\omega}$ to $M_{L, \epsilon}$. Thus, the manifold $M_{L, \epsilon}$, the hypersurfaces Σ_i , and the orthogeodesic segment ω are as desired. \square

Remark 4. The proof of Theorem 1 demonstrates that we may take the closed hyperbolic manifolds giving rise to a dense set of systoles in $(0, +\infty)$ to be quasi-arithmetic in the sense of Vinberg and all share the same adjoint trace field (indeed, the same Vinberg ambient algebraic group); see [20] and compare [8, Thm. 1.3]. On the other hand, by varying the form f in the proof, one easily produces a family of pairwise incommensurable closed hyperbolic manifolds in each dimension whose systoles remain dense in $(0, +\infty)$.

In fact, it is possible to achieve the latter without varying f ; one can select from any family of closed hyperbolic manifolds whose systoles are dense in $(0, +\infty)$ pairwise incommensurable manifolds whose systoles remain dense in $(0, +\infty)$. Indeed, given a closed hyperbolic manifold M , the set of all geodesic lengths of manifolds commensurable to M is closed and discrete in $[0, +\infty)$. When M is nonarithmetic, the latter follows from Margulis’s arithmeticity criterion [15, Thm. IX.6.5], and when M is arithmetic, from the fact that, for fixed $D > 0$ and $\mu > 1$, there are only finitely many monic integer polynomials of degree $\leq D$ and Mahler measure $\leq \mu$; see the discussion immediately following Conjecture 10.2 in [12].

Remark 5. Instead of appealing to the work of Belolipetsky–Thomson (loc. cit.) to find the manifold $M_{L,\epsilon}$ in the proof of Theorem 1, we could have instead used [6, Cor. 1.12] as in the proof of Theorem 4 in [9]. An interesting feature of the former approach is that it afforded us a manifold $M_{L,\epsilon}$ that is *congruence* arithmetic.

Proof of Theorem 2. Let $\mu = \lambda + \lambda^{-1}$. Let $k = \mathbb{Q}(\mu)$, and \mathcal{O}_k be the ring of integers of k . Following the proof of [10, Thm. 6.3], we define the form f in $n+1$ variables over k to be that given by the symmetric matrix

$$\begin{pmatrix} 1 & & \mu/2 \\ & I_{n-1} & \\ \mu/2 & & 1 \end{pmatrix}.$$

The form f is of signature $(n, 1)$, so that we may again identify \mathbb{H}^n with one of the two sheets of the hyperboloid $\{x \in \mathbb{R}^{n+1} | f(x) = -1\}$ and $\text{Isom}(\mathbb{H}^n)$ with $O'(f; \mathbb{R})$, where $O'(f; \mathbb{R})$ is the index-2 subgroup of $O(f; \mathbb{R})$ preserving \mathbb{H}^n . Moreover, the field k is totally real, and for any embedding $\sigma : k \rightarrow \mathbb{R}$ other than the identity (of which at least one exists), the form f^σ is positive definite. It follows that $\Gamma := O'(f; \mathcal{O}_k)$ is a uniform arithmetic lattice of $O'(f; \mathbb{R}) = \text{Isom}(\mathbb{H}^n)$, which contains the reflections

$$\tau_1 := \begin{pmatrix} -1 & & -\mu \\ & I_{n-1} & \\ & & 1 \end{pmatrix}, \quad \tau_2 := \begin{pmatrix} -\mu & & 1 - \mu^2 \\ & I_{n-1} & \\ 1 & & \mu \end{pmatrix}$$

whose respective fixed hyperplanes $H_1, H_2 \subset \mathbb{H}^n$ satisfy

$$\text{dist}_{\mathbb{H}^n}(H_1, H_2) = \log(\lambda)/2.$$

As in the proof of Theorem 1, by the aforementioned work of Belolipetsky–Thomson (loc. cit.), we may pass to a non-zero ideal $I \subset \mathcal{O}_k$ such that

- the principal congruence subgroup $\Gamma(I)$ is torsion-free, so that $M_{\log(\lambda)} := \Gamma(I) \backslash \mathbb{H}^n$ is a manifold;
- we have $\text{sys}(M_{\log(\lambda)}) \geq \log(\lambda)$;
- the H_i project to disjoint properly embedded 2-sided totally geodesic hypersurfaces Σ_1 and Σ_2 in $M_{\log(\lambda)}$;
- any orthogeodesic segment in $M_{\log(\lambda)}$ with endpoints on $\Sigma_1 \cup \Sigma_2$ has length $\geq \log(\lambda)/2$.

Now cutting $M_{\log(\lambda)}$ along the Σ_i and then doubling the resulting manifold along its boundary yields a (possibly disconnected) closed manifold M of systole precisely $\log(\lambda)$ each of whose components is arithmetic (since $\tau_i \in \Gamma$ for $i = 1, 2$). \square

Acknowledgements. We thank Misha Belolipetsky, Bram Petri, and Alan Reid for helpful comments. S.D. was supported by the Huawei Young Talents Program. This work was initiated during a visit of S.D. to Yale University in March 2024; S.D. thanks Sebastian Hurtado and the entire geometry/topology/dynamics team at Yale for their hospitality.

REFERENCES

- [1] I. AGOL, *Systoles of hyperbolic 4-manifolds*, arXiv preprint math/0612290, (2006).
- [2] I. AGOL, D. D. LONG, AND A. W. REID, *The Bianchi groups are separable on geometrically finite subgroups*, Ann. of Math. (2), 153 (2001), pp. 599–621.
- [3] M. V. BELOLIPETSKY AND S. A. THOMSON, *Systoles of hyperbolic manifolds*, Algebr. Geom. Topol., 11 (2011), pp. 1455–1469.

- [4] R. BENEDETTI AND C. PETRONIO, *Lectures on hyperbolic geometry*, Universitext, Springer-Verlag, Berlin, 1992.
- [5] N. BERGERON, *Premier nombre de Betti et spectre du laplacien de certaines variétés hyperboliques*, Enseign. Math. (2), 46 (2000), pp. 109–137.
- [6] N. BERGERON, F. HAGLUND, AND D. T. WISE, *Hyperplane sections in arithmetic hyperbolic manifolds*, J. Lond. Math. Soc. (2), 83 (2011), pp. 431–448.
- [7] A. BOREL AND HARISH-CHANDRA, *Arithmetic subgroups of algebraic groups*, Ann. of Math. (2), 75 (1962), pp. 485–535.
- [8] G. COSAC AND C. DÓRIA, *Closed geodesics on semi-arithmetic Riemann surfaces*, Math. Res. Lett., 29 (2022), pp. 961–1001.
- [9] S. DOUBA, *Systoles of hyperbolic hybrids*, arXiv preprint arXiv:2309.16051, (2023).
- [10] V. EMERY, J. G. RATCLIFFE, AND S. T. TSCHANTZ, *Salem numbers and arithmetic hyperbolic groups*, Transactions of the American Mathematical Society, 372 (2019), pp. 329–355.
- [11] M. FRACZYK AND L. L. PHAM, *Bottom of the length spectrum of arithmetic orbifolds*, Trans. Amer. Math. Soc., 376 (2023), pp. 4745–4764.
- [12] T. GELANDER, *Homotopy type and volume of locally symmetric manifolds*, Duke Math. J., 124 (2004), pp. 459–515.
- [13] D. H. LEHMER, *Factorization of certain cyclotomic functions*, Ann. of Math. (2), 34 (1933), pp. 461–479.
- [14] D. D. LONG, *Immersion and embeddings of totally geodesic surfaces*, Bull. London Math. Soc., 19 (1987), pp. 481–484.
- [15] G. A. MARGULIS, *Discrete subgroups of semisimple Lie groups*, vol. 17 of Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], Springer-Verlag, Berlin, 1991.
- [16] G. D. MOSTOW, *Quasi-conformal mappings in n -space and the rigidity of hyperbolic space forms*, Publications Mathématiques de l’IHÉS, 34 (1968), pp. 53–104.
- [17] W. D. NEUMANN AND A. W. REID, *Arithmetic of hyperbolic manifolds*, in Topology ’90 (Columbus, OH, 1990), vol. 1 of Ohio State Univ. Math. Res. Inst. Publ., de Gruyter, Berlin, 1992, pp. 273–310.
- [18] R. SALEM, *Algebraic numbers and Fourier analysis*, in Selected reprints, Wadsworth Math. Ser., Wadsworth, Belmont, CA, 1983, pp. iii+68.
- [19] P. SCOTT, *Subgroups of surface groups are almost geometric*, J. London Math. Soc. (2), 17 (1978), pp. 555–565.
- [20] S. THOMSON, *Quasi-arithmeticity of lattices in $PO(n, 1)$* , Geom. Dedicata, 180 (2016), pp. 85–94.

INSTITUT DES HAUTES ÉTUDES SCIENTIFIQUES, 35 ROUTE DE CHARTRES, 91440 BURES-SUR-YVETTE, FRANCE

Email address: `douba@ihes.fr`

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, NEW HAVEN, CT 06511, USA

Email address: `junzhi.huang@yale.edu`