ONE-BUBBLE NODAL BLOW-UP FOR ASYMPTOTICALLY CRITICAL STATIONARY SCHRÖDINGER-TYPE EQUATIONS

BRUNO PREMOSELLI AND FRÉDÉRIC ROBERT

ABSTRACT. We investigate in this work families $(u_{\varepsilon})_{\varepsilon>0}$ of sign-changing blowingup solutions of asymptotically critical stationary nonlinear Schrödinger equations of the following type:

$$\Delta_a u_\varepsilon + h_\varepsilon u_\varepsilon = |u_\varepsilon|^{p_\varepsilon - 2} u_\varepsilon$$

in a closed manifold (M, g), where h_{ε} converges in $C^1(M)$. Assuming that $(u_{\varepsilon})_{\varepsilon>0}$ blows-up as a single sign-changing bubble, we obtain necessary conditions for blow-up that constrain the localisation of blow-up points and exhibit a strong interaction between h, the geometry of (M, g) and the bubble itself. These conditions are new and are a consequence of the sign-changing nature of u_{ε} .

1. INTRODUCTION

1.1. Statement of the main results. Let $(M^n, g), n \ge 3$ be a smooth, connected and closed manifold, where closed means compact without boundary. We study in this paper sign-changing solutions $u \in C^2(M)$ of the equation

(1.1)
$$\Delta_a u + hu = |u|^{p-2} u \text{ in } M$$

where $\Delta_g = -\operatorname{div}_g(\nabla \cdot)$ is the Laplace-Beltrami operator, $h \in C^1(M)$ and $2 with <math>2^* = \frac{2n}{n-2}$. When $h \equiv \frac{n-2}{4(n-1)}S_g$, where S_g is the scalar curvature of (M,g), $p = 2^*$ and u > 0, (1.1) is the celebrated Yamabe equation. We let $H^1(M)$ be the completion of $C^{\infty}(M)$ for $u \mapsto ||u||_{H^1} := ||u||_2 + ||\nabla u||_2$. It has been known since the seminal work of Struwe [36] that families of solutions to (1.1) that are uniformly bounded in $H^1(M)$ may develop concentration phenomena in the form of "bubbles" (see also Druet-Hebey-Robert [12] for the case of a compact Riemannian manifold). These "bubbles" correspond to a loss of compactness of the solutions, and understanding the conditions in which they may appear and their mutual interactions has been the subject of many works in the last decades.

Despite the abundance of contributions for positive solutions of (1.1), the literature for sign-changing solutions is less developed. In the present work we take a step in this direction and study sign-changing solutions of (1.1) that blow-up as a single sign-changing bubble. In the sequel, we fix $r_g \in (0, i_g(M))$ where $i_g(M) > 0$ is the injectivity radius of (M, g). Throughout this paper $\chi \in C_c^{\infty}(\mathbb{R}^n)$ will denote a cutoff function such that $\chi(t) = 1$ if $|t| \leq r_g/2$ and $\chi(t) = 0$ if $|t| \geq r_g$. We say that a family $(B_{\varepsilon})_{\varepsilon>0}$ of functions in $H^1(M)$ is a family of bubbles if there exists

Date: April 24th, 2024.

The first author was supported by the FNRS CdR grant J.0135.19, the Fonds Thélam and an ARC Avancé 2020 grant.

a non-zero solution V to

(1.2)
$$\Delta_{\xi} V = |V|^{2^{\star}-2} V \text{ in } \mathbb{R}^n, \quad V \in D^{1,2}(\mathbb{R}^n).$$

where $D^{1,2}(\mathbb{R}^n)$ is the completion of $C_c^{\infty}(\mathbb{R}^n)$ for the norm $u \mapsto \|\nabla u\|_2$, and families of points $(x_{\varepsilon})_{\varepsilon>0}$ in M and of positive real numbers $(\mu_{\varepsilon})_{\varepsilon>0}$ such that

$$B_{\varepsilon}(x) - \chi(d_g(\cdot, x_{\varepsilon}))\mu_{\varepsilon}^{-\frac{n-2}{2}}V\left(\frac{\exp_{x_{\varepsilon}}^{-1}(\cdot)}{\mu_{\varepsilon}}\right) \to 0 \text{ in } H^1(M)$$

as $\varepsilon \to 0$. Here $\exp_{x_{\varepsilon}}$ is the exponential chart at x_{ε} and ξ is the Euclidean metric, so that $\Delta_{\xi} = -\sum_{i=1}^{n} \partial_i^2$. We will say that such a family $(B_{\varepsilon})_{\varepsilon>0}$ is centered at x_{ε} and of radius μ_{ε} and modeled on V, and V itself will sometimes be referred to as the bubble. A more precise definition is given in Definition 3.1 and properties of such families are investigated in Section 3 below. All the solutions considered in this paper are allowed to change sign unless mentioned otherwise.

Our main results establish a necessary condition for the existence of blowing-up solutions of (1.1) that blow-up with a single sign-changing bubble. We introduce one last notation as follows: if $B = (B_{\varepsilon})_{\varepsilon>0}$ is a family of bubbles centered at x_{ε} , of radius μ_{ε} and modeled on V, we define

$$\begin{aligned} \operatorname{Weyl}_{g} \otimes B &:= \frac{4n}{3(n-2)^{2}} \left(\int_{\mathbb{R}^{n}} |V|^{2^{\star}} dx \right)^{-1} \times \\ \operatorname{Weyl}_{g}(x_{0})_{i\alpha j\beta} \int_{\mathbb{R}^{n}} x^{\alpha} x^{\beta} \partial_{ij}^{2} V \left(\frac{n-2}{2} V + x^{l} \partial_{l} V \right) \, dx \end{aligned}$$

where $x_0 = \lim_{\varepsilon \to 0} x_{\varepsilon}$. We first state a result when $n \ge 5$.

Theorem 1.1. Let $(M^n, g), n \ge 5$ be a smooth, connected and closed manifold, let $h, (h_{\varepsilon})_{0 < \varepsilon \le 1} \in C^1(M)$ such that $\lim_{\varepsilon \to 0} h_{\varepsilon} = h$ in $C^1(M)$ and $\Delta_g + h$ is coercive. Let $(p_{\varepsilon})_{0 < \varepsilon \le 1} \in (2, 2^*]$ be such that $\lim_{\varepsilon \to 0} p_{\varepsilon} = 2^*$. Let $(u_{\varepsilon})_{0 < \varepsilon \le 1}$ be a family of solutions of

(1.3)
$$\Delta_g u_{\varepsilon} + h_{\varepsilon} u_{\varepsilon} = |u_{\varepsilon}|^{p_{\varepsilon}-2} u_{\varepsilon} \quad in \ M.$$

We assume that $(u_{\varepsilon})_{0 < \varepsilon < 1}$ satisfies

(1.4)
$$u_{\varepsilon} = cB_{\varepsilon} + o(1) \text{ in } H^{1}(M)$$

for some c > 0, where $B = (B_{\varepsilon})_{\varepsilon > 0}$ is a bubble centered at $(x_{\varepsilon})_{\varepsilon} \in M$ and with radius $(\mu_{\varepsilon})_{\varepsilon} \in (0, +\infty)$ and modeled on $V \in D^{1,2}(\mathbb{R}^n)$ as in Definition 3.1 below.

Then c = 1, $\lim_{\varepsilon \to 0} \mu_{\varepsilon}^{2^{\star} - p_{\varepsilon}} = 1$, the following limit exists:

$$\Lambda := \lim_{\varepsilon \to 0} \frac{2^* - p_\varepsilon}{\mu_\varepsilon^2} \ge 0,$$

and h satisfies:

(1.5)
$$h(x_0) - \frac{n-2}{4(n-1)} S_g(x_0) = \frac{(n-2)^2}{4n} \frac{\int_{\mathbb{R}^n} |V|^{2^*} dx}{\int_{\mathbb{R}^n} V^2 dx} \Big(\Lambda - Weyl_g \otimes B\Big),$$

where we have let $x_0 := \lim_{\varepsilon \to 0} x_{\varepsilon}$.

The new term $\operatorname{Weyl}_g \otimes B$ appearing in (1.5) is well-defined when $n \geq 5$ and we prove in Section 3 that it does not depend on the representative chosen for the family $(B_{\varepsilon})_{\varepsilon>0}$ appearing in (1.4) (see Proposition 3.1 below).

In dimensions 3 and 4 we prove the following analogue of Theorem 1.1:

Theorem 1.2. Let $(M^n, g), n = 3, 4$ be a smooth, connected and closed manifold, let $h, (h_{\varepsilon})_{0 < \varepsilon \leq 1}, (p_{\varepsilon})_{0 < \varepsilon \leq 1} \in (2, 2^*]$ be as in the statement of Theorem 1.1. Let $(u_{\varepsilon})_{0 < \varepsilon \leq 1}$ be a family of solutions of (1.3) satisfying again (1.4) for some c > 0, where $B = (B_{\varepsilon})_{\varepsilon > 0}$ is a bubble centered at $(x_{\varepsilon})_{\varepsilon} \in M$ and with radius $(\mu_{\varepsilon})_{\varepsilon} \in$ $(0, +\infty)$ and modeled on $V \in D^{1,2}(\mathbb{R}^n)$ as in Definition 3.1 below.

Then c = 1, $\lim_{\varepsilon \to 0} \mu_{\varepsilon}^{2^* - p_{\varepsilon}} = 1$, the following limits exist:

$$\Lambda := \lim_{\varepsilon \to 0} \begin{cases} \frac{6 - p_{\varepsilon}}{\mu_{\varepsilon}} & n = 3\\ \frac{4 - p_{\varepsilon}}{\mu_{\varepsilon}^2 \ln \frac{1}{\mu_{\varepsilon}}} & n = 4 \end{cases} \ge 0,$$

and h satisfies

(1.6)
$$\frac{\left(\int_{\mathbb{R}^4} |V|^2 V \, dx\right)^2}{\omega_3 \int_{\mathbb{R}^4} |V|^4 \, dx} \left(h(x_0) - \frac{1}{6} S_g(x_0)\right) = \Lambda \text{ when } n = 4$$

(1.7)
$$-6\frac{\left(\int_{\mathbb{R}^3} |V|^4 V \, dx\right)^2}{\int_{\mathbb{R}^3} |V|^6 \, dx} m_h(x_0) = \Lambda \text{ when } n = 3,$$

where as before we have let $x_0 := \lim_{\varepsilon \to 0} x_{\varepsilon}$. In (1.6) ω_3 is the area of the 3-sphere and in (1.7) $m_h(x_0) \in \mathbb{R}$ is the mass of the Green's function of $\Delta_g + h$ at $x_0 \in M$ (see Definition 3.3 below).

A few comments on Theorems 1.1 and 1.2 are in order. Assumption (1.4) implies that $(u_{\varepsilon})_{\varepsilon>0}$ blows-up as a single, possibly sign-changing bubble. In other words, (1.4) is a Struwe decomposition with a single bubbling profile. No assumption is made on the energy of V (except that it is finite) and V is not assumed to be nondegenerate in the sense of Duyckaerts-Kenig-Merle [13]. Assumption (1.4) should be regarded, insofar as sign-changing solutions are considered, as the analogue of the classical notion of isolated and simple blow-up point for positive solutions introduced in Schoen [35].

Theorems 1.1 and 1.2 are not new for *positive* solutions. It has been known since Obata [26] and Struwe [36] that when $(u_{\varepsilon})_{\varepsilon>0}$ is a family of *positive* solutions of (1.3) the bubble V is positive and radial and given by (2.2) below. In this case, and when $n \ge 5$, the additional term Weyl_g $\otimes B$ in (1.5) vanishes by the symmetries of Weyl_g (see Proposition 3.3 below), and condition (1.5) becomes

$$h(x_0) - \frac{n-2}{4(n-1)} S_g(x_0) = C(n)\Lambda \ge 0, \quad \text{with } \Lambda := \lim_{\varepsilon \to 0} \frac{2^* - p_\varepsilon}{\mu_\varepsilon^2}$$

for some dimensional constant C(n) > 0. When n = 3, 4, (1.6) and (1.7) show that

$$h(x_0) - \frac{1}{6}S_g(x_0) = C(4)\Lambda$$
 when $n = 4$ and $m_h(x_0) = -C(3)\Lambda$ when $n = 3$,

where Λ is as in the statement of the Theorem. When $p_{\varepsilon} = 2^{\star}$ for all ε , and for positive finite-energy solutions, these conditions have been known since the work of Li-Zhu [20] and Druet [10, 11].

If $(u_{\varepsilon})_{\varepsilon>0}$ is a blowing-up family of *sign-changing* solutions of (1.3) with a single bubbling profile as in (1.4), however, the bubble V is sign-changing in general and (1.5), (1.6) and (1.7) display new phenomena. When $n \ge 5$ the additional term $\operatorname{Weyl}_{a} \otimes B$ cannot be expected to vanish *a priori*. We construct indeed in Section 3

examples of manifolds (M, g) and of solutions V of (1.2) such that $\operatorname{Weyl}_g \otimes B \neq 0$ (see Proposition 3.4 below). The term $\operatorname{Weyl}_g \otimes B$ is thus a striking new feature of sign-changing blow-up that does not appear for positive solutions. When $n \geq 5$, condition (1.5) is entirely new and highlights a strong interaction between h, the geometry of (M, g) and the limiting bubble V itself. When n = 3, 4, (1.6) and (1.7) involve $\int_{\mathbb{R}^n} |V|^{2^*-2}V dx$. But if V is a sign-changing solution of (1.2) for $n \geq 3$, the latter integral may vanish if V decays strongly at infinity (see (2.9) below). When this is the case conditions (1.6) and (1.7) indicate that, in dimensions 3 and 4, the interaction between B_{ε} and the geometry of (M, g) takes place at a higher-order than it did for positive solutions.

1.2. Consequences of Theorem 1.1 on the blow-up picture of (1.3). Theorems 1.1 and 1.2 shed new light on the *stability* of the set of sign-changing solutions of (1.3). We say that equation (1.1), with $p = 2^*$, is stable if for every family $(h_{\varepsilon})_{\varepsilon>0}$ converging to h, every energy-bounded family of solutions $(u_{\varepsilon})_{\varepsilon>0}$ of (1.3) converges in $C^2(M)$, up to a subsequence. Druet [10] proved that, for *positive* solutions, stability holds provided $h \neq \frac{n-2}{4(n-1)}S_g$ everywhere in M (with a caveat in dimensions 3 and 6, see [10]). This result was generalised to the case of signchanging solutions, when $n \geq 7$ and when (M, g) is locally conformally flat, in Premoselli-Vétois [31]. When (M, g) is not locally conformally flat, and at least when $n \geq 5$, however, (1.5) shows that necessary conditions for a one bubble blowup involve h, the geometry of (M, g) and the possible limiting bubble V itself. The set of sign-changing solutions of (1.2) is still poorly understood and no classification result is known, but examples of large-energy sequences of solutions have been constructed by Del Pino-Musso-Pacard-Pistoia [7], Ding [9], Medina-Musso [22], Medina-Musso-Wei [23]. The wealth of solutions of (1.2) is the main obstacle to understanding the term $\operatorname{Weyl}_a \otimes B$ in (1.5). It seems therefore highly unlikely, even in the single-bubble case, that stability for sign-changing solutions of (1.3) can be enforced solely by a global assumption on h, as was the case for positive solutions.

Theorems 1.1 and 1.2, however, allow us to rule out limiting bubbles V that may appear in the blow-up in some cases. We start with an example in dimensions 3 and 4. If V is a solution of (1.2) it satisfies

$$V(x) = \frac{\lambda(V)}{|x|^{n-2}} + O(|x|^{1-n}) \quad \text{as } |x| \to +\infty,$$

where $(n-2)\omega_{n-1}\lambda(V) = \int_{\mathbb{R}^n} |V|^{2^*-2}Vdx$ (see (2.4) and (2.9) below). Bubbles decaying at infinity as $|x|^{2-n}$ satisfy $\lambda(V) > 0$: this is the case for the positive bubbles given by (2.2) and for the bubbles of [7]. An immediate corollary of Theorem 1.2 is as follows:

Corollary 1.3. Let (M^n, g) , n = 3, 4, be a smooth, closed, connected Riemannian manifold and $h, (h_{\varepsilon})_{0 < \varepsilon \leq 1} \in C^1(M)$ be such that $\lim_{\varepsilon \to 0} h_{\varepsilon} = h$ in $C^1(M)$ and $\Delta_g + h$ is coercive. Let $(p_{\varepsilon})_{0 < \varepsilon \leq 1} \in (2, 2^*]$ be such that $\lim_{\varepsilon \to 0} p_{\varepsilon} = 2^*$. Let $(u_{\varepsilon})_{0 < \varepsilon \leq 1}$ be a family of solutions of (1.3) satisfying (1.4), where B_{ε} is a family of bubbles centered at x_{ε} , of radius μ_{ε} and modeled on V. Assume that

- the mass of $\Delta_g + h$ is positive at every point of M if n = 3
- $h < \frac{1}{6}S_g$ everywhere in M if n = 4.

Then $\int_{\mathbb{R}^n} |V|^{2^*-2} V dx = 0$. In particular, $V(x) = O(|x|^{1-n})$ as $|x| \to +\infty$ and V is neither positive nor one of the bubbles of [7].

In dimensions $n \ge 5$ (1.5) does not provide any information on the decay of V at infinity, but we can still rule out the existence of symmetric bubbles in some cases. We recall that Ding [9] constructed a family of solutions of (1.2), possessing $O(p) \times O(n-p)$ symmetry for $2 \le p \le n-1$. We show that Ding solutions cannot appear as bubbles in a single-bubble blow-up on symmetric manifolds:

Corollary 1.4. Let $(M^n, g) = (\mathbb{S}^p \times \mathbb{S}^q, g_p \times g_q)$, where g_p and g_q are the round metrics on \mathbb{S}^p and \mathbb{S}^q , $p, q \ge 2$ and $n = p + q \ge 5$. Let $h, (h_{\varepsilon})_{0 < \varepsilon \le 1} \in C^1(M)$ be such that $\lim_{\varepsilon \to 0} h_{\varepsilon} = h$ in $C^1(M)$ and $\Delta_g + h$ is coercive. Let $(p_{\varepsilon})_{0 < \varepsilon \le 1} \in (2, 2^*]$ be such that $\lim_{\varepsilon \to 0} p_{\varepsilon} = 2^*$. Let $(u_{\varepsilon})_{0 < \varepsilon \le 1}$ be a family of solutions of (1.3) satisfying (1.4), where B_{ε} is a family of bubbles centered at x_{ε} , of radius μ_{ε} , modeled on V. Assume that $h \le \frac{n-2}{4(n-1)}S_g$ everywhere. Then

$$Weyl_q \otimes B \geq 0.$$

As a consequence, V cannot be a Ding solution.

Proof. Since $h \leq \frac{n-2}{4(n-1)}S_g$ everywhere, (1.5) shows that $\operatorname{Weyl}_g \otimes B \geq 0$. But as shown in Corollary 3.5 below, $\operatorname{Weyl}_g \otimes B < 0$ if V is a Ding solution. \Box

Corollaries 1.3 (when n = 3) and 1.4 both apply to one-bubble sign-changing blow-up for the sign-changing Yamabe equation:

$$\Delta_g u + \frac{n-2}{4(n-1)} S_g u = |u|^{2^*-2} u.$$

Solutions of this equation naturally appear as minimisers for the second conformal eigenvalue (see Ammann-Humbert [1]), and their stability properties shed new light on the set of possible minimisers (see Premoselli-Vétois [30]). Other stability results for sign-changing solutions are in Premoselli-Vétois [29, 31]. Examples of sign-changing families of solutions of (1.3) have been constructed in Bonheure-Casteras-Premoselli [5], Micheletti-Pistoia-Vétois [25], Pistoia-Vétois [27] and Robert-Vétois [33,34]. Deng-Musso-Wei [8] have considered an innovative construction of a family of sign-changing solutions $(u_{\varepsilon})_{\varepsilon>0}$ to the (1.3) with $h_{\varepsilon} \equiv h$ and $p_{\varepsilon} := 2^* - \varepsilon$ that blows-up like a single bubble as in (1.4). We discuss similar constructions in Section 6.

The structure of the paper is as follows. In Section 2 we introduce a few properties of sign-changing solutions of (1.2) and highlight the main differences with respect to the positive case. Section 3 contains the formal definition and properties of the families of bubbles that we investigate here. We also investigate the term $\text{Weyl}_g \otimes B$ there and construct examples of manifolds and bubbles where $\text{Weyl}_g \otimes B \neq 0$. Section 4 is the core of the analysis of this paper and we prove there that relation (1.4) can be improved into optimal pointwise bounds on u_{ε} . This is the content of Proposition 4.1 and is the main ingredient in the proof of Theorems 1.1 and 1.2. Finally, Theorems 1.1 and 1.2 are proven in Section 5.

Acknowledgements: the authors would like to thank P.-D. Thizy and J. Vétois for many fruitful discussions in the early version of this work.

2. Finite-energy nodal solutions of the Yamabe equation in \mathbb{R}^n

We let Σ be the set of non-zero finite-energy solutions of (1.2):

(2.1) $\Sigma = \{ V \in D^{1,2}(\mathbb{R}^n) \setminus \{0\}, \quad \Delta_{\xi} V = |V|^{2^* - 2} V \}.$

Regularity theory for critical equations (see eg Trudinger [37]) together with standard elliptic theory, shows that there exists $\alpha_n > 0$ such that $V \in C_{loc}^{3,\alpha_n}(\mathbb{R}^n)$ for all $V \in \Sigma$. If $V \in \Sigma$ is positive it is equal by Obata [26], up to translations and rescalings, to

(2.2)
$$B_0^+(x) = \left(1 + \frac{|x|^2}{n(n-2)}\right)^{-\frac{n-2}{2}}, \quad x \in \mathbb{R}^n.$$

By Caffarelli-Gidas-Spruck [6] this result remains true for all positive solutions of (1.2), without the assumption that they belong to $D^{1,2}(\mathbb{R}^n)$. Note also that, up to dilation, B_0^+ is the only non-zero *radial* solution of (1.2) which belongs to $D^{1,2}(\mathbb{R}^n)$ by a simple application of Pohozaev's identity.

If $V \in D^{1,2}(\mathbb{R}^n)$ we define its Kelvin transform as

(2.3)
$$V^*(x) = \frac{1}{|x|^{n-2}} V\left(\frac{x}{|x|^2}\right) \text{ for a.e. } x \in \mathbb{R}^n \setminus \{0\}.$$

It is well-known (see for instance Duyckaerts-Kenig-Merle [13, Proposition 3.1]) that $V \mapsto V^*$ defines an isometry of $D^{1,2}(\mathbb{R}^n)$ and of $L^{2^*}(\mathbb{R}^n)$ and that

$$\Delta_{\xi} V^*(x) = \frac{1}{|x|^{n+2}} \Delta_{\xi} V\left(\frac{x}{|x|^2}\right)$$

in $\mathbb{R}^n \setminus \{0\}$, provided V is of class C^2 . As a consequence we have $V^* \in \Sigma$ whenever $V \in \Sigma$ (see again [13]) and the regularity theory for (1.2) shows that $V^* \in C^3(\mathbb{R}^n)$ when $V \in \Sigma$. The following result provides a precise description of the behavior at infinity of the elements in Σ :

Lemma 2.1. Let $V \in \Sigma$. There exist $\lambda(V) \in \mathbb{R}$ and $\alpha(V) \in \mathbb{R}^n$ such that the following asymptotic expansion holds as $|x| \to +\infty$ and can be differentiated:

(2.4)
$$V(x) = \frac{\lambda(V)}{|x|^{n-2}} + \frac{\langle \alpha(V), x \rangle}{|x|^n} + O\left(\frac{1}{|x|^n}\right).$$

When we say that expansion (2.4) can be differentiated we mean that the following holds as $|x| \to +\infty$: for any $1 \le i \le n$,

(2.5)
$$\partial_i V(x) = -(n-2)\lambda(V)\frac{x_i}{|x|^n} + \frac{\alpha(V)_i |x|^2 - n\langle \alpha(V), x \rangle x_i}{|x|^{n+2}} + O\left(\frac{1}{|x|^{n+1}}\right).$$

Proof. Let V^* be, as before, the Kelvin transform of V. It is of class C^3 in \mathbb{R}^n and we can consider its Taylor expansion at 0: there exist $\lambda(V) \in \mathbb{R}$ and $\alpha(V) \in \mathbb{R}^n$ such that, as $y \to 0$, and for $1 \le i \le n$,

(2.6)
$$V^*(y) = \lambda(V) + \langle \alpha(V), y \rangle + O(|y|^2)$$
$$\partial_i V^*(y) = \alpha(V)_i + O(|y|)$$

hold. Since for any $x \neq 0$ we have $V(x) = \frac{1}{|x|^{n-2}} V^*\left(\frac{x}{|x|^2}\right)$, (2.4) follows from (2.6). To prove (2.5) we write that by the chain rule, for $x \neq 0$,

$$(2.7) \ \partial_i V(x) = -(n-2)\frac{x_i}{|x|^n} V^*\left(\frac{x}{|x|^2}\right) + \sum_{j=1}^n \frac{1}{|x|^{n-2}} \left(\frac{\delta_{ij}}{|x|^2} - 2\frac{x_i x_j}{|x|^4}\right) \partial_j V^*\left(\frac{x}{|x|^2}\right)$$

holds, and we conclude using again (2.6).

As a consequence of (2.4) and (2.5), there exists C = C(n, V) such that

$$(2.8) |V(x)| + (1+|x|)|\nabla V(x)| + (1+|x|)^2|\nabla^2 V(x)| \le C(1+|x|)^{2-n}$$

for all $x \in \mathbb{R}^n$. This can also be shown by the conformal invariance of (1.2), see e.g. Premoselli [28, Lemma 2.2]. As shown in Premoselli [28, Lemma 2.1], we have

(2.9)
$$\lambda(V) = \frac{1}{(n-2)\omega_{n-1}} \int_{\mathbb{R}^n} |V|^{2^*-2} V dx \quad \text{for all } V \in \Sigma.$$

If B_0^+ is as in (2.2) we have $\lambda(B_0^+) = (n(n-2))^{\frac{n-2}{2}}$. The solutions constructed in [7] also satisfy $\lambda(V) \neq 0$. As proven in Lemma 2.1, $\lambda(V) = V^*(0)$. As a consequence, examples where $\lambda(V) = 0$ are easily obtained by Kelvin-transforming sign-changing solutions of Σ at a point where they vanish.

Remark 2.2. An accurate inspection of the proof of Lemma 2.1 shows that we can state a slightly more precise result. Let $V \in \Sigma$.

• Assume that n = 3, 4. Then there exists $k \in \mathbb{N}$ and a non-zero homogeneous polynomial P of degree k such that

$$V(x) = \frac{P(x)}{|x|^{n-2+2k}} + O\left(\frac{1}{|x|^{n-1+k}}\right) \quad \text{as } |x| \to +\infty.$$

• Assume that n = 5. Then there exists $k \in \{0, ..., 4\}$ and a homogeneous polynomial P of degree k such that

$$V(x) = \frac{P(x)}{|x|^{3+2k}} + \begin{cases} O\left(\frac{1}{|x|^{4+k}}\right) & \text{if } k \le 3\\ O\left(\frac{1}{|x|^{7+\frac{1}{3}}}\right) & \text{if } k = 4 \end{cases} \quad \text{as } |x| \to +\infty.$$

• Assume finally that $n \ge 6$. Let $0 < \alpha < 1$ if n = 6 and $0 < \alpha \le \frac{4}{n-2}$ if $n \ge 7$. Then there exists $k \in \{0, \ldots, 3\}$ and a homogeneous polynomial P of degree k such that V satisfies

$$V(x) = \frac{P(x)}{|x|^{n-2+2k}} + O\left(\frac{1}{|x|^{n-2+k+\alpha}}\right) \quad \text{as } |x| \to +\infty.$$

All these expansions can be differentiated. In dimensions n = 3, 4, P is the first non-zero homogeneous polynomial in the Taylor expansion of V^* at 0, which always exists by classical finite continuation results (see e.g. Aronszajn [2]) since $V \in D^{1,2}(\mathbb{R}^n) \setminus \{0\}$. In dimensions $n \geq 5$, P may vanish. Examples of solutions in any dimension $n \geq 3$ for which the degree of P is larger or equal than 2 are still unknown and the question of their existence was raised in Duyckaerts-Kenig-Merle[13].

3. RIEMANNIAN PRELIMINARIES

3.1. Sign-changing families of bubbles in M. We define the notion of Riemannian bubble that we will consider in this work:

Definition 3.1. Let $(x_{\varepsilon})_{\varepsilon>0} \in M$ and $(\mu_{\varepsilon})_{\varepsilon>0} \in \mathbb{R}_{>0}$ be families of points in Mand positive real numbers, with $\mu_{\varepsilon} \to 0$ as $\varepsilon \to 0$. We let $x_0 := \lim_{\varepsilon \to 0} x_{\varepsilon}$. Let $V \in \Sigma$ be fixed and let $B = (B_{\varepsilon})_{\varepsilon>0}$ be a family of functions in $H^1(M)$. We say that B is a family of bubbles centered at x_{ε} with radius μ_{ε} and modeled on V if there exists a family of charts $(\varphi_{\varepsilon})_{\varepsilon \geq 0}$ such that $\varphi_{\varepsilon} : B_g(x_0, r_g) \to \mathbb{R}^n$ is normal at x_{ε} for all $\varepsilon > 0$, $\lim_{\varepsilon \to 0} \varphi_{\varepsilon} = \varphi_0$ in C^k for all k and such that

(3.1)
$$B_{\varepsilon} - \chi(d_g(\cdot, x_{\varepsilon}))\mu_{\varepsilon}^{-\frac{n-2}{2}}V\left(\frac{\varphi_{\varepsilon}(\cdot) - \varphi_{\varepsilon}(x_{\varepsilon})}{\mu_{\varepsilon}}\right) \to 0 \quad in \ H^1(M)$$

as $\varepsilon \to 0$.

Here Σ is as in (2.1). As the following simple result shows, Definition 3.1 is independent of the choice of the local chart up to an isometry. In the sequel, we let O(n) be the group of isometries of the Euclidean space \mathbb{R}^n :

Proposition 3.1. Let $(x_{\varepsilon})_{\varepsilon>0} \in M$ and $(\mu_{\varepsilon})_{\varepsilon>0} \in \mathbb{R}_{>0}$ be families of points in Mand positive real numbers, with $\mu_{\varepsilon} \to 0$ as $\varepsilon \to 0$, and let $B = (B_{\varepsilon})_{\varepsilon}$ and $\overline{B} = (\overline{B}_{\varepsilon})_{\varepsilon}$ be families of bubbles centered at x_{ε} with radius μ_{ε} and respectively modeled on $V, \overline{V} \in \Sigma$ as in Definition (3.1). Then

$$B_{\varepsilon} - \bar{B}_{\varepsilon} \to 0 \text{ in } H^1(M) \text{ as } \varepsilon \to 0 \quad \iff \quad \bar{V} = V \circ O$$

for some orthogonal mapping $O \in O(n)$.

Families of bubbles $(B_{\varepsilon})_{\varepsilon>0}$ and $(\bar{B}_{\varepsilon})_{\varepsilon>0}$ centered at x_{ε} with radius μ_{ε} and satisfying $B_{\varepsilon} - \bar{B}_{\varepsilon} \to 0$ in $H^1(M)$ as $\varepsilon \to 0$ will be called *equivalent* families.

Proof. Let $\varphi_{\varepsilon}, \bar{\varphi}_{\varepsilon} : B_g(x_0, r_g) \to \mathbb{R}^n$ be normal coordinate charts for g at x_{ε} associated to B and \bar{B} as in definition 3.1. If $x \in \mathbb{R}^n$ we have

(3.2)
$$\varphi_{\varepsilon} \circ \bar{\varphi}_{\varepsilon}^{-1}(\bar{\varphi}_{\varepsilon}(x_{\varepsilon}) + x) = \varphi_{\varepsilon}(x_{\varepsilon}) + O_{\varepsilon}x + O(|x|^2) \quad \text{as } x \to 0,$$

where we have let $O_{\varepsilon} = D\varphi_{\varepsilon}(x_{\varepsilon}) \circ (D\bar{\varphi}_{\varepsilon}(x_{\varepsilon}))^{-1}$. Since φ_{ε} and $\bar{\varphi}_{\varepsilon}$ are normal charts at x_{ε} , $O_{\varepsilon} \in O(n)$. Straightforward computations using (2.8) and (3.2) show that (3.3)

$$\chi(d_g(\cdot, x_{\varepsilon})) \left[\mu_{\varepsilon}^{-\frac{n-2}{2}} V\left(\frac{\varphi_{\varepsilon}(\cdot) - \varphi_{\varepsilon}(x_{\varepsilon})}{\mu_{\varepsilon}}\right) - \mu_{\varepsilon}^{-\frac{n-2}{2}} V\left(\frac{O_{\varepsilon}\left(\bar{\varphi}_{\varepsilon}(\cdot) - \bar{\varphi}_{\varepsilon}(x_{\varepsilon})\right)}{\mu_{\varepsilon}}\right) \right] \to 0$$

in $H^1(M)$ as $\varepsilon \to 0$. Since $B_{\varepsilon} - \bar{B}_{\varepsilon} \to 0$ in $H^1(M)$ by assumption we obtain

$$\chi(d_g(\cdot, x_{\varepsilon})) \left[\mu_{\varepsilon}^{-\frac{n-2}{2}} V\left(\frac{O_{\varepsilon}\left(\bar{\varphi}_{\varepsilon}(\cdot) - \bar{\varphi}_{\varepsilon}(x_{\varepsilon})\right)}{\mu_{\varepsilon}}\right) - \mu_{\varepsilon}^{-\frac{n-2}{2}} \bar{V}\left(\frac{\bar{\varphi}_{\varepsilon}(\cdot) - \bar{\varphi}_{\varepsilon}(x_{\varepsilon})}{\mu_{\varepsilon}}\right) - \right] \to 0$$

in $H^1(M)$ as $\varepsilon \to 0$. A simple scaling argument then shows that $V \circ O = \overline{V}$ where $O := \lim_{\varepsilon \to 0} O_{\varepsilon}$.

Equations (3.2) and (3.3) show that, up to composing V with an euclidean isometry, we can always assume that φ_{ε} is an exponential chart at x_{ε} in Definition 3.1. Choose, for any $p \in B_g(x_0, r_g)$, an identification of T_pM with \mathbb{R}^n . This defines an exponential map at p, that we will denote by $\exp_p : \mathbb{R}^n \to M$ in all of this paper. Mimicking the arguments in (3.2) and (3.3) we similarly have, for $V \in \Sigma$,

$$\chi(d_g(\cdot, x_{\varepsilon})) \left[\mu_{\varepsilon}^{-\frac{n-2}{2}} V\left(\frac{\varphi_{\varepsilon}(\cdot) - \varphi_{\varepsilon}(x_{\varepsilon})}{\mu_{\varepsilon}}\right) - \mu_{\varepsilon}^{-\frac{n-2}{2}} \left(V \circ O \right) \left(\frac{\exp_{x_{\varepsilon}}^{-1}(\cdot)}{\mu_{\varepsilon}}\right) \right] \to 0$$

for some $O \in O(n)$. We have thus shown that $B = (B_{\varepsilon})_{\varepsilon>0}$ is a family of bubbles centered at x_{ε} with radius μ_{ε} and modeled on $V \in \Sigma$ if and only if

(3.4)
$$B_{\varepsilon} - \chi(d_g(\cdot, x_{\varepsilon}))\mu_{\varepsilon}^{-\frac{n-2}{2}} \left(V \circ O \right) \left(\frac{\exp_{x_{\varepsilon}}^{-1}(\cdot)}{\mu_{\varepsilon}} \right) \to 0 \quad \text{in } H^1(M)$$

as $\varepsilon \to 0$, for some $O \in O(n)$. Any two exponential charts at x_{ε} are related by an isometry of \mathbb{R}^n . A consequence of (3.4) is that Definition 3.1 is independent of the choice of the exponential chart at x_{ε} up to composing V with an isometry. The classical example of a family of bubbles is the so-called family of standard bubbles:

(3.5)
$$B_{\varepsilon}(x) := \chi(d_g(x, x_{\varepsilon}))B_{\varepsilon}^+(x), \text{ where}$$
$$B_{\varepsilon}^+(x) := \left(\frac{\mu_{\varepsilon}}{\mu_{\varepsilon}^2 + \frac{d_g(x, x_{\varepsilon})^2}{n(n-2)}}\right)^{\frac{n-2}{2}} = \mu_{\varepsilon}^{-\frac{n-2}{2}}B_0^+\left(\frac{\exp_{x_{\varepsilon}}^{-1}(x)}{\mu_{\varepsilon}}\right),$$

where B_0^+ is given by (2.2). This example is peculiar: since B_0^+ is radially symmetrical (as a consequence of positivity), any exponential chart at x_{ε} yields the same family of bubbles, and the function B_{ε}^+ thus obtained is well-defined on the whole of M. Naively composing with an exponential chart does not, however, allow to globally pull-back a sign-changing element $V \in \Sigma$ in M without losing regularity at the cut-locus, hence the need for Definition 3.1.

The following proposition is straightforward:

Proposition 3.2. Let $(B_{\varepsilon})_{\varepsilon>0}$ be a family of bubble as in Definition 3.1. Then

- There exists C > 0 such that $||B_{\varepsilon}||_{H^1} \leq C$ for all $\varepsilon > 0$
- $\lim_{\varepsilon \to 0} \|B_{\varepsilon}\|_{L^2} = 0$
- $B_{\varepsilon} \to 0$ weakly in $H^1(M)$ as $\varepsilon \to 0$ $\lim_{\varepsilon \to 0} \|\nabla B_{\varepsilon}\|_2^2 = \int_{\mathbb{R}^n} |\nabla V|^2 dx.$

3.2. The product $\operatorname{Weyl}_g \otimes B$. We now investigate the properties of the term $\operatorname{Weyl}_q \otimes B$ appearing in the Statement of Theorem 1.1.

Definition 3.2. Assume that $n \geq 5$. Let $B = (B_{\varepsilon})_{\varepsilon>0}$ be a family of bubbles centered at x_{ε} with radius μ_{ε} and modeled on $V \in \Sigma$ as in Definition 3.1. We define

(3.6)
$$Weyl_g \otimes B := \frac{4n}{3(n-2)^2} \left(\int_{\mathbb{R}^n} |V|^{2^*} dx \right)^{-1} \times Weyl_g(x_0)_{i\alpha j\beta} \int_{\mathbb{R}^n} x^{\alpha} x^{\beta} \partial_{ij}^2 V\left(\frac{n-2}{2}V + x^l \partial_l V\right) dx$$

where $x_0 = \lim_{\varepsilon \to 0} x_{\varepsilon}$ and where the coordinates of the Weyl tensor are taken with respect to the chart \exp_{x_0} .

It is intended in (3.6) that repeated indices are summed over. That the second integral in (3.6) is finite follows from (2.8) since $n \ge 5$. Note that the term $\operatorname{Weyl}_q \otimes B$ depends on x_0 since the family $B = (B_{\varepsilon})_{\varepsilon} > 0$ does. We recall the symmetries of the Weyl tensor of g at any point $x \in M$:

$$\begin{aligned} \operatorname{Weyl}_g(x)_{ijk\ell} &= -\operatorname{Weyl}_g(x)_{jik\ell} = -\operatorname{Weyl}_g(x)_{ij\ell k} = \operatorname{Weyl}_g(x)_{k\ell ij} \\ \operatorname{Weyl}_g(x)_{iji\ell} &= 0 \text{ and } \operatorname{Weyl}_g(x)_{ijk\ell} + \operatorname{Weyl}_g(x)_{jki\ell} + \operatorname{Weyl}_g(x)_{kij\ell} = 0 \end{aligned}$$

for any $i, j, k, \ell \in \{1, ..., n\}$, where repeated indices are summed over. As a consequence of these symmetries, simple integration by parts (see for instance Hebey-Vaugon [17] or Lemma 3.2 in Mesmar-Robert [24]) show that

(3.7)

$$Weyl_g \otimes B = \frac{4n}{3(n-2)^2} \left(\int_{\mathbb{R}^n} |V|^{2^*} dx \right)^{-1} \times Weyl_g(x_0)_{i\alpha j\beta} \int_{\mathbb{R}^n} x^\alpha x^\beta \partial_i V \partial_j V dx.$$

Again the integral on the right is finite when $n \geq 5$ by (2.8). A consequence of (3.7) and of the tensorial nature of $\operatorname{Weyl}_g(x_0)$ is that $\operatorname{Weyl}_g \otimes B$ does not change if V is replaced by $V \circ O$ for some $O \in O(n)$, provided the coordinates of $\operatorname{Weyl}_g(x_0)$ are now computed in the chart $O \circ \exp_{x_0}^{-1}$. Hence $\operatorname{Weyl}_g \otimes B$ does not depend on the choice of the exponential chart at x_0 and only depends on the equivalence class of families of bubbles at x_0 in the sense of Proposition 3.1.

We describe two simple cases where $\operatorname{Weyl}_q \otimes B$ vanishes globally:

Proposition 3.3. Assume that $n \geq 5$ and let B be a family of bubbles centered at $(x_{\varepsilon})_{\varepsilon} \in M$, with radius $(\mu_{\varepsilon})_{\varepsilon} \in \mathbb{R}_{>0}$ and modeled on $V \in \Sigma$, and $x_0 = \lim_{\varepsilon \to 0} x_{\varepsilon}$. Then $Weyl_a \otimes B = 0$ in the following (non-exhaustive) situations:

- $Weyl_a(x_0) = 0$
- V is radial (and hence $V = B_0^+$ given by (2.2)).

Proof. By (3.6) the first point is trivial. If $V = B_0^+$ is radial, the second point follows from the radiality of B_0^+ , from the equivalent expression (3.7) and from the total antisymmetry of the Weyl tensor.

The next two results shows that $\operatorname{Weyl}_g\otimes B$ cannot be expected to vanish in general:

Proposition 3.4. Let (\mathbb{S}^p, g_p) and (\mathbb{S}^q, g_q) be the unit spheres of dimensions $p, q \geq 2$ endowed with their round metrics. Assume that $n = p + q \geq 5$ and that $V \in D^{1,2}(\mathbb{R}^n) \setminus \{0\}$ solves (1.2) and is invariant under the action of $O(p) \times O(q)$ but not under the action of O(n). Let $M = \mathbb{S}^p \times \mathbb{S}^q$ endowed with the metric $g = g_p \otimes g_q$. Let $B = (B_{\varepsilon})_{\varepsilon>0}$ be a family of bubbles in (M,g) modeled on V as in Definition 3.1, centered at $(x_{\varepsilon})_{\varepsilon>0} \in M$ and with radius $(\mu_{\varepsilon})_{\varepsilon>0} \in \mathbb{R}_{>0}$ and $x_0 = \lim_{\varepsilon \to 0} x_{\varepsilon}$. Then $Weyl_q \otimes B < 0$.

Proof. We let $M := \mathbb{S}^p \times \mathbb{S}^q$ where $p, q \geq 2$ and $p + q = n \geq 5$ and $g := g_p \otimes g_q$ be the product metric of the round metrics g_p and g_q respectively on \mathbb{S}^p and \mathbb{S}^q . We follow the geometric notations of Besse [4] and Hebey [15]. The Riemann tensors for each metric are

$$\operatorname{Rm}_{g_p} = \frac{1}{2}g_p \odot g_p$$
 and $\operatorname{Rm}_{g_q} = \frac{1}{2}g_q \odot g_q$,

where \odot is the Kulkarni-Nomizu product, given in coordinates by $(T \odot S)_{ijkl} = T_{ik}S_{jl} - T_{il}S_{jk} + T_{jl}S_{ik} - T_{jk}S_{il}$. Standard results on product manifolds give that, after the necessary projections, $\operatorname{Rm}_g = \frac{1}{2}g_p \odot g_p + \frac{1}{2}g_q \odot g_q$, $\operatorname{Ric}_g = (p-1)g_p + (q-1)g_q$ and $S_g = p(p-1) + q(q-1)$. The Weyl tensor of g is defined as

$$\operatorname{Weyl}_{g} = \operatorname{Rm}_{g} - \frac{1}{n-2}\operatorname{Ric}_{g} \odot g + \frac{S_{g}}{2(n-1)(n-2)}g \odot g.$$

If $x_0 = (x_0^p, x_0^q) \in \mathbb{S}^p \times \mathbb{S}^q$ we compute the coordinates of $\operatorname{Weyl}_g(x_0)$ in the chart obtained from the exponential chart on \mathbb{S}^p at x_0^p and the exponential chart on \mathbb{S}^q at x_0^q : this yields a normal chart on $\mathbb{S}^p \times \mathbb{S}^q$. The Latin letters i, j, k... refer to coordinates in \mathbb{S}^p , the Greek letter $\alpha, \beta, ...$ refer to \mathbb{S}^q . The previous observations show that, at any point around x_0 we have

$$(Weyl_g)_{ijkl} = C_1(g_{ik}g_{jl} - g_{il}g_{jk}) (Weyl_g)_{\alpha\beta\gamma\delta} = C_2(g_{\alpha\gamma}g_{\beta\delta} - g_{\alpha\delta}g_{\beta\gamma}) (Weyl_g)_{i\alphaj\beta} = -C_3g_{ij}g_{\alpha\beta} (Weyl_g)_{ij\alpha\beta} = 0,$$

with $C_1 = \frac{2q(q-1)}{(n-1)(n-2)}, C_2 = \frac{2p(p-1)}{(n-1)(n-2)}$ and $C_3 = \frac{2(p-1)(q-1)}{(n-1)(n-2)}$. The mixed terms $(\operatorname{Weyl}_g)_{ijk\alpha}$ and $(\operatorname{Weyl}_g)_{i\alpha\beta\gamma}$ are null, and the other terms are obtained via the symmetries of the Weyl tensor. We denote the coordinates in \mathbb{R}^n by (X, Y), where $X \in \mathbb{R}^p$ and $Y \in \mathbb{R}^q$. Let $V \in \Sigma$ be $O(p) \times O(q)$ invariant and let $C_{n,V} = \frac{4n}{3(n-2)^2} \left(\int_{\mathbb{R}^n} |V|^{2^*} dx \right)^{-1}$. Using (3.7) and the explicit expression of $\operatorname{Weyl}_g(x_0)$ we have

$$\begin{split} C_{n,V}^{-1} \mathrm{Weyl}_g \otimes B &= C_1 \int_{\mathbb{R}^n} \left[|X|^2 |\nabla_1 V|^2 - (X, \nabla_1 V)^2 \right] \, dX dY \\ &+ C_2 \int_{\mathbb{R}^n} \left[|Y|^2 |\nabla_2 V|^2 - (Y, \nabla_2 V)^2 \right] \, dX dY \\ &- C_3 \int_{\mathbb{R}^n} \left[|X|^2 |\nabla_2 V|^2 + |Y|^2 |\nabla_1 V|^2 - 2(X, \nabla_1 V)(Y, \nabla_2 V) \right] dX \, dY \end{split}$$

where $\nabla_1 V$ and $\nabla_2 V$ denote respectively the gradient of V with respect to $X \in \mathbb{R}^p$ and $Y \in \mathbb{R}^q$. The $O(p) \times O(q)$ invariance of V implies that V(X,Y) = V(|X|,|Y|), and as a consequence we have $\nabla_1 V(X,Y) = \partial_1 V(X,Y) \frac{X}{|X|}$ and $\nabla_2 V(X,Y) = \partial_2 V(X,Y) \frac{Y}{|Y|}$ where $\partial_1 V$ and $\partial_2 V$ are the partial derivatives of $(r_1, r_2) \mapsto V(r_1, r_2)$. We obtain in the end

$$\begin{aligned} C_{n,V}^{-1} \operatorname{Weyl}_{g} \otimes B &= -C_{3} \int_{\mathbb{R}^{n}} \left[|X|^{2} (\partial_{2}V)^{2} + |Y|^{2} (\partial_{1}V)^{2} - 2|X| |Y| \partial_{1}V \partial_{2}V \right] dX \, dY \\ &= -C_{3} \int_{\mathbb{R}^{n}} \left(|X| \partial_{2}V - |Y| \partial_{1}V \right)^{2} dX \, dY \leq 0. \end{aligned}$$

If $\operatorname{Weyl}_g \otimes B = 0$, then $r_2 \partial_1 V - r_1 \partial_2 V$ for all $r_1 = |X| > 0$ and $r_2 = |Y| > 0$. This shows that V = V(r), for $r^2 = r_1^2 + r_2^2$ when $r_1, r_2 > 0$, and this equality extends continuously to $\mathbb{R}^n \setminus \{0\}$. We then get that V is radially symmetrical, contradicting our assumption. Therefore $\operatorname{Weyl}_g \otimes B < 0$.

Corollary 3.5. There are manifolds (M, g) of dimension $n \ge 5$ and families of bubbles $B = (B_{\varepsilon})_{\varepsilon>0}$ as in Definition 3.1 such that $Weyl_q \otimes B < 0$.

Proof. We choose $p, q \geq 2$ such that $n := p + q \geq 5$. As before we denote by g_n the round metric in \mathbb{S}^n . It follows from the result of Ding [9] that for all L > 0, there exists $\tilde{V} \in C^2(\mathbb{S}^n)$ that solves $\Delta_{g_n}\tilde{V} + \frac{n(n-2)}{4}\tilde{V} = |\tilde{V}|^{2^*-2}\tilde{V}$, satisfies $\int_{\mathbb{S}^n} |\tilde{V}|^{2^*} dv_{g_n} \geq L$ and is $O(p) \times O(q+1)$ -invariant, that is $\tilde{V}(x,y) = \tilde{V}(|x|,|y|)$ for $(x,y) \in \mathbb{R}^p \times \mathbb{R}^{q+1} = \mathbb{R}^{n+1}$. Let $\pi_N : \mathbb{S}^n \setminus \{N\} \to \mathbb{R}^n$ be the stereographic

projection from the North pole. Let, for $x \in \mathbb{R}^n$,

$$V(x) := \left(\frac{2}{1+|x|^2}\right)^{\frac{n-2}{2}} \tilde{V}\big(\pi_N^{-1}(x)\big).$$

Then $V \in C^2(\mathbb{R}^n) \cap D^{1,2}(\mathbb{R}^n)$ solves $\Delta_{\xi} V = |V|^{2^*-2} V$ in \mathbb{R}^n , satisfies $\int_{\mathbb{R}^n} |V|^{2^*} dx \ge L$ and using the coordinate expression of the stereographic projection it is easily seen that it is $O(p) \times O(q)$ -invariant, that is V(x', x'') = V(|x'|, |x''|) for $(x', x'') \in \mathbb{R}^p \times \mathbb{R}^q = \mathbb{R}^n$. Up to taking L large enough, we get that V is not O(n)-invariant since by Obata [26] finite-energy radial solutions of (1.2) are translations and rescalings of B_0^+ given by (2.2) and all have fixed minimal Dirichlet energy. Let now $M := \mathbb{S}^p \times \mathbb{S}^q$ and $g := g_p \otimes g_q$ be the product metric of the round metrics on \mathbb{S}^p and \mathbb{S}^q . Let $B = (B_{\varepsilon})_{\varepsilon>0}$ be any family of bubbles in (M, g) modeled on V as in Definition 3.1, centered at $(x_{\varepsilon})_{\varepsilon} \in M$ and with radius $(\mu_{\varepsilon})_{\varepsilon} \in \mathbb{R}_{>0}$ and $x_0 = \lim_{\varepsilon \to 0} x_{\varepsilon}$, where $x_0 \in M$ is any point. For L large enough Proposition 3.4 applies and shows that Weyl_q $\otimes B < 0$.

3.3. The mass of the Green's function in 3D. We conclude this section by recalling the definition of the mass when n = 3, following Li-Zhu [20]:

Definition 3.3. Let (M,g) be a smooth, closed, connected 3-dimensional Riemannian manifold. Let $h \in C^1(M)$ be such that $\Delta_g + h$ is coercive. Let G_h be the Green's function for $\Delta_g + h$. Then for all $x_0 \in M$, there exists $m_h(x_0) \in \mathbb{R}$, denoted as the mass, such that

$$G_h(x, x_0) = \frac{1}{4\pi d_g(x, x_0)} + m_h(x_0) + o(1) \text{ as } x \to x_0.$$

When n = 3, $\Delta_g + h$ is coercive and $x_0 \in M$, It follows from the construction of the Green's function (see Aubin [3], Robert [32]) that there exists $\beta_{x_0} \in C^0(M) \cap$ $H_2^p(M)$ for all p < 3 such that

(3.8)
$$G_h(x, x_0) = \frac{\chi(x)}{4\pi d_g(x, x_0)} + \beta_{x_0}(x) \text{ for all } x \in M \setminus \{x_0\},$$

where $\chi \in C^{\infty}(M)$ is a cutoff function with $\chi(x) = 1$ in $B_g(x_0, \frac{r_g}{2})$ and $\chi(x) = 0$ in $M \setminus B_g(x_0, r_g)$. We have $(\Delta_g + h)\beta_{x_0} = -(\Delta_g + h)((4\pi)^{-1}\chi d_g(\cdot, x_0)^{-1}) = O(d_g(\cdot, x_0)^{-1})$ and standard elliptic theory then shows that (3.9)

$$|\beta_{x_0}(x)| \le C, |\nabla\beta_{x_0}(x)| \le C\left(1 + |\ln d_g(x_0, x)|\right) \text{ for all } x \in B_g\left(x_0, \frac{r_g}{2}\right) \setminus \{0\}.$$

It now follows from Definition 3.3 that

$$m_h(x_0) := \beta_{x_0}(x_0).$$

4. A priori pointwise estimates for families of bubbles

4.1. Statement of the results. In this section we will assume, as in the introduction, that $(u_{\varepsilon})_{0 < \varepsilon \leq 1}$ is a family of possibly sign-changing solutions of (1.3) satisfying (1.4):

$$u_{\varepsilon} = cB_{\varepsilon} + o(1)$$
 in $H^1(M)$

for some c > 0, where B_{ε} is a family of bubbles as in Definition 3.1, centered at x_{ε} with radius μ_{ε} . Here $(x_{\varepsilon})_{0 < \varepsilon \leq 1}$ and $(\mu_{\varepsilon})_{0 < \varepsilon \leq 1}$ are families of points in M and of positive numbers with $\mu_{\varepsilon} \to 0$ as $\varepsilon \to 0$. We let $V \in \Sigma$, where Σ is as in (2.1), be

the function that appears in (3.4). Under this assumption $(u_{\varepsilon})_{0<\varepsilon\leq 1}$ blows-up with only one possibly sign-changing bubble and a zero weak limit. Throughout this paper we will adopt the following conventions: for nonnegative families $(a_{\varepsilon})_{0<\varepsilon\leq 1}$ and $(b_{\varepsilon})_{0<\varepsilon\leq 1}$ of functions in M (or real numbers), we will write $a_{\varepsilon} \leq b_{\varepsilon}$, or equivalently $a_{\varepsilon} = O(b_{\varepsilon})$, when there exists a positive constant C such that

$$a_{\varepsilon}(x) \leq Cb_{\varepsilon}(x)$$
 for all $0 < \varepsilon \leq 1$ and any $x \in M$.

Since $\Delta_g + h_{\varepsilon}$ is coercive we let, for any $0 < \varepsilon \leq 1$, $G_{h_{\varepsilon}}$ be the positive Green's function of $\Delta_g + h_{\varepsilon}$ (see Robert [32]). Since $h_{\varepsilon} \to h$ in $C^1(M)$ as $\varepsilon \to 0$ and $\Delta_g + h$ is coercive there exists C > 0 such that

(4.1)
$$\left| d_g(x,y)^{n-2} G_{h_{\varepsilon}}(x,y) - \frac{1}{(n-2)\omega_{n-1}} \right| \le C d_g(x,y)$$

holds true for any $(x, y) \in M \times M \setminus D(M)$ and for any $0 < \varepsilon \leq 1$, where $D(M) = \{(x, x), x \in M\}$ and where ω_{n-1} is the area of the standard sphere \mathbb{S}^{n-1} . Define, for $x \in M$,

$$F_{\varepsilon}(x_{\varepsilon}, x) = (n-2)\omega_{n-1}d_g(x_{\varepsilon}, x)^{n-2}G_{h_{\varepsilon}}(x_{\varepsilon}, x)$$

We define, for any $x \in M$ and any $0 < \varepsilon \leq 1$:

(4.2)
$$B^{\mu_{\varepsilon},x_{\varepsilon}}(x) = \chi \left(d_g(x_{\varepsilon},x) \right) F_{\varepsilon}(x_{\varepsilon},x) \mu_{\varepsilon}^{-\frac{n-2}{2}} V\left(\frac{1}{\mu_{\varepsilon}} \exp_{x_{\varepsilon}}^{-1}(x)\right) \\ + \left(1 - \chi \left(d_g(x_{\varepsilon},x) \right) \right) (n-2) \omega_{n-1} \lambda(V) \mu_{\varepsilon}^{\frac{n-2}{2}} G_{h_{\varepsilon}}(x_{\varepsilon},x),$$

where $\lambda(V)$ is as in (2.9) and for $\chi \in C_c^{\infty}(\mathbb{R})$ with $\chi \equiv 1$ in $[0, \frac{i_g(M)}{4}]$ and $\chi \equiv 0$ in $[\frac{i_g(M)}{2}, +\infty)$, and where $i_g(M)$ is the injectivity radius of (M, g). By (4.1) and properties of the Green's function (see [32]) we have

$$F_{\varepsilon}(x_{\varepsilon}, x) = 1 + O(d_g(x_{\varepsilon}, x))$$
 and $|\nabla_x F_{\varepsilon}(x_{\varepsilon}, x)| = O(1)$ for $x \in M$.

It is therefore easily seen with (3.4) that

$$B^{\mu_{\varepsilon}, x_{\varepsilon}} = B_{\varepsilon} + o(1) \quad \text{in } H^1(M),$$

so that $(B^{\mu_{\varepsilon},x_{\varepsilon}})_{0<\varepsilon\leq 1}$ is a family of bubbles which is equivalent to $(B_{\varepsilon})_{0<\varepsilon\leq 1}$ in the sense of Proposition 3.1. The original assumption (1.4) implies that u_{ε} still satisfies

(4.3)
$$u_{\varepsilon} = cB^{\mu_{\varepsilon}, x_{\varepsilon}} + o(1) \quad \text{in } H^{1}(M)$$

for some c > 0. We will thus, in the subsequent analysis, work with the family $(B^{\mu_{\varepsilon},x_{\varepsilon}})_{0<\varepsilon\leq 1}$. If $(y_{\varepsilon})_{\varepsilon}$ is a family of points in M satisfying $\mu_{\varepsilon} \ll d_g(x_{\varepsilon},y_{\varepsilon}) \leq \frac{i_g(M)}{4}$ we also have, by (2.4), (4.1) and (4.2),

(4.4)
$$B^{\mu_{\varepsilon},x_{\varepsilon}}(y_{\varepsilon}) = F_{\varepsilon}(x_{\varepsilon},y_{\varepsilon})\mu_{\varepsilon}^{-\frac{n-2}{2}}V\left(\frac{1}{\mu_{\varepsilon}}\exp_{x_{\varepsilon}}^{-1}(y_{\varepsilon})\right)$$
$$= (n-2)\omega_{n-1}\mu_{\varepsilon}^{\frac{n-2}{2}}G_{h_{\varepsilon}}(x_{\varepsilon},y_{\varepsilon})\left(\lambda(V) + O\left(\frac{\mu_{\varepsilon}}{d_{g}(x_{\varepsilon},y_{\varepsilon})}\right)\right).$$

The function $B^{\mu_{\varepsilon},x_{\varepsilon}}$ thus interpolates the (rescaled) pull-back of V in M to its first-order expansion at infinity.

In this section we prove that (4.3) still holds true globally pointwise in M, up to an error term that is of the order of the positive standard bubble B_{ε}^+ given by (3.5). The main result of this section is the following:

Proposition 4.1. Let $(M^n, g), n \geq 3$ be a smooth, connected and closed manifold, let $(h_{\varepsilon})_{0 < \varepsilon \leq 1}$ be a family of C^1 functions in M converging in $C^1(M)$ towards h and let $(p_{\varepsilon})_{0 < \varepsilon \leq 1}$ be a family of real numbers satisfying $2 < p_{\varepsilon} \leq 2^*$ for all $0 < \varepsilon \leq 1$ and such that $\lim_{\varepsilon \to 0} p_{\varepsilon} = 2^*$. Assume that $\Delta_g + h$ is coercive. Let $(u_{\varepsilon})_{0 < \varepsilon \leq 1}$ be a family of solutions of (1.3) satisfying (1.4) for some c > 0.

Then c = 1 in (1.4) and there exists a family $(\sigma_{\varepsilon})_{0 < \varepsilon \leq 1}$ of positive real numbers with $\lim_{\varepsilon \to 0} \sigma_{\varepsilon} = 0$ such that,

(4.5)
$$\left\| \left| \frac{u_{\varepsilon} - B^{\mu_{\varepsilon}, x_{\varepsilon}}}{B_{\varepsilon}^{+}} \right| \right|_{L^{\infty}(M)} \le \sigma_{\varepsilon}$$

for any $0 < \varepsilon \leq 1$, where $B^{\mu_{\varepsilon}, x_{\varepsilon}}$ is as in (4.2) and B_{ε}^+ is as in (3.5).

The function $V \in \Sigma$ that appears in the definition of $B^{\mu_{\varepsilon},x_{\varepsilon}}$ is the one that defines the family $(B_{\varepsilon})_{0<\varepsilon\leq 1}$ and is given by (3.4). In particular, for any $x \in M$ and any $0 < \varepsilon \leq 1$ Proposition 4.1 shows that

$$\left|u_{\varepsilon}(x) - B^{\mu_{\varepsilon}, x_{\varepsilon}}(x)\right| \le \sigma_{\varepsilon} B_{\varepsilon}^{+}(x).$$

As a consequence, if $(y_{\varepsilon})_{0 < \varepsilon < 1}$ is any family of points in M we have

$$u_{\varepsilon}(y_{\varepsilon}) = B^{\mu_{\varepsilon}, x_{\varepsilon}}(y_{\varepsilon}) + o\left(B_{\varepsilon}^{+}(y_{\varepsilon})\right)$$

as $\varepsilon \to 0$.

Proposition 4.1 is sharp if $\lambda(V) \neq 0$ which, by (2.4), corresponds to a bubble that decays to infinity in \mathbb{R}^n as $|x|^{2-n}$. This is the case for the positive bubble B_0^+ given by (2.2) and for the sign-changing solutions of (1.2) constructed in DelPino-Musso-Pacard-Pistoia [7]. Proposition 4.1 for positive solutions has been known since the work of Li-Zhu [20] and Druet [10] and Druet-Hebey-Robert [12]] (see also [16]). In this section we prove Proposition 4.1 by following the approach in Ghoussoub-Mazumdar-Robert [14], that draws inspiration from the techniques in [12,16]. Since we consider sign-changing solutions and bubbles we have to adapt the existing techniques that have been developed for positive solutions. In the exactly critical case $p_{\varepsilon} \equiv 2^*$, Proposition 4.1 has already been proven in Premoselli [28] in the more general case where multiple sign-changing bubbles may appear.

4.2. Proof of Proposition 4.1. As a preliminary observation we claim that

(4.6)
$$c = \lim_{\varepsilon \to 0} \mu_{\varepsilon}^{-\frac{(n-2)^2}{8}(2^{\star}-p_{\varepsilon})} \in [1, +\infty).$$

We set $\tilde{U}_{\varepsilon}(x) := \mu_{\varepsilon}^{\frac{n-2}{2}} u_{\varepsilon}(\exp_{x_{\varepsilon}}(\mu_{\varepsilon}x))$ for $x \in B(0, \frac{i_g(M)}{2\mu_{\varepsilon}})$. It follows from (1.4) and (3.4) that $\tilde{U}_{\varepsilon} \to cV$ in $H^1_{loc}(\mathbb{R}^n)$. With a change of variable, equation (1.3) writes

$$\Delta_{g_{\varepsilon}}\tilde{U}_{\varepsilon} + \mu_{\varepsilon}^{2}h_{\varepsilon}(\exp_{x_{\varepsilon}}(\mu_{\varepsilon}x))\tilde{U}_{\varepsilon} = \mu_{\varepsilon}^{\frac{n-2}{2}(2^{\star}-p_{\varepsilon})}|\tilde{U}_{\varepsilon}|^{2^{\star}-2}\tilde{U}_{\varepsilon} \text{ in } B\left(0,\frac{i_{g}(M)}{2\mu_{\varepsilon}}\right),$$

where we have let $g_{\varepsilon}(x) = \exp_{x_{\varepsilon}}^{*} g(\mu_{\varepsilon} x)$. Passing to the limit $\varepsilon \to 0$ yields

$$\Delta_{\xi}(cV) = \left(\lim_{\varepsilon \to 0} \mu_{\varepsilon}^{\frac{n-2}{2}(2^{\star}-p_{\varepsilon})}\right) c^{2^{\star}-1} |V|^{2^{\star}-2} V \text{ weakly in } \mathbb{R}^{n}.$$

Since $\mu_{\varepsilon} \to 0$ as $\varepsilon \to 0$ and $p_{\varepsilon} \leq 2^*$ we have $\mu_{\varepsilon}^{2^*-p_{\varepsilon}} \leq 1$ for ε small enough. Since $V \in D^{1,2}(\mathbb{R}^n) \setminus \{0\}$ it cannot be harmonic, which shows that $\lim_{\varepsilon \to 0} \mu_{\varepsilon}^{\frac{n-2}{2}(2^*-p_{\varepsilon})} \in (0,1]$. Finally, V solves (1.2), which proves the claim.

As already remarked, we can assume that (4.3) holds. This implies that the family $(u_{\varepsilon})_{0<\varepsilon<1}$ blows-up, that is

$$\lim_{\varepsilon \to 0} \|u_{\varepsilon}\|_{L^{\infty}(M)} = +\infty$$

By (4.6) it is also easily seen that

(4.7)
$$\mu_{\varepsilon}^{\frac{n-2}{2}-\frac{2}{p_{\varepsilon}-2}} \to c \quad \text{as } \varepsilon \to 0.$$

Decomposition (4.3) with (4.2) shows in particular that $u_{\varepsilon} \to 0$ in $H^1(M \setminus B_g(x_{\varepsilon}, \delta))$ as $\varepsilon \to 0$, for any $\delta > 0$ fixed. The regularity theory of Trudinger [37], together with standard elliptic theory, then shows that

(4.8)
$$u_{\varepsilon} \to 0 \quad \text{in} \quad C^2_{loc}(M \setminus \{x_0\})$$

as $\varepsilon \to 0$, where we have let $x_0 = \lim_{\varepsilon \to 0} x_{\varepsilon}$. We prove Proposition 4.1 in several steps.

Step 1: Let, for any
$$x \in B(0, \frac{i_g(M)}{2\mu_{\varepsilon}})$$
,
$$U_{\varepsilon}(x) = \mu_{\varepsilon}^{\frac{2}{p_{\varepsilon}-2}} u_{\varepsilon} \left(\exp_{x_{\varepsilon}}(\mu_{\varepsilon}x) \right)$$

We claim that

(4.9)
$$U_{\varepsilon} \to V \quad \text{in } C^2_{loc}(\mathbb{R}^n)$$

as $\varepsilon \to 0$.

Proof of (4.9). Let, for $x \in B(0, \frac{i_g(M)}{2\mu_{\varepsilon}}), g_{\varepsilon}(x) = \exp_{x_{\varepsilon}}^* g(\mu_{\varepsilon} x)$. By (1.3) U_{ε} satisfies

(4.10)
$$\Delta_{g_{\varepsilon}} U_{\varepsilon} + \mu_{\varepsilon}^{2} \tilde{h}_{\varepsilon} U_{\varepsilon} = |U_{\varepsilon}|^{p_{\varepsilon}-2} U_{\varepsilon} \quad \text{in } B\left(0, \frac{\imath_{g}(M)}{2\mu_{\varepsilon}}\right)$$

where we have let $\tilde{h}_{\varepsilon}(x) = h_{\varepsilon} \left(\exp_{x_{\varepsilon}}(\mu_{\varepsilon}x) \right)$. Equation (4.3) together with (4.7) shows that, for any compact set $K \subset \mathbb{R}^n$, we have

$$\begin{split} \int_{K} |U_{\varepsilon} - V|^{p_{\varepsilon}} dv_{g_{\varepsilon}} &= \mu_{\varepsilon}^{\frac{2p_{\varepsilon}}{p_{\varepsilon}-2}-n} \int_{\exp_{x_{\varepsilon}}(\mu_{\varepsilon}K)} \left| u_{\varepsilon} - \mu_{\varepsilon}^{\frac{n-2}{2} - \frac{2}{p_{\varepsilon}-2}} B^{\mu_{\varepsilon}, x_{\varepsilon}} \right|^{p_{\varepsilon}} dv_{g} \\ &= \mu_{\varepsilon}^{\frac{2p_{\varepsilon}}{p_{\varepsilon}-2}-n} \int_{\exp_{x_{\varepsilon}}(\mu_{\varepsilon}K)} \left| u_{\varepsilon} - (c+o(1)) B^{\mu_{\varepsilon}, x_{\varepsilon}} \right|^{p_{\varepsilon}} dv_{g} \\ &\lesssim \int_{\exp_{x_{\varepsilon}}(\mu_{\varepsilon}K)} \left| u_{\varepsilon} - (c+o(1)) B^{\mu_{\varepsilon}, x_{\varepsilon}} \right|^{p_{\varepsilon}} dv_{g} \\ &= o(1) \end{split}$$

as $\varepsilon \to 0$, where the second line follows from (4.7) and in the third line we used that $\mu_{\varepsilon}^{\frac{2p_{\varepsilon}}{p_{\varepsilon}-2}-n} \lesssim 1$ since $\frac{2p_{\varepsilon}}{p_{\varepsilon}-2}-n \ge 0$. The local regularity theory of Trudinger [37] and standard elliptic theory using (4.10) then show that the convergence of U_{ε} towards V takes place in $C_{loc}^2(\mathbb{R}^n)$, which concludes the proof of (4.9).

By (4.3), (4.9) and Sobolev's inequality, we get

(4.11)
$$\lim_{R \to +\infty} \limsup_{\varepsilon \to 0} \|u_{\varepsilon}\|_{L^{2^{\star}}(M \setminus B_g(x_{\varepsilon}, R\mu_{\varepsilon}))} = 0.$$

Step 2: Let $0 < \delta < i_g(M)$ be fixed. We claim that

(4.12)
$$\lim_{R \to +\infty} \limsup_{\varepsilon \to 0} \max_{B_g(x_{\varepsilon}, \delta) \setminus B_g(x_{\varepsilon}, R\mu_{\varepsilon})} d_g(x_{\varepsilon}, x)^{\frac{2}{p_{\varepsilon}-2}} |u_{\varepsilon}(x)| = 0.$$

Proof. To prove (4.12) we will prove more generally that

(4.13)
$$\limsup_{\varepsilon \to 0} \max_{x \in M} \left(\mu_{\varepsilon} + d_g(x_{\varepsilon}, x) \right)^{\frac{2}{p_{\varepsilon} - 2}} \left| u_{\varepsilon} - cB^{\mu_{\varepsilon}, x_{\varepsilon}} \right| (x) = 0.$$

It is easily seen using (4.2) that (4.12) follows from (4.13). To prove (4.13) we proceed by contradiction and assume that, up to passing to a subsequence $(\varepsilon_k)_{k\geq 0}$ with $\lim_{k\to+\infty} \varepsilon_k = 0$, there exists $\eta_0 > 0$ and a sequence $(y_k)_{k\geq 0}$ of points of Msuch that

(4.14)
$$(\mu_{\varepsilon_k} + d_g(x_{\varepsilon_k}, y_k))^{\frac{2}{p_{\varepsilon_k} - 2}} |u_{\varepsilon_k} - cB^{\mu_{\varepsilon_k}, x_{\varepsilon_k}}|(y_k)$$
$$= \max_{x \in M} (\mu_{\varepsilon_k} + d_g(x_{\varepsilon_k}, x))^{\frac{2}{p_{\varepsilon_k} - 2}} |u_{\varepsilon_k} - cB^{\mu_{\varepsilon_k}, x_{\varepsilon_k}}|(x) \ge \eta_0.$$

By (4.7) and (4.9) we get that

= o(1)

(4.15)
$$\frac{d_g(x_{\varepsilon_k}, y_k)}{\mu_{\varepsilon_k}} \to +\infty$$

as $k \to +\infty$. By (4.2) and since $\frac{2}{p_{\varepsilon_k}-2} - \frac{n-2}{2} \ge 0$ we thus have

(4.16)
$$(\mu_{\varepsilon_k} + d_g(x_{\varepsilon_k}, y_k))^{\frac{n}{p_{\varepsilon_k} - 2}} |B^{\mu_{\varepsilon_k}, x_{\varepsilon_k}}|(y_k) \\ \lesssim \left(\frac{\mu_{\varepsilon_k}}{\mu_{\varepsilon} + d_g(x_{\varepsilon_k}, y_k)}\right)^{\frac{n-2}{2}} (\mu_{\varepsilon} + d_g(x_{\varepsilon_k}, y_k))^{\frac{2}{p_{\varepsilon_k} - 2} - \frac{n-2}{2}}$$

as $k \to +\infty$. Using (4.2) and (4.8), (4.14) also shows that $d_g(x_{\varepsilon_k}, y_k) \to 0$, and hence, with (4.16), that $|u_{\varepsilon_k}(y_k)| \to +\infty$ as $k \to +\infty$. We let $\nu_k = |u_{\varepsilon_k}(y_k)|^{-\frac{p_{\varepsilon_k}-2}{2}}$, so that $\nu_k \to 0$, and for $x \in B(0, \frac{i_g(M)}{2\nu_k})$ we let

$$w_k(x) = \nu_k^{\frac{2}{p_{\varepsilon_k} - 2}} u_{\varepsilon_k} \left(\exp_{y_k}(\nu_k x) \right)$$

By (1.3) w_k satisfies

(4.17)
$$\Delta_{g_k} w_k + \nu_k^2 \tilde{h}_k w_k = |w_k|^{p_{\varepsilon_k} - 2} w_k \quad \text{in } B\left(0, \frac{i_g(M)}{2\nu_k}\right),$$

where we have let $h_k(x) = h_{\varepsilon_k} \left(\exp_{y_k}(\nu_k x) \right)$ and $g_k = \exp_{y_k}^* g(\nu_k x)$. Independently, by (4.14), (4.15) and (4.16) we see that

(4.18)
$$d_g(x_{\varepsilon_k}, y_k) \gtrsim \nu_k.$$

With the latter, (4.14) shows that there exists $r_0 > 0$ small enough such that

$$||w_k||_{L^{\infty}(B(0,2r_0))} \le 2$$

for any k large enough. With (4.17) and standard elliptic theory, and since $|w_k(0)| = 1$, w_k converges in $C^2(\overline{B(0, r_0)})$ as $k \to +\infty$, up to a subsequence, to a non-zero solution w_0 of

$$\Delta_{\xi} w_0 = |w_0|^{2^* - 2} w_0 \quad \text{in } B(0, r_0),$$

that thus satisfies

(4.19)
$$\int_{B(0,r_0)} |w_0|^{2^*} dx > 0$$

Independently, (4.15) and (4.18) show that, up to assuming that r_0 is small enough, we have

(4.20)
$$\frac{1}{\mu_{\varepsilon_k}} d_g \left(x_{\varepsilon_k}, B_g(y_k, r_0 \nu_k) \right) \to +\infty$$

as $k \to +\infty$. We let $q_k = \frac{n}{2}p_{\varepsilon_k} - n$. As is easily seen, it satisfies $q_k \leq 2^*$ (since $p_{\varepsilon_k} \leq 2^*$), $q_k \to 2^*$ as $k \to +\infty$ and

$$\frac{2q_k}{p_{\varepsilon_k} - 2} = n$$

We then have, by the C^2 convergence of w_k towards w_0 and by Hölder's inequality,

$$\int_{B(0,r_0)} |w_0|^{2^\star} dx = \limsup_{k \to +\infty} \int_{B(0,r_0)} |w_k|^{q_k} dv_{g_k}$$
$$= \limsup_{k \to +\infty} \int_{B_g(y_k,r_0\nu_k)} |u_{\varepsilon_k}|^{q_k} dv_g$$
$$\lesssim \limsup_{k \to +\infty} \left(\int_{B_g(y_k,r_0\nu_k)} |u_{\varepsilon_k}|^{2^\star} dv_g \right)^{\frac{q_k}{2^\star}}$$
$$= 0.$$

where the last equality follows from (4.11) and (4.20). This is a contradiction with (4.19). Hence (4.14) cannot occur, which proves (4.13).

For
$$0 < \delta < i_g(M)$$
 we let
(4.21) $\eta_{\varepsilon}(\delta) = ||u_{\varepsilon}||_{L^{\infty}(M \setminus B_g(x_{\varepsilon}, \delta))}.$

By (4.8), $\eta_{\varepsilon}(\delta) \to 0$ as $\varepsilon \to 0$ for any $0 < \delta < i_g(M)$ fixed.

Step 3: We claim that for any $0 < \alpha < \frac{1}{2}$ there exists $0 < \delta_{\alpha} < i_g(M)$ and $C_{\alpha} > 0$ such that, for any $x \in M \setminus B_g(x_{\varepsilon}, \mu_{\varepsilon})$ and any $0 < \varepsilon \leq 1$

$$(4.22) |u_{\varepsilon}(x)| \le C_{\alpha} \left(\mu_{\varepsilon}^{\frac{2}{p_{\varepsilon}-2}(1-2\alpha)} d_g(x_{\varepsilon}, x)^{-\frac{4(1-\alpha)}{p_{\varepsilon}-2}} + \eta_{\varepsilon}(\delta_{\alpha}) d_g(x_{\varepsilon}, x)^{-\frac{4\alpha}{p_{\varepsilon}-2}} \right)$$

holds.

Proof of (4.22). Let $0 < \alpha < \frac{1}{2}$. For $x \in M \setminus B_g(x_{\varepsilon}, \mu_{\varepsilon})$ we let

$$\Phi_{\varepsilon}^{\alpha}(x) = \mu_{\varepsilon}^{\frac{2}{p_{\varepsilon}-2}(1-2\alpha)} G_{h_{\varepsilon}}(x_{\varepsilon}, x)^{\frac{4(1-\alpha)}{(n-2)(p_{\varepsilon}-2)}} + \eta_{\varepsilon}(\delta) G_{h_{\varepsilon}}(x_{\varepsilon}, x)^{\frac{4\alpha}{(n-2)(p_{\varepsilon}-2)}}$$

where $G_{h_{\varepsilon}}$ is the Green's function of $\Delta_g + h_{\varepsilon}$. It follows from properties of the Green's function that there exists $\delta'_{\alpha} > 0$ such that

$$d_g(x, x_{\varepsilon})^2 \left(\frac{|\nabla_x G_{h_{\varepsilon}}(x_{\varepsilon}, x)|^2}{G_{h_{\varepsilon}}(x_{\varepsilon}, x)^2} - h_{\varepsilon}(x) \right) \ge \frac{(n-2)^2}{4} \text{ for } d_g(x, x_{\varepsilon}) < \delta'_{\alpha}.$$

Let $\delta_{\alpha} \in (0, \delta'_{\alpha})$ be such that

$$2\|h\|_{L^{\infty}(M)}\delta_{\alpha}^{2} < (n-2)^{2}\alpha(1-\alpha)$$

and let $0 < \delta \le \delta_{\alpha}$ be fixed. We claim that there exists C > 0 such that, for any $0 < \varepsilon \le 1$,

(4.23)
$$\left\| \left| \frac{u_{\varepsilon}}{\Phi_{\varepsilon}^{\alpha}} \right| \right\|_{L^{\infty}(M \setminus B_g(x_{\varepsilon}, \mu_{\varepsilon}))} \le C$$

holds. By (4.1) it is easily seen that (4.22) follows from (4.23). We thus prove (4.23). We let, for any $0 < \varepsilon \leq 1$, $y_{\varepsilon} \in M \setminus B_g(x_{\varepsilon}, \mu_{\varepsilon})$ be such that

$$\frac{u_{\varepsilon}(y_{\varepsilon})}{\Phi_{\varepsilon}^{\alpha}(y_{\varepsilon})} \bigg| = \max_{x \in M \setminus B_g(x_{\varepsilon}, \mu_{\varepsilon})} \bigg| \frac{u_{\varepsilon}(x)}{\Phi_{\varepsilon}^{\alpha}(x)} \bigg|.$$

We proceed by contradiction and assume that, up to passing to a subsequence $(\varepsilon_k)_{k\geq 0}$ with $\lim_{k\to+\infty} \varepsilon_k = 0$, and up to replacing u_{ε_k} by $-u_{\varepsilon_k}$ (which still solves (1.3)), we have

(4.24)
$$\frac{u_{\varepsilon_k}(y_{\varepsilon_k})}{\Phi_{\varepsilon_k}^{\alpha}(y_{\varepsilon_k})} \to +\infty$$

as $k \to +\infty$. We first assume that, up to a subsequence,

(4.25)
$$\mu_{\varepsilon_k} = o(d_g(y_{\varepsilon_k}, x_{\varepsilon_k})) \quad \text{and} \quad d_g(x_{\varepsilon_k}, y_{\varepsilon_k}) \le \delta_\alpha$$

as $k \to +\infty$. By definition of y_{ε_k} we have

(4.26)
$$\frac{\Delta_g u_{\varepsilon_k}(y_{\varepsilon_k})}{u_{\varepsilon_k}(y_{\varepsilon_k})} \ge \frac{\Delta_g \Phi_{\varepsilon_k}^{\alpha}(y_{\varepsilon_k})}{\Phi_{\varepsilon_k}^{\alpha}(y_{\varepsilon_k})}$$

On the one side, straightforward computations show that

$$d_g(x_{\varepsilon_k}, y_{\varepsilon_k})^2 \frac{\Delta_g \Phi_{\varepsilon_k}^{\alpha}(y_{\varepsilon_k})}{\Phi_{\varepsilon_k}^{\alpha}(y_{\varepsilon_k})} \ge \frac{(n-2)^2}{2} \alpha(1-\alpha) > 0 \text{ for } d_g(x, x_{\varepsilon}) < \delta_{\alpha}$$

as $k \to +\infty$. On the other side, (1.3) and (4.25) show that

$$d_g(x_{\varepsilon_k}, y_{\varepsilon_k})^2 \frac{\Delta_g u_{\varepsilon_k}(y_{\varepsilon_k})}{u_{\varepsilon_k}(y_{\varepsilon_k})} \le d_g(x_{\varepsilon_k}, y_{\varepsilon_k})^2 |u_{\varepsilon_k}(y_{\varepsilon_k})|^{p_{\varepsilon_k}-2} + 2||h||_{L^{\infty}(M)} \delta_{\alpha}^2$$

as $k \to +\infty$. With (4.12), (4.26) and the choice of δ_{α} this yields a contradiction for k large enough, and shows that (4.25) cannot happen. Thus we either have, up to a subsequence, $\frac{d_g(x_{\varepsilon_k}, y_{\varepsilon_k})}{\mu_{\varepsilon_k}} \to D \in [1, +\infty)$ or $d_g(x_{\varepsilon_k}, y_{\varepsilon_k}) \ge \delta_{\alpha}$ as $k \to +\infty$. Using (4.9) in the first case and (4.21) in the second case we obtain a contradiction with (4.24). This proves (4.23) and thus (4.22).

Step 4: We claim that

(4.27)
$$|u_{\varepsilon}(x)| \lesssim B_{\varepsilon}^+(x)$$
 for any $x \in M$.

where B_{ε}^+ is defined in (3.5).

Proof of (4.27). We first prove a weaker statement. Let $0 < \delta < i_g(M)$ be fixed. We claim that there exists $C_{\delta} > 0$ such that, for any $0 < \varepsilon \leq 1$ and $x \in M$,

(4.28)
$$|u_{\varepsilon}(x)| \le C_{\delta} \left(B_{\varepsilon}^{+}(x) + \eta_{\varepsilon}(\delta) \right)$$

holds, where B_{ε}^+ is as in (3.5). Let $(y_{\varepsilon})_{0 < \varepsilon \leq 1}$ be any family of points in M. First, if $d_g(y_{\varepsilon}, x_{\varepsilon}) \geq \delta$, (4.28) follows from (4.21). We can thus assume that $y_{\varepsilon} \in B_g(x_{\varepsilon}, \delta)$ for all $0 < \varepsilon \leq 1$. A representation formula for u_{ε} at y_{ε} gives, with (1.3),

(4.29)
$$u_{\varepsilon}(y_{\varepsilon}) = \int_{B_{g}(x_{\varepsilon},\mu_{\varepsilon})} G_{h_{\varepsilon}}(y_{\varepsilon},y) |u_{\varepsilon}(y)|^{p_{\varepsilon}-2} u_{\varepsilon}(y) dy + \int_{M \setminus B_{g}(x_{\varepsilon},\mu_{\varepsilon})} G_{h_{\varepsilon}}(y_{\varepsilon},y) |u_{\varepsilon}(y)|^{p_{\varepsilon}-2} u_{\varepsilon}(y) dy$$

Straightforward computations with (4.7), (4.9) and (4.1) show that

(4.30)
$$\left| \int_{B_g(x_{\varepsilon},\mu_{\varepsilon})} G_{h_{\varepsilon}}(y_{\varepsilon},y) |u_{\varepsilon}(y)|^{p_{\varepsilon}-2} u_{\varepsilon}(y) dy \right| \lesssim B_{\varepsilon}^+(y_{\varepsilon})$$

(see e.g. [16, Proposition 6.1]). Straightforward computations using (4.7), (4.22) and (4.1) show that

(4.31)
$$\left| \int_{M \setminus B_g(x_{\varepsilon}, \mu_{\varepsilon})} G_{h_{\varepsilon}}(y_{\varepsilon}, y) |u_{\varepsilon}(y)|^{p_{\varepsilon}-2} u_{\varepsilon}(y) dy \right| \lesssim B_{\varepsilon}^+(y_{\varepsilon}) + \eta_{\varepsilon}(\delta)^{p_{\varepsilon}-1}.$$

Plugging (4.30) and (4.31) in (4.29) and since $\eta_{\varepsilon}(\delta) \to 0$ as $\varepsilon \to 0$ proves (4.28). We now claim that, for any $0 < \delta < i_q(M)$ fixed,

(4.32)
$$\eta_{\varepsilon}(\delta) \lesssim \mu_{\varepsilon}^{\frac{n-2}{2}}$$

1

holds for ε small enough. With (3.5), (4.28) and (4.32) this will conclude the proof of (4.27). We prove (4.32) by contradiction: we fix $0 < \delta < i_g(M)$ and, up to passing to a subsequence $(\varepsilon_k)_{k\geq 0}$ with $\lim_{k\to+\infty} \varepsilon_k = 0$ we assume that

(4.33)
$$\frac{\eta_{\varepsilon_k}(\delta)}{\mu_{\varepsilon_k}^{n-2}} \to +\infty$$

as $k \to +\infty$. We let, for $k \ge 0$ and for any $x \in M$,

$$\hat{u}_k(x) = \frac{u_{\varepsilon_k}(x)}{\eta_{\varepsilon_k}(\delta)},$$

and let $y_k \in M \setminus B_g(x_{\varepsilon_k}, \delta)$ be such that $|\hat{u}_k(y_k)| = 1$. By (3.5), (4.28) and (4.33) the family $(\hat{u}_k)_{k\geq 0}$ is uniformly bounded in compact subsets of $M \setminus \{x_0\}$, where $x_0 = \lim_{\varepsilon \to 0} x_{\varepsilon}$. By (1.3), standard elliptic theory and since $\eta_{\varepsilon_k}(\delta) \to 0$ as $k \to +\infty$, \hat{u}_k thus converges in $C^2_{loc}(M \setminus \{x_0\})$, up to a subsequence, to some function $\hat{u}_0 \in C^2(M \setminus \{x_0\})$ that satisfies $\Delta_g \hat{u}_0 + h \hat{u}_0 = 0$ in $M \setminus \{x_0\}$ and $|\hat{u}_0(y_0)| = 1$, where $y_0 = \lim_{k \to +\infty} y_k \in M \setminus B_g(x_0, \delta)$. Passing (4.28) to the limit pointwise shows in addition that we have

$$|\hat{u}_0(x)| \le C_\delta$$
 for all $x \in M \setminus \{x_0\}$.

Classical arguments thus show that the singularity of \hat{u}_0 at x_0 is removable and that \hat{u}_0 satisfies $\Delta_g \hat{u}_0 + h \hat{u}_0 = 0$ in M in a strong sense. This implies $\hat{u}_0 \equiv 0$ since $\Delta_g + h$ is coercive, which is a contradiction with $|\hat{u}_0(y_0)| = 1$. Thus (4.32) holds true and (4.27) is proven.

A consequence of (4.27), (1.3) and standard elliptic theory is the following estimate on the first and second derivatives of u_{ε} : for $0 \le k \le 2$,

(4.34)
$$|\nabla^k u_{\varepsilon}(x)| \lesssim \frac{\mu_{\varepsilon}^{\frac{n-2}{2}}}{(\mu_{\varepsilon} + d_g(x_{\varepsilon}, x))^{n-2+k}} \quad \text{for all } x \in M \text{ and } 0 < \varepsilon \le 1.$$

Estimate (4.34) follows from a scaling argument and from standard elliptic theory.

Step 5: we now claim that

(4.35)
$$2^{\star} - p_{\varepsilon} = \begin{cases} O(\mu_{\varepsilon}) & \text{if } n = 3\\ O\left(\mu_{\varepsilon}^{2} \ln \frac{1}{\mu_{\varepsilon}}\right) & \text{if } n = 4\\ O(\mu_{\varepsilon}^{2}) & \text{if } n \ge 5 \end{cases}$$

as $\varepsilon \to 0$. In particular $\lim_{\varepsilon \to 0} \mu_{\varepsilon}^{2^* - p_{\varepsilon}} = 1$.

Proof of (4.35). Let $0 < \delta < i_g(M)$ be fixed and let

$$\Omega_{\varepsilon} = B_g(x_{\varepsilon}, \delta).$$

We write a Pohozaev identity for u_{ε} on Ω_{ε} : let X be any smooth vector field in M. Integrating (1.3) by parts against $\langle X, \nabla u_{\varepsilon} \rangle$ classically shows that

$$(4.36) \qquad \int_{\Omega_{\varepsilon}} h_{\varepsilon} u_{\varepsilon} \langle X, \nabla u_{\varepsilon} \rangle dv_{g} + \int_{\Omega_{\varepsilon}} \left(\nabla X - \frac{1}{2} \operatorname{div}_{g} Xg \right) (\nabla u_{\varepsilon}, \nabla u_{\varepsilon}) dv_{g}$$
$$(4.36) \qquad + \int_{\partial\Omega_{\varepsilon}} \left(\frac{1}{2} \langle X, \nu \rangle |\nabla u_{\varepsilon}|_{g}^{2} - \langle X, \nabla u_{\varepsilon} \rangle \partial_{\nu} u_{\varepsilon} - \frac{1}{p_{\varepsilon}} \langle X, \nu \rangle |u_{\varepsilon}|_{p_{\varepsilon}}^{p_{\varepsilon}} \right) d\sigma_{g}$$
$$= -\frac{1}{p_{\varepsilon}} \int_{\Omega_{\varepsilon}} \operatorname{div}_{g} X |u_{\varepsilon}|_{p_{\varepsilon}}^{p_{\varepsilon}} dv_{g}$$

(see e.g. [16, Proposition 6.2]). Let now X_{ε} be the smooth vector field whose coordinates in the exponential chart at x_{ε} are $(X_{\varepsilon}(x))^i = x^i$. For $x \in \Omega_{\varepsilon}$ and any $1 \leq i, j \leq n$ we have $\nabla_i X^j_{\varepsilon}(x) = \delta^j_i + O(d_g(x_{\varepsilon}, x)^2)$. As a consequence,

$$\int_{\Omega_{\varepsilon}} \left(\nabla X - \frac{1}{2} \mathrm{div}_g X g \right) (\nabla u_{\varepsilon}, \nabla u_{\varepsilon}) dv_g = \int_{\Omega_{\varepsilon}} \left(1 - \frac{n}{2} + O(d_g(x_{\varepsilon}, \cdot)^2) \right) |\nabla u_{\varepsilon}|_g^2 dv_g$$
and

$$-\frac{1}{p_{\varepsilon}} \int_{\Omega_{\varepsilon}} \operatorname{div}_{g} X |u_{\varepsilon}|^{p_{\varepsilon}} dv_{g} = \int_{\Omega_{\varepsilon}} \left(-\frac{n}{p_{\varepsilon}} + O(d_{g}(x_{\varepsilon}, \cdot)^{2}) |u_{\varepsilon}|^{p_{\varepsilon}} dv_{g} \right).$$

Integrating (1.3) by parts shows independently that

$$\int_{\Omega_{\varepsilon}} |\nabla u_{\varepsilon}|_{g}^{2} dv_{g} = \int_{\Omega_{\varepsilon}} |u_{\varepsilon}|^{p_{\varepsilon}} dv_{g} - \int_{\Omega_{\varepsilon}} h_{\varepsilon} u_{\varepsilon}^{2} dv_{g} + \int_{\partial\Omega_{\varepsilon}} u_{\varepsilon} \partial_{\nu} u_{\varepsilon} d\sigma_{g},$$

so that (4.36) becomes

$$\begin{split} n\left(\frac{1}{2^{\star}}-\frac{1}{p_{\varepsilon}}\right) &\int_{\Omega_{\varepsilon}} |u_{\varepsilon}|^{p_{\varepsilon}} dv_{g} \\ &= \int_{\Omega_{\varepsilon}} \left(\frac{n-2}{2}u_{\varepsilon}^{2}+h_{\varepsilon}u_{\varepsilon}\langle X,\nabla u_{\varepsilon}\rangle\right) dv_{g} \\ &+ O\left(\int_{\Omega_{\varepsilon}} d_{g}(x_{\varepsilon},\cdot)^{2}|\nabla u_{\varepsilon}|_{g}^{2} dv_{g}+\int_{\Omega_{\varepsilon}} d_{g}(x_{\varepsilon},\cdot)^{2}|u_{\varepsilon}|^{p_{\varepsilon}} dv_{g}\right) \\ &+ \int_{\partial\Omega_{\varepsilon}} \left(\frac{1}{2}\langle X,\nu\rangle|\nabla u_{\varepsilon}|_{g}^{2}-\langle X,\nabla u_{\varepsilon}\rangle\partial_{\nu}u_{\varepsilon}-\frac{n-2}{2}u_{\varepsilon}\partial_{\nu}u_{\varepsilon} \\ &-\frac{1}{p_{\varepsilon}}\langle X,\nu\rangle|u_{\varepsilon}|^{p_{\varepsilon}}\right) d\sigma_{g}. \end{split}$$

Straightforward computations using (4.27) and (4.34) show that

$$\int_{\Omega_{\varepsilon}} d_g(x_{\varepsilon}, \cdot)^2 |\nabla u_{\varepsilon}|_g^2 dv_g + \int_{\Omega_{\varepsilon}} d_g(x_{\varepsilon}, \cdot)^2 |u_{\varepsilon}|^{p_{\varepsilon}} dv_g$$
$$+ \int_{\Omega_{\varepsilon}} |h_{\varepsilon} u_{\varepsilon} \langle X, \nabla u_{\varepsilon} \rangle |dv_g + \int_{\Omega_{\varepsilon}} |h_{\varepsilon} |u_{\varepsilon}^2 dv_g = \begin{cases} O(\mu_{\varepsilon}) & \text{if } n = 3\\ O(\mu_{\varepsilon}^2 \ln \frac{1}{\mu_{\varepsilon}}) & \text{if } n = 4\\ O(\mu_{\varepsilon}^2) & \text{if } n \geq 5 \end{cases}$$

20

By (4.27) and (4.34) we have

$$\int_{\partial\Omega_{\varepsilon}} \left(\frac{1}{2} \langle X, \nu \rangle |\nabla u_{\varepsilon}|_{g}^{2} - \langle X, \nabla u_{\varepsilon} \rangle \partial_{\nu} u_{\varepsilon} - \frac{n-2}{2} u_{\varepsilon} \partial_{\nu} u_{\varepsilon} - \frac{1}{p_{\varepsilon}} \langle X, \nu \rangle |u_{\varepsilon}|^{p_{\varepsilon}} \right) d\sigma_{g} = O(\mu_{\varepsilon}^{n-2})$$

Plugging the latter computations in (4.36) thus shows that

(4.37)
$$n\left(\frac{1}{2^{\star}} - \frac{1}{p_{\varepsilon}}\right) \int_{\Omega_{\varepsilon}} |u_{\varepsilon}|^{p_{\varepsilon}} dv_{g} = \begin{cases} O(\mu_{\varepsilon}) & \text{if } n = 3\\ O(\mu_{\varepsilon}^{2} \ln \frac{1}{\mu_{\varepsilon}}) & \text{if } n = 4\\ O(\mu_{\varepsilon}^{2}) & \text{if } n \ge 5 \end{cases}$$

Finally, using (4.7) and (4.9), Fatou's lemma shows that

$$\int_{\Omega_{\varepsilon}} |u_{\varepsilon}|^{p_{\varepsilon}} dv_g \ge c_0$$

for some positive $c_0 > 0$ independent of ε . Going back to (4.37) thus proves (4.35).

Step 6: End of the proof of Proposition 4.1. We are now in position to conclude the proof of Proposition 4.1. The equality c = 1 follows from (4.6) and (4.35) and we thus only need to prove (4.5). We proceed by contradiction and assume that (4.5) is false: up to passing to a subsequence $(\varepsilon_k)_{k\geq 0}$ with $\lim_{k\to+\infty} \varepsilon_k = 0$, there exists a sequence $(y_k)_{k\geq 0}$ of points of M such that

$$(4.38) |u_{\varepsilon_k}(y_k) - B^{\mu_{\varepsilon_k}, x_{\varepsilon_k}}(y_k)| \ge \eta_0 B^+_{\varepsilon_k}(y_k)$$

for some $\eta_0 > 0$, where $B^{\mu_{\varepsilon},x_{\varepsilon}}$ and B^+_{ε} are as in (4.2) and (3.5). Since by (4.35) we have c = 1, (4.9) shows with (4.38) that

(4.39)
$$\frac{d_g(x_{\varepsilon_k}, y_{\varepsilon_k})}{\mu_{\varepsilon_k}} \to +\infty$$

as $k \to +\infty$. We write a representation formula for u_{ε_k} in M: if $G_{h_{\varepsilon_k}}$ again denotes the Green's function for $\Delta_g + h_{\varepsilon_k}$, we have with (1.3):

(4.40)
$$u_{\varepsilon_k}(y_k) = \int_M G_{h_{\varepsilon_k}}(y_k, y) |u_{\varepsilon_k}(y)|^{p_{\varepsilon_k}-2} u_{\varepsilon_k}(y) dv_g(y).$$

Let R > 1 be fixed and let $x \in B(0, R)$. By (4.1) and (4.39) we have

$$(n-2)\omega_{n-1}d_g(x_{\varepsilon_k}, y_k)^{n-2}G_{h_{\varepsilon_k}}\left(\exp_{x_{\varepsilon_k}}(\mu_{\varepsilon_k}x), y_k)\right) \to 1 \quad \text{if } d_g(x_{\varepsilon_k}, y_k) \to 0$$

uniformly in B(0,R) as $k \to +\infty$. Since $h_{\varepsilon_k} \to h$ in $C^1(M)$ we also have

$$G_{h_{\varepsilon_k}}\left(y_k, \exp_{x_{\varepsilon_k}}(\mu_{\varepsilon_k}x)\right) = G_h(x_{\varepsilon_k}, y_k) + o(1) \quad \text{if } d_g(x_{\varepsilon_k}, y_k) \not\to 0$$

as $k \to +\infty$. Since y_k satisfies (4.39), and by (4.1), the dominated convergence theorem gives with the definition (4.4)

(4.41)

$$\int_{B_g(x_{\varepsilon_k}, R\mu_{\varepsilon_k})} G_{h_{\varepsilon_k}}(y_k, \cdot) |u_{\varepsilon_k}|^{p_{\varepsilon_k} - 2} u_{\varepsilon_k} dv_g$$

$$= (1 + o(1)) G_{h_{\varepsilon_k}}(x_{\varepsilon_k}, y_k) \mu_{\varepsilon_k}^{\frac{n-2}{2}} \int_{B(0,R)} |V|^{2^* - 2} V dx$$

$$= (1 + o(1)) G_{h_{\varepsilon_k}}(x_{\varepsilon_k}, y_k) (n - 2) \omega_{n-1} \mu_{\varepsilon_k}^{\frac{n-2}{2}} \left(\lambda(V) + O(R^{-2})\right)$$

$$= B^{\mu_{\varepsilon_k}, x_{\varepsilon_k}}(y_k) + o(B_{\varepsilon_k}(y_k)) + O(R^{-2}B_{\varepsilon_k}(y_k))$$

where $\lambda(V)$ is as in (2.4), where the third line follows from (2.8) and (2.9) and the last one from (4.2) and (4.39). Independently, straightforward computations show with (4.27) that

$$\left|\int_{M\setminus B_g(x_{\varepsilon_k},R\mu_{\varepsilon_k})} G_{h_{\varepsilon_k}}(y_k,\cdot)|u_{\varepsilon_k}|^{p_{\varepsilon_k}-2} u_{\varepsilon_k} dv_g\right| \lesssim R^{-2} B_{\varepsilon_k}^+(y_k) + o\left(B_{\varepsilon_k}^+(y_k)\right)$$

holds. Plugging the latter and (4.41) in (4.40) shows that, for any fixed R > 0 and for any k large enough we have

$$u_{\varepsilon_k}(y_k) = B^{\mu_{\varepsilon_k}, x_{\varepsilon_k}}(y_k) + o\left(B^+_{\varepsilon_k}(y_k)\right) + O\left(R^{-2}B^+_{\varepsilon_k}(y_k)\right)$$

By choosing R large enough and passing to a subsequence we get a contradiction with (4.38). This concludes the proof of Proposition 4.1. \Box

We conclude this section by showing an improvement of (4.34) when k = 1 that will be needed in the next section. Let $(h_{\varepsilon})_{0 < \varepsilon \leq 1}$, $(p_{\varepsilon})_{0 < \varepsilon \leq 1}$ be as in the statement of Proposition 4.1 and let $(u_{\varepsilon})_{0 < \varepsilon \leq 1}$ be a family of solutions of (1.3) satisfying (1.4) for some c > 0. In particular Proposition 4.1 applies to $(u_{\varepsilon})_{0 < \varepsilon \leq 1}$ and (4.5) holds. We claim that there exists a family of positive real numbers $(\sigma_{\varepsilon})_{0 < \varepsilon \leq 1}$, with $\sigma_{\varepsilon} \to 0$ as $\varepsilon \to 0$, and a positive constant C such that for any $x \in M$ and any $0 < \varepsilon \leq 1$ we have

(4.42)
$$\left| \nabla u_{\varepsilon}(x) - \nabla B^{\mu_{\varepsilon}, x_{\varepsilon}}(x) \right| \leq \sigma_{\varepsilon} \frac{\mu_{\varepsilon}^{\frac{n-2}{2}}}{(\mu_{\varepsilon} + d_g(x_{\varepsilon}, x))^{n-1}} + C\mu_{\varepsilon}^{\frac{n-2}{2}}$$

where $B^{\mu_{\varepsilon}, x_{\varepsilon}}$ is as in (4.2).

Proof of (4.42). Let $0 < 2\delta < i_g(M)$ be fixed. We define $w_{\varepsilon}(x) = u_{\varepsilon}(\exp_{x_{\varepsilon}}(x))$ for $x \in B(0, 2\delta)$. We first prove that there exists a family $(\sigma_{\varepsilon})_{0 < \varepsilon \leq 1}$, with $\sigma_{\varepsilon} \to 0$ as $\varepsilon \to 0$, and a constant C > 0 such that for any $x \in B(0, 2\delta)$ and any $0 < \varepsilon \leq 1$ we have

(4.43)
$$\left|\partial_i w_{\varepsilon}(x) - \mu_{\varepsilon}^{-\frac{n}{2}}(\partial_i V) \left(\frac{x}{\mu_{\varepsilon}}\right)\right| \le \sigma_{\varepsilon} \frac{\mu_{\varepsilon}^{\frac{n-2}{2}}}{(\mu_{\varepsilon} + |x|)^{n-1}} + C\mu_{\varepsilon}^{\frac{n-2}{2}}.$$

Let $x \in B(0, 2\delta)$. The representation formula (4.40) can be differentiated with respect to x: this gives, for $1 \le i \le n$,

$$\partial_i w_{\varepsilon}(x) = \int_M \partial_i G_{h_{\varepsilon}}\big(\exp_{x_{\varepsilon}}(x), y\big) |u_{\varepsilon}(y)|^{p_{\varepsilon}-2} u_{\varepsilon}(y) dv_g(y),$$

where we have let $\partial_i G_{h_{\varepsilon}}(\exp_{x_{\varepsilon}}(x), y) = \frac{\partial}{\partial x_i} (G_{h_{\varepsilon}}(\exp_{x_{\varepsilon}}(\cdot), y))(x)$. Standard results on the Green's functions (see e.g. [32]) show that there exists C > 0 such that for all ε small enough and all $x, y \in B(0, 2\delta)$,

$$(4.44) \left| \partial_i G_{h_{\varepsilon}} \left(\exp_{x_{\varepsilon}}(x), \exp_{x_{\varepsilon}}(y) \right) + \frac{1}{\omega_{n-1}} \frac{(x-y)_i}{|x-y|^n} \right| \le C|x-y|^{2-n} + C|x| \cdot |x-y|^{1-n}$$

holds. Let now, for $x \in \mathbb{R}^n$, $\tilde{V}_{\varepsilon}(x) = \mu_{\varepsilon}^{-\frac{n-2}{2}} V(\frac{x}{\mu_{\varepsilon}})$. Let $(z_{\varepsilon})_{0 < \varepsilon \leq 1}$ be any family of points in $B(0, 2\delta)$. Straightforward computations using (4.35) and (4.5) show that

$$\begin{aligned} \partial_i w_{\varepsilon}(z_{\varepsilon}) &= -\int_{B(0,2\delta)} \frac{1}{\omega_{n-1}} \frac{(z_{\varepsilon} - y)_i}{|z_{\varepsilon} - y|^n} |\tilde{V}_{\varepsilon}|^{p_{\varepsilon} - 2} \tilde{V}_{\varepsilon} dy + o\left(\frac{\mu_{\varepsilon}^{\frac{n-2}{2}}}{(\mu_{\varepsilon} + |z_{\varepsilon}|)^{n-1}}\right) + O(\mu_{\varepsilon}^{\frac{n-2}{2}}) \\ &= -\int_{B(0,2\delta)} \frac{1}{\omega_{n-1}} \frac{(z_{\varepsilon} - y)_i}{|z_{\varepsilon} - y|^n} |\tilde{V}_{\varepsilon}|^{2^* - 2} \tilde{V}_{\varepsilon} dy + o\left(\frac{\mu_{\varepsilon}^{\frac{n-2}{2}}}{(\mu_{\varepsilon} + |z_{\varepsilon}|)^{n-1}}\right) + O(\mu_{\varepsilon}^{\frac{n-2}{2}}) \\ &+ \int_{B(0,2\delta)} \frac{1}{\omega_{n-1}} \frac{(z_{\varepsilon} - y)_i}{|z_{\varepsilon} - y|^n} \left(|\tilde{V}_{\varepsilon}|^{2^* - 2} - |\tilde{V}_{\varepsilon}|^{p_{\varepsilon} - 2}\right) \tilde{V}_{\varepsilon} dy. \end{aligned}$$

Let us estimate the last term. Using the control (4.6), we get that

$$\begin{split} \left| \int_{B(0,2\delta)} \frac{1}{\omega_{n-1}} \frac{(z_{\varepsilon} - y)_{i}}{|z_{\varepsilon} - y|^{n}} \left(|\tilde{V}_{\varepsilon}|^{2^{*}-2} - |\tilde{V}_{\varepsilon}|^{p_{\varepsilon}-2} \right) \tilde{V}_{\varepsilon} dy \right| \\ &\leq \int_{B(0,2\delta)} |z_{\varepsilon} - y|^{1-n} \left| |\tilde{V}_{\varepsilon}|^{2^{*}-p_{\varepsilon}} - 1 \right| \cdot |\tilde{V}_{\varepsilon}|^{p_{\varepsilon}-1} dy \\ &\leq \int_{B(0,2\delta) \cap \{ |\tilde{V}_{\varepsilon}| < \mu_{\varepsilon}^{\frac{n-2}{2}} \}} |z_{\varepsilon} - y|^{1-n} \left| |\tilde{V}_{\varepsilon}|^{2^{*}-p_{\varepsilon}} - 1 \right| \cdot |\tilde{V}_{\varepsilon}|^{p_{\varepsilon}-1} dy \\ &+ \int_{B(0,2\delta) \cap \{ |\tilde{V}_{\varepsilon}| \ge \mu_{\varepsilon}^{\frac{n-2}{2}} \}} |z_{\varepsilon} - y|^{1-n} \left| |\tilde{V}_{\varepsilon}|^{2^{*}-p_{\varepsilon}} - 1 \right| \cdot |\tilde{V}_{\varepsilon}|^{p_{\varepsilon}-1} dy \\ &\leq C \mu_{\varepsilon}^{\frac{n-2}{2}(p_{\varepsilon}-1)} + \int_{B(0,2\delta) \cap \{ |\tilde{V}_{\varepsilon}| \ge \mu_{\varepsilon}^{\frac{n-2}{2}} \}} |z_{\varepsilon} - y|^{1-n} \left| |\tilde{V}_{\varepsilon}|^{2^{*}-p_{\varepsilon}} - 1 \right| \cdot |\tilde{V}_{\varepsilon}|^{p_{\varepsilon}-1} dy \end{split}$$

The control (2.8) and (4.6) yield

$$\left\{ |\tilde{V}_{\varepsilon}| \ge \mu_{\varepsilon}^{\frac{n-2}{2}} \right\} \Rightarrow \left| |\tilde{V}_{\varepsilon}|^{2^* - p_{\varepsilon}} - 1 \right| = o(1)$$

as $\varepsilon \to 0$, so that we get

$$\begin{aligned} \partial_i w_{\varepsilon}(z_{\varepsilon}) &= -\int_{\mathbb{R}^n} \frac{1}{\omega_{n-1}} \frac{(z_{\varepsilon} - y)_i}{|z_{\varepsilon} - y|^n} |\tilde{V}_{\varepsilon}|^{2^* - 2} \tilde{V}_{\varepsilon} dy + o\left(\frac{\mu_{\varepsilon}^{\frac{n-2}{2}}}{(\mu_{\varepsilon} + |z_{\varepsilon}|)^{n-1}}\right) + O(\mu_{\varepsilon}^{\frac{n-2}{2}}) \\ &= \partial_i \tilde{V}_{\varepsilon}(z_{\varepsilon}) + o\left(\frac{\mu_{\varepsilon}^{\frac{n-2}{2}}}{(\mu_{\varepsilon} + |z_{\varepsilon}|)^{n-1}}\right) + O(\mu_{\varepsilon}^{\frac{n-2}{2}}), \end{aligned}$$

where the last equality follows from a representation formula for (1.2) in \mathbb{R}^n , since $V \in \Sigma$. This proves (4.43). Estimate (4.42) now follows from (4.1), (4.43), (4.44) and the explicit expression of $B^{\mu_{\varepsilon}, x_{\varepsilon}}$ in (4.2).

Remark 4.2. We assumed for simplicity here that $\Delta_g + h$ is coercive. If $\Delta_g + h$ had a kernel and, more generally, if $(u_{\varepsilon})_{0<\varepsilon\leq 1}$ was a finite-energy family of solutions of (1.3) (without the one-bubble assumption (1.4)), an analogue of Proposition 4.1 would still hold. We refer for instance to Premoselli [28] where a generalisation of Proposition 4.1 was proven when $p_{\varepsilon} = 2^*$ for all $0 < \varepsilon \leq 1$, in the general multi-bubble case and when $\Delta_g + h$ is allowed to have a kernel. The proof of [28] allows to deal with any multi-bubble configuration and would still adapt to the asymptotically critical setting $p_{\varepsilon} \leq 2^*$, $p_{\varepsilon} \to 2^*$ as a careful inspection of the proof shows. In the one-bubble case (1.4) that we consider here, the proof of Proposition 4.1 that we gave in this section is shorter and more direct.

5. Proof of Theorems 1.1 and 1.2

Throughout this section we let $(M^n, g), n \geq 3$ be a smooth, connected and closed manifold, $(h_{\varepsilon})_{0 < \varepsilon \leq 1}$ be a family of C^1 functions in M converging in $C^1(M)$ towards h and let $(p_{\varepsilon})_{0 < \varepsilon \leq 1}$ be a family of real numbers satisfying $2 < p_{\varepsilon} \leq 2^*$ for all $0 < \varepsilon \leq 1$ and such that $\lim_{\varepsilon \to 0} p_{\varepsilon} = 2^*$. We assume that $\Delta_g + h$ is coercive. We assume again, as in Section 4, that $(u_{\varepsilon})_{0 < \varepsilon \leq 1}$ is a family of possibly sign-changing solutions of (1.3) satisfying (1.4):

$$u_{\varepsilon} = cB_{\varepsilon} + o(1)$$
 in $H^1(M)$,

for some c > 0, where B_{ε} is a family of bubbles as in Definition 3.1 centered at x_{ε} with radius μ_{ε} and modeled on $V \in \Sigma$. In particular the analysis of Section 4 applies: Proposition 4.1, (4.34) and (4.42) show that there exists a family $(\sigma_{\varepsilon})_{0 < \varepsilon \leq 1}$ with $\sigma_{\varepsilon} \to 0$ as $\varepsilon \to 0$ and a positive constant C such that, for any $x \in M$ and any $\varepsilon > 0$,

$$\begin{aligned} \left| u_{\varepsilon}(x) - B^{\mu_{\varepsilon}, x_{\varepsilon}}(x) \right| &\leq \sigma_{\varepsilon} B_{\varepsilon}^{+}(x), \\ (5.1) \qquad \left| \nabla u_{\varepsilon}(x) - \nabla B^{\mu_{\varepsilon}, x_{\varepsilon}}(x) \right| &\leq \sigma_{\varepsilon} \frac{\mu_{\varepsilon}^{\frac{n-2}{2}}}{(\mu_{\varepsilon} + d_{g}(x_{\varepsilon}, x))^{n-1}} + C\mu_{\varepsilon}^{\frac{n-2}{2}} \quad \text{and} \\ (\mu_{\varepsilon} + d_{g}(x_{\varepsilon}, x))^{k} \left| \nabla^{k} u_{\varepsilon}(x) \right| &\leq C B_{\varepsilon}^{+}(x) \quad \text{for } k = 1, 2 \end{aligned}$$

hold, where B_{ε}^+ and $B^{\mu_{\varepsilon},x_{\varepsilon}}$ are as in (3.5) and (4.2). Equation (4.35) also applies and shows that $2^* - p_{\varepsilon} = O(\mu_{\varepsilon})$. In this section we will prove Theorem 1.1 using (5.1). We first recall the classical conformal normal coordinates result of Lee-Parker

[18]. It states the existence of a positive function $\Lambda \in C^{\infty}(M \times M)$ such that, letting $\Lambda_y(x) = \Lambda(y, x)$ for any $y, x \in M$ and $g_y := \Lambda_y^{\frac{4}{n-2}}g$ we have

(5.2)

$$\Lambda_{y}(x) = 1 + O\left(d_{g_{y}}(y, x)^{2}\right),$$

$$\operatorname{Ric}_{g_{y}}(x) = O\left(d_{g_{y}}(y, x)\right)$$

$$S_{g_{y}} = O\left(d_{g_{y}}(y, x)^{2}\right) \quad \text{and}$$

$$\sqrt{|g_{y}|}(x) = 1 + O\left(d_{g_{y}}(y, x)\right)^{N}$$

as $d_{g_y}(y,x) \to 0$, for a fixed N large enough, where Ric_{g_y} denotes the Ricci tensor of g_y and where it is understood that $\sqrt{|g_y|}$ is computed with respect to the exponential map of g_y at y. For $\varepsilon > 0$ we let $g_{x_{\varepsilon}} = \Lambda_{x_{\varepsilon}}^{\frac{4}{n-2}}g$ and we let $\exp_{x_{\varepsilon}}^{g_{x_{\varepsilon}}}$ be the exponential chart of $g_{x_{\varepsilon}}$ at x_{ε} . Let $\delta > 0$ be such that $2\delta < \inf_{\xi \in M} i_{g_{\xi}}(M)$. We define g_{ε} and v_{ε} in $B(0, 2\delta) \subset \mathbb{R}^n$ as follows:

(5.3)
$$g_{\varepsilon} = \left(\exp_{x_{\varepsilon}}^{g_{x_{\varepsilon}}}\right)^* g_{x_{\varepsilon}} \quad \text{and} \\ v_{\varepsilon}(x) = \frac{u_{\varepsilon}}{\Lambda_{x_{\varepsilon}}} \circ \exp_{x_{\varepsilon}}^{g_{x_{\varepsilon}}}.$$

First, (5.2) shows that, for any fixed $x \in M$, we have

(5.4)
$$d_{g_x}(x,y) = d_g(x,y) + O\left(d_g(x,y)^3\right) \text{ as } d_g(x,y) \to 0.$$

The latter with (2.8), (4.2) and (5.1) implies that

(5.5)
$$\begin{aligned} \left| v_{\varepsilon}(x) - \tilde{V}_{\varepsilon}(x) \right| &\leq \sigma_{\varepsilon} \tilde{B}_{\varepsilon}^{+}(x) + C\mu_{\varepsilon}^{\frac{n-2}{2}}, \\ \left| \nabla v_{\varepsilon}(x) - \nabla \tilde{V}_{\varepsilon}(x) \right| &\leq \sigma_{\varepsilon} \frac{\mu_{\varepsilon}^{\frac{n-2}{2}}}{(\mu_{\varepsilon} + |x|)^{n-1}} + C\mu_{\varepsilon}^{\frac{n-2}{2}} \quad \text{and} \\ (\mu_{\varepsilon} + d_{g_{\varepsilon}}(x_{\varepsilon}, x))^{k} \left| \nabla^{k} v_{\varepsilon}(x) \right| &\leq C \tilde{B}_{\varepsilon}^{+}(x) \quad \text{for } k = 1, 2 \end{aligned}$$

for any $x \in B(0, 2\delta)$, where C > 0 is independent of ε and where we have let

$$\tilde{B}_{\varepsilon}^{+}(x) = \frac{\mu_{\varepsilon}^{\frac{n-2}{2}}}{\left(\mu_{\varepsilon}^{2} + \frac{|x|^{2}}{n(n-2)}\right)^{\frac{n-2}{2}}} \quad \text{and} \quad \tilde{V}_{\varepsilon}(x) = \mu_{\varepsilon}^{-\frac{n-2}{2}} V\left(\frac{x}{\mu_{\varepsilon}}\right).$$

Moreover, it follows from (4.9) that

(5.6)
$$\lim_{\varepsilon \to 0} \hat{v}_{\varepsilon} = V \text{ in } C^2_{loc}(\mathbb{R}^n), \text{ where } \hat{v}_{\varepsilon}(x) := \mu_{\varepsilon}^{\frac{n-2}{2}} v_{\varepsilon}(\mu_{\varepsilon} x)$$

Using (1.3) and the conformal invariance property of the conformal Laplacian it is easily seen that v_ε satisfies

(5.7)
$$\Delta_{\xi} v_{\varepsilon} + A_{\varepsilon}(v_{\varepsilon}) + \hat{h}_{\varepsilon} v_{\varepsilon} = \Lambda_{\varepsilon}^{p_{\varepsilon} - 2^{\star}} |v_{\varepsilon}|^{p_{\varepsilon} - 2} v_{\varepsilon} \quad \text{in } B(0, 2\delta),$$

where $\Lambda_{\varepsilon} := \Lambda_{x_{\varepsilon}} \circ \exp_{x_{\varepsilon}}^{g_{x_{\varepsilon}}}$, ξ denotes the Euclidean metric and where we have let $\Delta_{\xi} = -\sum_{i=1}^{n} \partial_i^2$,

(5.8)
$$\hat{h}_{\varepsilon}(x) = \left[\Lambda_{x_{\varepsilon}}^{2-2^{\star}}\left(h_{\varepsilon} - \frac{n-2}{4(n-1)}S_g\right)\right]\left(\exp_{x_{\varepsilon}}^{g_{x_{\varepsilon}}}(x)\right)$$

and

(5.9)
$$A_{\varepsilon}(v_{\varepsilon}) = (\Delta_{g_{\varepsilon}} - \Delta_{\xi}) v_{\varepsilon} + \frac{n-2}{4(n-1)} S_{g_{\varepsilon}} v_{\varepsilon}.$$

Integrating (5.7) against $\langle x, \nabla v_{\varepsilon} \rangle + \frac{n-2}{2} v_{\varepsilon}$ in $B(0, \delta)$ with respect to the Euclidean metric yields the following Pohozaev identity (see Marques [21, Lemma 2.1]):

$$\int_{\partial B(0,\delta)} \left(\frac{n-2}{2} v_{\varepsilon} \partial_{\nu} v_{\varepsilon} - \frac{\delta}{2} |\nabla v_{\varepsilon}|_{\xi}^{2} + \delta(\partial_{\nu} v_{\varepsilon})^{2} + \frac{1}{p_{\varepsilon}} \Lambda_{\varepsilon}^{p_{\varepsilon}-2^{\star}} \delta |v_{\varepsilon}|^{p_{\varepsilon}} \right) d\sigma$$

$$= \frac{1}{p_{\varepsilon}} \int_{B(0,\delta)} (p_{\varepsilon} - 2^{\star}) x^{k} \partial_{k} \Lambda_{\varepsilon} \Lambda_{x_{\varepsilon}}^{p_{\varepsilon}-1-2^{\star}} |v_{\varepsilon}|^{p_{\varepsilon}} dx$$

$$+ n \left(\frac{1}{p_{\varepsilon}} - \frac{1}{2^{\star}} \right) \int_{B(0,\delta)} \Lambda_{\varepsilon}^{p_{\varepsilon}-2^{\star}} |v_{\varepsilon}|^{p_{\varepsilon}} dx$$

$$+ \int_{B(0,\delta)} \left(x^{k} \partial_{k} v_{\varepsilon} + \frac{n-2}{2} v_{\varepsilon} \right) A_{\varepsilon}(v_{\varepsilon}) dx$$

$$+ \int_{B(0,\delta)} \left(x^{k} \partial_{k} v_{\varepsilon} + \frac{n-2}{2} v_{\varepsilon} \right) \hat{h}_{\varepsilon} v_{\varepsilon} dx.$$

In all the computations of this section it is intended that all the terms involving $p_{\varepsilon} - 2^*$ disappear in the case where $p_{\varepsilon} = 2^*$ for any $\varepsilon > 0$. We now estimate all the

integrals appearing in (5.10). First, straightforward computations using (5.2) and (5.5) show that

(5.11)
$$\frac{1}{p_{\varepsilon}} \int_{B(0,\delta)} (p_{\varepsilon} - 2^{\star}) x^{k} \partial_{k} \Lambda_{\varepsilon} \Lambda_{\varepsilon}^{p_{\varepsilon} - 1 - 2^{\star}} |v_{\varepsilon}|^{p_{\varepsilon}} dx + n \left(\frac{1}{p_{\varepsilon}} - \frac{1}{2^{\star}}\right) \int_{B(0,\delta)} \Lambda_{\varepsilon}^{p_{\varepsilon} - 2^{\star}} |v_{\varepsilon}|^{p_{\varepsilon}} dx = \frac{(n-2)^{2}}{4n} (2^{\star} - p_{\varepsilon}) \int_{\mathbb{R}^{n}} |V|^{2^{\star}} dx + o \left(|p_{\varepsilon} - 2^{\star}|\right)$$

and that

(5.12)
$$\int_{\partial B(0,\delta)} \left(\frac{n-2}{2} v_{\varepsilon} \partial_{\nu} v_{\varepsilon} - \frac{\delta}{2} |\nabla v_{\varepsilon}|_{\xi}^{2} + \delta (\partial_{\nu} v_{\varepsilon})^{2} + \frac{1}{p_{\varepsilon}} \Lambda_{\varepsilon}^{p_{\varepsilon}-2^{\star}} \delta |v_{\varepsilon}|^{p_{\varepsilon}} \right) d\sigma$$
$$= O(\mu_{\varepsilon}^{n-2}) = \begin{cases} o(\mu_{\varepsilon}^{2}) & \text{if } n \ge 5\\ O(\mu_{\varepsilon}^{2}) & \text{if } n = 4 \end{cases}$$

as $\varepsilon \to 0$. A simple integration by parts shows that

(5.13)
$$\int_{B(0,\delta)} \left(x^k \partial_k v_{\varepsilon} + \frac{n-2}{2} v_{\varepsilon} \right) \hat{h}_{\varepsilon} v_{\varepsilon} dx$$
$$= \frac{1}{2} \int_{\partial B(0,\delta)} \delta \hat{h}_{\varepsilon} v_{\varepsilon}^2 d\sigma - \int_{B(0,\delta)} \left(\hat{h}_{\varepsilon} + \frac{1}{2} x^k \partial_k \hat{h}_{\varepsilon} \right) v_{\varepsilon}^2 dx,$$

so that with (5.5), we get that

(5.14)
$$\int_{B(0,\delta)} \left(x^k \partial_k v_{\varepsilon} + \frac{n-2}{2} v_{\varepsilon} \right) \hat{h}_{\varepsilon} v_{\varepsilon} dx$$
$$= -\int_{B(0,\delta)} \left(\hat{h}_{\varepsilon} + \frac{1}{2} x^k \partial_k \hat{h}_{\varepsilon} \right) v_{\varepsilon}^2 dx + \begin{cases} o(\mu_{\varepsilon}^2) & \text{if } n \ge 5\\ O(\mu_{\varepsilon}^2) & \text{if } n = 4 \end{cases}$$
$$= -\int_{B(0,\delta)} \hat{h}_{\varepsilon} v_{\varepsilon}^2 dx + \begin{cases} o(\mu_{\varepsilon}^2) & \text{if } n \ge 5\\ O(\mu_{\varepsilon}^2) & \text{if } n = 4 \end{cases}$$

Let now $w \in C^{\infty}(B(0, 2\delta))$ be a smooth function. We have, for any $\varepsilon > 0$ and by (5.2),

$$\Delta_{g_{\varepsilon}}w - \Delta_{\xi}w = -\left(g_{\varepsilon}^{ij} - \delta^{ij}\right)\partial_{ij}^{2}w - \partial_{i}(g_{\varepsilon})^{ij}\partial_{j}w + O\left(|x|^{N-1}|\nabla w|_{\xi}\right)$$

where repeated indices are summed over. By (5.2) we have $\operatorname{Ric}_{g_{\varepsilon}}(x_{\varepsilon}) = 0$. Cartan's expansion of g_{ε} and the symmetries of the Riemann tensor show that for any $1 \leq j \leq n$

$$\partial_i (g_{\varepsilon})^{ij}(x) = -\frac{1}{3} \operatorname{Ric}_{g_{\varepsilon}}(x_{\varepsilon})_{pj} x^p + O(|x|^2) = O(|x|^2).$$

By (5.2) again we also have $S_g(x_{\varepsilon}) = 0$, so that $\operatorname{Rm}_{g_{\varepsilon}}(x_{\varepsilon}) = \operatorname{Weyl}_{g_{\varepsilon}}(x_{\varepsilon})$, where $\operatorname{Weyl}_{g_{\varepsilon}}$ is the (4,0) Weyl tensor of g_{ε} . Using again Cartan's expansion of g_{ε} we find that for any smooth function w in $B(0, 2\delta)$, we have

(5.15)
$$\Delta_{g_{\varepsilon}}w - \Delta_{\xi}w = \frac{1}{3} \operatorname{Weyl}_{g_{\varepsilon}}(x_{\varepsilon})_{ipqj} x^{p} x^{q} \partial_{ij}^{2} w + O\left(|x|^{3} |\nabla^{2} w|\right) + O\left(|x|^{2} |\nabla w|\right).$$

Applying the latter to v_{ε} and using (5.5) shows that

$$A_{\varepsilon}(v_{\varepsilon}) = \frac{1}{3} \operatorname{Weyl}_{g_{\varepsilon}}(x_{\varepsilon})_{ipqj} x^{p} x^{q} \partial_{ij}^{2} v_{\varepsilon} + O\left(\frac{\mu_{\varepsilon}^{\frac{n-2}{2}}}{(\mu_{\varepsilon} + |x|)^{n-3}}\right).$$

Integrating the latter against $x^k \partial_k v_{\varepsilon} + \frac{n-2}{2} v_{\varepsilon}$ thus gives, using again (5.5), that

(5.16)
$$\begin{aligned} \int_{B(0,\delta)} \left(x^k \partial_k v_{\varepsilon} + \frac{n-2}{2} v_{\varepsilon} \right) A_{\varepsilon}(v_{\varepsilon}) dx \\ &= \frac{1}{3} \operatorname{Weyl}_{g_{\varepsilon}}(x_{\varepsilon})_{ipqj} \int_{B(0,\delta)} x^p x^q \partial_{ij}^2 v_{\varepsilon} \left(x^k \partial_k v_{\varepsilon} + \frac{n-2}{2} v_{\varepsilon} \right) dx \\ &+ \begin{cases} o(\mu_{\varepsilon}^2) & \text{if } n \ge 5\\ O(\mu_{\varepsilon}^2) & \text{if } n = 4 \end{cases} \end{aligned}$$

where again repeated indices are summed over. We now separate between the cases n = 3, n = 4 and $n \ge 5$ and prove Theorems 1.1 and 1.2.

5.1. The case $n \ge 5$. With \hat{v}_{ε} and \hat{h}_{ε} as in (5.6) and (5.8) a change of variables gives

$$\int_{B(0,\delta)} \hat{h}_{\varepsilon} dx = \mu_{\varepsilon}^2 \int_{B(0,\mu_{\varepsilon}^{-1}\delta)} \hat{h}_{\varepsilon}(\mu_{\varepsilon} x) \hat{v}_{\varepsilon}^2 dx.$$

It follows from (5.5) that $|\hat{v}_{\varepsilon}(x)| \leq C(1+|x|)^{2-n} \in L^2(\mathbb{R}^n)$ for $n \geq 5$. Therefore, using (5.6) and Lebesgue's convergence theorem, we get that (5.14) yields

(5.17)
$$\int_{B(0,\delta)} \left(x^k \partial_k v_{\varepsilon} + \frac{n-2}{2} v_{\varepsilon} \right) \hat{h}_{\varepsilon} v_{\varepsilon} dx$$
$$= -\left(h(x_0) - \frac{n-2}{4(n-1)} S_g(x_0) \right) \int_{\mathbb{R}^n} V^2 dx \cdot \mu_{\varepsilon}^2 + o(\mu_{\varepsilon}^2)$$

as $\varepsilon \to 0$, where $x_0 = \lim_{\varepsilon \to 0} x_{\varepsilon}$. This limit exists up to passing to a subsequence as $\varepsilon \to 0$, which will be implicit from now on. Note that the integral $\int_{\mathbb{R}^n} V^2 dx$ is finite by (2.8) since $n \ge 5$.

We now deal with (5.16), still for $n \geq 5$. By construction g_{ε} converges towards $\left(\exp_{x_0}^{g_{x_0}}\right)^* g_{x_0}$ in $C^1(\overline{B(0,\delta)})$ as $\varepsilon \to 0$, so that $\operatorname{Weyl}_{g_{\varepsilon}}(x_{\varepsilon})_{ipqj} \to \operatorname{Weyl}_{g_{x_0}}(x_0)_{ipqj}$ as $\varepsilon \to 0$, where the components of $\operatorname{Weyl}_{g_{x_0}}$ are measured in the exponential chart of g_{x_0} at x_0 . Since g_{x_0} is conformal to g and since $\Lambda_{x_0}(x_0) = 1$ we have $g_{x_0}(x_0) = g(x_0)$, and by the conformal invariance of the Weyl tensor we also have

$$\operatorname{Weyl}_{g_{x_0}}(x_0)_{ipqj} = \operatorname{Weyl}_g(x_0)_{ipqj},$$

where the components of $\operatorname{Weyl}_g(x_0)$ in the right-hand side are measured in the exponential chart of g at x_0 . By (5.5) and the dominated convergence theorem we therefore obtain, with (5.16), that

$$(5.18) \qquad \int_{B(0,\delta)} \left(x^k \partial_k v_{\varepsilon} + \frac{n-2}{2} v_{\varepsilon} \right) A_{\varepsilon}(v_{\varepsilon}) dx$$
$$= \frac{1}{3} \operatorname{Weyl}_g(x_0)_{ipqj} \int_{\mathbb{R}^n} x^p x^q \partial_{ij}^2 V \left(x^k \partial_k V + \frac{n-2}{2} V \right) dx \cdot \mu_{\varepsilon}^2 + o(\mu_{\varepsilon}^2)$$
$$= -\frac{(n-2)^2}{4n} \left(\int_{\mathbb{R}^n} |V|^{2^*} dx \right) \operatorname{Weyl}_g \otimes B \cdot \mu_{\varepsilon}^2 + o(\mu_{\varepsilon}^2)$$

as $\varepsilon \to 0$, where we used the Definition 3.2 of Weyl_g $\otimes B$. Using (4.35) and plugging (5.11), (5.12), (5.14), (5.17), (5.18) in (5.10) proves (1.5) and concludes the proof of Theorem 1.1.

5.2. The case n = 4. It follows from (2.4), (4.4), (5.2) and (5.5) that

(5.19)
$$\lim_{R \to +\infty} \limsup_{\varepsilon \to 0} \sup_{x \in B(0, R^{-1}) \setminus B(0, R\mu_{\varepsilon})} \left| \frac{|x|^2 v_{\varepsilon}(x)}{\mu_{\varepsilon}} - \lambda(V) \right| \text{ and} \\\lim_{R \to +\infty} \limsup_{\varepsilon \to 0} \sup_{x \in B(0, R^{-1}) \setminus B(0, R\mu_{\varepsilon})} \left| \frac{|x|^3 \nabla v_{\varepsilon}(x)}{\mu_{\varepsilon}} + 2\lambda(V) \frac{x}{|x|} \right| = 0.$$

For any fixed R > 0 we have, by (5.5) and (5.19),

$$\int_{B(0,\delta)} \hat{h}_{\varepsilon} v_{\varepsilon}^2 dx = O(R^4 \mu_{\varepsilon}^2) + \left(\hat{h}_{\varepsilon}(0) + \varepsilon_R\right) \mu_{\varepsilon}^2 \int_{B(0,R^{-1}) \setminus B(0,R\mu_{\varepsilon})} \frac{\lambda(V)^2}{|x|^4} dx$$

where $\varepsilon_R \to 0$ as $R \to +\infty$. As a consequence,

$$\int_{B(0,\delta)} \hat{h}_{\varepsilon} v_{\varepsilon}^2 dx = \left(\lambda(V)^2 \omega_3 \hat{h}_{\varepsilon}(0) + o(1)\right) \mu_{\varepsilon}^2 \ln \frac{1}{\mu_{\varepsilon}}$$

as $\varepsilon \to 0$. By (5.8) and going back to (5.14) we therefore have, when n = 4,

(5.20)
$$\int_{B(0,\delta)} \left(x^k \partial_k v_{\varepsilon} + v_{\varepsilon} \right) \hat{h}_{\varepsilon} v_{\varepsilon} dx$$
$$= -\lambda (V)^2 \omega_3 \left(h(x_0) - \frac{1}{6} S_g(x_0) \right) \mu_{\varepsilon}^2 \ln \frac{1}{\mu_{\varepsilon}} + o\left(\mu_{\varepsilon}^2 \ln \frac{1}{\mu_{\varepsilon}} \right)$$

as $\varepsilon \to 0$, where as before we have let $x_0 = \lim_{\varepsilon \to 0} x_{\varepsilon}$. We are left with the term (5.16). Integrating by parts and using (5.5) shows that

(5.21)
$$\int_{B(0,\delta)} x^p x^q \partial_{ij}^2 v_{\varepsilon} \left(x^k \partial_k v_{\varepsilon} + \frac{n-2}{2} v_{\varepsilon} \right) dx$$
$$= \int_{B(0,\delta)} x^p x^q \partial_i v_{\varepsilon} \partial_j v_{\varepsilon} dx + O(\mu_{\varepsilon}^2).$$

Arguing as above we get by (5.5) and (5.19) that

$$\begin{split} \operatorname{Weyl}_{g}(x_{0})_{ipqj} & \int_{B(0,\delta)} x^{p} x^{q} \partial_{i} v_{\varepsilon} \partial_{j} v_{\varepsilon} \, dx \\ &= O(R^{4} \mu_{\varepsilon}^{2}) + \operatorname{Weyl}_{g}(x_{0})_{ipqj} \int_{B(0,R^{-1}) \setminus B(0,R\mu_{\varepsilon})} x^{p} x^{q} \partial_{i} v_{\varepsilon} \partial_{j} v_{\varepsilon} dx \\ &= 4\lambda(V)^{2} \mu_{\varepsilon}^{2} \operatorname{Weyl}_{g}(x_{0})_{ipqj} \int_{B(0,R^{-1}) \setminus B(0,R\mu_{\varepsilon})} x^{p} x^{q} \frac{x_{i}}{|x|^{4}} \frac{x_{j}}{|x|^{4}} \, dx \\ &+ O(R^{4} \mu_{\varepsilon}^{2}) + O\left(\varepsilon_{R} \mu_{\varepsilon}^{2} \ln \frac{1}{\mu_{\varepsilon}}\right) \\ &= O(R^{4} \mu_{\varepsilon}^{2}) + O\left(\varepsilon_{R} \mu_{\varepsilon}^{2} \ln \frac{1}{\mu_{\varepsilon}}\right) \end{split}$$

where as before $\lim_{R\to+\infty} \varepsilon_R = 0$ and where the last term involving $\operatorname{Weyl}_g(x_0)$ vanishes by antisymmetry of $\operatorname{Weyl}_a(x_0)$. Going back to (5.16) with (5.21) we have

proven that

(5.22)
$$\int_{B(0,\delta)} \left(x^k \partial_k v_{\varepsilon} + \frac{n-2}{2} v_{\varepsilon} \right) A_{\varepsilon}(v_{\varepsilon}) dx = o\left(\mu_{\varepsilon}^2 \ln \frac{1}{\mu_{\varepsilon}} \right)$$

as $\varepsilon \to 0$. Remark that by (2.9) we have $2\omega_3\lambda(V) = \int_{\mathbb{R}^4} |V|^2 V dx$. With this observation, using (4.35) and plugging (5.11), (5.12), (5.14), (5.20) and (5.22) in (5.10), we get (1.6), which proves Theorem 1.2 for n = 4.

5.3. The case n = 3. We are left with the case n = 3. Using (5.11) and (5.13) the Pohozaev identity (5.10) writes as

(5.23)
$$\int_{\partial B(0,\delta)} \left(\frac{1}{2} v_{\varepsilon} \partial_{\nu} v_{\varepsilon} - \frac{\delta}{2} |\nabla v_{\varepsilon}|_{\xi}^{2} + \delta (\partial_{\nu} v_{\varepsilon})^{2} + \frac{1}{p_{\varepsilon}} \Lambda_{x_{\varepsilon}}^{p_{\varepsilon}-6} \delta |v_{\varepsilon}|^{p_{\varepsilon}} \right) d\sigma$$
$$= \frac{1}{12} (6 - p_{\varepsilon}) \int_{\mathbb{R}^{n}} |V|^{6} dx + o \left(|p_{\varepsilon} - 6|\right)$$
$$+ \int_{B(0,\delta)} \left(x^{k} \partial_{k} v_{\varepsilon} + \frac{1}{2} v_{\varepsilon} \right) \tilde{A}_{\varepsilon}(x) dx$$

with

$$\tilde{A}_{\varepsilon}(x) := \left(\Delta_{g_{\varepsilon}} - \Delta_{\xi}\right) v_{\varepsilon} + \left(\frac{1}{8}S_{g_{\varepsilon}} + \hat{h}_{\varepsilon}\right) v_{\varepsilon},$$

where \hat{h}_{ε} is given by (5.8) and where v_{ε} is defined in (5.3). First, by using (5.5) together with (5.15), we see that

$$|\tilde{A}_{\varepsilon}(x)| \lesssim \frac{\mu_{\varepsilon}^{\frac{1}{2}}}{\mu_{\varepsilon} + |x|} \quad \text{for } x \in B(0, \delta).$$

As a consequence, straightforward computations show that

(5.24)
$$\left| \int_{B(0,\delta)} \left(x^k \partial_k v_{\varepsilon} + \frac{1}{2} v_{\varepsilon} \right) \tilde{A}_{\varepsilon}(x) \, dx \right| = O\left(\delta \mu_{\varepsilon} \right)$$

as $\varepsilon \to 0$. Independently, (5.5) shows that

(5.25)
$$\left| \int_{\partial B(0,\delta)} \frac{1}{p_{\varepsilon}} \Lambda_{x_{\varepsilon}}^{p_{\varepsilon}-6} \delta |v_{\varepsilon}|^{p_{\varepsilon}} d\sigma \right| = o(\mu_{\varepsilon})$$

as $\varepsilon \to 0$. We now compute the remaining boundary term. It follows from (4.4), (5.2), (5.5), (5.7) and standard elliptic theory that

(5.26)
$$\lim_{\varepsilon \to 0} \frac{v_{\varepsilon}(x)}{\mu_{\varepsilon}^{\frac{1}{2}}} = \lambda \hat{G}(x) \quad \text{in } C^2_{loc}(B_{2\delta}(0) \setminus \{0\}),$$

where we have let

$$\lambda := 4\pi\lambda(V)$$

/ \

and

(5.27)

$$\hat{G}(x) := \frac{G_h(x_0, \exp_{x_0}^{g_{x_0}}(x))}{\Lambda_{x_0}(\exp_{x_0}^{g_{x_0}}(x))} \quad \text{for all } x \in B_{2\delta}(0) \setminus \{0\}.$$

Here we recall that $\lambda(V)$ is the first term in the expansion at infinity of V (see (2.4)), G_h is the Green's function for the operator $\Delta_g + h$ in M, Λ_{x_0} is as in (5.2)

and $\exp_{x_0}^{g_{x_0}}$ is the exponential map at x_0 with respect to g_{x_0} . It first follows from (5.26) that

(5.28)
$$\lim_{\varepsilon \to 0} \mu_{\varepsilon}^{-1} \int_{\partial B(0,\delta)} \left(\frac{1}{2} v_{\varepsilon} \partial_{\nu} v_{\varepsilon} - \frac{\delta}{2} |\nabla v_{\varepsilon}|_{\xi}^{2} + \delta (\partial_{\nu} v_{\varepsilon})^{2} \right) d\sigma$$
$$= \lambda^{2} \int_{\partial B(0,\delta)} \left(\frac{1}{2} \hat{G} \partial_{\nu} \hat{G} - \frac{\delta}{2} |\nabla \hat{G}|_{\xi}^{2} + \delta (\partial_{\nu} \hat{G})^{2} \right) d\sigma.$$

By (5.2) and (5.4), and since $G_h(x_0, \cdot)$ satisfies the expansion (3.8), there exists a continuous function $\hat{\beta} : B(0, 2\delta) \to \mathbb{R}$ such that

$$\hat{G}(x) = \frac{1}{4\pi|x|} + \hat{\beta}(x) \quad \text{for all } x \in B_{2\delta}(0) \setminus \{0\}.$$

It follows from Definition 3.3 that $\hat{\beta}(0) = m_h(x_0)$ and (3.9) and (5.4) show that we have

$$|\hat{\beta}(x)| \le C; \ |\nabla\hat{\beta}(x)| \le C(1+|\ln|x||) \quad \text{for all } x \in B_{2\delta}(0) \setminus \{0\}.$$

Straightforward computations then show that

(5.29)
$$\int_{\partial B(0,\delta)} \left(\frac{1}{2} \hat{G} \partial_{\nu} \hat{G} - \frac{\delta}{2} |\nabla \hat{G}|_{\xi}^2 + \delta (\partial_{\nu} \hat{G})^2 \right) d\sigma = -\frac{1}{2} m_h(x_0) + \varepsilon_{\delta},$$

where $\lim_{\delta\to 0} \varepsilon_{\delta} = 0$. Using (4.35), plugging (5.24), (5.25), (5.28) and (5.29) in (5.23) and using (5.27) finally shows that

(5.30)
$$\lim_{\varepsilon \to 0} \frac{6 - p_{\varepsilon}}{\mu_{\varepsilon}} = \frac{-96\pi^2 \lambda(V)^2 m_h(x_0)}{\int_{\mathbb{R}^3} |V|^6 dx}$$

This proves (1.7) and concludes the proof of Theorem 1.2. \Box

6. A NON-BLOWUP SITUATION

In this last Section we show how Theorem 1.1 can be used to rule out the existence of one-bubble blowing-up solutions of equations like:

(6.1)
$$\Delta_g u_{\varepsilon} + h u_{\varepsilon} = |u_{\varepsilon}|^{2^* - 2 - \varepsilon} u_{\varepsilon}$$

at a given point, in some cases. In [8], Deng-Musso-Wei have proposed a creative approach to construct families of sign-changing solutions $(u_{\varepsilon})_{\varepsilon>0}$ to the equation (6.1) that blow-up like a single bubble as in (1.4). The bubble that the authors choose in [8] is one of the symmetric bubbles constructed in [7]. In their arguments, Deng-Musso-Wei have introduced the function

$$\varphi_{\ell} := h - c_n \left(1 + \frac{n-4}{3n} \ell \right) S_g,$$

where ℓ is a parameter associated to the familie of bubbles in [7]. The function φ_{ℓ} plays a crucial role in the results of [8]. We show below how, under some conditions, it is possible to use φ_{ℓ} to rule out one-bubble blow-up.

We let (M, g) be a smooth, closed, connected Riemannian manifold of dimension $n \ge 5$ that is locally conformally flat and has constant scalar curvature equal to -1. This is for instance satisfied if (M, g) is a closed hyperbolic manifold. We define

$$c_n = \frac{n-2}{4(n-1)}$$
 and we let $1 < t < 1 + \frac{n-4}{3n}$.

Let $\xi_0 \in M$ be any point.

Step 6.1: We claim that there exists $h \in C^{\infty}(M)$ such that

(6.2)
$$\begin{cases} h(\xi_0) = -tc_n \\ \nabla h(\xi_0) = 0 \\ \nabla^2 h(x_0) \text{ is positive-definite} \\ \Delta_g + h \text{ is coercive in } H^1(M) \end{cases}$$

Under (6.2), ξ_0 is in particular a local strict minimum point of h.

We prove the Claim. Let for instance $\chi \in C_c^{\infty}(\mathbb{R}^n)$ be a cutoff function with $0 \leq \chi \leq 1, \, \chi(x) = 1$ in B(0,1) and $\chi(x) = 0$ in $\mathbb{R}^n \setminus B(0,2)$. Let, for $0 < \delta < \frac{i_g(M)}{2}$ and $x \in M$,

$$h_{\delta}(x) = 1 - \chi \left(\frac{d_g(\xi_0, x)}{\delta}\right) + \chi \left(\frac{d_g(\xi_0, x)}{\delta}\right) \left(-tc_n + d_g(\xi_0, x)^2\right).$$

We claim that h_{δ} satisfies (6.2) for δ small enough (but fixed). Indeed, in $B_g(\xi_0, \delta)$, we have $h_{\delta}(x) = -tc_n + d_g(\xi_0, x)^2$, and straightforward computations show that

$$\|h_{\delta} - 1\|_{L^{\frac{n}{2}}(M)} \le C\delta^2$$

so that $\Delta_g + h_{\delta}$ is coercive for δ small enough. This proves the claim.

Step 6.2: We now let $h \in C^{\infty}(M)$ satisfy (6.2). For $\ell \geq 1$ and $\xi \in M$ we define

$$\varphi_{\ell}(\xi) = h(\xi) - c_n \Big(1 + \frac{n-4}{3n} \ell \Big) S_g(\xi) = h(\xi) + c_n \Big(1 + \frac{n-4}{3n} \ell \Big),$$

We claim that

(6.3)
$$\begin{cases} \Delta_g + h \text{ is coercive;} \\ \varphi_\ell(\xi_0) > 0 \text{ for all } l \ge 1; \\ \xi_0 \text{ is a } C^1 \text{-stable critical point of } \varphi_\ell \text{ uniformly in } \ell \ge 1. \end{cases}$$

We prove the claim. We have that

$$\varphi_{\ell}(\xi_0) = c_n \left(1 + \frac{n-4}{3n}\ell - t \right) \ge c_n \left(1 + \frac{n-4}{3n} - t \right) > 0$$

for any $\ell \geq 1$. It is easily seen with (6.2) that, for any $\ell \geq 1$, ξ_0 is a critical point of φ_ℓ satisfying $\varphi_\ell(\xi_0) > 0$ and $\varphi_\ell(\xi_0) \to +\infty$ as $\ell \to +\infty$. Since ξ_0 has non-zero degree, it is a C^1 -stable critical point of h (following the terminology of [19]). That is, there is $r_0 > 0$ that satisfies the following: for any $0 < r \leq r_0$ there exists $\alpha > 0$ such that any $\tilde{h} \in C^1(\overline{B_g(\xi_0, r)})$ satisfying $\|\tilde{h} - h\|_{C^1(\overline{B_g(\xi_0, r)})} \leq \alpha$ has at least one critical point in $B_g(\xi_0, r)$. By definition of φ_ℓ we observe that ξ_0 then remains a C^1 -stable critical point of φ_ℓ uniformly in $\ell \geq 1$: that is, for any $0 < r \leq r_0$ and α as above, and for any $\ell \geq 1$, any function $\tilde{\varphi} \in C^1(\overline{B_g(\xi_0, r)})$ satisfying $\|\tilde{\varphi} - \varphi_\ell\|_{C^1(\overline{B_g(\xi_0, r)})} \leq \delta$ has at least one critical point in $B_g(\xi_0, r)$. This proves (6.3).

Step 6.3: We claim that there is no family $(u_{\varepsilon})_{0 < \varepsilon \leq 1} \in C^2(M)$ of solutions of (6.1) such that

(6.4)
$$u_{\varepsilon} = B_{\varepsilon} + o(1) \text{ in } H^1(M),$$

where $B = (B_{\varepsilon})_{\varepsilon>0}$ is a bubble centered at $(\xi_{\varepsilon})_{\varepsilon} \in M$, $\lim_{\varepsilon \to 0} \xi_{\varepsilon} = \xi_0$ as in Definition 3.1.

We prove the claim by contradiction. Since (M, g) is locally conformally flat, then the Weyl tensor vanishes. It then follows from Theorem 1.1 of the present paper that if there is blow-up like in (6.4), then equation (1.5) in particular tells us that

(6.5)
$$h(\xi_0) + c_n = h(\xi_0) - c_n S_g(\xi_0) \ge 0.$$

But this is an obvious contradiction with (6.2) since $h(\xi_0) = -tc_n$ and t > 1. This proves the claim.

References

- B. Ammann and E. Humbert, *The second Yamabe invariant*, J. Funct. Anal. 235 (2006), no. 2, 377–412. MR2225458
- [2] N. Aronszajn, A unique continuation theorem for solutions of elliptic partial differential equations or inequalities of second order, J. Math. Pures Appl. (9) 36 (1957), 235-249. MR92067
- [3] T. Aubin, Équations du type Monge-Ampère sur les variétés kählériennes compactes, Bull. Sci. Math. (2) 102 (1978), no. 1, 63–95.
- [4] Arthur L. Besse, *Einstein manifolds*, Classics in Mathematics, Springer-Verlag, Berlin, 2008. Reprint of the 1987 edition. MR2371700
- [5] Denis Bonheure, Jean-Baptiste Castéras, and Bruno Premoselli, *Classification of radial blow*up at the first critical exponent for the lin-ni-takagi problem in the ball (2022). Preprint.
- [6] Luis A. Caffarelli, Basilis Gidas, and Joel Spruck, Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth, Comm. Pure Appl. Math. 42 (1989), no. 3, 271–297. MR982351 (90c:35075)
- Manuel del Pino, Monica Musso, Frank Pacard, and Angela Pistoia, Large energy entire solutions for the Yamabe equation, J. Differential Equations 251 (2011), no. 9, 2568–2597. MR2825341 (2012k:35176)
- [8] Shengbing Deng, Monica Musso, and Juncheng Wei, New type of sign-changing blow-up solutions for scalar curvature type equations, Int. Math. Res. Not. IMRN 13 (2019), 4159–4197. MR3978436
- Wei Yue Ding, On a conformally invariant elliptic equation on Rⁿ, Comm. Math. Phys. 107 (1986), no. 2, 331–335. MR863646
- [10] Olivier Druet, From one bubble to several bubbles: the low-dimensional case, J. Differential Geom. 63 (2003), no. 3, 399–473. MR2015469
- [11] _____, Compactness for Yamabe metrics in low dimensions, Int. Math. Res. Not. 23 (2004), 1143–1191. MR2041549 (2005b:53056)
- [12] Olivier Druet, Emmanuel Hebey, and Frédéric Robert, Blow-up theory for elliptic PDEs in Riemannian geometry, Mathematical Notes, vol. 45, Princeton University Press, Princeton, NJ, 2004. MR2063399 (2005g:53058)
- [13] Thomas Duyckaerts, Carlos Kenig, and Frank Merle, Solutions of the focusing nonradial critical wave equation with the compactness property, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 15 (2016), 731–808. MR3495446
- [14] Nassif Ghoussoub, Saikat Mazumdar, and Frédéric Robert, Multiplicity and stability of the Pohozaev obstruction for Hardy-Schrödinger equations with boundary singularity, Mem. Amer. Math. Soc. 285 (2023), no. 1415, v+126. MR4599209
- [15] Emmanuel Hebey, Introduction à l'analyse non linéaire sur les varyétés, Diderot Editeur Arts Sciences, 1997.
- [16] _____, Compactness and stability for nonlinear elliptic equations, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2014. MR3235821
- [17] Emmanuel Hebey and Michel Vaugon, From best constants to critical functions, Math. Z. 237 (2001), no. 4, 737–767.
- [18] John M. Lee and Thomas H. Parker, *The Yamabe problem*, Bull. Amer. Math. Soc. (N.S.) 17 (1987), no. 1, 37–91. MR888880 (88f:53001)
- [19] YanYan Li, On a singularly perturbed elliptic equation, Adv. Differential Equations 2 (1997), no. 6, 955–980. MR1606351
- [20] Yanyan Li and Meijun Zhu, Yamabe type equations on three-dimensional Riemannian manifolds, Commun. Contemp. Math. 1 (1999), no. 1, 1–50. MR1681811 (2000m:53051)

- [21] Fernando Coda Marques, A priori estimates for the Yamabe problem in the non-locally conformally flat case, J. Differential Geom. 71 (2005), no. 2, 315–346. MR2197144 (2006i:53046)
- [22] Maria Medina and Monica Musso, Doubling nodal solutions to the Yamabe equation in ℝⁿ with maximal rank, J. Math. Pures Appl. (9) **152** (2021), 145–188. MR4280834
- [23] Maria Medina, Monica Musso, and Juncheng Wei, Desingularization of Clifford torus and nonradial solutions to the Yamabe problem with maximal rank, J. Funct. Anal. 276 (2019), no. 8, 2470–2523. MR3926123
- [24] Hussein Mesmar and Frédéric Robert, Concentration analysis for elliptic critical equations with no boundary control: Ground-state blow-up, Discrete and Continuous Dynamical Systems
 S 17 (2024), no. 4, 1599–1620.
- [25] Anna Maria Micheletti, Angela Pistoia, and Jérôme Vétois, Blow-up solutions for asymptotically critical elliptic equations on Riemannian manifolds, Indiana Univ. Math. J. 58 (2009), no. 4, 1719–1746. MR2542977
- [26] Morio Obata, The conjectures on conformal transformations of Riemannian manifolds, J. Differential Geometry 6 (1971/72), 247–258. MR303464
- [27] Angela Pistoia and Jérôme Vétois, Sign-changing bubble towers for asymptotically critical elliptic equations on Riemannian manifolds, J. Differential Equations 254 (2013), no. 11, 4245–4278. MR3035432
- [28] Bruno Premoselli, A priori estimates for finite-energy sign-changing solutions blowing-up solutions of critical elliptic equations, Int. Math. Res. Not. IMRN (2021). To appear.
- [29] Bruno Premoselli and Jérôme Vétois, Compactness of sign-changing solutions to scalar curvature-type equations with bounded negative part, J. Differential Equations 266 (2019), no. 11, 7416–7458. MR3926106
- [30] _____, Sign-changing blow-up for the Yamabe equation at the lowest energy level, Adv. Math. 410 (2022), Paper No. 108769, 50. MR4509411
- [31] Bruno Premoselli and Jérôme Vétois, Stability and instability results for sign-changing solutions to second-order critical elliptic equations, Journal de Mathématiques Pures et Appliquées 167 (2022), 257–293.
- [32] Frédéric Robert, Existence et asymptotiques optimales des fonctions de Green des opérateurs elliptiques d'ordre deux, 2010. https://iecl.univ-lorraine.fr/files/2021/04/ConstrucGreen.pdf.
- [33] Frédéric Robert and Jérôme Vétois, Sign-changing blow-up for scalar curvature type equations, Comm. Partial Differential Equations 38 (2013), no. 8, 1437–1465. MR3169751
- [34] _____, Sign-changing solutions to elliptic second order equations: glueing a peak to a degenerate critical manifold, Calc. Var. Partial Differential Equations 54 (2015), no. 1, 693–716. MR3385177
- [35] Richard M. Schoen, Lecture notes from courses at stanford (1988). Written by D. Pollack, preprint.
- [36] Michael Struwe, A global compactness result for elliptic boundary value problems involving limiting nonlinearities, Math. Z. 187 (1984), no. 4, 511–517. MR760051
- [37] Neil S. Trudinger, Remarks concerning the conformal deformation of Riemannian structures on compact manifolds, Ann. Scuola Norm. Sup. Pisa (3) 22 (1968), 265–274. MR0240748

Bruno Premoselli, Université Libre de Bruxelles, Service d'Analyse, CP 218, Boulevard du Triomphe, B-1050 Bruxelles, Belgique.

Email address: bruno.premoselli@ulb.be

FRÉDÉRIC ROBERT, UNIVERSITÉ DE LORRAINE, CNRS, IECL, F-54000 NANCY, FRANCE *Email address:* frederic.robert@univ-lorraine.fr