

Group Vertex Magicness of H -join and Generalised Friendship Graph

S. Balamoorthy, S.V. Bharanedhar *

Department of Mathematics,
Central University of Tamil Nadu,
Thiruvavur 610005,
India.

April 29, 2024

Abstract

Let A be a non-trivial Abelian group. A graph G is A -vertex magic if there exists a labeling $l : V(G) \rightarrow A \setminus \{0\}$ and μ in A such that $w(v) = \sum_{u \in N_G(v)} l(u) = \mu$ for any vertex v of G , where $N_G(v) = \{u \in V(G) : vu \in E(G)\}$ is the open neighborhood of the vertex v in G . In this paper, we characterize \mathbb{Z} -vertex magic graphs. We prove that G is A -vertex magic if its reduced graph is A -vertex magic, where A is a non-trivial Abelian group. We also construct a family of A -vertex magic graphs using H -generalized join operation. In addition, we introduce a new type of labeling called A' -vertex magic labeling of graphs and characterize A -vertex magicness using A' -vertex magicness. We give a new procedure to embed any graph as an induced subgraph of an A -vertex magic graph and we construct infinite families of A -vertex magic graphs both of these procedure uses less number of vertices compared to the one given in (Sabeel et al. in Australas. J. Combin. 85(1) (2023), 49-60). Moreover, we generalize some results proved in (Sabeel et al. in Australas. J. Combin. 85(1) (2023), 49-60). Finally we completely classify generalised friendship graph using A -vertex magicness and group vertex magicness.

Keywords: A -vertex magic; Group vertex magic; H -join; Generalised Friendship graph;

1 Introduction

All the graphs considered in this paper are connected simple finite graphs and A denote a non-trivial Abelian group, not necessarily finite. Let $G = (V(G), E(G))$ be a graph,

*Corresponding Author.

E-Mail addresses: moorthybala545@gmail.com (S. Balamoorthy), bharanedhar3@gmail.com (S.V. Bharanedhar)

we denote by $V(G)$ the vertex set of G . A graph G is said to be A -vertex magic if there exists a labeling $l : V(G) \rightarrow A \setminus \{0\}$ and μ in A such that $w(v) = \sum_{u \in N_G(v)} l(u) = \mu$ for any vertex v of G , where $N_G(v) = \{u \in V(G) : vu \in E(G)\}$ is the open neighborhood of the vertex v in G . The element μ is called a magic constant of the labeling l . A graph G that admits such a labeling is called an A -vertex magic graph. If G is an A -vertex magic graph for every non-trivial Abelian group A , then G is called a group vertex magic graph. We denote the degree of a vertex v in a graph G by $d_G(v)$. If $v \in V(G)$ is such that $d_G(v) = 1$, then v is called a pendant vertex of G . In this case, the vertex to which v is adjacent is called a support vertex. A vertex w is called a strong support vertex if w is adjacent to at least two pendant vertices.

The concept of group vertex magic graphs was introduced by Kamatchi et al. [7] and they have studied some properties of this magicness for trees with diameter at most 4. The main motivation of group vertex magic graph is group magic graph, which was introduced by Lee et al. [8] and this work was further studied by Lee et al. [9], Low and Lee [10]. In [11], Sabeel et al. have completely characterized A -vertex magic trees of diameter at most 5, where A is a finite Abelian group. In [2], the present authors have given a necessary condition for a graph to be A -vertex magic and they have obtained results for the A -vertex magicness of product of graphs.

In this paper, we use group elements to label the vertices of a graph and we have extended the study of group vertex magicness of a graph by considering an arbitrary Abelian group. This work considers the vertex magicness of H -join and generalised friendship graphs. We use the following important results in group theory namely Cauchy's theorem, Sylow's first theorem and fundamental theorem of finite Abelian groups. We refer to Bondy and Murty [4] for graph theoretic terminology and notations. We refer to Herstein [5] for ideas in algebra.

Let H be a graph with vertex set $\{v_1, v_2, \dots, v_k\}$ and let G_1, G_2, \dots, G_k be arbitrary graphs. The H -join operation of the graphs G_1, G_2, \dots, G_k is denoted as $H[G_1, G_2, \dots, G_k]$, is obtained by replacing the vertex v_i of H by the graph G_i for $1 \leq i \leq k$ and every vertex of G_i is made adjacent with every vertex of G_j , whenever v_i is adjacent to v_j in H . The vertex set $V(G) = \bigcup_{i=1}^k V(G_i)$ and edge set

$$E(G) = \left(\bigcup_{i=1}^k E(G_i) \right) \cup \left(\bigcup_{v_i v_j \in E(H)} \{uv : u \in V(G_i), v \in V(G_j)\} \right) \text{ (see [12] and [3]).}$$

If $G \cong G_i$, for $1 \leq i \leq k$, then $H[G_1, G_2, \dots, G_k] \cong H[G]$, the *lexicographic product* of H and G .

Let R be a commutative ring an element x of R is said to be a unit if it has a multiplicative inverse. A non-zero element u of R is said to be a zero-divisor if its product with a non-zero element yields zero. Let $Z(R)$ denotes set of all zero-divisor of R . Given a ring R , we associate a simple graph $\Gamma(R)$ to R with vertex set the set of all zero-divisor $Z(R)$ of R and two distinct vertices u and v are adjacent if and only

if $uv = 0$. This graph $\Gamma(R)$ is called the zero-divisor graph of R .

A caterpillar is a tree in which the removal of pendant vertices yields a path.

Observation 1. *If G is a connected graph with $|V(G)| \leq 3$, then G is A -vertex magic, where $|A| > 2$.*

The graph P_4 is not A -vertex magic graph with minimum number of edges.

Theorem 1. *Let G be a \mathbb{Z} -vertex magic graph. Then there exists $k \in \mathbb{N}$ such that G is \mathbb{Z}_m -vertex magic, for all $m \geq k$.*

Proof. Assume that l is \mathbb{Z} -vertex magic labeling of G . Then $l(v_i) = k_i$, where $k_i \in \mathbb{Z}, i = 1, 2, \dots, n$. Let $k = \max\{|k_i|\} + 1$. Now, define $l' : V(G) \rightarrow \mathbb{Z}_m \setminus \{0\}$ by $l'(v_i) = l(v_i)a$, where $o(a) = m$ and $m \geq k$. Clearly, the labeling l' is \mathbb{Z}_m -vertex labeling of G , for all $m \geq k$. \square

Theorem 2. *Let $m \in \mathbb{N}$ and l is a \mathbb{Z}_p -vertex magic labeling of G , for all primes $p \geq m$. Then G is \mathbb{Z} -vertex magic.*

Proof. Let us assume l is \mathbb{Z}_p -vertex magic labeling of G , for all prime $p \geq m$. Clearly, l is $\mathbb{Z}_{p'}$ -vertex magic, where $p' = \min\{p : p \geq m\}$. Then $l(v_i) = k_i a$, where $k_i \in \mathbb{Z}$ and $o(a) = p'$. Now, define $l' : V(G) \rightarrow \mathbb{Z} \setminus \{0\}$ by $l'(v_i) = k_i a$, here a is generator of \mathbb{Z} . Suppose l' is not a \mathbb{Z} -vertex magic labeling of G , then there exists $v_i, v_j \in V(G)$ such that $w(v_i) \neq w(v_j)$, which implies $w(v_i) \neq w(v_j)$ is also true for all \mathbb{Z}_{p_1} , where p_1 is a prime number, such that $p_1 > \max\{\deg(v_i) : v_i \in V(G)\} \cdot p'$, which is a contradiction. \square

Let G be a simple graph. In [1] and [3], the authors define the following relation on $V(G)$. For any two vertices $u, v \in V(G)$, define $u \sim_G v$ if and only if $N_G(u) = N_G(v)$. It is easy to see that, the relation \sim_G is an equivalence relation on $V(G)$. Let $[u]$ be the equivalence class which contains u and S be the set of all equivalence classes of this relation \sim_G . The reduced graph H of G is defined as follows. The *reduced graph* H of G is the graph with vertex set $V(H) = S$ and two distinct vertices $[u]$ and $[v]$ are adjacent in H if and only if u and v are adjacent in G . Note that, if $V(H) = \{[u_1], [u_2], \dots, [u_k]\}$, then G is the H -join of $\langle [u_1] \rangle, \langle [u_2] \rangle, \dots, \langle [u_k] \rangle$, that is, $G \cong H[\langle [u_1] \rangle, \langle [u_2] \rangle, \dots, \langle [u_k] \rangle]$ ($\langle [u] \rangle$ denote the subgraph induced by $[u]$) and each $[u_i]$ is an independent subset of G , we take $|[u_i]| = m_i$, where $m_i \in \mathbb{N}$ for each i . Clearly, H is isomorphic to a subgraph of G induced by $\{u_1, u_2, \dots, u_k\}$.

Theorem 3. *If the reduced graph H of G is A -vertex magic, where $|A| > 2$ then G is A -vertex magic.*

Proof. Assume that H is A -vertex magic with the corresponding labeling l such that $l([u_i]) = a_i$, where $a_i \in A \setminus \{0\}$ and $[u_i] = \{u_1^i, \dots, u_{m_i}^i\}$, for $i = 1, \dots, k$. Define the

label on $V(G)$ as follows, suppose m_i is odd, then define

$$l'(u_j^i) = \begin{cases} a_i & \text{if } j = 1 \\ a & \text{if } j > 1 \text{ and } j \text{ is odd} \\ -a & \text{if } j \text{ is even,} \end{cases}$$

for some $a \in A \setminus \{0\}$. Suppose m_i is even, then define

$$l'(u_j^i) = \begin{cases} a_i + a & \text{if } j = 1 \\ a & \text{if } j > 1 \text{ and } j \text{ is odd} \\ -a & \text{if } j \text{ is even,} \end{cases}$$

where $a_i \neq -a$. Let $v \in V(G)$. Then $v \in [u_i]$, for some i . As $w(v)$ (corresponding label l' in G) is equal to $w([u_i])$ (corresponding label l in H), G is A -vertex magic. \square

The following result proved in [2] is a consequence of Theorem 3.

Corollary 1. (Theorem 3, [2]) Let $G = K_{n_1, n_2, \dots, n_m}$ be a complete m -partite graph. Then G is A -vertex magic, where $|A| > 2$.

Proof. By the relation \sim_G on $V(G)$, we see that K_m is the reduced graph of G . Since K_m is A -vertex magic, where $|A| > 2$ and hence the result follows. \square

The following result proved in [11] is an immediate consequence of Corollary 1.

Corollary 2. (Corollary 2.4, [11]) For any finite Abelian group A with $|A| \geq 3$, all tree of diameter 2 are A -vertex magic.

The following corollaries are an immediate consequence of Theorem 3.

Corollary 3. If any connected graph G has at most three equivalence classes under the relation \sim_G , then the graph G is A -vertex magic, where $|A| > 2$.

Proof. By Theorem 3 and observation 1, we get the required result. \square

Corollary 4. Let $G = K_{n_1, n_2, \dots, n_m}$ be a complete m -partite graph. The $G + K_n$ is A -vertex magic.

Proof. The reduced graph H of $G + K_n$ is regular graph. By Theorem 3, the result follows. \square

Theorem 4. Let H be a simple graph on k vertices and let $G_j = K_{n_j}^c$, for all $j = 1, 2, \dots, k$. The graph $G = H[G_1, G_2, \dots, G_k]$ is A -vertex magic, where $|A| > 2$, if $n_j > 1$, for each $j = 1, 2, \dots, k$.

Proof. Let $V(G) = \bigcup_{j=1}^k V(K_{n_j}^c)$ and $v_i^j \in V(K_{n_j}^c)$, where $i = 1, 2, \dots, n_j$. Assume that each $n_j > 1$. Suppose n_j is odd, define $l : V(G) \rightarrow A \setminus \{0\}$ by

$$l(v_i^j) = \begin{cases} a + b & \text{if } i = 1 \\ -b & \text{if } i = 2 \\ -a & \text{if } i > 1 \text{ and } i \text{ is odd} \\ a & \text{if } i > 2 \text{ and } i \text{ is even,} \end{cases}$$

where $a \neq -b$.

Suppose n_j is even, define $l : V(G) \rightarrow A \setminus \{0\}$ by

$$l(v_i^j) = \begin{cases} a & \text{if } i \text{ is odd} \\ -a & \text{if } i \text{ is even.} \end{cases}$$

Thus $w(v) = 0$, for all $v \in V(G)$. □

The following result proved in [2] is an immediate consequence of Theorem 4.

Corollary 5. (Theorem 15, [2]) *Let G be any simple graph on n vertices. The graph $G[K_m^c]$ is A -vertex magic, $|A| > 2$ if and only if $m > 1$.*

Consider the zero-divisor graph $\Gamma(R)$, where R is a finite commutative reduced ring with unity. It is well-known that such a ring has to be a product of finite fields [3]. That is, there exist k finite fields $F_{q_1}, F_{q_2}, \dots, F_{q_k}$ such that $R \cong F_{q_1} \times F_{q_2} \times \dots \times F_{q_k}$, where q_i 's are prime powers, say $q_i = p_i^{n_i}$, $n_i \in \mathbb{N}$.

Corollary 6. *Let $q_i \geq 3$ be a prime power, for $1 \leq i \leq k$ and $R \cong F_{q_1} \times F_{q_2} \times \dots \times F_{q_k}$. Then the graph $\Gamma(R)$ is A -vertex magic, where $|A| > 2$.*

Proof. An easy observation, if $v \in V(\Gamma(R))$, then there exists a vertex $u \in V(\Gamma(R))$ such that $u \neq v$ and $N(u) = N(v)$. Therefore, each equivalence class (\sim_G) contains at least two element in $V(\Gamma(R))$. By Theorem 4, R is A -vertex magic, where $|A| > 2$. □

Definition 1. *Let G be a graph and let H be its reduced graph. We say that H is A' magic if there exists a labeling $l : V(H) \rightarrow A$ such that 0 can also be used to label the vertices $[u] \in V(H)$ whenever $|[u]| \geq 2$ and $\sum_{[u] \in N([v])} l([u]) = \mu$, for all $[v] \in V(H)$.*

Lemma 1. *Let A be an Abelian group with at least three elements. If $n \geq 2$ and $a \in A$, then there exists a_1, a_2, \dots, a_n in $A \setminus \{0\}$ such that $a = \sum_{i=1}^n a_i$.*

Proof. Consider the following cases.

Case 1. $a = 0$

Let $b, c \in A \setminus \{0\}$ and $b \neq -c$. If n is odd, define

$$a_i = \begin{cases} b + c & \text{if } i = 1 \\ -c & \text{if } i = 2 \\ -b & \text{if } i > 1 \text{ and } i \text{ is odd} \\ b & \text{if } i > 2 \text{ and } i \text{ is even.} \end{cases}$$

If n is even, define

$$a_i = \begin{cases} b & \text{if } i \text{ is odd} \\ -b & \text{if } i \text{ is even.} \end{cases}$$

Case 2. $a \neq 0$

Let $a, b \in A \setminus \{0\}$ and $a \neq -b$. If n is odd, define

$$a_i = \begin{cases} a & \text{if } i \text{ is odd} \\ -a & \text{if } i \text{ is even.} \end{cases}$$

If n is even, define

$$a_i = \begin{cases} a + b & \text{if } i = 1 \\ -b & \text{if } i = 2 \\ -a & \text{if } i > 1 \text{ and } i \text{ is odd} \\ a & \text{if } i > 2 \text{ and } i \text{ is even.} \end{cases}$$

□

The following result proved in [11] is an immediate consequence of Lemma 1.

Corollary 7. (Lemma 2.1 in [11]) *Let A be a finite Abelian group with $|A| \geq 3$ and let $g \in A$. Then, for each $n \geq 2$, there exist a_1, a_2, \dots, a_n in $A \setminus \{0\}$ such that $g = a_1 + a_2 + \dots + a_n$.*

Theorem 5. *A graph G is A -vertex magic, where $|A| > 2$ if and only if its reduced graph H of G is A' magic.*

Proof. Assume that G is A -vertex magic and l is corresponding A -vertex magic labeling. Define $l' : V(H) \rightarrow A$ by $l'([v]) = \sum_{\substack{u \in V(G) \\ u \sim_G v}} l(u)$. Then

$$\begin{aligned} w([v]) &= \sum_{[u] \in N([v])} l'([u]) \\ &= \sum_{[u] \in N([v])} \left(\sum_{\substack{u' \in V(G) \\ u' \sim_G u}} l(u') \right) \\ &= \sum_{u \in N(v)} l(u) \\ &= w(v) \end{aligned}$$

Since G is A -vertex magic, we have H is A' magic.

Conversely, assume that the reduced graph H of G is A' magic and the corresponding labeling be f . Define $f' : V(G) \rightarrow A \setminus \{0\}$ by if $|[v]| = 1$, then $f'(v) = f([v])$ and if $|[v]| \geq 2$, then we label the vertices of the class using Lemma 1 in such a way that sum

of the labels of all vertices in this equivalence class is equal to $f([v])$ in the reduced graph H of G .

$$\begin{aligned} w(v) &= \sum_{u \in N(v)} f'(u) \\ &= \sum_{[u] \in N([v])} f([u]) \\ &= w([v]). \end{aligned}$$

□

The following corollaries are immediate consequence of Theorem 5.

Corollary 8. *Let A be an Abelian group with at least three elements. If G is a graph in which every non-pendant vertex is a strong support vertex, then G is A -vertex magic.*

Proof. Let v_1, v_2, \dots, v_k be non-pendant vertices of G and v_{i_j} be the pendant vertices of G , which are adjacent to v_i , $j = 1, 2, \dots, m_i$, where $m_i \geq 2$ for all $i = 1, 2, \dots, k$. Let H be the reduced graph of G . Define $l : V(H) \rightarrow A \setminus \{0\}$ by

$$l([v_m]) = \begin{cases} a & \text{if } d_G(v_m) > 2 \\ -(d_H([v_i]) - 1)a & \text{if } v_m = v_{i_j} \text{ for some } i, j \end{cases}$$

where $a \in A \setminus \{0\}$. Thus $w([v]) = a$, for all $[v] \in V(H)$. By Theorem 5, we get the required result. □

The following result proved in [11] is an immediate consequence of corollary 8.

Corollary 9. *(Theorem 2.2 in [11]) Let A be a finite Abelian group with $|A| \geq 3$. If G is a graph in which every non-pendant vertex is a strong support vertex, then G is A -vertex magic.*

Corollary 10. *A caterpillar T is A -vertex magic, where $|A| > 2$ if and only if T has no vertex of degree two.*

Proof. Let T be a caterpillar. Assume that T has a non-corner vertex with degree 2, let v_2 be such a vertex. Also, in the reduced graph H , $\deg([v_2]) = 2$ (see Figure 1).

Suppose H is A' -vertex magic, then $w([v_2]) = l([v_1]) + l([v_3])$. Let $[v'_1]$ be a equivalence class which contained all vertices only adjacent to v_1 in T . Since $w([v'_1]) = w([v_2])$ which implies $l([v_3]) = 0$. But we know $|[v_3]| = 1$, by Theorem 5 which is a contradiction to $l([v_3]) = 0$ in H . The same argument works for corner vertex also.

Conversely, let T be a caterpillar such that it has no vertex of degree 2. The reduced graph H of T is A' magic, for all Abelian groups A with $|A| > 2$ (see Figure 2). By Theorem 5, T is A -vertex magic. □



Figure 1:



Figure 2: A' -vertex magic labeling of reduced graph H of T

Remark 1. Given a graph G , under the relation \sim_G the vertex set $V(G)$ is partitioned into equivalence classes. For every equivalence class $[v_i]$, with $|[v_i]| = 1$, introduce a new vertex x_i in $[v_i]$. Now, join the new vertex x_i to all vertices in equivalence classes belongs to $N([v_i])$. Thus each equivalence class will have at least two vertices. Hence by Theorem 4 the resultant graph is A -vertex magic, for all $|A| > 2$.

In the above construction we newly add at most $|V(G)|$ vertices, the graph G is convert into A -vertex magic, for all $|A| > 2$. Therefore, we get the following corollary as an immediate consequence of the above construction.

Corollary 11. Any graph H' is an induced subgraph of A -vertex magic graph G , where $|A| > 2$.

The above corollary 11 is a generalisation of the following result proved in [11] which has been proved only for finite Abelian groups and it uses more number of vertices.

Corollary 12. (Corollary 2.3, [11]) Any graph G is an induced subgraph of an A -vertex magic graph H .

We now state a procedure to construct infinite classes of A -vertex magic graphs, when $|A| > 2$. Let G be a A -vertex magic graph with magic labeling l . As \sim_G is an equivalence relation, join a new vertex x in any one of the equivalence class $[v_i]$ of G and join by an edge x into each vertex in $N(v_i)$, the resulting graph is G' . Then

by Lemma 1, we can relabel the vertices in $[v_i]$ and x whose label sum is equal to $\sum_{v \in [v_i]} l(v)$, for the other vertices in G' we retain the labeling given in G by l .

The following result proved in [11] is an immediate consequence of the above construction.

Corollary 13. (Theorem 3.1, [11]) *Let $t \geq 2$. Suppose G is an A -vertex magic graph of order n with a labeling l and magic constant g . If there exists an edge uv in G with $l(u) = l(v) = g$, then the graph G' , obtained from G by subdividing the edge uv and by attaching t pendant vertices at the new vertex x , is an A -vertex magic graph of order $n + t + 1$ with the same magic constant g .*

The main difference between our construction and corollary 13 is, it deals with more number of vertices and it is proved under an additional assumption on the magic labeling. We just add only one new vertex and our construction is true for infinite Abelian groups also.

2 Group Vertex Magic labeling of Generalised Friendship Graph

In [6], Henning Fernau et al. defined the Generalised Friendship Graph as follows

Definition 2. *The friendship graph f_m is a collection of m triangles with a common vertex. The generalised friendship graph $f_{n,m}$ is a collection of m cycles (all of order n), meeting at a common vertex.*

Proposition 1. *Let $m \in \mathbb{N}$. The group A is isomorphic to the group $\prod \mathbb{Z}_{m_i}$. Now, $m_i \mid (2m - 1)$, for all i if and only if $o(a) \mid (2m - 1)$, for all $a \in A$.*

Proof. Let $A \cong \prod \mathbb{Z}_{m_i}$. Suppose that $m_i \mid (2m - 1)$, for all i . Let $a \in A$. Then $a = (a_1, a_2, \dots, a_i, \dots)$, where $a_i \in \mathbb{Z}_{m_i}$ for each i , which implies $o(a_i) \mid m_i$ for all i . Then $o(a_i) \mid (2m - 1)$. Now, $(2m - 1)a_i = 0$ for all i . Therefore, $(2m - 1)a = (0, 0, \dots, 0, \dots)$. Hence $o(a) \mid (2m - 1)$.

Conversely, assume that $o(a) \mid (2m - 1)$, for all $a \in A$. Since $(2m - 1)a = (0, 0, \dots, 0, \dots)$, which implies $(2m - 1)a_i = 0$ for all i . Therefore $o(a_i) \mid (2m - 1)$. Since this is true for every $a_i \in \mathbb{Z}_{m_i}$, it also hold for generator of \mathbb{Z}_{m_i} . Hence $m_i \mid (2m - 1)$, for all i . \square

Theorem 6. *Let $n \equiv 1, 3 \pmod{4}$. The generalised friendship graph $G = f_{n,m}$ is A -vertex magic if and only if $A \cong \prod \mathbb{Z}_{m_i}$, where $m_i \mid (2m - 1)$, for all i .*

Proof. Let V_1, V_2, \dots, V_{m+1} be a partition of $V(G)$ with $|V_i| = n - 1$, where $i = 1, 2, \dots, m$. Let $V_i = \{v_{i_1}, v_{i_2}, \dots, v_{i_{n-1}}\}$ and $V_{m+1} = \{v\}$. If G is A -vertex magic, then $l(v_{i_k}) = l(v_{i_{k+4}})$.

Case 1: $n = 3$.

Assume that $f_{3,m}$ is A -vertex magic with $l(v_{i_1}) = a, l(v_{i_2}) = b, l(v) = c$. Since $w(v_{i_1}) = w(v_{i_2})$, which implies $a = b$. Let $j = \{1, 2, \dots, m\}$ and $i \neq j$. Assume that $l(v_{j_1}) = a_1, l(v_{j_2}) = b_1$. Since $w(v_{i_1}) = w(v_{j_1})$, which implies $b = b_1$. Since $w(v_{i_2}) = w(v_{j_2})$, which implies $a = a_1$. Since $w(v_{i_1}) = w(v_{j_2})$, which implies $b = a_1$. Also, $w(v_{i_1}) = w(v)$, which implies $c = (2m - 1)a$. Hence $A \not\cong \prod \mathbb{Z}_{m_i}$, where $m_i \mid (2m - 1)$, for all i . Conversely, assume that $A \not\cong \prod \mathbb{Z}_{m_i}$, where $m_i \mid (2m - 1)$, for all i . By Proposition 1, A has an element a such that $o(a) \nmid (2m - 1)$. Define $l : V(G) \rightarrow A \setminus \{0\}$ by $l(v_{i_k}) = a$, for all i, k and $l(v) = (2m - 1)a$. Thus $w(v) = 2ma$, for all $v \in V(G)$.

Case 2: $n \equiv 3(\text{mod } 4)$ and $n > 3$.

Assume that $f_{n,m}$ is A -vertex magic with $l(v_{i_1}) = a, l(v_{i_2}) = b, l(v_{i_3}) = c, l(v_{i_4}) = d$ and $l(v) = e$. Since $w(v_{i_1}) = w(v_{i_{n-1}})$, which implies $b = a$. Since $w(v_{i_1}) = w(v_{i_3})$, which implies $e = d$. Also $w(v_{i_2}) = w(v_{i_{n-1}})$, which implies $c = e$. Let $j \in \{1, 2, \dots, m\}$ and $i \neq j$. Assume that $l(v_{j_1}) = a_1, l(v_{j_2}) = b_1, l(v_{j_3}) = c_1, l(v_{j_4}) = d_1$. By similar argument, we get $b_1 = a_1$ and $c_1 = d_1 = e$. Since $w(v_{i_1}) = w(v_{j_1})$, which implies $b = b_1$. Since $w(v_{i_{n-1}}) = w(v_{j_{n-1}})$, which implies $a = a_1$. Now, $w(v_{i_1}) = w(v)$, which implies $e = (2m - 1)a$. Hence $A \not\cong \prod \mathbb{Z}_{m_i}$, where $m_i \mid (2m - 1)$, for all i .

Conversely, assume that $A \not\cong \prod \mathbb{Z}_{m_i}$, where $m_i \mid (2m - 1)$, for all i . By Proposition 1, A has an element a such that $o(a) \nmid (2m - 1)$. Define $l : V(G) \rightarrow A \setminus \{0\}$ by

$$l(v_{i_k}) = \begin{cases} (2m - 1)a & \text{if } k \equiv 0, 3(\text{mod } 4) \\ a & \text{otherwise,} \end{cases}$$

for all i, k and $l(v) = (2m - 1)a$. Thus $w(v) = 2ma$, for all $v \in V(G)$.

Case 3: $n \equiv 1(\text{mod } 4)$.

Assume that $f_{n,m}$ is A -vertex magic with $l(v_{i_1}) = a, l(v_{i_2}) = b, l(v_{i_3}) = c, l(v_{i_4}) = d$ and $l(v) = e$. Since $w(v_{i_1}) = w(v_{i_{n-1}})$, which implies $b = c$. Since $w(v_{i_2}) = w(v_{i_3})$, which implies $a = d$. Also $w(v_{i_2}) = w(v_{i_{n-1}})$, which implies $a = e$. Let $j \in \{1, 2, \dots, m\}$ and $i \neq j$. Assume that $l(v_{j_1}) = a_1, l(v_{j_2}) = b_1, l(v_{j_3}) = c_1, l(v_{j_4}) = d_1$. By similar argument, we get $b_1 = c_1, a_1 = d_1 = e$, which implies $a_1 = a = d = d_1$. Since $w(v_{i_1}) = w(v_{j_1})$, which implies $b = b_1$. Since $w(v_{i_{n-1}}) = w(v_{j_{n-1}})$, which implies $c = c_1$. Also, $w(v) = w(v_{i_1})$, which implies $b = (2m - 1)a$. Hence $A \not\cong \prod \mathbb{Z}_{m_i}$, where $m_i \mid (2m - 1)$, for all i .

Conversely, assume that $A \not\cong \prod \mathbb{Z}_{m_i}$, where $m_i \mid (2m - 1)$. By Proposition 1, A has an element a such that $o(a) \nmid (2m - 1)$. Define $l : V(G) \rightarrow A \setminus \{0\}$ by

$$l(v_{i_k}) = \begin{cases} (2m - 1)a & \text{if } k \equiv 2, 3(\text{mod } 4) \\ a & \text{otherwise,} \end{cases}$$

for all i, k and $l(v) = a$. Thus $w(v) = 2ma$, for all $v \in V(G)$. □

Theorem 7. *If $n \equiv 0(\text{mod } 4)$, then the generalised friendship graph $f_{n,m}$ is group vertex magic.*

Proof. Let A be a non-trivial Abelian group. Let V_1, V_2, \dots, V_{m+1} be a partition of $V(G)$ with $|V_i| = n - 1$, where $i = 1, 2, \dots, m$. Let $V_i = \{v_{i_1}, v_{i_2}, \dots, v_{i_{n-1}}\}$ and $V_{m+1} = \{v\}$. Define $l : V(G) \rightarrow A \setminus \{0\}$ by

$$l(v_{i_k}) = \begin{cases} -a & \text{if } k \equiv 1, 2 \pmod{4} \\ a & \text{otherwise} \end{cases}$$

and $l(v) = a$. Thus $w(v) = 0$, for all $v \in V(G)$ (see Figure 3).




Figure 3: Group vertex magic labeling of $f_{n,m}$, where $n \equiv 0 \pmod{4}$

□

Theorem 8. Let $n \equiv 2 \pmod{4}$. The generalised friendship graph $G = f_{n,m}$ is group vertex magic if and only if $m \equiv 1 \pmod{3}$.

Proof. Let V_1, V_2, \dots, V_{m+1} be a partition of $V(G)$ with $|V_i| = n - 1$, where $i = 1, 2, \dots, m$. Let $V_i = \{v_{i_1}, v_{i_2}, \dots, v_{i_{n-1}}\}$ and $V_{m+1} = \{v\}$. Assume that $f_{n,m}$ is group vertex magic. By observation, we have $l(v_{i_k}) = l(v_{i_{k+4}})$. Assume that $l(v_{i_1}) = a, l(v_{i_2}) = b, l(v_{i_3}) = c, l(v_{i_4}) = d$ and $l(v) = e$. Since $w(v_{i_1}) = w(v_{i_{n-1}})$, which implies $b = d$. Since $w(v_{i_1}) = w(v_{i_3})$, which implies $e = d$. Also, $w(v_{i_2}) = w(v_{i_3})$, which implies $a + c = b + d$. Let $j \in \{1, 2, \dots, m\}$ and $i \neq j$. Assume that $l(v_{j_1}) = a_1, l(v_{j_2}) = b_1, l(v_{j_3}) = c_1, l(v_{j_4}) = d_1$. By similar argument, we get $b_1 = d_1 = e$ and $a_1 + c_1 = b_1 + d_1$. Since $w(v_{i_1}) = w(v_{j_1})$, which implies $b = b_1$. Since $w(v_{i_{n-1}}) = w(v_{j_{n-1}})$, which implies $d = d_1$. Let $a' \in \mathbb{Z}_3 \setminus \{0\}$. Assume that $l(v_{i_2}) = l(v_{i_4}) = l(v) = a'$. Since $w(v_{i_2}) = w(v_{i_3})$, which implies $l(v_{i_1}) = l(v_{i_3}) = a'$,

therefore $l(v) = a'$, for all v . Since $w(v_{i_1}) = w(v)$, which implies $2a' = 2ma'$, we get $m \equiv 1 \pmod{3}$.

Conversely, assume that $m \equiv 1 \pmod{3}$ and p is a prime number.

Case 1. $p = 2$.

If 2 divides $o(A)$, then define $l : V(G) \rightarrow A \setminus \{0\}$ by $l(v) = a$, for all $v \in V(G)$, where $o(a) = 2$. Thus $w(v) = 0$, for all $v \in V(G)$.

Case 2. $p = 3$.

If 3 divides $o(A)$, then define $l : V(G) \rightarrow A \setminus \{0\}$ by $l(v) = a$, for all $v \in V(G)$, where $o(a) = 3$. Thus $w(v) = 2a$, for all v .

Case 3. $p > 3$.

If p divides $o(A)$, then

Subcase 1. Assume that m is odd.

Define $l : V(G) \rightarrow A \setminus \{0\}$ by

$$l(v_{i_k}) = \begin{cases} -a & \text{if } i \text{ is even and } k \equiv 1 \pmod{4} \\ 3a & \text{if } i \text{ is even and } k \equiv 3 \pmod{4} \\ a & \text{otherwise} \end{cases}$$

and $l(v) = a$, where $o(a) = p$. Thus $w(v) = 2a$, for all v .

Subcase 2. Assume that m is even.

Define $l : V(G) \rightarrow A \setminus \{0\}$ by

$$l(v_{i_k}) = \begin{cases} 3a & \text{if } i = 1 \text{ and } k \equiv 1 \pmod{4} \\ -a & \text{if } i = 1 \text{ and } k \equiv 3 \pmod{4} \\ -2a & \text{if } i = 2 \text{ and } k \equiv 1 \pmod{4} \\ 4a & \text{if } i = 2 \text{ and } k \equiv 3 \pmod{4} \\ -a & \text{if } i \text{ is even, } i > 2 \text{ and } k \equiv 1 \pmod{4} \\ 3a & \text{if } i \text{ is even, } i > 2 \text{ and } k \equiv 3 \pmod{4} \\ a & \text{otherwise} \end{cases}$$

and $l(v) = a$, where $o(a) = p$. Thus $w(v) = 2a$, for all $v \in V(G)$ (see Figure 4).

By Theorem 2 and Case 3, we get G is \mathbb{Z} -vertex magic. \square

Theorem 9. Let $n \equiv 2 \pmod{4}$ and $m \not\equiv 1 \pmod{3}$. The generalised friendship graph $f_{n,m}$ is A -vertex magic if and only if $A \neq \mathbb{Z}_3$.

Proof. Let V_1, V_2, \dots, V_{m+1} be a partition of $V(G)$ with $|V_i| = n - 1$, where $i = 1, 2, \dots, m$. Let $V_i = \{v_{i_1}, v_{i_2}, \dots, v_{i_{n-1}}\}$ and $V_{m+1} = \{v\}$. The necessary condition follows from Theorem 8. Conversely, assume that $A \neq \mathbb{Z}_3$.

Case 1. 2 divides $o(A)$.

The proof follows from the labeling in case 1 of above theorem.




Figure 4: \mathbb{Z}_p -vertex magic of $f_{n,m}$, $p > 3$, $n \equiv 2(\text{mod } 4)$ and m is even

Case 2. 9 divides $o(A)$.

By Sylow's first theorem and fundamental theorem of finite Abelian groups, A has a subgroup isomorphic to either \mathbb{Z}_9 or $\mathbb{Z}_3 \times \mathbb{Z}_3$. Suppose A has a subgroup isomorphic to $\mathbb{Z}_3 \times \mathbb{Z}_3$.

Subcase 1. Assume that m is odd. Define $l : V(G) \rightarrow \mathbb{Z}_3 \times \mathbb{Z}_3 \setminus \{0\}$ by

$$l(v_{i_k}) = \begin{cases} (1, 0) & \text{if } i > 1, i \text{ is even and } k \equiv 1(\text{mod } 4) \\ (1, 2) & \text{if } i > 1, i \text{ is even and } k \equiv 3(\text{mod } 4) \\ (2, 0) & \text{if } i > 1, i \text{ is odd and } k \equiv 1(\text{mod } 4) \\ (0, 2) & \text{if } i > 1, i \text{ is odd and } k \equiv 3(\text{mod } 4) \\ (1, 1) & \text{otherwise} \end{cases}$$

for all i, k and $l(v) = (1, 1)$

Subcase 2. Assume that m is even. Define $l : V(G) \rightarrow \mathbb{Z}_3 \times \mathbb{Z}_3 \setminus \{0\}$ by

$$l(v_{i_k}) = \begin{cases} (1, 0) & \text{if } i = 1 \text{ and } k \equiv 1(\text{mod } 4) \\ (1, 2) & \text{if } i = 1 \text{ and } k \equiv 3(\text{mod } 4) \\ (0, 1) & \text{if } i \text{ is even and } k \equiv 1(\text{mod } 4) \\ (2, 1) & \text{if } i \text{ is even and } k \equiv 3(\text{mod } 4) \\ (0, 2) & \text{if } i > 2, i \text{ is odd and } k \equiv 1(\text{mod } 4) \\ (2, 0) & \text{if } i > 2, i \text{ is odd and } k \equiv 3(\text{mod } 4) \\ (1, 1) & \text{otherwise} \end{cases}$$

for all i, k and $l(v) = (1, 1)$. Thus $w(v) = (2, 2)$ for all $v \in V(G)$.

Suppose A has a subgroup isomorphic to \mathbb{Z} , \mathbb{Z}_9 or \mathbb{Z}_p , where $p > 3$, then the proof follows from Theorem 2 and the labeling in case 3 of above theorem. \square

3 Concluding remarks

In this paper we have proved the A -vertex magicness of several graphs by considering its reduced graph. These ideas helps us to embed every graph as an induced subgraph of an A -vertex magic graph and also to construct infinite families of A -vertex magic graphs in a much more simpler way.

Acknowledgements

The work of S. Balamoorthy is supported by a Junior Research Fellowship from CSIR-UGC, India(UGC-Ref.No.:1085/(CSIR-UGC NET JUNE 2019)). The authors would like to thank Dr T. Kavaskar, Central University of Tamil Nadu, Thiruvavur for the support and fruitful discussions.

References

- [1] D.F. Anderson, R. Levy and J. Shapiro, Zero-divisor graphs, von Neumann regular rings and Boolean algebras. J. Pure Appl. Algebra 180 (2003), 221–241.
- [2] S. Balamoorthy, S.V. Bharanedhar and N. Kamatchi, On the products of group vertex magic graphs, AKCE Int. J. Graphs Comb. 19(3) (2022), 268–275.
- [3] S. Balamoorthy, T. Kavaskar and Vinothkumar. K (2023). Wiener index of an ideal-based zero-divisor graph of commutative ring with unity. AKCE International Journal of Graphs and Combinatorics, 1–9. <https://doi.org/10.1080/09728600.2023.2263040>
- [4] J.A. Bondy and U.S.R. Murty, Graph Theory with Applications, American Elsevier Publishing Co., Inc., New York (1976).
- [5] I.N. Herstein, Topics in Algebra, John Wiley and Sons, 2nd ed., New York (2006).
- [6] H. Fernau, J.F. Ryanb and K.A. Sugeng, A sum labelling for the generalised friendship graph, Discrete math. 308 (2008), 734–740.
- [7] N. Kamatchi, K. Paramasivam, A.V. Prajeesh, K. Muhammed Sabeel and S. Arumugam, On group vertex magic graphs, AKCE Int. J. Graphs Combin. 17(1) (2020), 461–465.

- [8] S.M. Lee, H. Sun and I. Wen, On the group magic graphs, *J. Combin. Math. Combin. Comput.* 38 (2001), 197–207.
- [9] S.M. Lee, F.Saba, E. Salehi, and H. Sun, On the V_4 -magic graphs, *Congr. Numer.* 156 (2002), 59–67.
- [10] R.M. Low and S.M. Lee, On the products of group-magic graphs, *Australas. J. Combin.* 34 (2006), 41–48.
- [11] K.M. Sabeel, K. Paramasivam, A.V. Prajeesh, N. Kamatchi and S. Arumugam, A characterization of group vertex magic trees of diameter up to 5, *Australas. J. Combin.* 85(1) (2023), 49-60.
- [12] M. Saravanan, S.P. Murugan and G. Arunkumar, A generalization of Fiedler’s Lemma and the spectra of H-join of graphs, *Linear Algebra Appl.* 625 (2021), 20–43.